

*Pacific  
Journal of  
Mathematics*

**EXTREMELY CLOSED SUBGROUPS  
AND A VARIANT ON GLAUBERMAN'S  $Z^*$ -THEOREM**

HUNG P. TONG-VIET

Volume 329 No. 2

April 2024



## EXTREMELY CLOSED SUBGROUPS AND A VARIANT ON GLAUBERMAN'S $Z^*$ -THEOREM

HUNG P. TONG-VIET

**Let  $G$  be a finite group and let  $H$  be a subgroup of  $G$ . We say that  $H$  is extremely closed in  $G$  if  $\langle H, H^g \rangle \cap N_G(H) = H$  for all  $g \in G$ . We determine the structure of finite groups with an extremely closed abelian  $p$ -subgroup for some prime  $p$ . In particular, we show that if  $G$  contains such a subgroup  $H$ , then  $G = N_G(H) O_{p'}(G)$ . This is a variant on the celebrated Glauberman's  $Z^*$ -theorem.**

### 1. Introduction

It is an important problem in finite group theory to determine whether a finite group is simple or not. Many nonsimplicity criteria have been obtained in the literature. Among those is the celebrated Glauberman's  $Z^*$ -theorem. To state this theorem, we need some definitions. Let  $G$  be a finite group and let  $p$  be a prime. Let  $x \in G$  be a  $p$ -element and let  $P$  be a Sylow  $p$ -subgroup of  $G$  containing  $x$ . We say that  $x$  is isolated in  $P$  with respect to  $G$  if  $x^G \cap P = \{x\}$ , that is,  $x$  is not conjugate in  $G$  to any element in  $P$  except for  $x$  itself. Here,  $x^G$  denotes the conjugacy class of  $G$  containing  $x$ . We say that  $x$  is isolated in  $G$  if  $x$  is isolated in some Sylow  $p$ -subgroup of  $G$  containing it. Glauberman's  $Z^*$ -theorem [18] states that if  $x \in G$  is an isolated involution in  $G$ , then  $G = C_G(x) O_2(G)$ . The proof of this theorem depends on the modular representation theory and is independent of the classification of finite simple groups. Recall that for a prime  $p$ ,  $O_{p'}(G)$  is the largest normal  $p'$ -subgroup of  $G$ . Extending this fundamental theorem to all primes, Glauberman's  $Z_p^*$ -theorem states that if  $x \in G$  is an isolated  $p$ -element, then  $G = C_G(x) O_{p'}(G)$ . For various proofs of this theorem, see [1; 19; 27; 40]. Note that all of these proofs depend on the classification of finite simple groups. For many equivalent statements of this theorem, see [30]. Also, see [14] for some variant of Glauberman's  $Z_p^*$ -theorem.

In this paper, we introduce the so-called *extremely closed subgroup* and obtain some new factorization of finite groups similar to Glauberman's  $Z_p^*$ -theorem which gives some nonsimplicity criteria for finite groups. Let  $G$  be a finite group and let

*MSC2020:* 20D05, 20D15.

*Keywords:* extremely closed, strongly closed, weakly closed.

$H \leq M$  be subgroups of  $G$ . We say that  $H$  is extremely closed in  $M$  with respect to  $G$  if  $\langle H, H^g \rangle \cap M = H$  for all  $g \in G - M$ , and that  $H$  is extremely closed in  $G$  if  $H$  is extremely closed in  $N_G(H)$  with respect to  $G$ . Trivially, if  $H$  is a normal or self-normalizing subgroup of  $G$ , then  $H$  is extremely closed in  $G$ .

If  $G$  is a finite group, we write  $Z(G)$  for the center of  $G$  and  $\Phi(G)$  for the Frattini subgroup of  $G$ , that is, the intersection of all maximal subgroups of  $G$ . Furthermore, if  $H$  is a subgroup of  $G$ , then  $\langle H^G \rangle$  is the normal closure of  $H$  in  $G$ . We next compare our definition of extremely closed subgroups with other known embedding properties of subgroups of finite groups.

The first motivation for our definition comes from work of Flavell [10] on the generation of finite groups with maximal subgroups of maximal subgroups. In particular, a triple  $(G, M, H)$  with  $H \leq M \leq G$  is called a  $\gamma$ -triple if  $H < M < G$  and  $\langle H, g \rangle \cap M = H$  for all  $g \in G - M$ . If  $H$  is maximal in  $M$  and  $M$  is maximal in  $G$  and moreover,  $G$  cannot be generated by any two conjugates of  $H$ , then  $(G, M, H)$  is a  $\gamma$ -triple. Clearly, if  $(G, M, H)$  is a  $\gamma$ -triple, then  $H$  is extremely closed in  $M$  with respect to  $G$ . The converse is not true by the example below.

**Example 1.1.** Let  $G = S_4$  be the symmetric group of degree 4. Let  $H = \langle (1, 2, 3) \rangle$  and  $M = N_G(H) \cong S_3$ . Then  $\langle H^G \rangle = A_4$  and  $\langle H^G \rangle \cap M = H$ . So  $\langle H, H^g \rangle \cap M = H$  for all  $g \in G$ , and hence  $H$  is extremely closed in  $M$  with respect to  $G$ . However, let  $g = (1, 3, 2, 4) \in G - M$ . Then  $\langle H, g \rangle = G$  and so  $\langle H, g \rangle \cap M = M \neq H$ . Therefore  $(G, M, H)$  is not a  $\gamma$ -triple.

For the second motivation, following Hawkes and Humphreys [31], a subgroup  $M$  of a finite group  $G$  is said to have property CR (character restriction) if every irreducible complex character of  $M$  is the restriction of a character of  $G$ . In [31], the authors studied finite solvable groups with a CR-subgroup and the general cases were considered by Isaacs in [32]. One important property of a CR-subgroup  $M$  of a finite group  $G$  is that if  $H \trianglelefteq M$ , then  $\langle H^G \rangle \cap M = H$  [32, Proposition 1.1]. Berkovich [3] called a triple  $(G, M, H)$  with  $H \trianglelefteq M \leq G$  *special* in  $G$  if  $\langle H^G \rangle \cap M = H$ . (Li [36] calls  $H$  an NE-subgroup of  $G$  if  $(G, N_G(H), H)$  is special in  $G$ .) Isaacs [32] showed that if  $P$  is a Sylow  $p$ -subgroup of  $G$ , where  $p$  is a prime and assume that  $N_G(P)$  satisfies CR in  $G$ , then  $N_G(P)$  has a normal complement in  $G$ . This result was extended by Berkovich [3], where he showed that if both triples  $(G, N_G(P), P)$  and  $(G, N_G(P), \Phi(P))$  are special in  $G$ , then  $N_G(P)$  has a normal complement in  $G$ . This gives a character theory free proof of Isaacs' result mentioned earlier. Observe that if a triple  $(G, N_G(H), H)$  is special in  $G$ , then  $H$  is extremely closed in  $G$ . However, the converse is not true.

**Example 1.2.** Let  $G = P : H$  be a semidirect product of  $H$  and  $P$ , where  $H = \langle a \rangle$  is a cyclic group of order 2 and  $P \cong 3_+^{1+2}$  is an extraspecial group of order 27 with

exponent 3, so

$$P = \langle x, y, z \mid z = [x, y], x^3 = y^3 = z^3 = 1 = [x, z] = [y, z] \rangle$$

and  $H$  acts on  $P$  via  $x^a = x^{-1}$ ,  $y^a = y^{-1}$  and  $z^a = z$ . Then  $N_G(H) = C_G(H) = H \langle z \rangle$ , and  $\langle H^G \rangle = G$ . For every  $g \in G - N_G(H)$ , we have  $T = \langle H, H^g \rangle$  is a dihedral group of order 6, so  $N_T(H) = H$ , and hence  $H$  is extremely closed in  $G$  but  $(G, N_G(H), H)$  is not a special triple since  $\langle H^G \rangle \cap N_G(H) = H \langle z \rangle \neq H$ .

Finally, we mention the last inspiration for our new embedding property. Let  $H \leq M$  be subgroups of a finite group  $G$ . Recall that  $H$  is said to be *strongly closed* in  $M$  with respect to  $G$  if, whenever  $a^g \in M$ , where  $a \in H$ ,  $g \in G$ , then  $a^g \in H$ . This is equivalent to saying that  $M \cap H^g \leq H$  for all  $g \in G$ . Furthermore, we say that  $H$  is strongly closed in  $G$  if  $H$  is strongly closed in  $N_G(H)$  with respect to  $G$ . Noting that in [4],  $H$  is called an  $\mathcal{H}$ -subgroup of  $G$  if  $H$  is strongly closed in  $G$ . If  $H = \langle x \rangle$  is cyclic of order 2, then  $H$  is strongly closed in  $G$  if and only if  $x$  is isolated in  $G$ . Finite groups with a strongly closed  $p$ -subgroup are determined in [15; 16; 20]. It is easy to see that if  $H$  is extremely closed in  $G$ , then  $H$  is strongly closed in  $G$ .

**Example 1.3.** Let  $G = U_3(4)$ . By [20], if  $P$  is a Sylow 2-subgroup of  $G$ , then  $H = Z(P) = \Phi(P)$  is a strongly closed abelian 2-subgroup of  $G$ . Using GAP [17], we can find  $g \in G$  of order 15 such that  $\langle H, H^g \rangle \cong A_5 \cong \text{SU}_2(4)$ ,  $N_G(H) \cong P \langle g \rangle$  and  $\langle H, H^g \rangle \cap N_G(H) \cong A_4 \neq H$ . Thus  $H$  is not extremely closed in  $G$ .

Finally, we recall the definition of weakly closed subgroups. Let  $H \leq M$  be subgroups of a finite group  $G$ . We say that  $H$  is *weakly closed* in  $M$  with respect to  $G$  if, whenever  $H^g \subseteq M$ , where  $g \in G$ , then  $H^g = H$ . It is easy to see that if  $H$  is strongly closed in  $M$  with respect to  $G$ , then  $H$  is weakly closed in  $M$  with respect to  $G$ . Moreover, when  $H$  is cyclic of prime order, these two concepts coincide. We know that if  $H \triangleleft M \leq G$  and  $\langle H^G \rangle \cap M = H$ , then  $H$  is extremely closed in  $M$  with respect to  $G$ . In our first result, we show that in certain cases, the converse holds.

**Theorem 1.4.** *Let  $G$  be a finite group and let  $H \leq M$  be subgroups of  $G$ . Suppose that  $\langle H, H^g \rangle \cap M = H$  for all  $g \in G$ . If  $H$  is maximal in  $M$  and  $M$  is maximal in  $G$ , then  $\langle H^G \rangle \cap M = H$ .*

Let  $H \leq M$  be subgroups of a finite group  $G$ . Recall that  $H$  is called a *weak second maximal subgroup* of  $G$  if there exists a maximal subgroup  $M$  of  $G$  such that  $H$  is maximal in  $M$ . Moreover,  $H$  is called a *second maximal subgroup* of  $G$  if  $H \neq G$  and it is maximal in every maximal subgroup of  $G$  containing it. Flavell [10] shows that if  $G$  is a finite nonabelian simple group and  $H$  is a weak second maximal subgroup of  $G$ , then  $G = \langle H, g \rangle$  for some  $g \in G$ . And in [9], it is shown that

if  $H$  is a second maximal subgroup of a finite nonabelian simple group  $G$ , then  $G = \langle H, H^g \rangle$  for some  $g \in G$ . As a corollary to Theorem 1.4, we obtain the following generation result for finite nonabelian simple groups.

**Corollary 1.5.** *Let  $G$  be a finite nonabelian simple group. Let  $M$  be a maximal subgroup of  $G$  and let  $H$  be a normal subgroup of  $M$  of prime index. Then  $G = \langle H, H^g \rangle$  for some  $g \in G$ .*

Note that if  $G = A_5$ ,  $M = S_3$  and  $H$  is a Sylow 2-subgroup of  $M$ , then  $M$  is a maximal subgroup of  $G$  and  $|M : H| = 3$  is a prime, but  $G$  cannot be generated by any two conjugates of  $H$ . So we cannot drop the hypothesis that  $H \trianglelefteq M$  in Corollary 1.5. We should mention that Flavell [13] asks whether a finite nonabelian simple group can be generated by two conjugates of a self-normalizing subgroup.

Let  $G$  be a finite group and let  $p$  be a prime. We now focus on extremely closed  $p$ -subgroups. Let  $H$  be an extremely closed  $p$ -subgroup of  $G$ . We first assume that  $H = \langle x \rangle$  is cyclic of order  $p$ . If  $p = 2$ , then it is not hard to see that  $x$  is isolated in  $G$  or equivalently  $H$  is strongly closed in  $G$  and so  $G = C_G(x) O_{2'}(G)$  by Glauberman's  $Z^*$ -theorem. In particular,  $G$  is not simple. When  $p$  is odd, we note that there exists a simple group with a weakly closed or strongly closed subgroup of order  $p$ , for instance, when a Sylow  $p$ -subgroup of  $G$  is cyclic. However, when  $H$  is extremely closed in  $G$ , the subgroups generated by any two distinct conjugates of  $H$  are Frobenius groups. By applying a result due to B. Fischer [8] concerning Frobenius automorphisms, we can show that  $(G, N_G(H), H)$  is special in  $G$ , and then we obtain a factorization similar to that of Glauberman's  $Z_p^*$ -theorem.

**Theorem 1.6.** *Let  $G$  be a finite group and let  $p$  be an odd prime divisor of the order of  $G$ . Let  $H$  be a cyclic subgroup of order  $p$  of  $G$ . If  $\langle H, H^g \rangle \cap N_G(H) = H$  for all  $g \in G$ , then  $\langle H^G \rangle \cap N_G(H) = H$ . In particular,  $G = N_G(H) O_{p'}(G)$ .*

It would be nice to have a proof of Theorem 1.6 which does not rely on the classification of finite simple groups. For an arbitrary prime  $p$ , if  $H$  is an abelian extremely closed  $p$ -subgroup of a finite group  $G$ , we also obtain a similar factorization  $G = N_G(H) O_{p'}(G)$  as in Theorem 1.6. For even prime, the proof depends only on the classification of finite groups with an abelian strongly closed subgroup by Goldschmidt [20] which is independent of the classification of finite simple groups. For odd primes, we make use of a result due to Guest [25] on the characterization of solvable radical of finite groups and the classification of finite groups with strongly closed subgroups by Flores and Foote [15].

**Theorem 1.7.** *Let  $G$  be a finite group and let  $p$  be prime. Let  $H$  be an abelian  $p$ -subgroup of  $G$ . If  $\langle H, H^g \rangle \cap N_G(H) = H$  for all  $g \in G$ , then  $G = N_G(H) O_{p'}(G)$ .*

Remark that we cannot drop the hypothesis that  $H$  is abelian when  $p = 2$  in Theorem 1.7, since the simple group  $G = L_2(17)$  has a self-normalizing Sylow 2-subgroup  $P$  which is nonabelian and so  $P$  is clearly an extremely closed 2-subgroup of  $G$ . Note that if  $|H| = 2$ , then Theorem 1.7 is just Glauberman's  $Z^*$ -theorem. Example 1.2 above shows that an abelian extremely closed 2-subgroup  $H$  may not satisfy the condition  $\langle H^G \rangle \cap N_G(H) = H$ . However, this holds true for odd primes. We obtain the following as a corollary to Theorem 1.7.

**Corollary 1.8.** *Let  $G$  be a finite group and let  $p$  be an odd prime. Let  $H$  be an abelian  $p$ -subgroup of  $G$ . If  $\langle H, H^g \rangle \cap N_G(H) = H$  for all  $g \in G$ , then  $\langle H^G \rangle \cap N_G(H) = H$ .*

Recall that for a finite group  $G$ , the solvable radical of  $G$ , denoted by  $R(G)$ , is the largest normal solvable subgroup of  $G$ . Theorem 1.7 and Corollary 1.8 now yield the following.

**Corollary 1.9.** *Let  $G$  be a finite group and let  $p$  be a prime. If  $H$  is an extremely closed abelian  $p$ -subgroup of  $G$ , then  $H \subseteq R(G)$ .*

By an application of Burnside's normal  $p$ -complement theorem and the solvability of finite groups admitting a fixed point free coprime group action, if  $H$  satisfies the hypothesis of the corollary, then  $\langle H, H^g \rangle$  is solvable for all  $g \in G$ . Thus if  $x \in H$ , then  $\langle x, x^g \rangle$  is solvable for all  $g \in G$ . By the main results in [21; 25], if  $p \geq 5$ , then  $x \in R(G)$  and hence  $H \leq R(G)$ . Thus the above corollary only provides new result when  $p = 2$  or 3.

In general, if  $x$  is a  $p$ -element and  $\langle x, x^g \rangle$  is  $p$ -solvable for all  $g \in G$ , then it is not true that  $x \in R_p(G)$ , where  $R_p(G)$  is the  $p$ -solvable radical of  $G$ , that is,  $R_p(G)$  is the largest normal  $p$ -solvable subgroup of  $G$ . For a counterexample, consider  $G = U_3(3)$  and  $x \in G$  a transvection, so  $x$  has order 3 and the conjugacy class of  $G$  containing  $x$  has size 56, then we can check that  $\langle x, x^g \rangle$  is isomorphic to either  $\langle x \rangle$  or  $SL_2(3)$  for every  $g \in G$ . Clearly  $\langle x, x^g \rangle$  is 3-nilpotent and hence it is 3-solvable for every  $g \in G$ . There is also a counterexample when  $p = 2$  since if  $x \in G$  is an involution, then  $\langle x, x^g \rangle$  is 2-nilpotent for every  $g \in G$ . Recall that a finite group  $G$  is  $p$ -nilpotent if it has a normal  $p$ -complement for some prime  $p$ . On the other hand, it is proved in [11] that if  $P$  is a Sylow  $p$ -subgroup of a finite group  $G$  for some prime  $p$ , then  $G$  is  $p$ -solvable if and only if  $\langle P, P^g \rangle$  is  $p$ -solvable for all  $g \in G$ . Generalizing this result, we can prove the following.

**Theorem 1.10.** *Let  $G$  be a finite group and let  $p$  be a prime. Let  $P$  be a Sylow  $p$ -subgroup of  $G$ . Then  $G$  is  $p$ -solvable if and only if  $\langle P, P^g \rangle$  is  $p$ -solvable for all  $g \in G$ .*

Our notation is standard. For finite group theory, we follow [22] and [35] and for finite simple groups, we follow the notation in [34].

For the organization of the paper, we collect some useful results in Section 2. We will prove Theorems 1.4–1.7 and the corollaries in Section 3 and the last theorem will be proved in Section 4.

## 2. Preliminaries

Let  $G$  be a finite group. Recall that the Fitting subgroup of  $G$ , denoted by  $F(G)$ , is the largest nilpotent normal subgroup of  $G$ . The layer of  $G$ , denoted by  $E(G)$ , is the product of all components of  $G$ , where a component of  $G$  is a subnormal quasisimple subgroup of  $G$ . A finite group  $L$  is quasisimple if  $L$  is perfect and  $L/Z(L)$  is a nonabelian simple group. The generalized Fitting subgroup of  $G$ , denoted by  $F^*(G)$ , is defined by  $F^*(G) = F(G)E(G)$ . As usual, if  $H \leq G$ , then  $N_G(H)$  and  $C_G(H)$  denote the normalizer and centralizer of  $H$  in  $G$ , respectively. Finally, a finite group  $G$  is almost simple with socle  $S$  if there exists a finite nonabelian simple group  $S$  such that  $S \trianglelefteq G \leq \text{Aut}(S)$ .

Recall that a subgroup  $H$  of  $G$  is called *pronormal* (resp. *abnormal*) in  $G$  if for any  $g \in G$ ,  $H^g = H^u$  for some  $u \in \langle H, H^g \rangle$ , (resp.  $\langle H, H^g \rangle = \langle H, g \rangle$ ). The first lemma is obvious, for completeness, we will include a proof here.

**Lemma 2.1.** *Let  $G$  be a finite group. Let  $H$  be a pronormal subgroup of  $G$  and let  $N \trianglelefteq G$ . Then the following hold.*

- (i) *If  $N \trianglelefteq G$  and  $P \in \text{Syl}_p(N)$ , then  $P$  is pronormal in  $G$ .*
- (ii)  *$N_G(H)$  is abnormal in  $G$ .*
- (iii) *If  $H \leq N \trianglelefteq G$ , then  $G = N_G(H)N$ .*
- (iv) *If  $H \leq L \leq G$ , then  $H$  is pronormal in  $L$ .*
- (v) *If  $H$  is subnormal in  $K$ , where  $K \leq G$ , then  $H \trianglelefteq K$ .*
- (vi) *If  $L = \langle H^G \rangle$ , then  $\langle H^L \rangle = L$ .*

*Proof.* (i) Let  $g \in G$ . Since  $P \leq N \trianglelefteq G$ , we have  $\langle P, P^g \rangle \leq N$ . As  $P \in \text{Syl}_p(N)$ , it follows that  $P \in \text{Syl}_p(\langle P, P^g \rangle)$  and hence by Sylow's theorem,  $P^g = P^u$  for some  $u \in \langle P, P^g \rangle$ . Thus  $P$  is pronormal in  $G$ .

(ii) Assume that  $H$  is pronormal in  $G$ . Let  $M = N_G(H)$  and let  $g \in G$ . By definition,  $H^g = H^u$ , for some  $u \in \langle H, H^g \rangle$ , whence  $gu^{-1} \in N_G(H) = M$ . Since  $u \in \langle H, H^g \rangle \leq \langle M, M^g \rangle$ , we have  $g \in \langle M, M^g \rangle$  and so  $\langle M, M^g \rangle = \langle M, g \rangle$ . Hence  $M$  is abnormal in  $G$ .

(iii) Let  $g \in G$ . We have  $H^g = H^u$ , where  $u \in \langle H, H^g \rangle \leq N$  as  $H \leq N \trianglelefteq G$ . Thus  $gu^{-1} \in N_G(H)$  and so  $g \in N_G(H)N$ .

(iv) This is obvious.

(v) By (iv), it suffices to show that if  $H$  is subnormal and pronormal in  $G$  then  $H \trianglelefteq G$ . In fact, we only need to prove the following: if  $H \trianglelefteq K \trianglelefteq G$  and  $H$  is pronormal in  $G$  then  $H \trianglelefteq G$ . By applying (iii), we have  $G = N_G(H)K$ . However, as  $H \trianglelefteq K$ ,  $K \leq N_G(H)$  and so  $G = N_G(H)$ .

(vi) Since  $H \leq L = \langle H^G \rangle \trianglelefteq G$ ,  $G = N_G(H)L$  by (iii) and thus

$$L = \langle H^G \rangle = \langle H^{N_G(H)L} \rangle = \langle H^L \rangle \leq L.$$

Therefore  $L = \langle H^L \rangle$ . □

We next deduce some properties of extremely closed subgroups.

**Lemma 2.2.** *Let  $G$  be a finite group and let  $H$  be a  $p$ -subgroup of  $G$  for some prime  $p$ . Let  $N \trianglelefteq G$  and assume that  $H$  is extremely closed in  $G$ . Let  $\bar{G} = G/N$ . Then the following hold.*

- (i) *For every  $g \in G$ , we have  $N_{\langle H, H^g \rangle}(H) = H$  and  $H \in \text{Syl}_p(\langle H, H^g \rangle)$ .*
- (ii)  *$H$  is pronormal in  $G$ .*
- (iii) *If  $H \leq L$ , then  $H$  is extremely closed in  $L$ .*
- (iv)  *$N_{\bar{G}}(\bar{H}) = \overline{N_G(H)}$ .*
- (v) *If  $H$  is abelian, then  $\langle H, H^g \rangle = HO_{p'}(\langle H, H^g \rangle)$ , for every  $g \in G$ .*
- (vi)  *$\bar{H}$  is extremely closed in  $\bar{G}$ .*
- (vii) *If  $H \leq Q \leq G$ , where  $Q$  is a  $p$ -group, then  $N_G(Q) \leq N_G(H)$ . In particular, if  $H \leq P \in \text{Syl}_p(G)$ , then  $P \leq N_G(H)$ .*

*Proof.* (i) Let  $g \in G$  and let  $T = \langle H, H^g \rangle$ . We have  $H = T \cap N_G(H) = N_T(H)$  and so  $H$  is a Sylow  $p$ -subgroup of  $T$  by Sylow's theorem.

(ii) Let  $g \in G$ . As above, let  $T = \langle H, H^g \rangle$ . From part (i),  $H$  is a Sylow  $p$ -subgroup of  $T$  and since  $|H^g| = |H|$  and  $H^g \leq T$ ,  $H^g$  is also a Sylow  $p$ -subgroup of  $T$ . By Sylow's theorem,  $H^g = H^u$  for some  $u \in T$ . Thus  $H$  is pronormal in  $G$ .

(iii) Let  $g \in L$ . Then  $\langle H, H^g \rangle \cap N_L(H) = \langle H, H^g \rangle \cap N_G(H) \cap L = H \cap L = H$ .

(iv) It suffices to show that  $N_G(HN) \leq N_G(H)N$ . Let  $g \in N_G(HN)$ . Then  $H^g \leq HN$  and hence  $T = \langle H, H^g \rangle \leq HN$ . Since  $H$  is pronormal in  $G$  by (ii), we have  $H^g = H^u$ , for some  $u \in T \leq HN$ . Thus  $gu^{-1} \in N_G(H)$  whence  $g \in N_G(H)HN = N_G(H)N$ .

(v) Assume that  $H$  is abelian. Let  $g \in G$  and let  $T = \langle H, H^g \rangle$ . By (i),  $H$  is a self-normalizing abelian Sylow  $p$ -subgroup of  $T$ . The result now follows from Burnside's normal  $p$ -complement theorem ([22, Theorem 7.4.3]).

(vi) Applying (iv), we need to show that  $\langle H, H^g \rangle N \cap N_G(H)N = HN$ , for all  $g \in G$ . Let  $T = \langle H, H^g \rangle$ . By Dedekind's Modular law, we have

$$TN \cap N_G(H)N = N(TN \cap N_G(H)).$$

Hence, it suffices to show that  $TN \cap N_G(H) \leq HN$ . Let  $y = xn \in TN \cap N_G(H)$ , where  $y \in N_G(H)$ ,  $x \in T$  and  $n \in N$ . We have  $H = H^y = H^{xn}$ , it implies that  $H^x = H^{n^{-1}}$ . By (i), we have  $N_T(H) = H$ , and so by (ii) and Lemma 2.1(ii),  $H$  is abnormal in  $T$ . Thus  $x \in \langle H, H^x \rangle = \langle H, H^{n^{-1}} \rangle \leq HN$ . Therefore,  $y = xn \in HN$ .  
 (vii) Since  $H$  is pronormal and subnormal in  $N_G(Q)$ , by Lemma 2.1(v),  $H$  is normal in  $N_G(Q)$ . The remaining claim is obvious.  $\square$

**Lemma 2.3.** *Let  $G$  be a finite group, let  $N \trianglelefteq G$  and let  $H$  be an extremely closed  $p$ -subgroup of  $G$  for some prime  $p$ . Let  $P$  be a Sylow  $p$ -subgroup of  $G$  containing  $H$  and let  $Q = H \cap N$ . Then the following hold.*

- (i)  $H$  is strongly closed in  $P$  with respect to  $G$ .
- (ii)  $Q$  is strongly closed in  $P \cap N$  with respect to  $N$ .
- (iii) If  $N \leq N_G(H)$ , then  $Q \trianglelefteq G$ .

*Proof.* Observe that  $R := P \cap N \in \text{Syl}_p(N)$  and by Lemma 2.2(vii),  $P \leq N_G(H)$ .

- (i) For  $g \in G$ , we have  $H^g \cap P \leq \langle H, H^g \rangle \cap N_G(H) = H$ . So  $H$  is strongly closed in  $P$  with respect to  $G$ .
- (ii) For  $n \in N$ , we have  $Q^n \leq H^n$  and  $R \leq P \leq N_G(H)$  and so

$$Q^n \cap R \leq \langle H, H^n \rangle \cap N_G(H) = H.$$

Furthermore, as  $Q \leq N \trianglelefteq G$  and  $R \leq N$ , we have  $Q^n \cap R \leq N$ . Hence we obtain  $Q^n \cap R \leq H \cap N = Q$ .

- (iii) Assume that  $N \leq N_G(H)$ . For each  $g \in G$ , we have

$$Q^g = Q^g \cap N \leq \langle H, H^g \rangle \cap N_G(H) \cap N = H \cap N = Q.$$

Hence  $Q \trianglelefteq G$  as wanted.  $\square$

We next quote some results that we will need for the proofs of the main theorems.

**Lemma 2.4.** *Suppose that  $G$  is a finite group with  $F(G) = 1$ . Let  $L$  be a component of  $G$ . If  $x \in G$  such that  $x \notin N_G(L)$  and  $x^2 \notin C_G(L)$  then there exists an element  $g \in G$  such that  $\langle x, x^g \rangle$  is not solvable.*

*Proof.* This is Lemma 1 in [25].  $\square$

**Lemma 2.5.** *Let  $G$  be a finite almost simple group with socle  $L$ . Suppose that  $x \in G$  is an element of order  $p$ , where  $p$  is an odd prime. Then one of the following holds:*

- (i)  $\langle x, x^g \rangle$  is not solvable for some  $g \in G$ ;
- (ii)  $p = 3$  and  $L$  is a finite simple group of Lie type defined over  $\mathbb{F}_3$ , a finite field with 3 elements, or  $L \cong U_n(2)$ ,  $n \geq 4$ . Moreover, the Lie rank of  $L$  is at least 2 unless  $L \cong U_3(3)$ .

*Proof.* This is Theorem  $A^*$  in [25]. □

Recall that a triple  $(G, M, H)$  with  $H \trianglelefteq M \leq G$  is called a  $W$ -triple if  $M \cap M^g \leq H$  for all  $g \in G - M$ .

**Lemma 2.6.** *Let  $G$  be a finite group and  $H \trianglelefteq M \leq G$ . Then  $(G, M, H)$  is a  $W$ -triple if and only if*

$$N_G(D) \leq M \text{ for all subgroups } D \leq M \text{ with } D \not\leq H.$$

*Proof.* This is Lemma 2.3 in [10]. □

**Lemma 2.7** (Wielandt's theorem). *Let  $G$  be a finite group. If  $(G, M, H)$  is a  $W$ -triple, then  $G$  contains a normal subgroup  $K$  such that  $G = MK$  and  $M \cap K = H$ . In particular, the triple  $(G, M, H)$  is special in  $G$ .*

*Proof.* The first claim is in [42] or [38, Exercise 1, p. 347] and the second is in [3, Lemma 9]. □

An automorphism  $\theta$  of a finite group is Frobenius if each nontrivial power of  $\theta$  is fixed point free.

**Lemma 2.8.** *Let  $G$  be a finite group and let  $D$  be a conjugacy class of  $G$  containing elements of order  $> 2$ . Assume that  $G = \langle D \rangle$ . Then some element of  $D$  induces a Frobenius automorphism on  $G'$  if and only if each pair of distinct elements in  $D$  generates a Frobenius group.*

*Proof.* This is [8, Satz I]. □

We also need the following results.

**Lemma 2.9.** *Let  $G$  be a finite group.*

- (i) *Let  $\pi$  be a set of odd primes and suppose that the  $\pi$ -group  $P$  acts as a group of automorphisms on the solvable finite  $\pi'$ -group  $G$ . Then*

$$C_{[G,P]}(P) = \langle C_{[g,P]}(P) : g \in G \rangle.$$

- (ii) *Let  $\alpha$  be a coprime automorphism of odd order of  $G$ . Then*

$$C_{[G, \langle \alpha \rangle]}(\alpha) = \langle C_{[g, \langle \alpha \rangle]}(\alpha) : g \in [G, \langle \alpha \rangle] \rangle.$$

*Proof.* The first claim is [12, Theorem A] and the second is [14, Theorem 2]. □

We will use the next result repeatedly.

**Lemma 2.10.** *Let  $\pi$  be a nonempty set of primes. Let  $Q$  be a finite  $\pi$ -group which acts fixed point freely on a finite  $\pi'$ -group  $R$ , that is,  $C_R(Q) = 1$ , then  $R$  is solvable.*

*Proof.* For a proof, see Theorem 2.3 in [29]. □

We also need the following consequence of Burnside's normal  $p$ -complement theorem.

**Lemma 2.11.** *Let  $G$  be a finite group and let  $p$  be a prime. Let  $H$  be an abelian  $p$ -subgroup of  $G$ . Assume that  $\langle H^G \rangle \cap N_G(H) = H$ . Then  $G = N_G(H) O_{p'}(G)$  and  $\langle H^G \rangle$  is solvable. Moreover,  $N_G(H)$  has a normal complement in  $G$ , which is  $O_{p'}(\langle H^G \rangle)$ .*

*Proof.* Let  $H$  be an abelian  $p$ -subgroup of  $G$ . Assume that  $\langle H^G \rangle \cap N_G(H) = H$ . Let  $L = \langle H^G \rangle \trianglelefteq G$ . Then  $N_L(H) = L \cap N_G(H) = H$ , so  $H$  is a self-normalizing abelian Sylow  $p$ -subgroup of  $L$ . In particular,  $H \leq Z(N_L(H))$ ; hence by Burnside's normal  $p$ -complement theorem [22, Theorem 7.4.3]  $L = HO_{p'}(L)$ . By Frattini's argument

$$G = N_G(H)L = N_G(H)O_{p'}(L) = N_G(H)O_{p'}(G).$$

The last equality holds since  $O_{p'}(L) \leq O_{p'}(G)$ .

Let  $L = \langle H^G \rangle$ . Then  $N_L(H) = H$ , that is,  $H$  is a self-normalizing cyclic Sylow  $p$ -subgroup of  $L$ . By Burnside's normal  $p$ -complement theorem,  $L$  has a normal  $p$ -complement  $K$ , and hence  $C_K(H) = 1$ . By Lemma 2.10,  $K$  is solvable and thus  $L = HK$  is solvable as well.

We now show that  $O_{p'}(L)$  is a normal complement to  $N_G(H)$  in  $G$ . To see this, observe that  $L = HO_{p'}(L)$  and  $H \cap O_{p'}(L) = 1$ . Note that  $O_{p'}(L) \trianglelefteq G$ . Thus it suffices to show that  $N_G(H) \cap O_{p'}(L) = 1$ . Indeed, we have

$$N_G(H) \cap O_{p'}(L) = N_G(H) \cap L \cap O_{p'}(L) = H \cap O_{p'}(L) = 1. \quad \square$$

Finally, we also need the following solvability result.

**Lemma 2.12.** *Let  $G$  be a finite group and let  $H$  be an abelian  $p$ -subgroup of  $G$  for some prime  $p$ . If  $H$  is extremely closed in  $G$ , then  $\langle H, H^g \rangle$  is solvable for all  $g \in G$ .*

*Proof.* Let  $g \in G$  and let  $T = \langle H, H^g \rangle$ . By Lemma 2.2(v), we have  $T = HO_{p'}(T)$ . Since  $N_T(H) = H$ ,  $H$  acts fixed point freely and coprimely on  $O_{p'}(T)$ , the claim now follows from Lemma 2.10.  $\square$

### 3. Extremely closed abelian $p$ -subgroups

We are now ready to prove the main theorems. We first prove Theorem 1.4.

*Proof of Theorem 1.4.* Assume that  $M$  is a maximal subgroup of  $G$ ,  $H$  is a maximal subgroup of  $M$  and that  $\langle H, H^g \rangle \cap M = H$  for all  $g \in G$ . We will show that  $\langle H^G \rangle \cap M = H$ . The hypothesis implies that  $H \trianglelefteq M$  and  $|M : H|$  is a prime.

Clearly, if  $H \trianglelefteq G$ , then the conclusion holds. So, let  $M = N_G(H)$ . Suppose that there exists a subgroup  $D \leq M$  with  $D \not\leq H$  and  $N_G(D) \not\leq M$ . Take  $g \in N_G(D) - M$ . Then  $M = DH$  and  $M^g = DH^g$ . As  $M \neq M^g$ ,  $G = D\langle H, H^g \rangle$ . We have

$$\langle H^G \rangle = \langle H^{D\langle H, H^g \rangle} \rangle \leq \langle H, H^g \rangle \leq \langle H^G \rangle$$

and so  $\langle H^G \rangle = \langle H, H^g \rangle$ . Thus  $\langle H^G \rangle \cap M = \langle H, H^g \rangle \cap M = H$  and we are done.

So, we can assume that whenever  $D \leq M$  with  $D \not\leq H$  then  $N_G(D) \leq M$ . By Lemma 2.6,  $(G, M, H)$  is a  $W$ -triple and so the result follows from Lemma 2.7.  $\square$

*Proof of Corollary 1.5.* Let  $G$  be a finite nonabelian simple group. Let  $M$  be a maximal subgroup of  $G$  and let  $H$  be a normal subgroup of  $M$  such that  $|M : H| = p$  is a prime. Suppose by contradiction that  $G \neq \langle H, H^g \rangle$  for all  $g \in G$ . Since  $G$  is nonabelian simple, either  $\langle H^G \rangle = G$  or  $H = 1$ . If  $H = 1$ , then  $M$  is a self-normalizing cyclic subgroup of  $G$  of prime order  $p$  and Burnside's normal  $p$ -complement theorem implies that  $G$  has a normal  $p$ -complement, a contradiction.

So  $H \neq 1$  and  $\langle H^G \rangle = G$ . In particular,  $M = N_G(H)$  and  $\langle H^G \rangle \cap M = M > H$ . Since  $H$  is maximal in  $M$  and  $M$  is maximal in  $G$ , Theorem 1.4 implies that there exists  $g \in G$  such that  $\langle H, H^g \rangle \cap M > H$ . The maximality of  $H$  in  $M$  implies that  $M \leq \langle H, H^g \rangle$ . Hence  $M = \langle H, H^g \rangle$  and  $N_G(H^g) = M^g \neq M$  as  $g \notin M$ . Let  $a \in M^g - M$ , then  $H^a H^g = M^a$  is a subgroup and  $H^g \langle H, H^a \rangle = G$ . Now  $G = \langle H^G \rangle = H^{\langle H, H^a \rangle} \leq \langle H, H^a \rangle$ , contradicting our assumption. Thus  $G = \langle H, H^g \rangle$  for some  $g \in G$ .  $\square$

*Proof of Theorem 1.6.* Suppose we have proven that  $\langle H^G \rangle \cap N_G(H) = H$ . By Lemma 2.11, we have  $G = N_G(H) O_{p'}(G)$  and  $\langle H^G \rangle$  is solvable.

It remains to show that if  $H$  is extremely closed in  $G$ , then  $\langle H^G \rangle \cap N_G(H) = H$ . Let  $G$  be a counterexample to the claim with minimal order. Then  $H$  is a cyclic group of odd prime order  $p$ ,  $\langle H, H^g \rangle \cap N_G(H) = H$  for all  $g \in G$  but  $\langle H^G \rangle \cap N_G(H) \neq H$ . Furthermore, since  $|H| = p$  is prime,  $H^g = H$  or  $\langle H, H^g \rangle$  is a Frobenius group for all  $g \in G$ .

(1) We first claim that  $\langle H^G \rangle = G$ . Suppose by contradiction that  $L := \langle H^G \rangle < G$ . By Lemma 2.2(ii),  $H$  is pronormal in  $G$  and hence  $L = \langle H^L \rangle$  by Lemma 2.1(vi). By Lemma 2.2(iii),  $H$  is extremely closed in  $L$  and so by the minimality of  $G$ ,  $\langle H^L \rangle \cap N_L(H) = H$ . However, as  $L = \langle H^L \rangle$ , we have  $N_L(H) = H$ , and thus

$$\langle H^G \rangle \cap N_G(H) = L \cap N_G(H) = N_L(H) = H.$$

This contradiction proves the claim.

(2) Assume that  $N_G(H) = C_G(H)$ . Write  $H = \langle x \rangle$ . Then  $\langle x, x^g \rangle$  is a Frobenius group for all  $g \in G$ . Hence  $x$  acts as a Frobenius automorphism on  $G'$  by Lemma 2.8 and so  $G'H$  is a Frobenius group (as  $H$  has prime order). In particular, we have  $G' \leq O_{p'}(G)$  and  $G = N_G(H) O_{p'}(G)$  in this case. In addition,  $\langle H^G \rangle = G'H$  and  $\langle H^G \rangle \cap N_G(H) = H$ , a contradiction.

(3) Assume that  $N_G(H) > C_G(H)$ . Let  $M$  be a maximal normal subgroup of  $G$ . Then  $G/M$  is a simple group. Since  $G = \langle H^G \rangle$ , we have  $H \not\leq M$ . Assume that  $M > 1$ . Then  $|G/M| < |G|$ . Lemma 2.2(vi) implies that  $HM/M$  is extremely closed in  $G/M$ . Hence, as  $G/M$  is simple,  $G = HM$ . Now  $N_G(H) = N_M(H)H = C_G(H)$ , which is a contradiction. Hence  $M = 1$  and  $G$  is a nonabelian simple group.

(4) By Lemma 2.12,  $\langle x, x^g \rangle$  is solvable for all  $g \in G$  and so by Lemma 2.5,  $p = 3$  and  $G$  is a finite simple group of Lie type defined over  $\mathbb{F}_3$  or  $G \cong U_n(2)$  with  $n \geq 4$ . Furthermore, except for  $U_3(3)$ , the Lie rank of  $G$  is strictly greater than 1. Now it is easy to see that a Sylow 3-subgroup of  $G$  is nonabelian. Let  $P$  be a Sylow 3-subgroup of  $G$  containing  $x$ . Then  $H = \langle x \rangle$  is isolated in  $P$  with respect to  $G$ , that is,  $H$  does not conjugate in  $G$  to any subgroup in  $P - H$ . By [23, Theorem 4.250], we deduce that  $G \cong U_3(3)$ . By [25, Theorem A\*],  $x$  is a transvection. Now for any conjugate  $x^g$  of  $x$  different from  $x$  and  $x^{-1}$ , we have that  $T := \langle x, x^g \rangle \cong \text{SL}_2(3)$ . However,  $\text{SL}_2(3)$  is not a Frobenius group.  $\square$

We will need the following result which is a consequence of Theorem A\* in [25] and Theorem 1.2 in [15].

**Proposition 3.1.** *Let  $G$  be a finite group and let  $p$  be an odd prime. Let  $H$  be a nontrivial abelian  $p$ -subgroup of  $G$ . Assume that  $G$  has a unique minimal normal subgroup  $N$  which is nonabelian such that  $G = HN$ . Then  $H$  is not extremely closed in  $G$ .*

*Proof.* Suppose by contradiction that  $H$  is an abelian extremely closed  $p$ -subgroup of  $G$ . By Lemma 2.12, we have  $\langle H, H^g \rangle$  is solvable for all  $g \in G$ . In particular, if  $x \in H$ , then  $\langle x, x^g \rangle$  is solvable for all  $g \in G$ .

By the uniqueness of  $N$ , we have  $G = \langle H^G \rangle = HN$ . We first show that  $G$  is an almost simple group with socle  $S$ . Let  $W \trianglelefteq N$  be a component of  $G$ . Assume that  $W \neq N$ . As  $W \trianglelefteq N$ ,  $N \leq N_G(W)$  and so  $N_G(W) = N_H(W)N$ . Since  $W < N$ ,  $N_G(W) < G$  and so  $N_H(W) < H$ . Let  $x \in H - N_H(W)$ . Then  $x \notin N_G(W)$  and  $x^2 \notin C_G(W)$  since  $p$  is odd, whence  $x$  and  $W$  satisfy the hypothesis of Lemma 2.4 and hence  $\langle x, x^g \rangle$  is not solvable for some  $g \in G$ , a contradiction. Thus  $W \trianglelefteq G$  and so  $G = HN$ , where  $N$  is a nonabelian simple group which is also a minimal normal subgroup of  $G$ . Thus  $G$  is an almost simple group with socle  $S$  as wanted.

Let  $x \in H$  be an element of order  $p$ . Then  $\langle x, x^g \rangle$  is solvable for all  $g \in G$ . By Lemma 2.5,  $p = 3$  and  $S$  is a finite simple group of Lie type defined over  $\mathbb{F}_3$  or  $S \cong U_n(2)$ ,  $n \geq 4$ . If  $|H| = 3$ , then  $\langle H^G \rangle$  is solvable by Theorem 1.6 and Lemma 2.11, which is impossible. Thus we may assume that  $|H| \geq 9$ . Since  $G = HS$ , we have  $G/S \cong H/(H \cap S)$  is a 3-group and thus if  $G \neq S$ , then elements in  $H - S$  induce outer automorphisms of  $S$  of 3-power order.

(i) Assume  $S \cong U_n(2)$ ,  $n \geq 4$  or  $G \neq S$ . If  $S \cong U_n(2)$ , then since  $n \geq 4$ , we have  $|\text{Out}(U_n(2))| = 2(n, 3)$ . If  $G \neq S$  and  $S \not\cong U_n(2)$ , then  $S \cong D_4(3)$  or  ${}^3D_4(3)$  since  $S$  has no nontrivial field automorphism so  $\text{Out}(S)$  contains diagonal automorphisms and possibly graph automorphisms only. In all cases, the Sylow 3-subgroup of  $\text{Out}(S)$  has order at most 3. Hence  $|G : S| = |H : S \cap H| \leq 3$ .

Since  $|H| \geq 9$ , if  $G$  is not simple then  $A = H \cap S > 1$ . By Lemma 2.3(ii),  $A$  is strongly closed in  $S$ . If  $G = S = U_n(2)$ , then  $A = H$  is strongly closed in  $S$ . In

both cases,  $S$  contains a nontrivial abelian 3-subgroup  $A$  which is strongly closed in  $S$ . Since  $S$  is simple,  $S = \langle A^S \rangle$ . As a Sylow 3-subgroup of  $S$  is nonabelian,  $A$  cannot be a Sylow 3-subgroup of  $S$ ; however this contradicts [15, Theorem 1.2(i)].

(ii)  $G = S$  is a finite simple group of Lie type defined over  $\mathbb{F}_3$ . The possibilities for  $G$  can be read off from [25, Table 1] and note the correction in [37, Remark 5.2].

Let  $P$  be a Sylow  $p$ -subgroup of  $G$  containing  $H$ , and let  $M$  be a maximal parabolic subgroup of  $G$  containing the Borel subgroup  $B := N_G(P)$ . Let  $R = O_p(M)$ . Then  $M = N_G(R)$ . By induction, we have  $\langle H^M \rangle \cap N_M(H) = H$ , and hence  $\langle H^M \rangle = HU$ , where  $U = O_{p'}(\langle H^M \rangle)$  and  $C_U(H) = 1$ . Therefore,  $U$  is a solvable normal  $p'$ -subgroup of  $M$  since  $\langle H^M \rangle \trianglelefteq M$ . As  $H \leq \langle H^M \rangle \trianglelefteq M$  and  $H$  is pronormal in  $M$ , we have  $M = N_M(H)U$ . However, as  $M$  is a parabolic subgroup of a finite simple group of Lie type  $G$ ,  $F^*(M) = O_p(M)$  (Corollaries 3.1.4 and 3.1.5 in [24]), so  $U = 1$ . Hence  $H \trianglelefteq M$  and  $M = N_G(H)$ . Thus  $M$  contains every maximal parabolic subgroup of  $G$  that contains the Borel subgroup  $B$ . However this happens only when the Lie rank of  $G$  is 1. Therefore  $G \cong U_3(3)$ . (We can also use Theorem 1.2 in [15] to arrive at this conclusion.)

From the Atlas [7], we have  $P \cong 3_+^{1+2}$  is an extraspecial group of order 27 and exponent 3. Thus  $|H| = 9$ . Since  $G$  has Lie rank 1, the Borel subgroup  $B$  of  $G$  is a maximal subgroup of  $G$ . Hence  $B = N_G(H)$ . Let  $g \in G - B$  and  $T = \langle H, H^g \rangle$ . We know that  $T = HO_{3'}(T)$  is solvable, so  $T < G$  and thus  $T$  lies in some maximal subgroup of  $G$  whose order must be divisible by  $|H| = 9$ . Inspecting the list of maximal subgroups of  $G$  in the Atlas [7], the only maximal overgroups of  $H$  and  $T$  in  $G$  are the Borel subgroups  $B$  and its  $G$ -conjugates. Hence  $\langle H, H^g \rangle \leq B^t$  for some  $t \in G$ . Note that  $B = P : W$ , where  $W = \langle y \rangle$  is a cyclic group of order 8. Since  $B^t$  has a normal Sylow 3-subgroup, we must have that  $H \leq P^t \trianglelefteq B^t$  and hence  $H$  is subnormal in  $B^t$ . Since  $H$  is pronormal in  $G$  and hence in  $B^t$ , we have  $H \trianglelefteq B^t$  or  $B^t = N_G(H) = B$ . Therefore,  $\langle H, H^g \rangle \leq B$  for all  $g \in G$ . However, this implies that  $\langle H^G \rangle \leq B$ , which is a contradiction.  $\square$

*Proof of Theorem 1.7.* Let  $G$  be a counterexample to Theorem 1.7 with minimal order. Then  $\langle H, H^g \rangle \cap N_G(H) = H$  for all  $g \in G$  but  $G \neq N_G(H)O_{p'}(G)$ , where  $H$  is an abelian  $p$ -subgroup of  $G$ .

(1)  $\langle H^G \rangle = G$ . Let  $L = \langle H^G \rangle$ . Assume  $L \neq G$ . By Lemmas 2.2(ii) and 2.1(iii), we have  $G = N_G(H)L$ . By Lemma 2.2(iii) and the minimality of  $G$ , we have  $L = N_L(H)O_{p'}(L)$ . As  $L \trianglelefteq G$ ,  $O_{p'}(L) \leq O_{p'}(G)$ , and hence

$$G = N_G(H)L = N_G(H)O_{p'}(L) = N_G(H)O_{p'}(G).$$

This contradiction shows that  $L = G$ .

(2)  $O_{p'}(G) = 1$ . Suppose by contradiction that  $N = O_{p'}(G) \neq 1$ . It follows from Lemma 2.2(vi) that the hypothesis carries over to  $\bar{G} = G/N$ , and so by the minimality

of  $G$ , we obtain  $\bar{G} = N_{\bar{G}}(\bar{H}) O_{p'}(\bar{G})$ . But  $O_{p'}(\bar{G}) = O_{p'}(G/O_{p'}(G)) = 1$ , and hence  $\bar{G} = N_{\bar{G}}(\bar{H})$  by Lemma 2.2(iv). Therefore,  $G = N_G(H)N = N_G(H) O_{p'}(G)$ , which is a contradiction. This proves the claim.

(3)  $O_p(G) = 1$ . Assume  $O_p(G) \neq 1$ . Let  $N \leq O_p(G)$  be a minimal normal subgroup of  $G$  and let  $U = H \cap N$ . Observe first that  $O_p(G) \leq N_G(H)$  since  $P \leq N_G(H)$  for any  $H \leq P \in \text{Syl}_p(G)$ . It follows that  $N \leq N_G(H)$ . We will show that  $N \leq Z(G)$ . Suppose first that  $U \neq 1$ . Lemma 2.3(iii) implies that  $U \trianglelefteq G$ . By the minimality of  $N$ , we have  $U = N$ , and so  $N \leq H$ . As  $H$  is abelian,  $H \leq C_G(N) \trianglelefteq G$ . By (1), we have  $C_G(N) = G$ , and  $N \leq Z(G)$ . If  $U = H \cap N = 1$ , then as  $H$  and  $N$  are both normal in  $N_G(H)$ , we have  $[H, N] \leq H \cap N = 1$  and thus  $H \leq C_G(N) \trianglelefteq G$ , so  $N \leq Z(G)$ .

By Lemma 2.2(vi), the hypothesis carries over to  $\bar{G} = G/N$ , and so by the minimality of  $G$ , we obtain  $\bar{G} = N_{\bar{G}}(\bar{H}) O_{p'}(\bar{G})$ . Let  $K \leq G$  be such that  $N \leq K$  and  $K/N = O_{p'}(G/N)$ . Then  $N \trianglelefteq K \trianglelefteq G$  and  $|K : N|$  is odd. By the Schur-Zassenhaus theorem,  $K = NT$  where  $T$  is Hall  $p'$ -subgroup of  $G$ . Moreover, since  $N$  is central in  $G$ ,  $T \trianglelefteq K$  and hence  $T \trianglelefteq G$  as it is characteristic in  $K \trianglelefteq G$ . It follows that  $T \leq O_{p'}(G) = 1$  and thus  $G = N_G(H)K = N_G(H)N = N_G(H)$ . This contradiction proves the claim.

(4) Let  $N$  be a minimal normal subgroup of  $G$ . We claim that  $G = HN$  and  $N$  is the unique minimal normal subgroup of  $G$ . By (2) and (3),  $N \cong S^k$  for some finite nonabelian simple group  $S$  with  $p \mid |S|$  and some integer  $k \geq 1$ . Let  $M = HN$ . Suppose that  $M < G$ . By the minimality of  $G$ , we have  $M = N_M(H) O_{p'}(M)$ . Hence  $O_{p'}(M) \leq N$ , and so  $O_{p'}(M) = 1$ . We deduce that  $H \trianglelefteq M$ . As  $H$  is an abelian  $p$ -group and  $N \cong S^k$ , we have  $H \cap N = 1$  and hence  $[H, N] \leq H \cap N = 1$ . Thus  $H \leq C_G(N) \trianglelefteq G$ . By (1), we have  $C_G(N) = G$ , and then  $N \leq Z(G)$ . This contradiction shows that  $G = HN$ . Since  $G/N \cong H/H \cap N$  is abelian,  $N$  must be a unique minimal normal subgroup of  $G$  as wanted.

If  $p$  is odd, then Proposition 3.1 yields a contradiction. Thus for the remaining, we assume that  $p = 2$ .

(5) We next claim that  $G$  is finite nonabelian simple group. By (4),  $F^*(G) = N$  and  $H$  is a strongly closed abelian 2-subgroup of  $G$ . Now by [20, Theorem A], we have  $F^*(G) = G$  and so  $N = G$ . It follows that  $G = S$  is simple. By [20, Theorem A] again,  $G$  is isomorphic to one of the following groups:

- (i)  $L_2(2^n)$ ,  $n \geq 3$ ;  ${}^2B_2(2^{2n+1})$ ,  $n \geq 1$ ; or  $U_3(2^n)$ ,  $n \geq 2$ .
- (ii)  $L_2(q)$ ,  $q \equiv 3, 5 \pmod{8}$ .
- (iii)  ${}^2G_2(3^{2n+1})$ ,  $n \geq 1$ ; or  $J_1$ , the first Janko group.

By Glauberman's  $Z^*$ -theorem, we may assume that  $|H| \geq 4$ .

(6) The final contradiction. We now consider each case above separately.

(a) Assume  $G$  is isomorphic to one of the groups in (i). Let  $H \leq P \in \text{Syl}_2(G)$  and let  $B = PT$  be the Borel subgroup of  $G$  containing  $P$ . By [20, (3.2)],  $H = Z(P)$  is a noncyclic elementary abelian 2-group,  $P$  is a  $T.I$  subgroup of  $G$ , and  $B$  is the unique maximal subgroup of  $G$  containing  $P$ . It follows that  $B = N_G(H)$ . Observe that for any  $1 \neq x \in H$ ,  $P \leq C_G(x) \leq B$ , as  $P$  is uniquely contained in  $B$ . For  $g \in G - B$ , let  $T = \langle H, H^g \rangle$ . Then  $N_T(H) = H$  and then by Burnside's normal  $p$ -complement theorem, we have  $T = HU$ , where  $U = O_{2'}(T) \trianglelefteq T$ . As  $H$  is noncyclic abelian,  $U = \langle C_U(a) : 1 \neq a \in H \rangle$  [35, 8.3.4]. However, as  $C_G(a) \leq B$  for all  $1 \neq a \in H$ , we have  $U \leq B$  and so  $T = \langle H, H^g \rangle \leq B$ . From the hypothesis, we must have  $T = T \cap B = H$ , and hence  $H^g = H$ . This implies that  $g \in B$ , contradicting the choice of  $g$ .

(b) Assume that  $G \cong L_2(5) \cong A_5$ . By [20, (3.4)],  $H \in \text{Syl}_2(G)$ ,  $N_G(H) \cong A_4$  and  $N_G(H)$  is the unique maximal subgroup of  $G$  containing  $C_G(a)$ , for all  $1 \neq a \in H$ . Take  $g \in G - N_G(H)$ , and let  $T = \langle H, H^g \rangle$ . Then  $T = HU$ , where  $U = O_{2'}(T)$ . As  $H$  is noncyclic,  $U = \langle C_U(a) : 1 \neq a \in H \rangle \leq N_G(H)$ . This leads to a contradiction as in the previous case.

(c) Assume  $G$  is isomorphic to one of the groups in (ii) with  $q \geq 11$ . By [20, (3.4)], we have  $H \in \text{Syl}_2(G)$  and  $N_G(H) \cong A_4$ . Clearly,  $G$  contains a maximal subgroup  $M$  isomorphic to the dihedral group  $D_{q\pm 1}$  such that  $M$  does not contain  $H$ . Assume  $M$  is generated by two involutions  $a, b$ . We can choose  $a, b$  such that  $b \in H$ . Now  $G = \langle M, H \rangle \leq \langle a, H \rangle$ , and hence  $G = \langle a, H \rangle$ . By Sylow's theorem, there exists some  $g \in G$  such that  $a \in H^g$ . Thus  $G = \langle a, H \rangle \leq \langle H^g, H \rangle$ , and then  $G = \langle H, H^g \rangle$ , which contradicts the hypothesis that  $\langle H, H^g \rangle \cap N_G(H) = H$ .

(d) Finally, assume that  $G$  is isomorphic to one of the groups in (iii). By [20, (3.4)], we have  $H \in \text{Syl}_2(G)$ . Now  $H$  contains an involution  $t$  such that  $C_G(t) = \langle t \rangle \times L$ , where  $L \cong L_2(q)$ ,  $q \equiv 3, 5 \pmod{8}$  and  $[G : C_G(t)]$  is odd [41]. As  $H \leq C_G(t) < G$ , by the minimality of  $G$ , we obtain  $C_G(t) = (C_G(t) \cap N_G(H)) O_{2'}(C_G(t))$ . However, as  $C_G(t) = \langle t \rangle \times L$ , where  $L$  is nonabelian simple, it follows that  $O_{2'}(C_G(t)) = 1$  whence  $C_G(t) \leq N_G(H)$ . Hence  $C_G(t) = N_G(H)$  since  $C_G(t)$  is maximal in  $G$  by [20, (3.4)]. It follows that  $H \trianglelefteq C_G(t)$  and then  $H \cap L \trianglelefteq L$ , where  $H \cap L \in \text{Syl}_2(L)$ , which contradicts the simplicity of  $L$ . □

*Proof of Corollary 1.8.* Let  $G$  be a counterexample to the corollary with minimal order. Then we have that  $H$  is an abelian  $p$ -subgroup for some odd prime  $p$  and  $\langle H, H^g \rangle \cap N_G(H) = H$  for all  $g \in G$  but  $\langle H^G \rangle \cap N_G(H) \neq H$ . By Theorem 1.7, we have  $G = N_G(H) O_{p'}(G)$ . Now we have that

$$\langle H^G \rangle = \langle H^{N_G(H) O_{p'}(G)} \rangle = \langle H^{O_{p'}(G)} \rangle \leq H O_{p'}(G).$$

Let  $L = \langle H^G \rangle$ . Then  $L = H(L \cap O_{p'}(G)) = H O_{p'}(L)$  and so  $L$  has a normal  $p$ -complement. Let  $L < G$ . By the minimality of  $G$ , we have  $\langle H^L \rangle \cap N_L(H) = H$ .

However,  $L = \langle H^L \rangle$  by Lemmas 2.2(ii) and 2.1(vi) and thus  $N_L(H) = H$  or equivalently  $\langle H^G \rangle \cap N_G(H) = H$ , a contradiction. Therefore, we can assume that  $G = \langle H^G \rangle = HO_{p'}(G)$ .

Let  $U = O_{p'}(G)$  and let  $Q = [U, H]$ . By [35, 8.2.7],  $U = QC_U(H)$  and  $Q = [Q, H]$ . Furthermore,  $N_G(H) = C_G(H) = HC_U(H)$  and thus  $G = HC_U(H)Q$ . As  $G = \langle H^G \rangle = H^{HC_U(H)Q} \leq HQ$ , we obtain  $G = HQ$  whence  $U = Q$ .

Let  $N \leq U$  be a minimal normal subgroup of  $G$ . Since  $HN/N$  is an abelian extremely closed  $p$ -subgroup of  $G/N$  by Lemma 2.2(vi) and  $G/N = \langle (HN/N)^{G/N} \rangle$ , we have  $N_G(HN) = HN$  and hence by Lemma 2.2 (iv),  $N_G(H)N = HN$ . Moreover,  $C_{U/N}(HN/N) = 1$ . It follows that  $U/N$  is solvable by Lemma 2.10.

Assume that  $N$  is abelian. Then  $U$  is solvable. Let  $1 \neq u \in U$  and let  $T = \langle H, H^u \rangle$ . Then  $T = H[u, H]$ , where  $[u, H] = O_{p'}(T)$  and  $C_{[u, H]}(H) = 1$  as  $H$  is self-normalizing in  $T$ . By Lemma 2.9(i),  $C_{[U, H]}(H) = \langle C_{[u, H]}(H) : u \in U \rangle = 1$ . Thus  $C_U(H) = C_{[U, H]}(H) = 1$  and so  $N_G(H) = C_G(H) = HC_U(H) = H$ . Therefore,  $\langle H^G \rangle \cap N_G(H) = H$ , which is a contradiction.

Assume that  $N \cong S^k$ , where  $S$  is a nonabelian simple group and  $k \geq 1$  is an integer. Assume that  $K = HN < G$ . By the minimality of  $G$ , we have  $\langle H^K \rangle \cap N_K(H) = H$ . Thus  $\langle H^K \rangle = HQ$  and  $C_Q(H) = 1$ , where  $Q = O_{p'}(\langle H^K \rangle)$ . As  $Q$  is characteristic in  $\langle H^K \rangle \trianglelefteq K$ , it follows that  $Q \trianglelefteq K$ . Since  $|K : N|$  is a power of  $p$  and  $Q$  is a  $p'$ -group, we must have  $Q \leq N$  and hence  $Q \trianglelefteq N$ . By [35, 1.7.5],  $Q$  is isomorphic to a direct product of the nonabelian simple group  $S$ , so  $Q$  is not solvable or  $Q = 1$ . If the former case holds, then since  $C_Q(H) = 1$  and  $Q$  is a  $p'$ -group, Lemma 2.10 implies that  $Q$  is solvable, which is a contradiction as it is a direct product of copies of  $S$ . Therefore,  $Q = 1$  and hence  $H \trianglelefteq K$ . It follows that  $R = H \cap N \trianglelefteq N$ . However, as  $N \cong S^k$  and  $H \cap N$  is a normal  $p$ -subgroup of  $N$ , we must have  $H \cap N = 1$ . Hence  $[H, N] \leq H \cap N = 1$  and so  $H \leq C_G(N) \trianglelefteq G$ . Since  $G = \langle H^G \rangle$ , we have  $C_G(N) = G$  or  $N \leq Z(G)$ , a contradiction. Therefore  $G = HN$  and since  $G/N$  is solvable,  $N$  is a unique minimal normal subgroup of  $G$ . Now Proposition 3.1 yields a contradiction. □

*Proof of Corollary 1.9.* Let  $H$  be an extremely closed abelian  $p$ -subgroup of  $G$  for some prime  $p$ . Assume first that  $p = 2$ . By Theorem 1.7,  $G = N_G(H)O_2(G)$  which implies that  $HO_2(G) \trianglelefteq G$  and clearly  $HO_2(G)$  is solvable and thus  $H \subseteq R(G)$ . Assume now that  $p$  is odd. By Corollary 1.8, we have  $N_G(H) \cap \langle H^G \rangle = H$  and thus  $\langle H^G \rangle$  is solvable by Lemma 2.11. □

#### 4. A $p$ -solvability criterion

Let  $p$  be a prime. A finite group  $G$  is said to be a minimal non- $p$ -solvable group if  $G$  is not  $p$ -solvable but every proper subgroup of  $G$  is  $p$ -solvable. A minimal simple group is a nonabelian finite simple groups whose all proper subgroups are solvable.

Observe that minimal non-2-solvable simple groups are exactly the minimal simple groups and these groups are classified by Thompson in [39].

**Lemma 4.1.** *Every minimal simple group is isomorphic to one of the following simple groups:*

- (1)  $L_2(2^r)$ ,  $r$  is a prime.
- (2)  $L_2(3^r)$ ,  $r$  is an odd prime.
- (3)  $L_2(r)$ ,  $r > 3$  is a prime such that  $5 \mid r^2 + 1$ .
- (4)  ${}^2B_2(2^r)$ ,  $r$  is an odd prime.
- (5)  $L_3(3)$ .

*Proof.* This is [39, Corollary 1]. □

The next result classifies minimal non-3-solvable simple groups.

**Lemma 4.2.** *Let  $G$  be a finite nonabelian simple group. Assume that every proper subgroup of  $G$  is 3-solvable. Then  $G$  is isomorphic to a minimal simple group or to the Suzuki group  ${}^2B_2(q)$  with  $q = 2^{2m+1}$ ,  $m \geq 1$ .*

*Proof.* This is Lemma 5.3 in [28]. □

Finally, we need the classification of finite non- $p$ -solvable simple groups for any primes  $p \geq 5$ .

**Lemma 4.3.** *Let  $G$  be a finite nonabelian simple group and let  $p \geq 5$  be a prime dividing  $|G|$ . Assume that every proper subgroup of  $G$  is  $p$ -solvable. Then one of the following holds.*

- (1)  $G = L_2(p)$ .
- (2)  $G = A_p$ .
- (3)  $G = L_2(q)$  with  $p \mid q^2 - 1$ .
- (4)  $G = L_n(q)$ ,  $n \geq 3$  is odd, and  $p$  divides  $q^n - 1$  but not  $\prod_{i=1}^{n-1} (q^i - 1)$ .
- (5)  $G = U_n(q)$ ,  $n \geq 3$  is odd, and  $p$  divides  $q^n - (-1)^n$  but not  $\prod_{i=1}^{n-1} (q^i - (-1)^i)$ .
- (6)  $G = {}^2B_2(q)$  with  $q = 2^{2m+1}$ ,  $m \geq 1$ .
- (7)  $G = {}^2G_2(q)$  and  $p \mid (q^2 - q + 1)$ , where  $q = 3^{2m+1}$ ,  $m \geq 1$ .
- (8)  $G = {}^2F_4(q)$  with  $q = 2^{2m+1}$ ,  $m \geq 1$  and  $p \mid (q^4 - q^2 + 1)$ .
- (9)  $G = {}^3D_4(q)$  and  $p \mid (q^4 - q^2 + 1)$ .
- (10)  $G = E_8(q)$  and  $p$  divides  $(q^{30} - 1)$  but not  $\prod_{i \in \{8, 14, 18, 20, 24\}} (q^i - 1)$ .
- (11)  $(G, p)$  is one of the following:  $(M_{23}, 23)$ ,  $(J_1, 7 \text{ or } 19)$ ,  $(Ly, 37 \text{ or } 67)$ ,  $(J_4, 29 \text{ or } 43)$ ,  $(Fi'_{24}, 29)$ ,  $(B, 47)$  or  $(M, 41 \text{ or } 59 \text{ or } 71)$ .

*Proof.* This is Lemma 5.4 in [28]. □

Let  $q$  be a prime power and let  $n \geq 2$  be an integer. A prime divisor  $p$  of  $q^n - 1$  is called a primitive prime divisor or ppd of  $q^n - 1$  if  $p$  does not divide  $q^k - 1$  for all integers  $k$  with  $1 \leq k < n$ . Zsigmondy's theorem [43] states that such a ppd  $p$  exists unless  $(n, q) = (6, 2)$  or  $n = 2$  and  $q$  is a Mersenne prime. Now if  $n > 1$  is an integer and  $p$  is a prime, then the  $p$ -part of  $n$ , denoted by  $n_p$ , is the largest power of  $p$  dividing  $n$ . We refer the reader to [5; 34] for the description of maximal subgroups of finite simple groups of Lie type.

**Proposition 4.4.** *Let  $G$  be a finite nonabelian simple group and  $p$  be a prime dividing  $|G|$ . Assume that every proper subgroup of  $G$  is  $p$ -solvable. Let  $P$  be a Sylow  $p$ -subgroup of  $G$ . Then  $G = \langle P, P^g \rangle$  for some  $g \in G$ .*

*Proof.* Let  $G$  be a finite nonabelian simple group and let  $p$  be a prime dividing  $|G|$ . Let  $P \in \text{Syl}_p(G)$ . Assume that every proper subgroup of  $G$  is  $p$ -solvable. Let  $x \in P$  with  $|x| = p$ . Then  $\langle x, x^g \rangle \leq \langle P, P^g \rangle$  for all  $g \in G$  and thus  $\langle x, x^g \rangle$  is  $p$ -solvable for all  $g \in G$ .

(a) If  $p = 2$ , then every finite nonabelian simple group  $G$  can be generated by two Sylow 2-subgroups by Theorem A in [26]. So, we may assume that  $p > 2$ .

(b) If  $G$  is a finite nonabelian simple group of Lie type in characteristic  $p$ , then  $G$  is generated by two Sylow  $p$ -subgroups by Proposition 2.5 in [6].

(c) Assume that  $p = 3$ . By Lemma 4.2, since 3 divides  $|G|$ ,  $G$  is a minimal simple group. By part (b), we only need to consider the cases when  $G$  is isomorphic to  $L_2(2^r)$ ,  $r$  is a prime, or  $L_2(r)$ ,  $r > 3$  is a prime and  $5 \mid r^2 + 1$ .

If  $G \cong L_2(4)$ , then we can check by using GAP [17] that there exists  $g \in G$  such that  $G = \langle P, P^g \rangle$ . Assume next that  $G \cong L_2(q)$ ,  $q = 2^r$  or  $r$ , where  $r$  is an odd prime. By [25, Theorem A\*], there exists an element  $x \in G$  of order 3 such that  $\langle x, x^g \rangle$  is nonsolvable for some  $g \in G$ . Since  $G$  is minimal simple, we must have  $G = \langle x, x^g \rangle$  and thus  $G = \langle P, P^g \rangle$ .

(d) Assume that  $p \geq 5$ . By part (b), and Lemma 4.3,  $G$  is one of the groups listed in (2)–(11) in that lemma. We now consider each case in turn.

(1) Assume  $G = A_p$ . Here  $|P| = p$ . Without loss of generality, take  $P = \langle x \rangle$ , where  $x = (1, 2, \dots, p)$  is a  $p$ -cycle in  $A_p$ . Let  $y = (1, 2, p, p-1, p-2, \dots, 3) \in A_p$  be another  $p$ -cycle. Then  $xy = (1, 3, 2)$  and clearly

$$\langle x, y \rangle = \langle (xy)^{-1}, x \rangle = \langle (1, 2, 3), (1, 2, \dots, p) \rangle = A_p$$

(see, e.g., [33, Theorem B]). Hence  $G$  is generated by two Sylow  $p$ -subgroups.

(2) Assume  $G = L_2(q)$  with  $p \mid q^2 - 1$ . If  $q \leq 11$ , we can check using GAP [17] that the result holds. Assume  $q \geq 13$ . Inspecting the argument in [25, Section 5.1.2], if  $x$  is any element of order  $p$ , then we can find  $g \in G$  such that  $\langle x, x^g \rangle \cong L_2(q)$  and hence  $\langle P, P^g \rangle = G$ .

(3) Assume  $G = L_n(q)$ ,  $n \geq 3$  is odd, and  $p$  divides  $q^n - 1$  but not  $\prod_{i=1}^{n-1} (q^i - 1)$ . Write  $q = s^f$ , where  $s$  is a prime and  $f \geq 1$  is an integer. In this case  $p$  is a ppd of  $q^n - 1$ . Hence  $P \in \text{Syl}_p(G)$  is cyclic of order  $(q^n - 1)_p$ . Since  $n \equiv 1 \pmod p$  and  $n \geq 3$  is odd, we have  $p \geq 2n + 1$  and  $p \nmid n$ .

Assume that  $t$  is a prime divisor of  $n$  and write  $n = tm$  for some integer  $m \geq 1$ . Assume that  $m > 1$ . Then  $G$  has a  $\mathcal{C}_3$ -subgroup  $H$  of type  $\text{GL}_m(q^t)$  (see [34, Table 3.5A]) which is maximal and contains a Sylow  $p$ -subgroup of  $G$ . Since  $n \geq 3$  is odd,  $m, t \geq 3$  and so  $q^t \geq 2^t \geq 8$ . Therefore,  $H$  is not  $p$ -solvable.

Thus we can assume that  $n = t$  is an odd prime. If  $P$  lies in a unique maximal subgroup  $H$  of  $G$ , then  $H$  is of type  $\text{GL}_1(q^n)$  by [2, Table B]. We can choose  $g \in G - H$  such that  $P^g \not\leq H$  and hence  $G = \langle P, P^g \rangle$ . Assume that  $P$  lies in some other maximal subgroup  $M$  of  $G$  not of type  $\text{GL}_1(q^n)$ . As in the proof of Case 3 of Proposition 6.2 in [2],  $M \in \mathcal{C}_5$  is a subfield subgroup of type  $\text{GL}_n(q_0)$ , where  $q = q_0^k$ ,  $k$  is an odd prime and  $(q_0^n - 1)_p = (q^n - 1)_p$  or  $M \in \mathcal{S}$  is almost simple with socle  $S \cong L_2(p)$  and  $n = \frac{1}{2}(p - 1)$ . However, in both cases,  $M$  is not  $p$ -solvable.

(4) Assume  $G = U_n(q)$ ,  $n \geq 3$  is odd, and  $p$  divides  $q^n + 1$  but not  $\prod_{i=1}^{n-1} (q^i - (-1)^i)$ . Write  $q = s^f$  where  $s$  is a prime and  $f \geq 1$ .

(a) Assume that  $n = 3$ . If  $q = 3$ , then  $p = 7$ . In this case,  $|P| = 7$  and  $P$  lies in  $L_2(7)$  which is not 7-solvable. Similarly, if  $q = 5$ , then  $p = 7$  and  $|P| = 7$  and  $P$  lies in  $A_7$ .

First, let  $q$  be a prime. Let  $H$  be a maximal subgroup of  $G$  containing  $P$ . By the proof of [2, Proposition 6.3], either  $P$  lies in a unique maximal subgroup of  $G$  and we are done or  $P$  is contained in  $L_2(7)$  and  $p = 7$ ; however,  $L_2(7)$  is not 7-solvable.

Assume  $q = s^f$  with  $f > 1$ . In this case, if  $P$  is not contained in a unique maximal subgroup, then  $P$  can be contained in a subfield subgroup of type  $\text{GU}_3(q_0)$  with  $q = q_0^k$ , and  $k$  is an odd prime (see [5, Table 8.5]). However such a maximal subgroup is not  $p$ -solvable.

(b) Assume  $n \geq 5$ . Then  $p$  is a ppd of  $q^{2n} - 1$ . Hence  $p \geq 2n + 1$ .

Assume that  $n = tm$ , where  $t$  is a prime divisor of  $n$  and  $m > 1$ . Since  $n \geq 5$  is odd,  $t, m \geq 3$ . Then  $G$  has a maximal subgroup of type  $\text{GU}_m(q^t)$  and contains a Sylow  $p$ -subgroup of  $G$ . Since  $q^t \geq 2^t \geq 8$ , such a maximal subgroup is not  $p$ -solvable.

Therefore,  $n = t \geq 5$  is a prime. Argue as in case (3), if  $P$  lies in a unique maximal subgroup of  $G$ , then the conclusion holds. As in the proof of Proposition 6.4 in [2],  $P$  lies in a subfield subgroup  $H$  of type  $\text{GU}_n(q_0)$ , where  $q_0^k = q$  and  $k \geq 3$  is a prime, or of type  $O_n(q)$  or  $H$  is an almost simple group with socle  $L_2(p)$  with  $n = \frac{1}{2}(p - 1)$  and  $7 \leq p \equiv 3 \pmod 4$ . However, in all cases, these maximal subgroups are not  $p$ -solvable.

(5)  $G = {}^2B_2(q)$  with  $q = 2^{2m+1}$ ,  $m \geq 1$ . Then  $|G| = q^2(q-1)(q+s+1)(q-s+1)$ , where  $s = \sqrt{2q} = 2^{m+1}$ . The maximal subgroups of  $G$  are listed in [5, Table 8.16]. Since  $p$  is not the characteristic of  $G$ ,  $p > 2$  and  $p \mid q-1$  or  $p \mid q \pm s + 1$ .

Assume first that  $p \mid q-1$ . Then  $P$  lies in maximal subgroups of the form  $[q^2] : (q-1)$  and  $D_{2(q-1)}$ . It follows that  $\langle P, p^g \rangle$  is solvable for all  $g \in G$ . Let  $x \in P$  with  $|x| = p \geq 5$ . Then  $\langle x, x^g \rangle$  is solvable for all  $g \in G$ . However, this is impossible in view of Theorem A\* in [25].

Assume that  $p \mid q \pm s + 1$ . In this case,  $P$  lies in a maximal subgroup of the form  $(q \pm s + 1) : 4$  or a subfield subgroup of the form  ${}^2B_2(q_0)$ , where  $q_0^k = q$ ,  $k \geq 3$  is a prime and  $q_0 > 2$ . Clearly, the subfield subgroup is not  $p$ -solvable (if it contains  $P$ ). Hence  $P$  lies in a unique maximal subgroup of  $G$  and the result follows.

(6)  $G = {}^2G_2(q)$  with  $q = 3^{2m+1}$ ,  $m \geq 1$  and  $p \mid q^2 - q + 1$ . We can use the same argument as in the previous case using [5, Table 8.43].

(7)  $G = {}^2F_2(q)$  with  $q = 2^{2m+1}$ ,  $m \geq 1$  and  $p \mid q^4 - q^2 + 1$ . In this case,  $p$  is a ppd of  $q^{12} - 1$ . Using the argument in Proposition 7.2 in [2], either  $P$  lies in a unique maximal subgroup or it lies in a subfield subgroup  ${}^2F_2(q_0)$ , which is not  $p$ -solvable.

(8)  $G = {}^3D_4(q)$  and  $p \mid q^4 - q^2 + 1$ . We can use the argument in Proposition 7.3 in [2] to obtain the conclusion as in the previous case.

(9)  $G = E_8(q)$  and  $p$  divides  $(q^{30} - 1)$  but not  $\prod_{i \in \{8, 14, 18, 20, 24\}} (q^i - 1)$ . In this case,  $p$  is a ppd of  $q^{30} - 1$ . From Proposition 7.10 in [2], either  $P$  lies in a unique maximal subgroup and the result follows or  $P$  can lie in a maximal exotic local subgroup  $2^{5+10} \cdot L_5(2)$  when  $|P| = p = 31$  or  $P$  lies in an almost simple group. In the last two possibilities, clearly, these maximal subgroups are not  $p$ -solvable.

(10)  $(G, p)$  is one of the following:  $(M_{23}, 23)$ ,  $(J_1, 7 \text{ or } 19)$ ,  $(Ly, 37 \text{ or } 67)$ ,  $(J_4, 29 \text{ or } 43)$ ,  $(Fi'_{24}, 29)$ ,  $(B, 47)$  or  $(M, 41 \text{ or } 59 \text{ or } 71)$ .

By [2, Table D],  $P$  lies in the unique maximal subgroup of  $G$  and the result follows except for the case  $(G, p) = (J_1, 7)$ . By the Atlas [7], the maximal subgroups of  $J_1$  containing a Sylow 7-subgroup are isomorphic to either  $2^3 : 7 : 3$  or  $7 : 6$ . Thus  $\langle x, x^g \rangle$  is solvable for all  $g \in J_1$ , where  $x \in P$  with  $|x| = 7$ . However, this contradicts Theorem A\* in [25].  $\square$

**Remark 4.5.** It is conjectured in [6] that if  $G$  is a finite nonabelian simple group and if  $r$  and  $s$  are prime divisors of  $|G|$ , then  $G$  can be generated by a Sylow  $r$ -subgroup and a Sylow  $s$ -subgroup. The previous proposition is just a special case of this conjecture when  $r = s = p$  and  $G$  is a minimal non- $p$ -solvable simple group.

*Proof of Theorem 1.10.* Let  $G$  be a finite group and let  $p$  be a prime. Let  $P$  be a Sylow  $p$ -subgroup of  $G$ . If  $G$  is  $p$ -solvable, then every subgroup of  $G$  is  $p$ -solvable. Therefore, it suffices to show that if  $\langle P, P^g \rangle$  is  $p$ -solvable for all  $g \in G$ , then  $G$  is

$p$ -solvable. Suppose not and let  $G$  be a counterexample with minimal order. Then  $\langle P, P^g \rangle$  is  $p$ -solvable for all  $g \in G$  but  $G$  is not solvable.

We first claim that every proper subgroup of  $G$  is  $p$ -solvable and thus  $G$  is a minimal non- $p$ -solvable group. Let  $H$  be a proper subgroup of  $G$  and let  $Q$  be a Sylow  $p$ -subgroup of  $H$ . Then  $Q \leq P^t$  for some  $t \in G$ . Now for every  $h \in H$ , we have

$$\langle Q, Q^h \rangle \leq \langle P^t, (P^t)^h \rangle = \langle P, P^{tht^{-1}} \rangle^t.$$

Since  $\langle P, P^{tht^{-1}} \rangle$  is  $p$ -solvable,  $\langle Q, Q^h \rangle$  is  $p$ -solvable. Therefore, by the minimality of  $|G|$ ,  $H$  is  $p$ -solvable.

By Proposition 4.4, we know that  $G$  is not a nonabelian simple group. Let  $N$  be a proper nontrivial normal subgroup of  $G$ . Now  $PN/N$  is a Sylow  $p$ -subgroup of  $G/N$  and it satisfies the hypothesis of the theorem. Since  $|G/N| < |G|$ ,  $G/N$  is  $p$ -solvable. As in the previous claim,  $N$  is also  $p$ -solvable and thus  $G$  is  $p$ -solvable as well. This final contradiction proves the theorem.  $\square$

### Acknowledgements

The author is grateful to the referee for numerous comments and suggestions that have significantly improved the exposition of the paper. The referee has simplified the proofs of both Theorem 1.4 and Corollary 1.5 as well as shortened the proof of Theorem 1.6 significantly. The author also thanks Chris Schroeder for careful reading of several versions of the paper.

### References

- [1] O. D. Artemovich, "Isolated elements of prime order in finite groups", *Ukrain. Mat. Zh.* **40**:3 (1988), 397–400, 408. MR Zbl
- [2] B. Baumeister, T. C. Burness, R. M. Guralnick, and H. P. Tong-Viet, "On the maximal overgroups of Sylow subgroups of finite groups", *Adv. Math.* **444** (2024), art. id. 109632. MR Zbl
- [3] Y. Berkovich, "Subgroups with the character restriction property and related topics", *Houston J. Math.* **24**:4 (1998), 631–638. MR Zbl
- [4] M. Bianchi, A. G. B. Mauri, M. Herzog, and L. Verardi, "On finite solvable groups in which normality is a transitive relation", *J. Group Theory* **3**:2 (2000), 147–156. MR Zbl
- [5] J. N. Bray, D. F. Holt, and C. M. Roney-Dougal, *The maximal subgroups of the low-dimensional finite classical groups*, London Mathematical Society Lecture Note Series **407**, Cambridge University Press, Cambridge, 2013. MR Zbl
- [6] T. Breuer and R. M. Guralnick, "Finite groups can be generated by a  $\pi'$ -subgroup and a  $\pi'$ -subgroup", *Israel J. Math.* **257**:1 (2023), 55–69. MR Zbl
- [7] J. H. Conway, R. T. Curtis, S. P. Norton, R. A. Parker, and R. A. Wilson, *Atlas of finite groups*, Oxford University Press, Eynsham, 1985. MR Zbl
- [8] B. Fischer, "Frobeniusautomorphismen endlicher Gruppen", *Math. Ann.* **163** (1966), 273–298. MR Zbl

- [9] P. Flavell, “Generating finite groups with conjugates of a subgroup”, *Comm. Algebra* **23**:11 (1995), 3947–3959. MR Zbl
- [10] P. Flavell, “Generating finite groups with maximal subgroups of maximal subgroups”, *J. Algebra* **177**:2 (1995), 372–384. MR Zbl
- [11] P. Flavell, “A characterisation of  $p$ -soluble groups”, *Bull. London Math. Soc.* **29**:2 (1997), 177–183. MR Zbl
- [12] P. Flavell, “The fixed points of coprime action”, *Arch. Math. (Basel)* **75**:3 (2000), 173–177. MR Zbl
- [13] P. Flavell, “Generating finite groups with conjugates of a subgroup, II”, *J. Algebra* **232**:2 (2000), 578–616. MR Zbl
- [14] P. Flavell and G. R. Robinson, “Fixed points of coprime automorphisms and generalizations of Glauberman’s  $Z^*$ -theorem”, *J. Algebra* **226**:2 (2000), 714–718. MR Zbl
- [15] R. J. Flores and R. M. Foote, “Strongly closed subgroups of finite groups”, *Adv. Math.* **222**:2 (2009), 453–484. MR Zbl
- [16] R. Foote, “A characterization of finite groups containing a strongly closed 2-subgroup”, *Comm. Algebra* **25**:2 (1997), 593–606. MR
- [17] The GAP Group, *GAP (version 4.7.5)*, 2014, available at <https://www.gap-system.org/>. Zbl
- [18] G. Glauberman, “Central elements in core-free groups”, *J. Algebra* **4** (1966), 403–420. MR Zbl
- [19] G. Glauberman, R. Guralnick, J. Lynd, and G. Navarro, “Centers of Sylow subgroups and automorphisms”, *Israel J. Math.* **240**:1 (2020), 253–266. MR Zbl
- [20] D. M. Goldschmidt, “2-fusion in finite groups”, *Ann. of Math. (2)* **99** (1974), 70–117. MR Zbl
- [21] N. Gordeev, F. Grunewald, B. Kunyavskiĭ, and E. Plotkin, “Baer–Suzuki theorem for the solvable radical of a finite group”, *C. R. Math. Acad. Sci. Paris* **347**:5-6 (2009), 217–222. MR Zbl
- [22] D. Gorenstein, *Finite groups*, 2nd ed., Chelsea Publishing Co., New York, 1980. MR Zbl
- [23] D. Gorenstein, *Finite simple groups: an introduction to their classification*, Plenum Publishing Corp., New York, 1982. MR Zbl
- [24] D. Gorenstein, R. Lyons, and R. Solomon, *The classification of the finite simple groups, I, Chapter A: Almost simple  $K$ -groups*, Mathematical Surveys and Monographs **40.3**, American Mathematical Society, Providence, RI, 1998. MR Zbl
- [25] S. Guest, “A solvable version of the Baer–Suzuki theorem”, *Trans. Amer. Math. Soc.* **362**:11 (2010), 5909–5946. MR Zbl
- [26] R. M. Guralnick, “Generation of simple groups”, *J. Algebra* **103**:1 (1986), 381–401. MR Zbl
- [27] R. M. Guralnick and G. R. Robinson, “On extensions of the Baer–Suzuki theorem”, *Israel J. Math.* **82**:1-3 (1993), 281–297. MR Zbl
- [28] R. M. Guralnick and P. H. Tiep, “Lifting in Frattini covers and a characterization of finite solvable groups”, *J. Reine Angew. Math.* **708** (2015), 49–72. MR Zbl
- [29] R. M. Guralnick, G. Malle, and G. Navarro, “Self-normalizing Sylow subgroups”, *Proc. Amer. Math. Soc.* **132**:4 (2004), 973–979. MR Zbl
- [30] R. M. Guralnick, H. P. Tong-Viet, and G. Tracey, “Weakly subnormal subgroups and variations of the Baer–Suzuki theorem”, preprint, 2024. arXiv 2402.00804
- [31] T. Hawkes and J. Humphreys, “A character-theoretic criterion for the existence of normal complements to subgroups of finite groups”, *J. Algebra* **94**:2 (1985), 382–387. MR Zbl

- [32] I. M. Isaacs, "Subgroups with the character restriction property", *J. Algebra* **100**:2 (1986), 403–420. MR Zbl
- [33] I. M. Isaacs and T. Zieschang, "Generating symmetric groups", *Amer. Math. Monthly* **102**:8 (1995), 734–739. MR Zbl
- [34] P. Kleidman and M. Liebeck, *The subgroup structure of the finite classical groups*, London Mathematical Society Lecture Note Series **129**, Cambridge University Press, Cambridge, 1990. MR Zbl
- [35] H. Kurzweil and B. Stellmacher, *The theory of finite groups: an introduction*, Springer, New York, 2004. MR Zbl
- [36] S. Li, "On minimal non-PE-groups", *J. Pure Appl. Algebra* **132**:2 (1998), 149–158. MR Zbl
- [37] C. Parker and J. Saunders, "Squares of conjugacy classes and a variant on the Baer–Suzuki Theorem", preprint, 2022. arXiv 2210.06962
- [38] M. Suzuki, *Group theory, II*, Grundle Math. Wissen. **248**, Springer, New York, 1986. MR Zbl
- [39] J. G. Thompson, "Nonsolvable finite groups all of whose local subgroups are solvable", *Bull. Amer. Math. Soc.* **74** (1968), 383–437. MR Zbl
- [40] H. P. Tong-Viet, "Conjugacy classes of  $p$ -elements and normal  $p$ -complements", *Pacific J. Math.* **308**:1 (2020), 207–222. MR Zbl
- [41] J. H. Walter, "Finite groups with abelian Sylow 2-subgroups of order 8", *Invent. Math.* **2** (1967), 332–376. MR Zbl
- [42] H. Wielandt, "Über die Existenz von Normalteilern in endlichen Gruppen", *Math. Nachr.* **18** (1958), 274–280. MR Zbl
- [43] K. Zsigmondy, "Zur Theorie der Potenzreste", *Monatsh. Math. Phys.* **3**:1 (1892), 265–284. MR Zbl

Received April 15, 2024. Revised May 10, 2024.

HUNG P. TONG-VIET  
DEPARTMENT OF MATHEMATICS AND STATISTICS  
BINGHAMTON UNIVERSITY  
BINGHAMTON, NY 13902-6000  
UNITED STATES  
htongvie@binghamton.edu



# PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

[msp.org/pjm](http://msp.org/pjm)

## EDITORS

Don Blasius (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[blasius@math.ucla.edu](mailto:blasius@math.ucla.edu)

Matthias Aschenbrenner  
Fakultät für Mathematik  
Universität Wien  
Vienna, Austria  
[matthias.aschenbrenner@univie.ac.at](mailto:matthias.aschenbrenner@univie.ac.at)

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Atsushi Ichino  
Department of Mathematics  
Kyoto University  
Kyoto 606-8502, Japan  
[atsushi.ichino@gmail.com](mailto:atsushi.ichino@gmail.com)

Robert Lipshitz  
Department of Mathematics  
University of Oregon  
Eugene, OR 97403  
[lipshitz@uoregon.edu](mailto:lipshitz@uoregon.edu)

Kefeng Liu  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Dimitri Shlyakhtenko  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[shlyakht@ipam.ucla.edu](mailto:shlyakht@ipam.ucla.edu)

Paul Yang  
Department of Mathematics  
Princeton University  
Princeton NJ 08544-1000  
[yang@math.princeton.edu](mailto:yang@math.princeton.edu)

Ruixiang Zhang  
Department of Mathematics  
University of California  
Berkeley, CA 94720-3840  
[ruixiang@berkeley.edu](mailto:ruixiang@berkeley.edu)

## PRODUCTION

Silvio Levy, Scientific Editor, [production@msp.org](mailto:production@msp.org)

---

See inside back cover or [msp.org/pjm](http://msp.org/pjm) for submission instructions.

---

The subscription price for 2024 is US \$645/year for the electronic version, and \$875/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).


---

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2024 Mathematical Sciences Publishers

# PACIFIC JOURNAL OF MATHEMATICS

Volume 329    No. 2    April 2024

---

Concentration inequalities for Paley–Wiener spaces SYED HUSAIN and FRIEDRICH LITTMANN	201
Characterizing the Fourier transform by its properties MATEUSZ KRUKOWSKI	217
Reduction types of CM curves MENTZELOS MELISTAS	233
The local character expansion as branching rules: nilpotent cones and the case of $SL(2)$ MONICA NEVINS	259
Extremely closed subgroups and a variant on Glauber’s $Z^*$ -theorem HUNG P. TONG-VIET	303
Vishik equivalence and similarity of quasilinear $p$ -forms and totally singular quadratic forms KRISTÝNA ZEMKOVÁ	327
$RLL$ -realization of two-parameter quantum affine algebra in type $D_n^{(1)}$ RUSHU ZHUANG, NAIHONG HU and XIAO XU	357