

*Pacific
Journal of
Mathematics*

***RLL-REALIZATION OF TWO-PARAMETER
QUANTUM AFFINE ALGEBRA IN TYPE $D_n^{(1)}$***

RUSHU ZHUANG, NAIHONG HU AND XIAO XU

Volume 329 No. 2

April 2024

RLL -REALIZATION OF TWO-PARAMETER QUANTUM AFFINE ALGEBRA IN TYPE $D_n^{(1)}$

RUSHU ZHUANG, NAIHONG HU AND XIAO XU

We obtain the basic R -matrix of the two-parameter quantum group $U = U_{r,s}(\mathfrak{so}_{2n})$ via its weight representation theory and determine its R -matrix with spectral parameters for the two-parameter quantum affine algebra $U = U_{r,s}(\widehat{\mathfrak{so}}_{2n})$. Using the Gauss decomposition of the R -matrix realization of $U = U_{r,s}(\mathfrak{so}_{2n})$, we study the commutation relations of the Gaussian generators and finally arrive at its RLL -formalism of the Drinfeld realization of two-parameter quantum affine algebra $U = U_{r,s}(\widehat{\mathfrak{so}}_{2n})$.

1. Introduction	357
2. Preliminaries	359
3. Basic R -matrix	362
4. Faddeev–Reshetikhin–Takhtajan realization of $U_{r,s}(\mathfrak{so}_{2n})$	367
5. The algebra $\mathcal{U}(\widehat{\mathcal{R}})$ and its Gauss decomposition	375
6. Drinfeld realization of $U_{r,s}(\widehat{\mathfrak{so}}_{2n})$	379
Acknowledgements	394
References	394

1. Introduction

Quantum groups were independently discovered by Drinfeld [6; 8] and Jimbo [15], who showed that the universal enveloping algebra $U(\mathfrak{g})$ of any Kac–Moody algebra \mathfrak{g} admits a certain q -deformation $U_q(\mathfrak{g})$ as a Hopf algebra. Their construction is given in terms of Chevalley generators and q -Serre relations. For the Yangian algebra $Y(\mathfrak{g})$ and the quantum affine algebra $U_q(\widehat{\mathfrak{g}})$ of complex simple Lie algebra \mathfrak{g} , Drinfeld [7] gave their new realizations, which are quantizations of the loop realizations of the classical loop and affine Lie algebras. Faddeev, Reshetikhin and Takhtajan [21]

Hu is the corresponding author. This work is supported by the NNSF of China (Grant 12171155), and in part by the Science and Technology Commission of Shanghai Municipality (Grant 22DZ2229014).
MSC2020: primary 16T25, 17B37, 81R50; secondary 17B67.

Keywords: basic R -matrix, Drinfeld realization, RLL formulation, quantum affine algebra, Gauss decomposition.

presented the RLL -realizations of $U_q(\mathfrak{g})$ [19] of the classical simple Lie algebras \mathfrak{g} by means of solutions of the quantum Yang–Baxter equation

$$R_{12}R_{13}R_{23} = R_{23}R_{13}R_{12},$$

where $R_{12} = R \otimes I$, etc., and $R \in \text{End}(\mathbb{C}^n \otimes \mathbb{C}^n)$. This realization is a natural analog of the matrix realizations of the classical Lie algebras, which originated from the quantum inverse scattering method developed by the St. Petersburg school. Later on, the R -matrix realization of quantum loop algebra $U_q(\mathfrak{g} \otimes \mathbb{C}[t, t^{-1}])$ using a solution of the quantum Yang–Baxter equation with spectral parameters $z, w \in \mathbb{C}$,

$$R_{12}(z)R_{13}(zw)R_{23}(w) = R_{23}(w)R_{13}(zw)R_{12}(z),$$

where $R(z)$ is a rational function of z with values in $\text{End}(\mathbb{C}^n \otimes \mathbb{C}^n)$ was given by Faddeev, Reshetikhin and Takhtajan in [9].

As we know [10], the affine Kac–Moody algebra $\widehat{\mathfrak{g}}$ admits a natural realization as a central extension of the loop algebra $\mathfrak{g} \otimes \mathbb{C}[t, t^{-1}]$, when \mathfrak{g} is a simple Lie algebra. In [20], Reshetikhin and Semenov-Tian-Shansky proved that the central extension can be viewed as the affine analog of the construction in [21]. In [5], by using the Gauss decomposition, Ding and Frenkel showed that the R -matrix realization and Drinfeld realization of quantum affine algebra $U_q(\widehat{\mathfrak{gl}}_n)$ are isomorphic. Recently, Jing, Liu and Molev [17; 18] presented the isomorphisms between the R -matrix and Drinfeld presentations of one-parameter quantum affine algebras $U_q(\widehat{\mathfrak{g}})$ of affine types $B_n^{(1)}, C_n^{(1)}, D_n^{(1)}$.

On the other hand, in 2001, Benkart and Witherspoon [1], motivated by the two-parameters (r, s) -Serre relations satisfied by the up-down operators defined on posets, reobtained the two-parameter quantum enveloping algebras $U_{r,s}(\mathfrak{g})$ corresponding to the general linear Lie algebra \mathfrak{gl}_n and the special linear Lie algebra \mathfrak{sl}_n , which were earlier defined by Takeuchi [22] in 1990. Bergeron, Gao, Hu in [2] found the defining structures of two-parameter quantum groups $U_{r,s}(\mathfrak{g})$ of orthogonal and symplectic Lie algebras, which are realized as the Drinfeld doubles and established their weight representation theory of category $\mathcal{O}^{r,s}$ [3]. Much research has been done for the other types including the affine types (see [4] and references therein). Hu, Rosso and Zhang [12; 13] defined and initiated the study of the vertex representations of Drinfeld realizations of two-parameter quantum affine algebras of untwisted types and constructed the quantum affine Lyndon bases. The RLL -realization of the two-parameter quantum affine algebra $U_{r,s}(\widehat{\mathfrak{gl}}_n)$ has been given by Jing and Liu [16] under the name of RTT -realization.

A natural open question is how to work out the RLL -realizations for the quantum affine algebras $U_{r,s}(\widehat{\mathfrak{g}})$ of types $B_n^{(1)}, C_n^{(1)}, D_n^{(1)}$. The difficulty lies in that there has been no information about two-parameter basic R -matrices in the corresponding cases for many years, let alone their Yang–Baxterizations. Recently we have

overcome such obstacles and solved these problems (also see the subsequent preprints [14; 23]).

The current paper is the first to give the RLL-realization of the two-parameter quantum affine algebra $U_{r,s}(\widehat{\mathfrak{so}}_{2n})$ by the Reshetikhin and Semenov-Tian-Shansky method. We show that Drinfeld’s construction can be naturally established in the Gaussian decomposition of a matrix composed of elements of the quantum affine algebra. The organization of the paper is as follows. In Section 2, we recall the basic results. In Section 3, we give the \widehat{R} -matrix of the two-parameter quantum group $U_{r,s}(\mathfrak{so}_{2n})$. In Section 4, we give the isomorphism between Faddeev–Reshetikhin–Takhtajan and Drinfeld–Jimbo definitions of $U_{r,s}(\mathfrak{so}_{2n})$, and further give the spectral parameter dependent R -matrix $\widehat{R}(z)$ as the Yang–Baxterization of the basic R -matrix we obtained. In Sections 5–6, we study the commutation relations between Gaussian generators and give the Drinfeld realization of $U_{r,s}(\widehat{\mathfrak{so}}_{2n})$ (modified version of [12]).

2. Preliminaries

In [2], let $\mathbb{K} = \mathbb{Q}(r, s)$ be a ground field of rational functions in r, s , where r, s are algebraically independent indeterminates. Assume Φ is a finite root system of type D_n with Π a base of simple roots. Regard Φ as a subset of a Euclidean space $E = \mathbb{R}^n$ with an inner product (\cdot, \cdot) . Let $\epsilon_1, \dots, \epsilon_n$ denote an orthonormal basis of E , and suppose $\Pi = \{\alpha_i = \epsilon_i - \epsilon_{i+1} \mid 1 \leq i < n\} \cup \{\alpha_n = \epsilon_{n-1} + \epsilon_n\}$ and $\Phi = \{\pm\epsilon_i \pm \epsilon_j \mid 1 \leq i \neq j \leq n\}$. In this case, set $r_i = r^{(\alpha_i, \alpha_i)/2}$ and $s_i = s^{(\alpha_i, \alpha_i)/2}$, so that $r_1 = \dots = r_n = r$ and $s_1 = \dots = s_n = s$.

The Cartan matrix of D_n is

$$(2-1) \quad D_n = (a_{ij})_{n \times n} = \begin{pmatrix} 2 & -1 & 0 & \cdots & 0 & 0 & 0 \\ -1 & 2 & -1 & \cdots & 0 & 0 & 0 \\ 0 & -1 & 2 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 2 & -1 & -1 \\ 0 & 0 & 0 & \cdots & -1 & 2 & 0 \\ 0 & 0 & 0 & \cdots & -1 & 0 & 2 \end{pmatrix}.$$

The quantum structural constant matrix is of the form

$$(2-2) \quad (\langle w'_i, w_j \rangle)_{n \times n} = \begin{pmatrix} rs^{-1} & r^{-1} & 1 & \cdots & 1 & 1 & 1 \\ s & rs^{-1} & r^{-1} & \cdots & 1 & 1 & 1 \\ 1 & s & rs^{-1} & \cdots & 1 & 1 & 1 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 1 & 1 & 1 & \cdots & rs^{-1} & r^{-1} & r^{-1} \\ 1 & 1 & 1 & \cdots & s & rs^{-1} & r^{-1}s^{-1} \\ 1 & 1 & 1 & \cdots & s & rs & rs^{-1} \end{pmatrix}.$$

In [2], let $U_{r,s}(\mathfrak{so}_{2n})$ be the unital associative algebra over $\mathbb{Q}(r, s)$ generated by $e_i, f_i, w_i^{\pm 1}, w_i^{\prime \pm 1}$ ($1 \leq i \leq n$), subject to the defining relations (D1)–(D6):

(D1) $w_i^{\pm 1}, w_i^{\prime \pm 1}$ all commute with one another and $w_i w_i^{-1} = w_i' w_i'^{-1} = 1$.

(D2) For $1 \leq i \leq n, 1 \leq j \leq n$, and $1 \leq k (\neq n-1) \leq n$, we have

$$\begin{aligned} w_j e_i &= r^{(\epsilon_j, \alpha_i)} s^{(\epsilon_{j+1}, \alpha_i)} e_i w_j, & w_j f_i &= r^{-(\epsilon_j, \alpha_i)} s^{-(\epsilon_{j+1}, \alpha_i)} f_i w_j, \\ w_j' e_i &= s^{(\epsilon_j, \alpha_i)} r^{(\epsilon_{j+1}, \alpha_i)} e_i w_j', & w_j' f_i &= s^{-(\epsilon_j, \alpha_i)} r^{-(\epsilon_{j+1}, \alpha_i)} f_i w_j', \\ w_n e_k &= r^{(\epsilon_{n-1}, \alpha_k)} s^{-(\epsilon_n, \alpha_k)} e_k w_n, & w_n f_k &= r^{-(\epsilon_{n-1}, \alpha_k)} s^{(\epsilon_n, \alpha_k)} f_k w_n, \\ w_n' e_k &= s^{(\epsilon_{n-1}, \alpha_k)} r^{-(\epsilon_n, \alpha_k)} e_k w_n', & w_n' f_k &= s^{-(\epsilon_{n-1}, \alpha_k)} r^{(\epsilon_n, \alpha_k)} f_k w_n'. \end{aligned}$$

(D3)

$$\begin{aligned} w_n e_{n-1} &= r^{(\epsilon_n, \alpha_{n-1})} s^{-(\epsilon_{n-1}, \alpha_{n-1})} e_{n-1} w_n, \\ w_n f_{n-1} &= r^{-(\epsilon_n, \alpha_{n-1})} s^{(\epsilon_{n-1}, \alpha_{n-1})} f_{n-1} w_n, \\ w_n' e_{n-1} &= s^{(\epsilon_n, \alpha_{n-1})} r^{-(\epsilon_{n-1}, \alpha_{n-1})} e_{n-1} w_n', \\ w_n' f_{n-1} &= s^{-(\epsilon_n, \alpha_{n-1})} r^{(\epsilon_{n-1}, \alpha_{n-1})} f_{n-1} w_n'. \end{aligned}$$

(D4) For $1 \leq i, j \leq n$, we have

$$[e_i, f_j] = \delta_{ij} \frac{w_i - w_i'^{-1}}{r - s}.$$

(D5) For any $1 \leq i \neq j \leq n$ but $(i, j) \notin \{(n-1, n), (n, n-1)\}$ with $a_{ij} = 0$, we have

$$\begin{aligned} [e_i, e_j] &= 0, & e_{n-1} e_n &= r s e_n e_{n-1}, \\ [f_i, f_j] &= 0, & f_n f_{n-1} &= r s f_{n-1} f_n. \end{aligned}$$

(D6) For $1 \leq i < j \leq n$ with $a_{ij} = -1$, we have (r, s) -Serre relations:

$$\begin{aligned} e_i^2 e_j - (r + s) e_i e_j e_i + r s e_j e_i^2 &= 0, \\ f_j f_i^2 - (r + s) f_i f_j f_i + r s f_i^2 f_j &= 0, \\ e_j^2 e_i - (r^{-1} + s^{-1}) e_j e_i e_j + r^{-1} s^{-1} e_i e_j^2 &= 0, \\ f_i f_j^2 - (r^{-1} + s^{-1}) f_j f_i f_j + r^{-1} s^{-1} f_j^2 f_i &= 0. \end{aligned}$$

Proposition 2.1. *The algebra $U_{r,s}(\mathfrak{so}_{2n})$ is a Hopf algebra with comultiplication Δ , counit ε and antipode S such that*

$$\begin{aligned} \Delta(e_i) &= e_i \otimes 1 + w_i \otimes e_i, & \Delta(f_i) &= 1 \otimes f_i + f_i \otimes w_i', \\ \varepsilon(e_i) &= 0, & \varepsilon(f_i) &= 0, \\ S(e_i) &= -w_i^{-1} e_i, & S(f_i) &= -f_i w_i'^{-1}, \end{aligned}$$

and w_i, w_i' are group-like elements for any $i \in I$.

Definition 2.2. We define the linear mapping $f : \Lambda \times \Lambda \rightarrow k^*$ as

$$f(\lambda, \mu) = \langle w'_\mu, w_\lambda \rangle^{-1},$$

which satisfies

$$f(\lambda + \nu, \mu) = f(\lambda, \mu)f(\nu, \mu), \quad f(\lambda, \mu + \nu) = f(\lambda, \mu)f(\lambda, \nu).$$

Definition 2.3. Let M, M' be finite-dimensional U -modules. Define an isomorphism of U -modules $\tilde{f} : M \otimes M' \rightarrow M \otimes M'$ as

$$\tilde{f}(m \otimes m') = f(\lambda, \mu)m \otimes m',$$

where $m \in M_\lambda, m' \in M'_\mu$ and $\lambda, \mu \in \Lambda$.

Corollary 2.4 [3]. $U_{r,s}(\mathfrak{g}) \cong U_{r,s}(\mathfrak{n}^-) \otimes U^0 \otimes U_{r,s}(\mathfrak{n})$, as vector spaces. In particular, it induces $U_q(\mathfrak{g}) \cong U_q(\mathfrak{n}^-) \otimes U_0 \otimes U_q(\mathfrak{n})$, as vector spaces.

Let $Q = \mathbb{Z}\Phi$ denote the root lattice and set $Q^+ = \sum_{i=1}^n \mathbb{Z}_{\geq 0}\alpha_i$. Then for any $\zeta = \sum_{i=1}^n \zeta_i \alpha_i \in Q$, we denote

$$\omega_\zeta = \omega_1^{\zeta_1} \cdots \omega_n^{\zeta_n}, \quad \omega'_\zeta = (\omega'_1)^{\zeta_1} \cdots (\omega'_n)^{\zeta_n}.$$

Then $U_{r,s}(\mathfrak{n}^\pm) = \bigoplus_{\eta \in Q^+} U_{r,s}^{\pm\eta}(\mathfrak{n}^\pm)$ Q^\pm -graded, where

$$U_{r,s}^\eta(\mathfrak{n}^\pm) = \{a \in U_{r,s}(\mathfrak{n}^\pm) \mid \omega_\zeta a \omega_\zeta^{-1} = \langle \omega'_\eta, \omega_\zeta \rangle a, \omega'_\zeta a \omega'_\zeta^{-1} = \langle \omega'_\zeta, \omega_\eta \rangle^{-1} a\},$$

for $\eta \in Q^+ \cup Q^-$.

Lemma 2.5 [3]. Set $d_\zeta = \dim_{\mathbb{K}} U_{r,s}^{+\zeta}(\mathfrak{n}^+)$. Consider the basis $\{u_k^\zeta\}_{k=1}^{d_\zeta}$ for $U_{r,s}^{+\zeta}(\mathfrak{n}^+)$, and $\{v_k^\zeta\}_{k=1}^{d_\zeta}$ is the dual basis for $U_{r,s}^{-\zeta}(\mathfrak{n}^-)$ with respect to the pairing. Now let

$$(2-3) \quad \Theta_\zeta = \sum_{k=1}^{d_\zeta} v_k^\zeta \otimes u_k^\zeta \in U \otimes U.$$

All but finitely many terms in this sum will act as multiplication by 0 on any weight space M_λ of $M \in \mathcal{O}$. $\Theta = \sum_{\zeta \in Q^+} \Theta_\zeta$ is a well-defined operator on such $M \otimes M$.

Theorem 2.6 [3]. For M, M' to be finite-dimensional modules of $U_{r,s}(\mathfrak{g})$, the map

$$R_{M',M} = \Theta \circ \tilde{f} \circ P : M' \otimes M \rightarrow M \otimes M'$$

must be an isomorphism of $U_{r,s}(\mathfrak{g})$ -module, where $P(m \otimes m') = m' \otimes m, m \in M, m' \in M'$.

Theorem 2.7 [3]. For M, M', M'' to be finite-dimensional modules of $U_{r,s}(\mathfrak{g})$, the map must satisfy

$$\Theta_{12}^f \circ \Theta_{13}^f \circ \Theta_{23}^f = \Theta_{23}^f \circ \Theta_{13}^f \circ \Theta_{12}^f.$$

Equivalently, we have

$$R_{12} \circ R_{23} \circ R_{12} = R_{23} \circ R_{12} \circ R_{23}.$$

Remark 2.8. Denote $\Theta_{12}^f = (\Theta_{M,M'} \circ \tilde{f}) \otimes 1$, $\Theta_{23}^f = 1 \otimes (\Theta_{M',M''} \circ \tilde{f})$, $\Theta_{13}^f = (1 \otimes P) \circ (\Theta^f \otimes 1) \circ (1 \otimes P)$. Also, $\Theta \circ \tilde{f}$ is the solution of the quantum Yang–Baxter equation.

3. Basic R -matrix

In this section, we consider the R -matrix $R := R_{V,V}$ for the vector representation $T_1 = T_V$ of the Drinfeld–Jimbo algebra $U = U_{r,s}(\mathfrak{so}_{2n})$.

Definition 3.1. The vector representation T_1 of the Drinfeld–Jimbo algebra U is the irreducible type 1 representation with highest weight $\lambda = (1, 0, \dots, 0)$ with respect to the ordered sequence $\alpha_1, \dots, \alpha_n$ of simple roots.

Consider a $2n$ -dimensional vector space V over \mathbb{K} with basis $\{v_j \mid 1 \leq j \leq 2n\}$. We define an action of the generators of $U = U_{r,s}(\mathfrak{so}_{2n})$ by specifying their matrices relative to this basis:

Lemma 3.2. Let E_{kl} be the $2n \times 2n$ matrix with 1 in the (k, l) -position and 0 elsewhere. The vector representation T_1 of $U_{r,s}(\mathfrak{so}_{2n})$ is described by the following list:

$$(B1) \quad \begin{aligned} T_1(e_i) &= E_{i,i+1} - r^{-\frac{1}{2}}s^{-\frac{1}{2}}E_{(i+1)',i'}, & T_1(e_n) &= E_{n,n+2} - r^{-\frac{1}{2}}s^{-\frac{1}{2}}E_{(n+2)',n'}, \\ T_1(f_i) &= E_{i+1,i} - r^{-\frac{1}{2}}s^{-\frac{1}{2}}E_{i',(i+1)'}, & T_1(f_n) &= E_{n+2,n} - r^{-\frac{1}{2}}s^{-\frac{1}{2}}E_{n',(n+2)'}, \end{aligned}$$

$$(B2) \quad \begin{aligned} T_1(w_i) &= rE_{i,i} + sE_{i+1,i+1} + s^{-1}E_{(i+1)',(i+1)'} + r^{-1}E_{i',i'} + \sum_{j \neq \{i,i+1,i',(i+1)'\}} E_{j,j}, \\ T_1(w'_i) &= sE_{i,i} + rE_{i+1,i+1} + r^{-1}E_{(i+1)',(i+1)'} + s^{-1}E_{i',i'} + \sum_{j \neq \{i,i+1,i',(i+1)'\}} E_{j,j}, \end{aligned}$$

$$(B3) \quad \begin{aligned} T_1(w_n) &= s^{-1}E_{n-1,n-1} + rE_{n,n} + r^{-1}E_{n',n'} + sE_{(n-1)',(n-1)'} + r^{-1}s^{-1}E_{1,1} \\ &\quad + r^{-1}s^{-1} \sum_{1 \leq j \leq n-2} E_{j,j} + rs \sum_{1 \leq j \leq n-2} E_{j',j'}, \\ T_1(w'_n) &= r^{-1}E_{n-1,n-1} + sE_{n,n} + s^{-1}E_{n',n'} + rE_{(n-1)',(n-1)'} + r^{-1}s^{-1}E_{1,1} \\ &\quad + r^{-1}s^{-1} \sum_{1 \leq j \leq n-2} E_{j,j} + rs \sum_{1 \leq j \leq n-2} E_{j',j'}, \end{aligned}$$

where $1 \leq i, j \leq n - 1$, and $i' = 2n + 1 - i$.

Proof. By straightforward calculations one checks that the preceding formulas define a weight representation T_1 of the algebra $U_{r,s}(\mathfrak{so}_{2n})$ on the vector space \mathbb{C}^{2n} . For the basis vector $\mathbf{e}_1 = (1, 0, \dots, 0)$, we easily verify that $T_1(E_j)\mathbf{e}_1 = 0$, $T_1(w_1)\mathbf{e}_1 = r\mathbf{e}_1$, $T_1(w_i)\mathbf{e}_1 = \mathbf{e}_1$, $T_1(w_n)\mathbf{e}_1 = r^{-1}s^{-1}\mathbf{e}_1$, $T_1(w'_1)\mathbf{e}_1 = s\mathbf{e}_1$, $T_1(w'_i)\mathbf{e}_1 = \mathbf{e}_1$, and $T_1(w'_n)\mathbf{e}_1 = rs\mathbf{e}_1$, for $1 \leq j \leq n$ and $2 \leq i \leq n-1$. Hence T_1 is the type 1 representation with highest weight $\lambda = \alpha_1 + \alpha_2 + \dots + \frac{1}{2}(\alpha_{n-1} + \alpha_n) = \epsilon_1$. Thus T_1 is indeed the vector representation of $U_{r,s}(\mathfrak{so}_{2n})$. \square

We illustrate through the following lemmas that the module $V \otimes V$ is decomposed into the direct sum of three simple submodules, $S^o(V \otimes V)$, $S'(V \otimes V)$ and $\Lambda(V \otimes V)$. These modules are defined and proved by the following three lemmas.

Lemma 3.3. *The module $S^o(V \otimes V)$ generated by $\sum_{i=1}^{2n} a_i v_{i'} \otimes v_i$ is simple, and a_i satisfies*

$$a_i = \begin{cases} (rs^{-1})^{\frac{n-i}{2}} & \text{if } 1 \leq i \leq n, \\ 1 & \text{if } i = n+1, \\ (rs^{-1})^{\frac{n-i+1}{2}} & \text{if } n+2 \leq i \leq 2n. \end{cases}$$

Proof. The operators e_k, f_k, w_k, w'_k act on $\sum_{i=1}^{2n} a_i v_{i'} \otimes v_i$, then the following computations show that $S^o(V \otimes V)$ is a simple module:

$$\begin{aligned} & e_k \cdot \left(\sum_{i=1}^{2n} a_i v_{i'} \otimes v_i \right) \\ &= (e_k \otimes 1 + w_k \otimes e_k) \cdot \left(\sum_{i=1}^{2n} a_i v_{i'} \otimes v_i \right) \\ &= (a_{(k+1)'} - a_k r^{\frac{1}{2}} s^{-\frac{1}{2}}) v_k \otimes v_{(k+1)'} + (a_{k+1} s^{-1} - r^{-\frac{1}{2}} s^{-\frac{1}{2}} a_k) v_{(k+1)'} \otimes v_k \\ &= 0, \end{aligned}$$

$$\begin{aligned} & e_n \cdot \left(\sum_{i=1}^{2n} a_i v_{i'} \otimes v_i \right) \\ &= (e_n \otimes 1 + w_n \otimes e_n) \cdot \left(\sum_{i=1}^{2n} a_i v_{i'} \otimes v_i \right) \\ &= (a_n r^{\frac{1}{2}} s^{-\frac{1}{2}} - a_{(n+2)'}) v_n \otimes v_{(n+2)'} + (r^{-\frac{1}{2}} s^{-\frac{1}{2}} a_n - a_{(n+2)'} s^{-1}) v_{(n+2)'} \otimes v_n \\ &= 0. \end{aligned}$$

Moreover,

$$w_k \cdot \left(\sum_{i=1}^{2n} a_i v_{i'} \otimes v_i \right) = \sum_{i=1}^{2n} a_i v_{i'} \otimes v_i, \quad w'_k \cdot \left(\sum_{i=1}^{2n} a_i v_{i'} \otimes v_i \right) = \sum_{i=1}^{2n} a_i v_{i'} \otimes v_i,$$

and for $1 \leq k \leq n$. By similar calculations as above, we know that the operators f_k act trivially on $\sum_{i=1}^{2n} a_i v_{i'} \otimes v_i$. So, it is easy to see that $S^o(V \otimes V)$ is simple. \square

Lemma 3.4. *The simple module $S'(V \otimes V)$ is defined as follows:*

- (i) $v_i \otimes v_i, 1 \leq i \leq 2n,$
- (ii) $v_i \otimes v_j + sv_j \otimes v_i, 1 \leq i \leq n$ and $i + 1 \leq j \leq n$ or $i' + 1 \leq j \leq 2n,$
- (iii) $v_i \otimes v_j + r^{-1}v_j \otimes v_i, 1 \leq i \leq n - 1, n + 1 \leq j \leq 2n - i$ or $n + 1 \leq i \leq 2n - 1,$
 $i + 1 \leq j \leq 2n,$
- (iv) $v_i \otimes v_{i'} + r^{-1}sv_{i'} \otimes v_i - (r^{-1}s)^{\frac{1}{2}}(v_{(i+1)' } \otimes v_{i+1} + v_{i+1} \otimes v_{(i+1)' }), 1 \leq i \leq n - 1,$

where $v_1 \otimes v_1$ is the highest weight vector.

Proof. Operators e_k, f_k act on vectors in (i)–(iv). The following computations show that $S'(V \otimes V)$ is a simple module, because for case (i), we have

$$e_k \cdot (v_i \otimes v_i) = \begin{cases} \delta_{i,k+1}(v_k \otimes v_{k+1} + sv_{k+1} \otimes v_k), \\ -\delta_{i,k'}(rs)^{-\frac{1}{2}}(v_{i-1} \otimes v_i + r^{-1}v_i \otimes v_{i-1}), \end{cases}$$

or

$$f_k \cdot (v_i \otimes v_i) = \begin{cases} \delta_{i,k}(v_k \otimes v_{k+1} + sv_{k+1} \otimes v_k), \\ -\delta_{i,(k+1)' } (rs)^{-\frac{1}{2}}(v_i \otimes v_{i+1} + r^{-1}v_{i+1} \otimes v_i), \end{cases}$$

where $1 \leq k \leq n - 1,$ and for $k = n,$ we have

$$e_n \cdot (v_i \otimes v_i) = \begin{cases} \delta_{i,n+2}(v_n \otimes v_{n+2} + sv_{n+2} \otimes v_n), \\ -\delta_{i,n+1}(rs)^{-\frac{1}{2}}(v_{n-1} \otimes v_{n+1} + r^{-1}v_{n+1} \otimes v_{n-1}), \end{cases}$$

$$f_n \cdot (v_i \otimes v_i) = \begin{cases} \delta_{i,n}(v_n \otimes v_{n+2} + sv_{n+2} \otimes v_n), \\ -\delta_{i,n-1}(rs)^{-\frac{1}{2}}(v_{n-1} \otimes v_{n+1} + r^{-1}v_{n+1} \otimes v_{n-1}). \end{cases}$$

For case (ii), we have

$$e_k \cdot (v_i \otimes v_j + sv_j \otimes v_i) = e_k(v_i) \otimes v_j + sw_k(v_j) \otimes e_k(v_i) + se_k(v_j) \otimes v_i + w_k(v_i) \otimes e_k(v_j),$$

so we get

$$e_k \cdot (v_i \otimes v_j + sv_j \otimes v_i) = \begin{cases} \delta_{i+j,2n+2}(v_i \otimes v_{i'} + r^{-1}sv_{i'} \otimes v_i \\ - (r^{-1}s)^{\frac{1}{2}}v_{(i+1)' } \otimes v_{i+1} + v_{i+1} \otimes v_{(i+1)' }), \\ \delta_{i,k+1}(v_{i-1} \otimes v_j + sv_j \otimes v_{i-1}), \\ \delta_{j,k'}(v_i \otimes v_{j-1} + sv_{j-1} \otimes v_i), \\ \delta_{i,k}\delta_{k+1,j}(v_i \otimes v_i). \end{cases}$$

For operators $f_k,$ we have

$$f_k \cdot (v_i \otimes v_j + sv_j \otimes v_i) = v_i \otimes f_k(v_j) + sv_j \otimes f_k(v_i) + f_k(v_i) \otimes w'_k(v_j) + sf_k(v_j) \otimes w'_k(v_i),$$

so we have

$$f_k \cdot (v_i \otimes v_j + sv_j \otimes v_i) = \begin{cases} \delta_{i+j,2n}(v_i \otimes v_{i'} + r^{-1}sv_{i'} \otimes v_i \\ -(r^{-1}s)^{\frac{1}{2}}v_{(i+1)'} \otimes v_{i+1} + v_{i+1} \otimes v_{(i+1)'}), \\ \delta_{j,(k+1)'}(v_i \otimes v_{j+1} + sv_{j+1} \otimes v_i), \\ \delta_{i,k}(v_{i+1} \otimes v_j + sv_j \otimes v_{i+1}), \\ \delta_{i,(k+1)'}\delta_{j,k'}(v_{i+1} \otimes v_{i+1}), \end{cases}$$

where $1 \leq k \leq n - 1$. For $k = n$, and $n' + 2 \leq l \leq 2n$, we have

$$f_n \cdot (v_i \otimes v_j + sv_j \otimes v_i) = \begin{cases} v_{n-1} \otimes v_{(n-1)'} + r^{-1}sv_{(n-1)'} \otimes v_{n-1} \\ -(r^{-1}s)^{\frac{1}{2}}(v_n \otimes v_{n'} + v_{n'} \otimes v_n), \\ \delta_{i,n}(v_{n+2} \otimes v_l + r^{-1}v_l \otimes v_{n+2}), \\ \delta_{i,n-1}(v_{n+1} \otimes v_l + r^{-1}v_l \otimes v_{n+1}), \\ v_l \otimes v_l. \end{cases}$$

For case (iii), we have

$$e_k \cdot (v_i \otimes v_j + r^{-1}v_j \otimes v_i) = e_k(v_i) \otimes v_j + r^{-1}w_k(v_j) \otimes e_k(v_i) + r^{-1}e_k(v_j) \otimes v_i + w_k(v_i) \otimes e_k(v_j),$$

so we get

$$e_k \cdot (v_i \otimes v_j + r^{-1}v_j \otimes v_i) = \begin{cases} \delta_{i+j,2n+2}(v_i \otimes v_{i'} + r^{-1}sv_{i'} \otimes v_i \\ -(r^{-1}s)^{\frac{1}{2}}(v_{(i+1)'} \otimes v_{i+1} + v_{i+1} \otimes v_{(i+1)'})), \\ \delta_{i,k+1}(v_{i-1} \otimes v_j + r^{-1}v_j \otimes v_{i-1}), \\ \delta_{j,k'}(v_i \otimes v_{j-1} + r^{-1}v_{j-1} \otimes v_i), \\ \delta_{i,k}\delta_{k+1,j}(v_i \otimes v_i). \end{cases}$$

For operators f_k , we have

$$f_k \cdot (v_i \otimes v_j + r^{-1}v_j \otimes v_i) = v_i \otimes f_k(v_j) + r^{-1}v_j \otimes f_k(v_i) + f_k(v_i) \otimes w'_k(v_j) + r^{-1}f_k(v_j) \otimes w'_k(v_i),$$

so we have

$$f_k \cdot (v_i \otimes v_j + r^{-1}v_j \otimes v_i) = \begin{cases} \delta_{i+j,2n}(v_i \otimes v_{i'} + r^{-1}sv_{i'} \otimes v_i \\ -(r^{-1}s)^{\frac{1}{2}}(v_{(i+1)'} \otimes v_{i+1} + v_{i+1} \otimes v_{(i+1)'})), \\ \delta_{j,(k+1)'}(v_i \otimes v_{j+1} + r^{-1}v_{j+1} \otimes v_i), \\ \delta_{i,k}(v_{i+1} \otimes v_j + r^{-1}v_j \otimes v_{i+1}), \\ \delta_{i,(k+1)'}\delta_{j,k'}(v_{i+1} \otimes v_{i+1}), \end{cases}$$

where $1 \leq k \leq n - 1$. For $k = n$, and $n' + 2 \leq l \leq 2n$, we have

$$f_n \cdot (v_i \otimes v_j + r^{-1}v_j \otimes v_i) = \begin{cases} v_{n-1} \otimes v_{(n-1)'} + r^{-1}sv_{(n-1)'} \otimes v_{n-1} \\ \quad - (r^{-1}s)^{\frac{1}{2}}(v_n \otimes v_{n'} + v_{n'} \otimes v_n), \\ \delta_{i,n}(v_{n+2} \otimes v_l + sv_l \otimes v_{n+2}), \\ \delta_{i,n-1}(v_{n+1} \otimes v_l + sv_l \otimes v_{n+1}), \\ \quad v_l \otimes v_l. \end{cases}$$

Therefore, it is easy to see that $S'(V \otimes V)$ is simple. □

Lemma 3.5. *The simple module $\Lambda(V \otimes V)$ is defined as follows:*

- (i) $v_i \otimes v_j - rv_j \otimes v_i, 1 \leq i \leq n$ and $i + 1 \leq j \leq n$ or $i' + 1 \leq j \leq 2n,$
- (ii) $v_i \otimes v_j - s^{-1}v_j \otimes v_i, 1 \leq i \leq n - 1, n + 1 \leq j \leq 2n - i,$ or $n + 1 \leq i \leq 2n - 1,$
 $i + 1 \leq j \leq 2n,$
- (iii) $-(rs)^{-\frac{1}{2}}v_i \otimes v_{i'} - s^{-1}v_{(i+1)'} \otimes v_{i+1} + r^{-1}v_{i+1} \otimes v_{(i+1)'} + (rs)^{-\frac{1}{2}}v_{i'} \otimes v_i,$
 $1 \leq i \leq n - 1,$

where $v_1 \otimes v_2 - rv_2 \otimes v_1$ is the highest weight vector.

Proof. We can check this lemma by repeating similar calculations to those in Lemma 3.4. □

Lemma 3.6. *The decomposition of $U_{r,s}(\mathfrak{so}_{2n})$ -module $V \otimes V$ is*

$$V \otimes V = S^o(V \otimes V) \oplus S'(V \otimes V) \oplus \Lambda(V \otimes V).$$

Proof. In [11], Hu and Pei proved that as braided tensor categories, the categories $\mathcal{O}^{r,s}$ of finite-dimensional weight $U_{r,s}(\mathfrak{g})$ -modules (of type 1) and \mathcal{O}^q are monoidally equivalent. Referring to the book by Klimyk and Schmüdgen [19], $U_q(\mathfrak{so}_{2n})$ -module $V \otimes V$ is completely reducible and can be decomposed into the direct sum of three simple modules. □

Proposition 3.7. *The minimum polynomial of $R = R_{V,V}$ on $V \otimes V$ is*

$$(t - r^{-\frac{1}{2}}s^{\frac{1}{2}}) \cdot (t + r^{\frac{1}{2}}s^{-\frac{1}{2}}) \cdot (t - r^{\frac{2n-1}{2}}s^{-\frac{2n-1}{2}}).$$

Proof. It follows from the definition of R that $R(v_1 \otimes v_1) = r^{-\frac{1}{2}}s^{\frac{1}{2}}v_1 \otimes v_1$ and $R(v_1 \otimes v_2 - rv_2 \otimes v_1) = -r^{\frac{1}{2}}s^{-\frac{1}{2}}(v_1 \otimes v_2 - rv_2 \otimes v_1)$. By the proceeding lemmas, $S'(V \otimes V)$ and $\Lambda(V \otimes V)$ are simple, and in fact, $v_1 \otimes v_1$ and $v_1 \otimes v_2 - rv_2 \otimes v_1$ are the highest weight vectors. In particular, each is a cyclic module generated by its highest weight vector, respectively, $R(a_1v_{1'} \otimes v_1) = a_1r^{\frac{1}{2}}s^{-\frac{1}{2}}v_1 \otimes v_{1'} + \dots,$ and $v_1 \otimes v_{1'}$ only occurs in $R(a_1v_{1'} \otimes v_1)$. So we have the desired result

$$R\left(\sum_{i=1}^{2n} a_i v_{i'} \otimes v_i\right) = r^{\frac{2n-1}{2}}s^{-\frac{2n-1}{2}}\left(\sum_{i=1}^{2n} a_i v_{i'} \otimes v_i\right). \quad \square$$

Theorem 3.8. *The braiding R -matrix $R = R_{V,V}$ acts as*

$$\begin{aligned}
 R = & r^{-\frac{1}{2}}s^{\frac{1}{2}} \sum_{i=1}^{2n} E_{ii} \otimes E_{ii} + r^{\frac{1}{2}}s^{-\frac{1}{2}} \sum_{i=1}^{2n} E_{i'i'} \otimes E_{i'i} \\
 & + r^{-\frac{1}{2}}s^{-\frac{1}{2}} \left\{ \sum_{\substack{1 \leq i \leq n-1 \\ i+1 \leq j \leq n}} E_{ij} \otimes E_{ji} + \sum_{\substack{1 \leq i \leq n-1 \\ i'+1 \leq j \leq 2n}} E_{ij} \otimes E_{ji} + \sum_{j=n+2}^{2n} E_{nj} \otimes E_{jn} \right. \\
 & \qquad \qquad \qquad \left. + \sum_{\substack{n+1 \leq i \leq 2n-1 \\ i+1 \leq j \leq 2n}} E_{ji} \otimes E_{ij} + \sum_{\substack{1 \leq i \leq n-1 \\ n+1 \leq j \leq 2n-i}} E_{ji} \otimes E_{ij} \right\} \\
 & + r^{\frac{1}{2}}s^{\frac{1}{2}} \left\{ \sum_{\substack{1 \leq i \leq n-1 \\ i+1 \leq j \leq n}} E_{ji} \otimes E_{ij} + \sum_{\substack{1 \leq i \leq n-1 \\ i'+1 \leq j \leq 2n}} E_{ji} \otimes E_{ij} + \sum_{j=n+2}^{2n} E_{jn} \otimes E_{nj} \right. \\
 & \qquad \qquad \qquad \left. + \sum_{\substack{n+1 \leq i \leq 2n-1 \\ i+1 \leq j \leq 2n}} E_{ij} \otimes E_{ji} + \sum_{\substack{1 \leq i \leq n-1 \\ n+1 \leq j \leq 2n-i}} E_{ij} \otimes E_{ji} \right\} \\
 & + (r^{-\frac{1}{2}}s^{\frac{1}{2}} - r^{\frac{1}{2}}s^{-\frac{1}{2}}) \left\{ \sum_{i < j} E_{jj} \otimes E_{ii} - \sum_{i > j} (r^{-\frac{1}{2}}s^{\frac{1}{2}})^{(\rho_i - \rho_j)} E_{i'j'} \otimes E_{i'j} \right\},
 \end{aligned}$$

where

$$\rho_i = \begin{cases} n - i & \text{if } 1 \leq i \leq n \\ 0 & \text{if } n + 1 \\ n - i + 1 & \text{if } n + 2 \leq i \leq 2n \end{cases}.$$

Proof. We need to check that the braiding R -matrix R acts on $S^o(V \otimes V)$, $S'(V \otimes V)$ as multiplication by $r^{\frac{2n-1}{2}}s^{-\frac{2n-1}{2}}$, $r^{-\frac{1}{2}}s^{\frac{1}{2}}$, and on $\Lambda(V \otimes V)$ as multiplication by $-r^{\frac{1}{2}}s^{-\frac{1}{2}}$. By straightforward calculations, one checks that the expression formula of the basic R -matrix is correct. \square

Remark 3.9. Consider the matrix $\widehat{R} = P \circ R$, where $P = \sum_{i,j} E_{ij} \otimes E_{ji}$, and R satisfying the braiding relations on the tensor power $V^{\otimes k}$:

$$\begin{aligned}
 R_i \circ R_{i+1} \circ R_i &= R_{i+1} \circ R_i \circ R_{i+1}, \\
 R_i \circ R_j &= R_j \circ R_i,
 \end{aligned}$$

where $1 \leq i < k$, $|i - j| \geq 2$, $R_i = id_V^{i-1} \otimes R \otimes id_V^{k-1}$.

4. Faddeev–Reshetikhin–Takhtajan realization of $U_{r,s}(\mathfrak{so}_{2n})$

In this section, we give an isomorphism between Faddeev–Reshetikhin–Takhtajan and Drinfeld–Jimbo definitions of $U_{r,s}(\mathfrak{so}_{2n})$, and the spectral parameter dependent $R(z)$. Let \mathcal{B} (resp. \mathcal{B}') denote the subalgebra of $U_{r,s}(\mathfrak{so}_{2n})$ generated by $e_i, w_i^{\pm 1}$ (resp. $f_i, w_i^{\pm 1}$), $1 \leq i \leq n$.

Definition 4.1. $U(\widehat{R})$ is an associative algebra with unit. It has generators l_{ij}^+, l_{ji}^- , $1 \leq i \leq 2n$. Let $L^\pm = (l_{ij}^\pm)$, $1 \leq i, j \leq 2n + 1$, with $l_{ij}^+ = l_{ji}^- = 0$, and $l_{ii}^- l_{ii}^+ = l_{ii}^+ l_{ii}^-$ for $1 \leq j < i \leq 2n + 1$. The defining relations are given in matrix form as follows:

$$(4-1) \quad \widehat{R}L_1^\pm L_2^\pm = L_2^\pm L_1^\pm \widehat{R}, \quad \widehat{R}L_1^+ L_2^- = L_2^- L_1^+ \widehat{R},$$

where $L_1^\pm = L^\pm \otimes 1$, $L_2^\pm = 1 \otimes L^\pm$.

Since L^\pm are upper and lower triangular, respectively, and the diagonal elements of these matrix are invertible, L^\pm have inverse $(L^\pm)^{-1}$ as matrices with elements in $U(\widehat{R})$. The relations between L_1^\pm and L_2^\pm immediately imply the following theorem.

Theorem 4.2. *The mapping ϕ_n between $U(\widehat{R})$ and $U_{r,s}(\mathfrak{so}_{2n})$ is an algebraic homomorphism.*

Proof. We check the theorem for the case of $n = 4$. Let us consider L^\pm ,

$$L^+ = \begin{pmatrix} l_{11}^+ & l_{12}^+ & \cdots & l_{18}^+ \\ 0 & l_{22}^+ & \ddots & \vdots \\ \vdots & \ddots & \ddots & l_{78}^+ \\ 0 & \cdots & 0 & l_{88}^+ \end{pmatrix}_{8 \times 8}, \quad L^- = \begin{pmatrix} l_{11}^- & 0 & \cdots & 0 \\ l_{21}^- & l_{22}^- & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ l_{81}^- & \cdots & l_{87}^- & l_{88}^- \end{pmatrix}_{8 \times 8}.$$

Then for the generators $L_1^\pm, L_2^\pm, \widehat{R}$, we have that

$$L_1^+ = \begin{pmatrix} l_{11}^+ I_8 & l_{12}^+ I_8 & \cdots & l_{18}^+ I_8 \\ 0 & l_{22}^+ I_8 & \ddots & \vdots \\ \vdots & \ddots & \ddots & l_{78}^+ I_8 \\ 0 & \cdots & 0 & l_{88}^+ I_8 \end{pmatrix}_{64 \times 64}, \quad L_1^- = \begin{pmatrix} l_{11}^- I_8 & 0 & \cdots & 0 \\ l_{21}^- I_8 & l_{22}^- I_8 & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ l_{81}^- I_8 & \cdots & l_{87}^- I_8 & l_{88}^- I_8 \end{pmatrix}_{64 \times 64},$$

$$L_2^\pm = \begin{pmatrix} L^\pm & 0 & \cdots & 0 \\ 0 & L^\pm & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & L^\pm \end{pmatrix}_{64 \times 64}, \quad \widehat{R} = \begin{pmatrix} A_{11} & A_{12} & \cdots & A_{18} \\ 0 & A_{22} & \ddots & \vdots \\ \vdots & \ddots & \ddots & A_{78} \\ 0 & \cdots & 0 & A_{88} \end{pmatrix}_{64 \times 64},$$

$$A_{11} = \begin{pmatrix} A'_{11} & 0 \\ 0 & \mathcal{Q}A'^{-1}_{11}\mathcal{Q} \end{pmatrix}_{8 \times 8}, \quad A'_{11} = \begin{pmatrix} r^{-\frac{1}{2}}s^{\frac{1}{2}} & 0 & 0 & 0 \\ 0 & r^{\frac{1}{2}}s^{\frac{1}{2}} & 0 & 0 \\ 0 & 0 & r^{\frac{1}{2}}s^{\frac{1}{2}} & 0 \\ 0 & 0 & 0 & r^{\frac{1}{2}}s^{\frac{1}{2}} \end{pmatrix}_{4 \times 4},$$

$$A_{22} = \begin{pmatrix} A'_{22} & 0 \\ 0 & \mathcal{Q}A'^{-1}_{22}\mathcal{Q} \end{pmatrix}_{8 \times 8}, \quad A'_{22} = \begin{pmatrix} r^{-\frac{1}{2}}s^{-\frac{1}{2}} & 0 & 0 & 0 \\ 0 & r^{-\frac{1}{2}}s^{\frac{1}{2}} & 0 & 0 \\ 0 & 0 & r^{\frac{1}{2}}s^{\frac{1}{2}} & 0 \\ 0 & 0 & 0 & r^{\frac{1}{2}}s^{\frac{1}{2}} \end{pmatrix}_{4 \times 4},$$

$$A_{33} = \begin{pmatrix} A'_{33} & 0 \\ 0 & QA'_{33}Q \end{pmatrix}_{8 \times 8}, \quad A'_{33} = \begin{pmatrix} r^{-\frac{1}{2}}s^{-\frac{1}{2}} & 0 & 0 & 0 \\ 0 & r^{-\frac{1}{2}}s^{-\frac{1}{2}} & 0 & 0 \\ 0 & 0 & r^{-\frac{1}{2}}s^{\frac{1}{2}} & 0 \\ 0 & 0 & 0 & r^{\frac{1}{2}}s^{\frac{1}{2}} \end{pmatrix}_{4 \times 4},$$

$$A_{44} = \begin{pmatrix} A'_{44} & 0 \\ 0 & QA'_{44}Q \end{pmatrix}_{8 \times 8}, \quad A'_{44} = \begin{pmatrix} r^{-\frac{1}{2}}s^{-\frac{1}{2}} & 0 & 0 & 0 \\ 0 & r^{-\frac{1}{2}}s^{-\frac{1}{2}} & 0 & 0 \\ 0 & 0 & r^{-\frac{1}{2}}s^{-\frac{1}{2}} & 0 \\ 0 & 0 & 0 & r^{-\frac{1}{2}}s^{\frac{1}{2}} \end{pmatrix}_{4 \times 4},$$

where $Q = \sum_{i=1}^4 E_{5-i,i}$, $A_{i'i'} = A_{ii}^{-1}$ and

$$A_{ij} = (r^{-\frac{1}{2}}s^{\frac{1}{2}} - r^{\frac{1}{2}}s^{-\frac{1}{2}})\{E_{ji} - (r^{-\frac{1}{2}}s^{\frac{1}{2}})^{(\rho_{i'} - \rho_j)}E_{i'j'}\},$$

$1 \leq i < j \leq 8$, $E_{ij} \in M(8, \mathbb{K})$, where the multiplication between matrices \widehat{R} , L_1^\pm and L_2^\pm is matrix multiplication. From the equation $\widehat{R}L_1^+L_2^+ = L_2^+L_1^+\widehat{R}$, we can derive the following calculations:

$$\widehat{R}L_1^+L_2^+(v_1 \otimes v_j) = L_2^+L_1^+\widehat{R}(v_1 \otimes v_j) \Rightarrow \begin{cases} l_{11}^+l_{12}^+ = rl_{12}^+l_{11}^+, & l_{11}^+l_{23}^+ = l_{23}^+l_{11}^+, \\ l_{11}^+l_{34}^+ = l_{34}^+l_{11}^+, & l_{11}^+l_{35}^+ = r^{-1}s^{-1}l_{35}^+l_{11}^+, \end{cases}$$

where $1 \leq j \leq 7$;

$$\widehat{R}L_1^+L_2^+(v_2 \otimes v_j) = L_2^+L_1^+\widehat{R}(v_2 \otimes v_j) \Rightarrow \begin{cases} l_{22}^+l_{12}^+ = sl_{12}^+l_{22}^+, & l_{22}^+l_{23}^+ = rl_{23}^+l_{22}^+, \\ l_{22}^+l_{34}^+ = l_{34}^+l_{22}^+, & l_{22}^+l_{35}^+ = (rs)^{-1}l_{35}^+l_{22}^+, \\ l_{12}^+l_{34}^+ = l_{34}^+l_{12}^+, & l_{12}^+l_{35}^+ = (rs)^{-1}l_{35}^+l_{12}^+, \end{cases}$$

and we have

$$(4-2) \quad l_{12}^+l_{23}^+ + (r^{-1} - s^{-1})l_{22}^+l_{13}^+ = l_{23}^+l_{12}^+,$$

$$(4-3) \quad l_{12}^+l_{13}^+ = rl_{13}^+l_{12}^+,$$

where $1 \leq j \leq 8$ ($j \neq 7$);

$$\widehat{R}L_1^+L_2^+(v_3 \otimes v_j) = L_2^+L_1^+\widehat{R}(v_3 \otimes v_j) \Rightarrow \begin{cases} l_{33}^+l_{12}^+ = l_{12}^+l_{33}^+, & l_{33}^+l_{23}^+ = sl_{23}^+l_{33}^+, \\ l_{33}^+l_{34}^+ = rl_{34}^+l_{33}^+, & l_{33}^+l_{35}^+ = s^{-1}l_{35}^+l_{33}^+, \end{cases}$$

and we have

$$(4-4) \quad l_{23}^+l_{34}^+ + (r^{-1} - s^{-1})l_{33}^+l_{24}^+ = l_{34}^+l_{23}^+,$$

$$(4-5) \quad l_{23}^+l_{35}^+ + (r^{-1} - s^{-1})l_{33}^+l_{25}^+ = (rs)^{-1}l_{35}^+l_{23}^+,$$

$$(4-6) \quad l_{23}^+l_{24}^+ = rl_{24}^+l_{23}^+,$$

$$(4-7) \quad l_{23}^+l_{25}^+ = s^{-1}l_{25}^+l_{23}^+,$$

$$(4-8) \quad l_{13}^+l_{23}^+ = s^{-1}l_{23}^+l_{13}^+,$$

where $1 \leq j \leq 8$ ($j \neq 6$);

$$\widehat{R}L_1^+L_2^+(v_4 \otimes v_j) = L_2^+L_1^+\widehat{R}(v_4 \otimes v_j) \Rightarrow \begin{cases} l_{44}^+l_{12}^+ = l_{12}^+l_{44}^+, & l_{44}^+l_{23}^+ = l_{23}^+l_{44}^+, \\ l_{44}^+l_{34}^+ = sl_{34}^+l_{44}^+, & l_{44}^+l_{35}^+ = rl_{35}^+l_{44}^+, \end{cases}$$

and we have

$$(4-9) \quad l_{24}^+l_{34}^+ = s^{-1}l_{34}^+l_{24}^+,$$

$$(4-10) \quad l_{25}^+l_{35}^+ = s^{-1}l_{35}^+l_{25}^+,$$

where $1 \leq j \leq 8$ ($j \neq 5$). In particular, we get

$$\widehat{R}L_1^+L_2^+(v_4 \otimes v_5) = L_2^+L_1^+\widehat{R}(v_4 \otimes v_5),$$

$$\widehat{R}L_1^+L_2^+(v_5 \otimes v_4) = L_2^+L_1^+\widehat{R}(v_5 \otimes v_4),$$

then we obtain

$$(4-11) \quad l_{34}^+l_{35}^+ + rs^{-1}(r^{-\frac{1}{2}}s^{\frac{1}{2}} - r^{\frac{1}{2}}s^{-\frac{1}{2}})l_{36}^+l_{33}^+ = rs^{-1}l_{35}^+l_{34}^+,$$

$$(4-12) \quad r^{-1}sl_{35}^+l_{34}^+ + (r^{-\frac{1}{2}}s^{\frac{1}{2}} - r^{\frac{1}{2}}s^{-\frac{1}{2}})l_{36}^+l_{33}^+ = l_{34}^+l_{35}^+.$$

By (4-2), (4-3) and (4-8), we get

$$l_{12}^{+2}l_{23}^+ + rs^{-1}l_{23}^+l_{12}^{+2} = (rs^{-1} + 1)l_{12}^+l_{23}^+l_{12}^+,$$

$$l_{23}^{+2}l_{12}^+ + r^{-1}sl_{12}^+l_{23}^{+2} = (r^{-1}s + 1)l_{23}^+l_{12}^+l_{23}^+.$$

By (4-4), (4-6) and (4-9), we get

$$l_{23}^{+2}l_{34}^+ + rs^{-1}l_{34}^+l_{23}^{+2} = (rs^{-1} + 1)l_{23}^+l_{34}^+l_{23}^+,$$

$$l_{34}^{+2}l_{23}^+ + r^{-1}sl_{23}^+l_{34}^{+2} = (r^{-1}s + 1)l_{34}^+l_{23}^+l_{34}^+.$$

By (4-5), (4-7) and (4-10), we get

$$s^2l_{23}^{+2}l_{35}^+ + (rs)^{-1}l_{35}^+l_{23}^{+2} = (r^{-1}s + 1)l_{23}^+l_{35}^+l_{23}^+,$$

$$(rs)^{-1}l_{35}^+l_{23}^{+2} + s^2l_{23}^+l_{35}^{+2} = (r^{-1}s + 1)l_{35}^+l_{23}^+l_{35}^+.$$

By (4-11) and (4-12), we get

$$l_{34}^+l_{35}^+ = l_{35}^+l_{34}^+.$$

For the equation $\widehat{R}L_1^-L_2^- = L_2^-L_1^-\widehat{R}$, we can repeat the similar calculation process as above. Then we define a morphism $\phi_4 : U(\widehat{R}) \rightarrow U_{r,s}(\mathfrak{so}_8)$:

$$l_{11}^+ \mapsto (w'_1w'_2w'_3{}^{\frac{1}{2}}w'_4{}^{\frac{1}{2}})^{-1}, \quad l_{12}^+ \mapsto (r-s)e_1l_{11}^+,$$

$$l_{22}^+ \mapsto (w'_2w'_3{}^{\frac{1}{2}}w'_4{}^{\frac{1}{2}})^{-1}, \quad l_{23}^+ \mapsto (r-s)e_2l_{22}^+,$$

$$\begin{aligned}
 l_{33}^+ &\mapsto (w'_3 w'_4)^{-\frac{1}{2}}, & l_{34}^+ &\mapsto (r-s)e_3 l_{33}^+, \\
 l_{44}^+ &\mapsto (w_3^{-1} w'_4)^{-\frac{1}{2}}, & l_{35}^+ &\mapsto (r-s)e_4 l_{33}^+, \\
 l_{11}^- &\mapsto (w_1 w_2 (w_3 w_4)^{\frac{1}{2}})^{-1}, & l_{21}^- &\mapsto -(r-s)l_{11}^- f_1, \\
 l_{22}^- &\mapsto (w_2 (w_3 w_4)^{\frac{1}{2}})^{-1}, & l_{32}^- &\mapsto -(r-s)l_{22}^- f_2, \\
 l_{33}^- &\mapsto (w_3 w_4)^{-\frac{1}{2}}, & l_{43}^- &\mapsto -(r-s)l_{33}^- f_3, \\
 l_{44}^- &\mapsto (w_3^{-1} w_4)^{-\frac{1}{2}}, & l_{53}^- &\mapsto -(r-s)l_{33}^- f_4, \\
 l_{i'i'}^+ &\mapsto (l_{ii}^+)^{-1}, & l_{i'i'}^- &\mapsto (l_{ii}^-)^{-1},
 \end{aligned}$$

where $1 \leq i \leq 4$. It is obvious that ϕ_4 still preserves the algebra structure, the relations in \mathcal{B} and \mathcal{B}' , respectively. Next, we need to ensure that ϕ_4 preserves the cross relations of \mathcal{B} and \mathcal{B}' . Considering the equation $\widehat{R}L_1^+ L_2^- = L_2^- L_1^+ \widehat{R}$, we have:

$$\widehat{R}L_1^+ L_2^-(v_1 \otimes v_j) = L_2^- L_1^+ \widehat{R}(v_1 \otimes v_j) \Rightarrow \begin{cases} l_{11}^+ l_{21}^- = r^{-1} l_{21}^- l_{11}^+, & l_{11}^+ l_{32}^- = l_{32}^- l_{11}^+, \\ l_{11}^+ l_{43}^- = l_{43}^- l_{11}^+, & l_{11}^+ l_{53}^- = r s l_{53}^- l_{11}^+, \end{cases}$$

where $1 \leq j \leq 7$;

$$\widehat{R}L_1^+ L_2^-(v_2 \otimes v_j) = L_2^- L_1^+ \widehat{R}(v_2 \otimes v_j) \Rightarrow \begin{cases} l_{22}^+ l_{21}^- = s^{-1} l_{21}^- l_{22}^+, & l_{22}^+ l_{32}^- = r^{-1} l_{32}^- l_{22}^+, \\ l_{22}^+ l_{43}^- = l_{43}^- l_{22}^+, & l_{22}^+ l_{53}^- = r s l_{53}^- l_{22}^+, \\ r s l_{12}^+ l_{21}^- - l_{21}^- l_{12}^+ = (s-r)(l_{22}^+ l_{11}^- - l_{22}^+ l_{11}^+), \\ l_{32}^- l_{12}^+ = r l_{12}^+ l_{32}^-, & l_{43}^- l_{12}^+ = l_{12}^+ l_{43}^-, \\ l_{12}^+ l_{53}^- = r s l_{53}^- l_{12}^+, & l_{11}^- l_{12}^+ = s l_{12}^+ l_{11}^-, \\ l_{22}^- l_{12}^+ = r l_{12}^+ l_{22}^-, & l_{33}^- l_{12}^+ = l_{12}^+ l_{33}^-, \\ l_{44}^- l_{12}^+ = l_{12}^+ l_{44}^-, \end{cases}$$

where $1 \leq j \leq 8$ ($j \neq 7$);

$$\widehat{R}L_1^+ L_2^-(v_3 \otimes v_j) = L_2^- L_1^+ \widehat{R}(v_3 \otimes v_j) \Rightarrow \begin{cases} l_{33}^+ l_{21}^- = l_{21}^- l_{33}^+, & l_{33}^+ l_{32}^- = s^{-1} l_{32}^- l_{33}^+, \\ l_{33}^+ l_{43}^- = r^{-1} l_{43}^- l_{33}^+, & l_{33}^+ l_{53}^- = s l_{53}^- l_{33}^+, \\ l_{23}^+ l_{21}^- = s^{-1} l_{21}^- l_{23}^+, \\ r s l_{23}^+ l_{32}^- - l_{32}^- l_{23}^+ = (s-r)(l_{33}^- l_{22}^+ - l_{33}^- l_{22}^-), \\ l_{23}^+ l_{43}^- = r^{-1} l_{43}^- l_{23}^+, & l_{23}^+ l_{53}^- = s l_{53}^- l_{23}^+, \\ l_{23}^+ l_{11}^- = l_{11}^- l_{23}^+, & l_{23}^+ l_{22}^- = s^{-1} l_{22}^- l_{23}^+, \\ l_{23}^+ l_{33}^- = r^{-1} l_{33}^- l_{23}^+, & l_{23}^+ l_{44}^- = l_{44}^- l_{23}^+, \end{cases}$$

where $1 \leq j \leq 8$ ($j \neq 6$);

$$\widehat{R}L_1^+L_2^-(v_4 \otimes v_j) = L_2^-L_1^+\widehat{R}(v_4 \otimes v_j) \Rightarrow \begin{cases} l_{34}^+l_{11}^- = l_{11}^-l_{34}^+, & l_{34}^+l_{22}^- = l_{22}^-l_{34}^+, \\ l_{34}^+l_{33}^- = s^{-1}l_{33}^-l_{34}^+, & l_{34}^+l_{44}^- = r^{-1}l_{44}^-l_{34}^+, \\ l_{34}^+l_{21}^- = l_{21}^-l_{34}^+, & l_{34}^+l_{32}^- = s^{-1}l_{32}^-l_{34}^+, \\ rs l_{34}^+l_{43}^- - l_{43}^-l_{34}^+ = (s-r)(l_{44}^-l_{33}^+ - l_{44}^+l_{33}^-), \\ l_{34}^+l_{53}^- = l_{53}^-l_{34}^+, \\ l_{44}^+l_{21}^- = l_{21}^-l_{44}^+, & l_{44}^+l_{32}^- = l_{32}^-l_{44}^+, \\ l_{44}^+l_{43}^- = s^{-1}l_{43}^-l_{44}^+, & l_{44}^+l_{53}^- = sl_{53}^-l_{44}^+, \end{cases}$$

where $1 \leq j \leq 8$ ($j \neq 5$);

$$\widehat{R}L_1^+L_2^-(v_5 \otimes v_j) = L_2^-L_1^+\widehat{R}(v_5 \otimes v_j) \Rightarrow \begin{cases} l_{35}^+l_{11}^- = rsl_{11}^-l_{35}^+, & l_{35}^+l_{22}^- = rsl_{22}^-l_{35}^+, \\ l_{35}^+l_{33}^- = rl_{33}^-l_{35}^+, & l_{35}^+l_{44}^- = r^{-1}l_{44}^-l_{35}^+, \\ l_{35}^+l_{21}^- = rsl_{21}^-l_{35}^+, & l_{35}^+l_{32}^- = rl_{32}^-l_{35}^+, \\ l_{35}^+l_{43}^- = l_{43}^-l_{35}^+, \\ rsl_{35}^+l_{53}^- - l_{53}^-l_{35}^+ = (s-r)(l_{55}^-l_{33}^+ - l_{55}^+l_{33}^-), \end{cases}$$

where $1 \leq j \leq 8$. Now we proceed to the case of general n , restricting the generating relations (4-1) to $E_{ij} \otimes E_{kl}$, $2 \leq i, j, k, l \leq 2n - 1$, by induction, we get all commutation relations except those between $l_{11}^\pm, l_{12}^\pm, l_{21}^\pm$ and l_{ii}^\pm, l_{ij}^\pm . Repeating similar computations as above, we have the following relations:

(B1) The $l_{11}^{\pm 1}, l_{ii}^{\pm 1}$ all commute with one another and $l_{11}^{\pm 1}(l_{11}^{\pm 1})^{-1} = l_{ii}^{\pm 1}(l_{ii}^{\pm 1})^{-1} = 1$.

(B2) For $3 \leq i \leq n$, we have

$$\begin{aligned} l_{ii}^+l_{12}^+ &= l_{12}^+l_{ii}^+, & l_{ii}^-l_{12}^+ &= l_{12}^+l_{ii}^-, \\ l_{ii}^+l_{21}^- &= l_{21}^-l_{ii}^+, & l_{ii}^-l_{21}^- &= l_{21}^-l_{ii}^-, \\ l_{22}^+l_{12}^+ &= s^{-1}l_{12}^+l_{22}^+, & l_{22}^+l_{21}^- &= sl_{21}^-l_{22}^+, \\ l_{22}^-l_{12}^+ &= r^{-1}l_{12}^+l_{22}^-, & l_{22}^-l_{21}^- &= rl_{21}^-l_{22}^-. \end{aligned}$$

(B3) For $1 \leq i \leq n$, we have

$$\begin{aligned} l_{11}^+l_{i,i+1}^+ &= l_{i,i+1}^+l_{11}^+, & l_{11}^+l_{n-1,n+1}^+ &= (rs)^{-1}l_{n-1,n+1}^+l_{11}^+, \\ l_{11}^-l_{i+1,i}^- &= l_{i+1,i}^-l_{11}^-, & l_{11}^-l_{n+1,n-1}^- &= (rs)^{-1}l_{n+1,n-1}^-l_{11}^-, \\ l_{11}^-l_{i,i+1}^+ &= l_{i,i+1}^+l_{11}^-, & l_{11}^-l_{n-1,n+1}^+ &= rsl_{n-1,n+1}^+l_{11}^-, \\ l_{11}^+l_{i+1,i}^- &= l_{i+1,i}^-l_{11}^+, & l_{11}^+l_{n+1,n-1}^- &= rsl_{n+1,n-1}^-l_{11}^+. \end{aligned}$$

$$(B4) \quad \begin{aligned} (l_{12}^+)^2 l_{23}^+ + r s^{-1} l_{23}^+ (l_{12}^+)^2 &= (r s^{-1} + 1) l_{12}^+ l_{23}^+ l_{12}^+, \\ (l_{23}^+)^2 l_{12}^+ + r^{-1} s l_{12}^+ (l_{23}^+)^2 &= (r^{-1} s + 1) l_{23}^+ l_{12}^+ l_{23}^+, \\ (l_{21}^-)^2 l_{32}^- + r s^{-1} l_{32}^- (l_{21}^-)^2 &= (r s^{-1} + 1) l_{21}^- l_{32}^- l_{21}^-, \\ (l_{32}^-)^2 l_{21}^- + r^{-1} s l_{21}^- (l_{32}^-)^2 &= (r^{-1} s + 1) l_{32}^- l_{21}^- l_{32}^-. \end{aligned}$$

(B5) For $3 \leq i \leq n - 1$, we have

$$\begin{aligned} l_{12}^+ l_{i,i+1}^+ &= l_{i,i+1}^+ l_{12}^+, & l_{12}^+ l_{n-1,n+1}^+ &= (r s)^{-1} l_{n-1,n+1}^+ l_{12}^+, \\ l_{21}^- l_{i,i+1}^+ &= l_{i,i+1}^+ l_{21}^-, & l_{21}^- l_{n-1,n+1}^+ &= (r s)^{-1} l_{n-1,n+1}^+ l_{21}^-, \\ l_{12}^+ l_{i+1,i}^- &= l_{i+1,i}^- l_{12}^+, & l_{12}^+ l_{n+1,n-1}^- &= r s l_{n+1,n-1}^- l_{12}^+, \\ l_{21}^- l_{i+1,i}^- &= l_{i+1,i}^- l_{21}^-, & l_{21}^- l_{n+1,n-1}^- &= (r s)^{-1} l_{n+1,n-1}^- l_{21}^-. \end{aligned}$$

We give explicit expressions of the L -functionals l_{ij}^\pm in terms of the generators of $U_{r,s}(\mathfrak{so}_{2n})$. Define $\phi_n : U(\widehat{R}) \rightarrow U_{r,s}(\mathfrak{so}_{2n})$ as follows:

$$\begin{aligned} l_{ii}^+ &\mapsto (w'_{\beta_i})^{-1}, & l_{i,i+1}^+ &\mapsto (r - s) e_i l_{ii}^+, \\ l_{ii}^- &\mapsto (w_{\beta_i})^{-1}, & l_{i+1,i}^- &\mapsto -(r - s) l_{ii}^- f_i, \\ l_{nn}^+ &\mapsto (w'_{\beta_n})^{-1}, & l_{n-1,n+1}^+ &\mapsto (r - s) e_n l_{n-1,n-1}^+, \\ l_{nn}^- &\mapsto (w_{\beta_n})^{-1}, & l_{n+1,n-1}^- &\mapsto -(r - s) l_{n-1,n-1}^- f_n, \\ l_{i'i'}^+ &\mapsto w'_{\beta_i}, & l_{i'i'}^- &\mapsto w_{\beta_i}, \end{aligned}$$

where $\beta_i = \alpha_i + \dots + \alpha_{n-2} + \frac{1}{2}(\alpha_{n-1} + \alpha_n)$, $\beta_n = \frac{1}{2}(\alpha_n - \alpha_{n-1})$, $1 \leq i \leq n - 1$. By induction, we can prove that ϕ_n still preserves the structure of algebra $U_{r,s}(\mathfrak{so}_{2n})$. \square

Theorem 4.3. $\phi_n : U(\widehat{R}) \rightarrow U_{r,s}(\mathfrak{so}_{2n})$ is an algebraic isomorphism.

Proof. It is easy to check that the image of ϕ_n contains all generators of $U_{r,s}(\mathfrak{so}_{2n})$. Therefore, ϕ_n is surjective.

It remains to show that ϕ_n is injective. To this end, we need to construct an algebra homomorphism $\psi_n : U_{r,s}(\mathfrak{so}_{2n}) \rightarrow U(\widehat{R})$,

$$\begin{aligned} e_i &\mapsto \frac{1}{r-s} l_{i,i+1}^+ (l_{ii}^+)^{-1}, & f_i &\mapsto \frac{1}{s-r} (l_{ii}^-)^{-1} l_{i+1,i}^-, \\ w'_i &\mapsto (l_{ii}^+)^{-1} l_{i+1,i+1}^+, & w_i &\mapsto (l_{ii}^-)^{-1} l_{i+1,i+1}^-, \\ e_n &\mapsto \frac{1}{r-s} l_{n-1,n+1}^+ (l_{n-1,n-1}^+)^{-1}, & f_n &\mapsto \frac{1}{s-r} (l_{n-1,n-1}^-)^{-1} l_{n+1,n-1}^-, \\ w'_n &\mapsto (l_{nn}^+)^{-1} l_{n-1,n-1}^+, & w_n &\mapsto (l_{nn}^-)^{-1} l_{n-1,n-1}^-, \end{aligned}$$

which satisfies $\psi_n \circ \phi_n = \text{id}$.

To prove that ψ_n still preserves the algebra structure of $U(\widehat{R})$ is completely similar to that of Theorem 4.2. Hence, ϕ_n is injective. (For a similar proof in the one-parameter setting, one can refer to Section 8.5 of [17]). \square

Proposition 4.4. *For the braiding R -matrix $R = R_{VV}$, the spectral parameter dependent $R(z)$ is given by*

$$\begin{aligned}
R(z) &= \sum_{i=1}^{2n} E_{ii} \otimes E_{ii} \\
&+ \frac{rs(z-1)}{rz-s} \left\{ \sum_{\substack{1 \leq i \leq n-1 \\ i+1 \leq j \leq n}} E_{ij} \otimes E_{ji} + \sum_{\substack{1 \leq i \leq n-1 \\ i'+1 \leq j \leq 2n}} E_{ij} \otimes E_{ji} + \sum_{j=n+2}^{2n} E_{nj} \otimes E_{jn} \right. \\
&\quad \left. + \sum_{\substack{n+1 \leq i \leq 2n-1 \\ i+1 \leq j \leq 2n}} E_{ji} \otimes E_{ij} + \sum_{\substack{1 \leq i \leq n-1 \\ n+1 \leq j \leq 2n-i}} E_{ji} \otimes E_{ij} \right\} \\
&+ \frac{z-1}{rz-s} \left\{ \sum_{\substack{1 \leq i \leq n-1 \\ i+1 \leq j \leq n}} E_{ji} \otimes E_{ij} + \sum_{\substack{1 \leq i \leq n-1 \\ i'+1 \leq j \leq 2n}} E_{ji} \otimes E_{ij} + \sum_{j=n+2}^{2n} E_{jn} \otimes E_{nj} \right. \\
&\quad \left. + \sum_{\substack{n+1 \leq i \leq 2n-1 \\ i+1 \leq j \leq 2n}} E_{ij} \otimes E_{ji} + \sum_{\substack{1 \leq i \leq n-1 \\ n+1 \leq j \leq 2n-i}} E_{ij} \otimes E_{ji} \right\} \\
&+ \frac{r-s}{rz-s} \left\{ z \sum_{\substack{i < j \\ i' \neq j'}} E_{jj} \otimes E_{ii} + \sum_{\substack{i > j \\ i \neq j'}} E_{jj} \otimes E_{ii} \right\} \\
&+ \frac{1}{(z-r^{1-n}s^{n-1})(rz-s)} \sum_{i,j=1}^{2n} d_{ij}(z) E_{ij'} \otimes E_{i'j},
\end{aligned}$$

$$\text{where } d_{ij}(z) = \begin{cases} (s-r)z\{(r^{-\frac{1}{2}}s^{\frac{1}{2}})^{\rho_j-\rho_i}(z-1) - \delta_{ij'}[z-(rs^{-1})^{1-n}]\}, & \text{if } i > j, \\ (s-r)\{(r^{-\frac{1}{2}}s^{\frac{1}{2}})^{\rho_j-\rho_i+2n-2}(z-1) - \delta_{ij'}[z-(rs^{-1})^{1-n}]\}, & \text{if } i < j, \\ s[z-(rs^{-1})^{2-n}](z-1), & \text{if } i = j. \end{cases}$$

Remark 4.5. Consider the \widehat{R} -matrix $\widehat{R}(z) = P \circ R(z)$, where P is defined as in Remark 3.9:

$$\begin{aligned}
\widehat{R}(z) &= \sum_{i=1}^{2n} E_{ii} \otimes E_{ii} \\
&+ \frac{rs(z-1)}{rz-s} \left\{ \sum_{\substack{1 \leq i \leq n-1 \\ i+1 \leq j \leq n}} E_{jj} \otimes E_{ii} + \sum_{\substack{1 \leq i \leq n-1 \\ i'+1 \leq j \leq 2n}} E_{jj} \otimes E_{ii} + \sum_{j=n+2}^{2n} E_{jj} \otimes E_{nn} \right. \\
&\quad \left. + \sum_{\substack{n+1 \leq i \leq 2n-1 \\ i+1 \leq j \leq 2n}} E_{ii} \otimes E_{jj} + \sum_{\substack{1 \leq i \leq n-1 \\ n+1 \leq j \leq 2n-i}} E_{ii} \otimes E_{jj} \right\}
\end{aligned}$$

$$\begin{aligned}
 & + \frac{z-1}{rz-s} \left\{ \sum_{\substack{1 \leq i \leq n-1 \\ i+1 \leq j \leq n}} E_{ii} \otimes E_{jj} + \sum_{\substack{1 \leq i \leq n-1 \\ i'+1 \leq j \leq 2n}} E_{ii} \otimes E_{jj} + \sum_{j=n+2}^{2n} E_{nn} \otimes E_{jj} \right. \\
 & \qquad \qquad \qquad \left. + \sum_{\substack{n+1 \leq i \leq 2n-1 \\ i+1 \leq j \leq 2n}} E_{jj} \otimes E_{ii} + \sum_{\substack{1 \leq i \leq n-1 \\ n+1 \leq j \leq 2n-i}} E_{jj} \otimes E_{ii} \right\} \\
 & + \frac{r-s}{rz-s} \left\{ z \sum_{\substack{i < j \\ i' \neq j'}} E_{ij} \otimes E_{ji} + \sum_{\substack{i > j \\ i' \neq j'}} E_{ij} \otimes E_{ji} \right\} + \sum_{i,j=1}^{2n} c_{ij}(z) E_{i'j'} \otimes E_{ij},
 \end{aligned}$$

where

$$c_{ij}(z) = \frac{d_{ij}(z)}{(z - r^{1-n} s^{n-1})(rz - s)}.$$

It is easy to check that $\widehat{R}(z)$ satisfies the quantum Yang–Baxter equation

$$\widehat{R}_{12}(z) \widehat{R}_{13}(zw) \widehat{R}_{23}(w) = \widehat{R}_{23}(w) \widehat{R}_{13}(zw) \widehat{R}_{12}(z),$$

and the unitary condition

$$(4-13) \quad \widehat{R}_{21}(z) \widehat{R}(z^{-1}) = \widehat{R}(z^{-1}) \widehat{R}_{21}(z) = 1.$$

5. The algebra $\mathcal{U}(\widehat{R})$ and its Gauss decomposition

Definition 5.1. The algebra $\mathcal{U}(\widehat{R})$ is an associative algebra with generators $l_{kl}^{\pm}[\mp m]$ ($m \in \mathbb{Z}_+ \setminus \{0\}$), and $l_{kl}^+[0] = l_{lk}^-[0] = 0$, $1 \leq l \leq k \leq n$ and the central element c via $r^{\frac{c}{2}}$ or $s^{\frac{c}{2}}$. Let $l_{ij}^{\pm}(z) = \sum_{m=0}^{\infty} l_{ij}^{\pm}[\mp m] z^{\pm m}$, and $L^{\pm}(z) = \sum_{i,j=1}^n E_{ij} \otimes l_{ij}^{\pm}(z)$. Then the relations are given by the following matrix equations on $\text{End}(V^{\otimes 2}) \otimes \mathcal{U}(\widehat{R})$:

$$(5-1) \quad l_{ii}^+[0], l_{ii}^-[0] \text{ are invertible and } l_{ii}^+[0] l_{ii}^-[0] = l_{ii}^-[0] l_{ii}^+[0],$$

$$(5-2) \quad \widehat{R}\left(\frac{z}{w}\right) L_1^{\pm}(z) L_2^{\pm}(w) = L_2^{\pm}(w) L_1^{\pm}(z) \widehat{R}\left(\frac{z}{w}\right),$$

$$(5-3) \quad \widehat{R}\left(\frac{z_+}{w_-}\right) L_1^+(z) L_2^-(w) = L_2^-(w) L_1^+(z) \widehat{R}\left(\frac{z_-}{w_+}\right),$$

where $z_+ = zr^{\frac{c}{2}}$ and $z_- = zs^{\frac{c}{2}}$. Here (5-2) is expanded in the direction of either $\frac{z}{w}$ or $\frac{w}{z}$, and (5-3) is expanded in the direction of $\frac{z_-}{w_+}$.

Remark 5.2. From (5-3) and the unitary condition of \widehat{R} -matrix (4-13), we have

$$(5-4) \quad \widehat{R}\left(\frac{z_{\pm}}{w_{\mp}}\right) L_1^{\pm}(z) L_2^{\mp}(w) = L_2^{\mp}(w) L_1^{\pm}(z) \widehat{R}\left(\frac{z_{\mp}}{w_{\pm}}\right).$$

So the relations of generating series (5-2), (5-3) are equivalent to

$$(5-5) \quad L_1^\pm(z)^{-1} L_2^\pm(w)^{-1} \widehat{R}\left(\frac{z}{w}\right) = \widehat{R}\left(\frac{z}{w}\right) L_2^\pm(w)^{-1} L_1^\pm(z)^{-1},$$

$$(5-6) \quad L_1^\pm(z)^{-1} L_2^\mp(w)^{-1} \widehat{R}\left(\frac{z_\pm}{w_\mp}\right) = \widehat{R}\left(\frac{z_\pm}{w_\mp}\right) L_2^\mp(w)^{-1} L_1^\pm(z)^{-1}.$$

They are also equivalent to

$$(5-7) \quad L_2^\pm(w)^{-1} \widehat{R}\left(\frac{z}{w}\right) L_1^\pm(z) = L_1^\pm(z) \widehat{R}\left(\frac{z}{w}\right) L_2^\pm(w)^{-1},$$

$$(5-8) \quad L_2^\mp(w)^{-1} \widehat{R}\left(\frac{z_\pm}{w_\mp}\right) L_1^\pm(z) = L_1^\pm(z) \widehat{R}\left(\frac{z_\pm}{w_\mp}\right) L_2^\mp(w)^{-1}.$$

Remark 5.3. Here we present the specific matrix expression formulas for (5-2) and (5-3), and reveal the differences between type $D_n^{(1)}$ and type $A_n^{(1)}$. For $D_n^{(1)}$, write

$$L^\pm(z) = \begin{pmatrix} l_{11}^\pm(z) & l_{12}^\pm(z) & \cdots & l_{1,2n}^\pm(z) \\ l_{21}^\pm(z) & l_{22}^\pm(z) & \ddots & \vdots \\ \vdots & \ddots & \ddots & l_{2n-1,2n}^\pm(z) \\ l_{2n,1}^\pm(z) & \cdots & l_{2n,2n-1}^\pm(z) & l_{2n,2n}^\pm(z) \end{pmatrix}_{2n \times 2n},$$

then for the generators $L_1^\pm(z)$, $L_2^\pm(z)$, $\widehat{R}(z)$, we have that

$$L_1^\pm(z) = \begin{pmatrix} l_{11}^\pm(z) I_{2n} & \cdots & l_{1,2n}^\pm(z) I_{2n} \\ \vdots & \ddots & \vdots \\ l_{2n,1}^\pm(z) I_{2n} & \cdots & l_{2n,2n}^\pm(z) I_{2n} \end{pmatrix}_{4n^2 \times 4n^2},$$

$$L_2^\pm(z) = \begin{pmatrix} L^\pm(z) & 0 & \cdots & 0 \\ 0 & L^\pm(z) & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & L^\pm(z) \end{pmatrix}_{4n^2 \times 4n^2},$$

$$\widehat{R}(z) = \begin{pmatrix} B_{11}(z) & \cdots & B_{1,2n}(z) \\ \vdots & \ddots & \vdots \\ B_{2n,1}(z) & \cdots & B_{2n,2n}(z) \end{pmatrix}_{4n^2 \times 4n^2},$$

$$B_{ll}(z) = \begin{pmatrix} a_{l1}(z) & 0 & \cdots & 0 \\ 0 & a_{l2}(z) & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & a_{l,2n}(z) \end{pmatrix}_{2n \times 2n},$$

where $B_{ll}(z)$ is a diagonal matrix, and a_{lj} is the coefficient of element $E_{ll} \otimes E_{jj}$ in $\widehat{R}(z)$, and $B_{ij}(z) = b_{ij}(z)E_{ji} + c_{i'j'}(z)E_{i'j'}$, where $b_{ij}(z)$ is the coefficient of element $E_{ij} \otimes E_{ji}$ in $\widehat{R}(z)$, and $c_{ij}(z)$ is the coefficient of element $E_{i'j'} \otimes E_{ij}$ in $\widehat{R}(z)$.

The multiplication between matrices $\widehat{R}\left(\frac{z}{w}\right)$, $L_1^\pm(z)$, $L_2^\pm(w)$ is matrix multiplication. From

$$\widehat{R}\left(\frac{z}{w}\right)L_1^\pm(z)L_2^\pm(w) = L_2^\pm(w)L_1^\pm(z)\widehat{R}\left(\frac{z}{w}\right),$$

we can derive the following calculation:

$$\widehat{R}\left(\frac{z}{w}\right)L_1^\pm(z)L_2^\pm(w) = \begin{pmatrix} M_{11} & \cdots & M_{1,2n} \\ \vdots & \ddots & \vdots \\ M_{2n,1} & \cdots & M_{2n,2n} \end{pmatrix}_{4n^2 \times 4n^2}, \quad M_{ij} \in M(2n, \mathbb{K}),$$

$$L_2^\pm(w)L_1^\pm(z)\widehat{R}\left(\frac{z}{w}\right) = \begin{pmatrix} M'_{11} & \cdots & M'_{1,2n} \\ \vdots & \ddots & \vdots \\ M'_{2n,1} & \cdots & M'_{2n,2n} \end{pmatrix}_{4n^2 \times 4n^2}, \quad M'_{ij} \in M(2n, \mathbb{K}).$$

We only give two types of matrix expressions that will be used later. Taking $M_{ij} = M'_{ij}$, where $1 \leq i, j \leq n$, consider M_{ij} :

$$(5-9) \quad \begin{pmatrix} a_{i1}\left(\frac{z}{w}\right)l_{ij}^\pm(z) & & & & b_{i1}\left(\frac{z}{w}\right)l_{1j}^\pm(z) & & & & \\ & \ddots & & & \vdots & & & & \\ & & a_{ii}\left(\frac{z}{w}\right)l_{ij}^\pm(z) & & & & & & \\ & & \vdots & & \ddots & & & & \\ c_{i'1}\left(\frac{z}{w}\right)l_{i'j}^\pm(z) & \cdots & c_{i'i}\left(\frac{z}{w}\right)l_{i'j}^\pm(z) & \cdots & c_{i'i'}\left(\frac{z}{w}\right)l_{ij}^\pm(z) & \cdots & c_{i'1'}\left(\frac{z}{w}\right)l_{1j}^\pm(z) & & \\ & & \vdots & & & & \ddots & & \\ & & b_{i1'}\left(\frac{z}{w}\right)l_{1j}^\pm(z) & & & & & a_{i1'}\left(\frac{z}{w}\right)l_{ij}^\pm(z) & \end{pmatrix} L^\pm(w),$$

and $1 \leq i \leq n, 1+n \leq j, M_{ij}$:

$$(5-10) \quad \begin{pmatrix} a_{i1}\left(\frac{z}{w}\right)l_{ij}^\pm(z) & & & & b_{i1}\left(\frac{z}{w}\right)l_{1j}^\pm(z) & & & & \\ & \ddots & & & \vdots & & & & \\ c_{i'1}\left(\frac{z}{w}\right)l_{i'j}^\pm(z) & \cdots & c_{i'i}\left(\frac{z}{w}\right)l_{i'j}^\pm(z) & \cdots & c_{i'i'}\left(\frac{z}{w}\right)l_{i'j}^\pm(z) & \cdots & c_{i'1'}\left(\frac{z}{w}\right)l_{1j}^\pm(z) & & \\ & & \ddots & & \vdots & & & & \\ & & & & a_{ii}\left(\frac{z}{w}\right)l_{ij}^\pm(z) & & & & \\ & & & & \vdots & & \ddots & & \\ & & & & b_{i1'}\left(\frac{z}{w}\right)l_{1j}^\pm(z) & & & a_{i1'}\left(\frac{z}{w}\right)l_{ij}^\pm(z) & \end{pmatrix} L^\pm(w),$$

where the elements in the i' -th row except for the element at position (i', i') are all zero for type $A_n^{(1)}$. Consider M'_{ij} , for $1 \leq i, j \leq n$:

$$(5-11) \quad L^\pm(w) \begin{pmatrix} a_{j1}(\frac{z}{w})l_{ij}^\pm(z) & & & c_{1j'}(\frac{z}{w})l_{i1'}^\pm(z) & & & \\ & \ddots & & \vdots & & & \\ b_{1j}(\frac{z}{w})l_{i1}^\pm(z) \cdots a_{jj}(\frac{z}{w})l_{ij}^\pm(z) \cdots c_{jj'}(\frac{z}{w})l_{ij'}^\pm(z) \cdots b_{1'j}(\frac{z}{w})l_{i1'}^\pm(z) & & \ddots & \vdots & & & \\ & & & \vdots & & & \\ & & & c_{j'j'}(\frac{z}{w})l_{ij'}^\pm(z) & & & \\ & & & \vdots & & \ddots & \\ & & & c_{1'j'}(\frac{z}{w})l_{i1'}^\pm(z) & & & a_{j1'}(\frac{z}{w})l_{ij}^\pm(z) \end{pmatrix},$$

moreover, $1 \leq i \leq n, 1 + n \leq j$:

$$(5-12) \quad L^\pm(w) \begin{pmatrix} a_{j1}(\frac{z}{w})l_{ij}^\pm(z) & c_{1j'}(\frac{z}{w})l_{i1'}^\pm(z) & & & & & \\ & \ddots & & \vdots & & & \\ & & c_{j'j'}(\frac{z}{w})l_{ij'}^\pm(z) & & & & \\ & & \vdots & \ddots & & & \\ b_{1j}(\frac{z}{w})l_{i1}^\pm(z) \cdots c_{j'j'}(\frac{z}{w})l_{ij'}^\pm(z) \cdots a_{jj}(\frac{z}{w})l_{ij}^\pm(z) \cdots b_{1'j}(\frac{z}{w})l_{i1'}^\pm(z) & & & & \ddots & & \\ & & \vdots & & & \ddots & \\ & & c_{1'j'}(\frac{z}{w})l_{i1'}^\pm(z) & & & & a_{j1'}(\frac{z}{w})l_{ij}^\pm(z) \end{pmatrix},$$

where the elements in the j' -th column except for the element at position (j', j') are all zero for type $A_n^{(1)}$.

Definition 5.4. Let $X = (x_{ij})_{i,j=1}^n$ be a sequence matrix over a ring with identity. Denote by X^{ij} the submatrix obtained from X by deleting the i -th row and j -th column. Suppose that the matrix X^{ij} is invertible. The (i, j) -th quasideterminant $|X|_{ij}$ of X is defined by

$$(5-13) \quad |X|_{ij} = \begin{vmatrix} x_{11} & \cdots & x_{1j} & \cdots & x_{1n} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ x_{i1} & \cdots & x_{ij} & \cdots & x_{in} \\ \vdots & \ddots & \vdots & \ddots & \vdots \\ x_{n1} & \cdots & x_{nj} & \cdots & x_{nn} \end{vmatrix} = x_{ij} - r_i^j (X^{ij})^{-1} c_j^i,$$

where r_i^j is the row matrix obtained from the i -th row of X by deleting the element x_{ij} , and c_j^i is the column matrix obtained from the j -th column of X by deleting the element x_{ij} .

Proposition 5.5. $L^\pm(z)$ have the following unique decomposition

$$(5-14) \quad L^\pm(z) = F^\pm(z)K^\pm(z)E^\pm(z),$$

by applying the Gauss decomposition to $L^\pm(z)$, where we introduce matrices with $N \times N$, and $N = 2n$,

$$(5-15) \quad F^\pm(z) = \begin{pmatrix} 1 & 0 & \cdots & 0 \\ f_{21}^\pm(z) & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ f_{N1}^\pm(z) & \cdots & f_{N,N-1}^\pm(z) & 1 \end{pmatrix},$$

$$(5-16) \quad E^\pm(z) = \begin{pmatrix} 1 & e_{12}^\pm(z) & \cdots & e_{1N}^\pm(z) \\ 0 & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & e_{N-1,N}^\pm(z) \\ 0 & \cdots & 0 & 1 \end{pmatrix},$$

$$(5-17) \quad K^\pm(z) = \begin{pmatrix} k_1^\pm(z) & 0 & \cdots & 0 \\ 0 & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & k_N^\pm(z) \end{pmatrix}.$$

Their entries are found by the quasideterminant formulas

$$(5-18) \quad k_m^\pm(z) = \begin{vmatrix} l_{11}^\pm(z) & \cdots & l_{1,m-1}^\pm(z) & l_{1m}^\pm(z) \\ \vdots & \ddots & \vdots & \vdots \\ l_{m1}^\pm(z) & \cdots & l_{m,m-1}^\pm(z) & l_{mm}^\pm(z) \end{vmatrix}$$

for $1 \leq m \leq 2n$, $k_m^\pm(z) = \sum_{t \in \mathbb{Z}_+} k_m^\pm(\mp t) z^{\pm t}$;

$$(5-19) \quad e_{ij}^\pm(z) = k_i^\pm(z)^{-1} \begin{vmatrix} l_{11}^\pm(z) & \cdots & l_{1,i-1}^\pm(z) & l_{1j}^\pm(z) \\ \vdots & \ddots & \vdots & \vdots \\ l_{i1}^\pm(z) & \cdots & l_{i,i-1}^\pm(z) & l_{ij}^\pm(z) \end{vmatrix}$$

for $1 \leq i < j \leq 2n$; $e_{ij}^\pm(z) = \sum_{m \in \mathbb{Z}_+} e_{ij}^\pm(\mp m) z^{\pm m}$;

$$(5-20) \quad f_{ji}^\pm(z) = \begin{vmatrix} l_{11}^\pm(z) & \cdots & l_{1,i-1}^\pm(z) & l_{1i}^\pm(z) \\ \vdots & \ddots & \vdots & \vdots \\ l_{j1}^\pm(z) & \cdots & l_{j,i-1}^\pm(z) & l_{ji}^\pm(z) \end{vmatrix} k_i^\pm(z)^{-1}$$

for $1 \leq i < j \leq 2n$, $f_{ji}^\pm(z) = \sum_{m \in \mathbb{Z}_+} f_{ji}^\pm(\mp m) z^{\pm m}$.

6. Drinfeld realization of $U_{r,s}(\widehat{\mathfrak{so}}_{2n})$

In this section, we will study the commutation relations between the Gaussian generators and give the Drinfeld realization of $U_{r,s}(\widehat{\mathfrak{so}}_{2n})$.

Theorem 6.1. *In the algebra $\mathcal{U}(\widehat{R})$, we have*

$$\begin{aligned} X_i^+(z) &= e_{i,i+1}^+(z_+) - e_{i,i+1}^-(z_-), & X_n^+(z) &= e_{n-1,n+1}^+(z_+) - e_{n-1,n+1}^-(z_-), \\ X_i^-(z) &= f_{i+1,i}^+(z_-) - f_{i+1,i}^-(z_+), & X_n^-(z) &= f_{n+1,n-1}^+(z_-) - f_{n+1,n-1}^-(z_+). \end{aligned}$$

For the generators $\{k_i^\pm(z), X_j^\pm(z), k_{n+1}^\pm(z), X_n^\pm(z) \mid 1 \leq i \leq n, 1 \leq j \leq n-1\}$, the relations between $k_i^\pm(z)$ and $X_j^\pm(z)$ are the same as those in $U_{r,s}(\widehat{\mathfrak{gl}}_n)$. The other relations are those involving $k_i^\pm(z)$ and $k_{n+1}^\pm(w)$, that is,

$$\begin{aligned} k_i^\pm(z)k_{n+1}^\pm(w) &= k_{n+1}^\pm(w)k_i^\pm(z), \\ k_{n+1}^+(z)k_{n+1}^-(w) &= k_{n+1}^-(w)k_{n+1}^+(z), \\ k_i^\mp(z)k_{n+1}^\pm(w) \frac{z_\mp - w_\pm}{rz_\mp - sw_\pm} &= \frac{z_\pm - w_\mp}{rz_\pm - sw_\mp} k_{n+1}^\pm(w)k_i^\mp(z). \end{aligned}$$

The relations involving $k_t^\pm(w)$ ($1 \leq t \leq n+1$) and $X_n^\pm(z)$ are

$$\begin{aligned} k_l^\pm(w)X_n^\pm(z) &= rsX_n^\pm(z)k_l^\pm(w), \\ rsk_l^\pm(w)X_n^\mp(z) &= X_n^\mp(z)k_l^\pm(w), \quad 1 \leq l \leq n-2, \\ (rw - sz_\pm)k_{n-1}^\pm(w)X_n^+(z) &= rs(w - z_\pm)X_n^+(z)k_{n-1}^\pm(w), \\ rs(w - z_\mp)k_{n-1}^\pm(w)X_n^-(z) &= (rw - sz_\mp)X_n^-(z)k_{n-1}^\pm(w), \\ k_n^\pm(w)X_n^+(z) &= \frac{wr - sz_\pm}{w - z_\pm} X_n^+(z)k_n^\pm(w), \\ X_n^-(z)k_n^\pm(w) &= \frac{wr - sz_\mp}{w - z_\mp} k_n^\pm(w)X_n^-(z), \\ k_{n+1}^\pm(w)X_n^+(z) &= \frac{rs(w - z_\pm)}{sw - rz_\pm} X_n^+(z)k_{n+1}^\pm(w), \\ X_n^-(z)k_{n+1}^\pm(w) &= \frac{rs(w - z_\mp)}{sw - rz_\mp} k_{n+1}^\pm(w)X_n^-(z). \end{aligned}$$

The relations involving $k_{n+1}^\pm(z)$ and $X_l^\pm(w)$ ($1 \leq l \leq n-1$) are

$$\begin{aligned} k_{n+1}^\pm(z)X_l^\mp(w) &= X_l^\mp(w)k_{n+1}^\pm(z), \\ k_{n+1}^\pm(z)X_l^\pm(w) &= X_l^\pm(w)k_{n+1}^\pm(z), \quad 1 \leq l \leq n-2, \\ k_{n+1}^\pm(w)X_{n-1}^+(z) &= \frac{rs(z_\pm - w)}{z_\pm s - rw} X_{n-1}^+(z)k_{n+1}^\pm(w), \\ X_{n-1}^-(z)k_{n+1}^\pm(w) &= \frac{rs(z_\mp - w)}{z_\mp s - rw} k_{n+1}^\pm(w)X_{n-1}^-(z). \end{aligned}$$

For the relations involving $X_n^\pm(z)$ and $X_l^\pm(z)$ ($1 \leq l \leq n$), we have

$$\begin{aligned}
 X_n^\pm(w)X_l^\pm(z) &= X_l^\pm(z)X_n^\pm(w), \\
 X_n^\pm(w)X_l^\mp(z) &= X_l^\mp(z)X_n^\pm(w), \quad 1 \leq l \leq n-3, \\
 X_n^+(w)X_{n-2}^+(z) &= \frac{rz-sw}{z-w}X_{n-2}^+(z)X_n^+(w), \\
 X_n^-(w)X_{n-2}^-(z) &= \frac{z-w}{rz-sw}X_{n-2}^-(z)X_n^-(w), \\
 X_n^\pm(w)X_{n-2}^\mp(z) &= X_{n-2}^\mp(z)X_n^\pm(w), \\
 X_n^\pm(w)X_{n-1}^\pm(z) &= (rs)^{\pm 1}X_{n-1}^\pm(z)X_n^\pm(w), \\
 X_{n-1}^\pm(w)X_n^\mp(z) &= X_n^\mp(z)X_{n-1}^\pm(w), \\
 X_n^+(z)X_n^+(w) &= \frac{zr-ws}{zs-rw}X_n^+(w)X_n^+(z), \\
 X_n^-(z)X_n^-(w) &= \frac{zs-wr}{zr-sw}X_n^-(w)X_n^-(z), \\
 [X_n^+(z), X_l^-(w)] &= (s^{-1} - r^{-1})\delta_{nl} \left\{ \delta\left(\frac{z_-}{w_+}\right)k_{n+1}^-(w_+)k_n^-(w_+)^{-1} \right. \\
 &\quad \left. - \delta\left(\frac{z_+}{w_-}\right)k_{n+1}^+(z_+)k_n^+(z_+)^{-1} \right\},
 \end{aligned}$$

and the following (r, s) -Serre relations hold in $\mathcal{U}(\widehat{R})$:

$$(6-1) \quad \left\{ X_{n-2}^-(z_1)X_{n-2}^-(z_2)X_n^-(w) - (r+s)X_{n-2}^-(z_1)X_n^-(w)X_{n-2}^-(z_2) \right. \\ \left. + rsX_n^-(w)X_{n-2}^-(z_1)X_{n-2}^-(z_2) \right\} + \{z_1 \leftrightarrow z_2\} = 0,$$

$$(6-2) \quad \left\{ X_n^+(z_1)X_n^+(z_2)X_{n-2}^+(w) - (r+s)X_n^+(z_1)X_{n-2}^+(w)X_n^+(z_2) \right. \\ \left. + rsX_{n-2}^+(w)X_n^+(z_1)X_n^+(z_2) \right\} + \{z_1 \leftrightarrow z_2\} = 0,$$

$$(6-3) \quad \left\{ rsX_{n-2}^+(z_1)X_{n-2}^+(z_2)X_n^+(w) - (r+s)X_{n-2}^+(z_1)X_n^+(w)X_{n-2}^+(z_2) \right. \\ \left. + X_n^+(w)X_{n-2}^+(z_1)X_{n-2}^+(z_2) \right\} + \{z_1 \leftrightarrow z_2\} = 0,$$

$$(6-4) \quad \left\{ rsX_n^-(z_1)X_n^-(z_2)X_{n-2}^-(w) - (r+s)X_n^-(z_1)X_{n-2}^-(w)X_n^-(z_2) \right. \\ \left. + X_{n-2}^-(w)X_n^-(z_1)X_n^-(z_2) \right\} + \{z_1 \leftrightarrow z_2\} = 0,$$

where the formal delta function $\delta(z) = \sum_{n \in \mathbb{Z}} z^n$.

Proof. The proof is based on the induction on n . We consider first the case $n = 4$,

$$L^\pm(z) = \begin{pmatrix} l_{11}^\pm(z) & l_{12}^\pm(z) & \cdots & l_{18}^\pm(z) \\ l_{21}^\pm(z) & l_{22}^\pm(z) & \ddots & \vdots \\ \vdots & \ddots & \ddots & l_{78}^\pm(z) \\ l_{81}^\pm(z) & \cdots & l_{87}^\pm(z) & l_{88}^\pm(z) \end{pmatrix}_{8 \times 8}.$$

Observe (5-9) and (5-11) and restrict them to $E_{ij} \otimes E_{kl}$, $1 \leq i, j, k, l \leq 4$, then

$$\begin{aligned} \widehat{R}_1\left(\frac{z}{w}\right)\widetilde{L}_1^\pm(z)\widetilde{L}_2^\pm(w) &= \widetilde{L}_2^\pm(w)\widetilde{L}_1^\pm(z)\widehat{R}_1\left(\frac{z}{w}\right), \\ \widehat{R}_1\left(\frac{z_+}{w_-}\right)\widetilde{L}_1^\pm(z)\widetilde{L}_2^\pm(w) &= \widetilde{L}_2^\pm(w)\widetilde{L}_1^\pm(z)\widehat{R}_1\left(\frac{z_-}{w_+}\right), \\ \widetilde{L}^\pm(z) &= \begin{pmatrix} l_{11}^\pm(z) & l_{12}^\pm(z) & l_{13}^\pm(z) & l_{14}^\pm(z) \\ l_{21}^\pm(z) & l_{22}^\pm(z) & l_{23}^\pm(z) & l_{24}^\pm(z) \\ l_{31}^\pm(z) & l_{32}^\pm(z) & l_{33}^\pm(z) & l_{34}^\pm(z) \\ l_{41}^\pm(z) & l_{42}^\pm(z) & l_{43}^\pm(z) & l_{44}^\pm(z) \end{pmatrix}_{4 \times 4}, \end{aligned}$$

and

$$\begin{aligned} \widehat{R}_1\left(\frac{z}{w}\right) &= \sum_{i=1}^4 E_{ii} \otimes E_{ii} + \frac{s(w-z)}{sw-zr} \left(\sum_{i>j} E_{ii} \otimes E_{jj} + s^{-1} \sum_{i<j} E_{ii} \otimes E_{jj} \right) \\ &\quad + \frac{(s-r)w}{sw-zr} \left(\sum_{i>j} E_{ij} \otimes E_{ji} + \frac{z}{w} \sum_{i<j} E_{ij} \otimes E_{ji} \right). \end{aligned}$$

Jing and Liu [16] gave the following spectral parameter dependent $\widehat{R}_A\left(\frac{z}{w}\right)$ of $U_{r,s}(\widehat{\mathfrak{g}}_n)$, in particular, setting $n = 4$,

$$\begin{aligned} \widehat{R}_A\left(\frac{z}{w}\right) &= \sum_{i=1}^4 E_{ii} \otimes E_{ii} + \frac{w-z}{w-zrs^{-1}} \left(\sum_{i>j} E_{ii} \otimes E_{jj} + s^{-1} \sum_{i<j} E_{ii} \otimes E_{jj} \right) \\ &\quad + \frac{(1-rs^{-1})w}{w-zrs^{-1}} \left(\sum_{i>j} E_{ij} \otimes E_{ji} + \frac{z}{w} \sum_{i<j} E_{ij} \otimes E_{ji} \right), \end{aligned}$$

we get $\widehat{R}_A\left(\frac{z}{w}\right) = \widehat{R}_1\left(\frac{z}{w}\right)$. Thereby, we can directly have the relations of generators $\{X_1^\pm(z), X_2^\pm(z), X_3^\pm(z), k_1^\pm(z), k_2^\pm(z), k_3^\pm(z), k_4^\pm(z)\}$.

Next, we need to obtain the relations between the remaining Gauss generators. First, using the Gauss decomposition, we write down $L^\pm(z)$ and $L^\pm(z)^{-1}$:

$$\begin{aligned} L^\pm(z) &= \begin{pmatrix} k_1^\pm(z) & k_1^\pm(z)e_{12}^\pm(z) & \cdots \\ f_{21}^\pm(z)k_1^\pm(z) & \vdots & \vdots \\ \vdots & \vdots & \vdots \end{pmatrix}, \\ L^\pm(z)^{-1} &= \begin{pmatrix} \cdots & \cdots & \cdots \\ \cdots & \cdots & -e_{N-1,N}^\pm(z)k_N^\pm(z)^{-1} \\ \cdots & -k_N^\pm(z)^{-1}f_{N,N-1}^\pm(z) & k_N^\pm(z)^{-1} \end{pmatrix}. \end{aligned}$$

Then using the generating series relations (5-2) and (5-4), we can complete our proof by using the following lemmas. □

Lemma 6.2. *The following equations hold in $\mathcal{U}(\widehat{R})$:*

$$(6-5) \quad k_5^\pm(w)X_i^\pm(z) = X_i^\pm(z)k_5^\pm(w),$$

$$(6-6) \quad k_5^\pm(w)X_i^\mp(z) = X_i^\mp(z)k_5^\pm(w),$$

$$(6-7) \quad k_j^\pm(z)k_5^\pm(w) = k_5^\pm(w)k_j^\pm(z),$$

$$(6-8) \quad k_j^\pm(z)k_5^\mp(w) \frac{z_\pm - w_\mp}{rz_\pm - sw_\mp} = \frac{z_\mp - w_\pm}{rz_\mp - sw_\pm} k_5^\mp(w)k_j^\pm(z),$$

where $1 \leq i \leq 2$ and $1 \leq j \leq 3$.

Proof. Due to the observation made in formulas (5-9) and (5-11), the relations between the Gaussian generators mentioned above follow from those for the quantum affine algebra $U_{r,s}(\widehat{\mathfrak{gl}}_5)$; see [16] □

Lemma 6.3.

$$(6-9) \quad k_1^\pm(w)X_4^+(z) = rsX_4^+(z)k_1^\pm(w),$$

$$(6-10) \quad rsk_1^\pm(w)X_4^-(z) = X_4^-(z)k_1^\pm(w),$$

$$(6-11) \quad k_2^\pm(w)X_4^+(z) = rsX_4^+(z)k_2^\pm(w),$$

$$(6-12) \quad rsk_2^\pm(w)X_4^-(z) = X_4^-(z)k_2^\pm(w),$$

$$(6-13) \quad X_4^\pm(w)X_1^\pm(z) = X_1^\pm(z)X_4^\pm(w),$$

$$(6-14) \quad X_4^\pm(w)X_1^\mp(z) = X_1^\mp(z)X_4^\pm(w).$$

Proof. We only give details for one case of (6-9), (6-11), (6-13) and (6-14), the remaining relations are verified in a similar way. By (5-9) and (5-11), taking the equations $M_{11} = M'_{11}$, we get

$$a_{15} \left(\frac{z}{w} \right) k_3^\pm(w)e_{3,5}^\pm(w)l_{11}^\pm(z) = a_{13} \left(\frac{z}{w} \right) l_{11}^\pm(z)k_3^\pm(w)e_{3,5}^\pm(w),$$

and the relations between $k_3^\pm(w)e_{3,5}^\pm(w)$ and $l_{11}^\mp(z)$ we have

$$a_{15} \left(\frac{z_\pm}{w_\pm} \right) k_3^\pm(w)e_{3,5}^\pm(w)l_{11}^\mp(z) = a_{13} \left(\frac{z_\pm}{w_\mp} \right) l_{11}^\mp(z)k_3^\pm(w)e_{3,5}^\pm(w),$$

so that $k_1^+(z)X_4^+(w) = rsX_4^+(w)k_1^+(z)$. Now apply $M_{12} = M'_{12}$ to obtain

$$a_{25} \left(\frac{z}{w} \right) k_3^\pm(w)e_{3,5}^\pm(w)l_{12}^\pm(z) = a_{13} \left(\frac{z}{w} \right) l_{12}^\pm(z)k_3^\pm(w)e_{3,5}^\pm(w),$$

$$a_{25} \left(\frac{z_\mp}{w_\pm} \right) k_3^\pm(w)e_{3,5}^\pm(w)l_{12}^\mp(z) = a_{13} \left(\frac{z_\pm}{w_\mp} \right) l_{12}^\mp(z)k_3^\pm(w)e_{3,5}^\pm(w).$$

So we get $X_4^+(w)X_1^+(z) = X_1^+(z)X_4^+(w)$. Since $M_{21} = M'_{21}$, we get

$$a_{23} \left(\frac{z}{w} \right) k_3^\pm(w)e_{3,5}^\pm(w)l_{21}^\pm(z) = a_{23} \left(\frac{z}{w} \right) l_{21}^\pm(z)k_3^\pm(w)e_{3,5}^\pm(w).$$

Furthermore,

$$a_{15} \left(\frac{z_{\pm}}{w_{\pm}} \right) k_3^{\pm}(w) e_{3,5}^{\pm}(w) l_{21}^{\mp}(z) = a_{23} \left(\frac{z_{\pm}}{w_{\mp}} \right) l_{21}^{\mp}(z) k_3^{\pm}(w) e_{3,5}^{\pm}(w),$$

thus we prove a case of relation in (6-14). Since $M_{22} = M'_{22}$, the formula is

$$a_{25} \left(\frac{z}{w} \right) k_3^{\pm}(w) e_{3,5}^{\pm}(w) l_{22}^{\pm}(z) = a_{23} \left(\frac{z}{w} \right) l_{22}^{\pm}(z) k_3^{\pm}(w) e_{3,5}^{\pm}(w),$$

on the other hand, we have

$$a_{25} \left(\frac{z_{\pm}}{w_{\pm}} \right) k_3^{\pm}(w) e_{3,5}^{\pm}(w) l_{22}^{\mp}(z) = a_{23} \left(\frac{z_{\pm}}{w_{\mp}} \right) l_{22}^{\mp}(z) k_3^{\pm}(w) e_{3,5}^{\pm}(w).$$

As a final step, use the relations between $l_{21}^{\pm}(z)$ and $X_4^+(w)$ and those between $e_{12}^{\mp}(z)$ and $X_4^+(w)$, to come to the relation

$$k_2^+(z) X_4^+(w) = rs X_4^+(w) k_2^+(z),$$

as required. □

Lemma 6.4.

$$(6-15) \quad (rw - sz_{\pm}) k_3^{\pm}(w) X_4^+(z) = rs(w - z_{\pm}) X_4^+(z) k_3^{\pm}(w),$$

$$(6-16) \quad rs(w - z_{\mp}) k_3^{\pm}(w) X_4^-(z) = (rw - sz_{\mp}) X_4^-(z) k_3^{\pm}(w),$$

$$(6-17) \quad X_4^{\pm}(w) X_2^{\mp}(z) = X_2^{\mp}(z) X_4^{\pm}(w),$$

$$(z - w) X_4^+(w) X_2^+(z) = X_2^+(z) X_4^+(w) (rz - sw),$$

$$(6-18) \quad (rz - sw) X_4^-(w) X_2^-(z) = X_2^-(z) X_4^-(w) (z - w).$$

Proof. The arguments for both formulas are quite similar so we only give a proof of one case of (6-17) and (6-18), taking the equations $M_{13} = M'_{13}$, we get

$$(6-19) \quad a_{35} \left(\frac{z}{w} \right) l_{15}^{\pm}(w) l_{13}^{\pm}(z) + b_{53} \left(\frac{z}{w} \right) l_{13}^{\pm}(w) l_{15}^{\pm}(z) \\ = l_{13}^{\pm}(z) l_{15}^{\pm}(w),$$

$$(6-20) \quad a_{35} \left(\frac{z}{w} \right) l_{25}^{\pm}(w) l_{13}^{\pm}(z) + b_{53} \left(\frac{z}{w} \right) l_{23}^{\pm}(w) l_{15}^{\pm}(z) \\ = a_{12} \left(\frac{z}{w} \right) l_{13}^{\pm}(z) l_{25}^{\pm}(w) + b_{12} \left(\frac{z}{w} \right) l_{23}^{\pm}(z) l_{15}^{\pm}(w),$$

$$(6-21) \quad a_{35} \left(\frac{z}{w} \right) l_{35}^{\pm}(w) l_{13}^{\pm}(z) + b_{53} \left(\frac{z}{w} \right) l_{33}^{\pm}(w) l_{15}^{\pm}(z) \\ = a_{13} \left(\frac{z}{w} \right) l_{13}^{\pm}(z) l_{35}^{\pm}(w) + b_{13} \left(\frac{z}{w} \right) l_{33}^{\pm}(z) l_{15}^{\pm}(w).$$

Using $f_{32}^\pm(w)f_{21}^\pm(w) \cdot (6-19) - f_{32}^\pm(w)(6-20) - f_{31}^\pm(w) \cdot (6-19) + (6-21)$, through a lot of calculations, we can obtain

$$\begin{aligned}
 (6-22) \quad & a_{35} \left(\frac{z}{w} \right) k_3^\pm(w) e_{35}^\pm(w) l_{13}^\pm(z) + b_{53} \left(\frac{z}{w} \right) k_3^\pm(w) l_{15}^\pm(z) \\
 & \quad + a_{13} \left(\frac{z}{w} \right) f_{32}^\pm(w) l_{13}^\pm(z) k_2^\pm(w) e_{25}^\pm(w) \\
 & = a_{13} \left(\frac{z}{w} \right) l_{13}^\pm(z) f_{32}^\pm(w) k_2^\pm(w) e_{25}^\pm(w) + a_{13} \left(\frac{z}{w} \right) l_{13}^\pm(z) k_3^\pm(w) e_{35}^\pm(w) \\
 & \quad + b_{13} \left(\frac{z}{w} \right) k_3^\pm(w) l_{11}^\pm(z) e_{15}^\pm(w).
 \end{aligned}$$

Taking the equations $M_{23} = M'_{23}$, we have

$$\begin{aligned}
 (6-23) \quad & a_{35} \left(\frac{z}{w} \right) l_{15}^\pm(w) l_{23}^\pm(z) + b_{53} \left(\frac{z}{w} \right) l_{13}^\pm(w) l_{25}^\pm(z) \\
 & \quad = a_{21} \left(\frac{z}{w} \right) l_{23}^\pm(z) l_{15}^\pm(w) + b_{21} \left(\frac{z}{w} \right) l_{13}^\pm(z) l_{25}^\pm(w),
 \end{aligned}$$

$$\begin{aligned}
 (6-24) \quad & a_{35} \left(\frac{z}{w} \right) l_{25}^\pm(w) l_{23}^\pm(z) + b_{53} \left(\frac{z}{w} \right) l_{23}^\pm(w) l_{25}^\pm(z) \\
 & \quad = l_{23}^\pm(z) l_{25}^\pm(w),
 \end{aligned}$$

$$\begin{aligned}
 (6-25) \quad & a_{35} \left(\frac{z}{w} \right) l_{35}^\pm(w) l_{23}^\pm(z) + b_{53} \left(\frac{z}{w} \right) l_{33}^\pm(w) l_{25}^\pm(z) \\
 & \quad = a_{23} \left(\frac{z}{w} \right) l_{23}^\pm(z) l_{35}^\pm(w) + b_{23} \left(\frac{z}{w} \right) l_{33}^\pm(z) l_{25}^\pm(w).
 \end{aligned}$$

A similar calculation, we come to the relation

$$\begin{aligned}
 (6-26) \quad & a_{35} \left(\frac{z}{w} \right) k_3^\pm(w) e_{35}^\pm(w) l_{23}^\pm(z) + b_{53} \left(\frac{z}{w} \right) k_3^\pm(w) l_{25}^\pm(z) \\
 & = b_{21} \left(\frac{z}{w} \right) \{ f_{32}^\pm(w) f_{21}^\pm(w) l_{13}^\pm(z) k_2^\pm(w) e_{25}^\pm(w) - f_{31}^\pm(w) l_{13}^\pm(z) k_2^\pm(w) e_{25}^\pm(w) \} \\
 & \quad + a_{23} \left(\frac{z}{w} \right) \{ l_{23}^\pm(z) f_{32}^\pm(w) k_2^\pm(w) e_{25}^\pm(w) + l_{23}^\pm(z) k_3^\pm(w) e_{35}^\pm(w) \} \\
 & \quad + b_{23} \left(\frac{z}{w} \right) \{ l_{33}^\pm(z) k_2^\pm(w) e_{25}^\pm(w) + l_{21}^\pm(z) k_3^\pm(w) e_{15}^\pm(w) \} \\
 & \quad \quad - f_{32}^\pm(w) l_{23}^\pm(z) k_2^\pm(w) e_{25}^\pm(w).
 \end{aligned}$$

Furthermore, by using $-f_{21}^\pm(z) \cdot (6-22) + (6-26)$, we get

$$\begin{aligned}
 & a_{35} \left(\frac{z}{w} \right) k_3^\pm(w) e_{35}^\pm(w) k_2^\pm(z) e_{23}^\pm(z) + b_{53} \left(\frac{z}{w} \right) k_3^\pm(w) k_2^\pm(z) e_{25}^\pm(z) \\
 & \quad = b_{23} \left(\frac{z}{w} \right) k_2^\pm(z) k_3^\pm(w) e_{25}^\pm(w) + a_{23} \left(\frac{z}{w} \right) k_2^\pm(z) e_{23}^\pm(z) k_3^\pm(w) e_{35}^\pm(w),
 \end{aligned}$$

and taking into account the relations between $e_{3,5}^{\mp}(w)$ and $e_{23}^{\pm}(z)$, we have

$$\begin{aligned} a_{35} \left(\frac{z_{\pm}}{w_{\pm}} \right) k_3^{\mp}(w) e_{35}^{\mp}(w) k_2^{\pm}(z) e_{23}^{\pm}(z) + b_{53} \left(\frac{z_{\pm}}{w_{\pm}} \right) k_3^{\mp}(w) k_2^{\pm}(z) e_{25}^{\pm}(z) \\ = b_{23} \left(\frac{z_{\pm}}{w_{\mp}} \right) k_2^{\pm}(z) k_3^{\mp}(w) e_{25}^{\mp}(w) + a_{23} \left(\frac{z_{\pm}}{w_{\mp}} \right) k_2^{\pm}(z) e_{23}^{\pm}(z) k_3^{\mp}(w) e_{35}^{\mp}(w). \end{aligned}$$

Therefore, we can arrive at $(z-w)X_4^+(w)X_2^+(z) = X_2^+(z)X_4^+(w)(rz-sw)$. Now turn to (6-24). Taking the equation $M_{31} = M'_{31}$, we get

$$\begin{aligned} (6-27) \quad a_{15} \left(\frac{z_{\pm}}{w_{\pm}} \right) k_3^{\mp}(w) e_{35}^{\mp}(w) l_{31}^{\pm}(z) \\ = \left\{ l_{31}^{\pm}(z) - b_{32} \left(\frac{z_{\pm}}{w_{\mp}} \right) f_{32}^{\mp}(w) l_{21}^{\pm}(z) \right. \\ \left. + b_{31} \left(\frac{z_{\pm}}{w_{\mp}} \right) (f_{32}^{\mp}(w) f_{21}^{\mp}(w) - f_{31}^{\mp}(w)) l_{11}^{\pm}(z) \right\} k_3^{\mp}(w) e_{35}^{\mp}(w), \end{aligned}$$

$$\begin{aligned} (6-28) \quad a_{15} \left(\frac{z}{w} \right) k_3^{\pm}(w) e_{35}^{\pm}(w) l_{31}^{\pm}(z) \\ = \left\{ l_{31}^{\pm}(z) - b_{32} \left(\frac{z}{w} \right) f_{32}^{\pm}(w) l_{21}^{\pm}(z) \right. \\ \left. + b_{31} \left(\frac{z}{w} \right) (f_{32}^{\pm}(w) f_{21}^{\pm}(w) - f_{31}^{\pm}(w)) l_{11}^{\pm}(z) \right\} k_3^{\pm}(w) e_{35}^{\pm}(w). \end{aligned}$$

Using $M_{32} = M'_{32}$, consider the relations

$$\begin{aligned} (6-29) \quad a_{15} \left(\frac{z_{\pm}}{w_{\pm}} \right) k_3^{\mp}(w) e_{35}^{\mp}(w) l_{32}^{\pm}(z) \\ = \left\{ l_{32}^{\pm}(z) - b_{32} \left(\frac{z_{\pm}}{w_{\mp}} \right) f_{32}^{\mp}(w) l_{22}^{\pm}(z) \right. \\ \left. + b_{31} \left(\frac{z_{\pm}}{w_{\mp}} \right) (f_{32}^{\mp}(w) f_{21}^{\mp}(w) - f_{31}^{\mp}(w)) l_{12}^{\pm}(z) \right\} k_3^{\mp}(w) e_{35}^{\mp}(w), \end{aligned}$$

$$\begin{aligned} (6-30) \quad a_{15} \left(\frac{z}{w} \right) k_3^{\pm}(w) e_{35}^{\pm}(w) l_{32}^{\pm}(z) \\ = \left\{ l_{32}^{\pm}(z) - b_{32} \left(\frac{z}{w} \right) f_{32}^{\pm}(w) l_{22}^{\pm}(z) \right. \\ \left. + b_{31} \left(\frac{z}{w} \right) (f_{32}^{\pm}(w) f_{21}^{\pm}(w) - f_{31}^{\pm}(w)) l_{12}^{\pm}(z) \right\} k_3^{\pm}(w) e_{35}^{\pm}(w). \end{aligned}$$

By $-(6-27) \cdot e_{12}^{\pm}(z) + (6-29)$ and $-(6-28) \cdot e_{12}^{\pm}(z) + (6-30)$, we can obtain

$$\begin{aligned} a_{15} \left(\frac{z}{w} \right) k_3^{\pm}(w) e_{35}^{\pm}(w) f_{32}^{\pm}(z) k_2^{\pm}(z) \\ = \left\{ f_{32}^{\pm}(z) k_2^{\pm}(z) - b_{32} \left(\frac{z}{w} \right) f_{32}^{\pm}(w) k_2^{\pm}(z) \right\} k_3^{\pm}(w) e_{35}^{\pm}(w), \\ a_{15} \left(\frac{z_{\pm}}{w_{\pm}} \right) k_3^{\mp}(w) e_{35}^{\mp}(w) f_{32}^{\pm}(z) k_2^{\pm}(z) \\ = \left\{ f_{32}^{\pm}(z) k_2^{\pm}(z) - b_{32} \left(\frac{z_{\pm}}{w_{\mp}} \right) f_{32}^{\mp}(w) k_2^{\pm}(z) \right\} k_3^{\mp}(w) e_{35}^{\mp}(w), \end{aligned}$$

and therefore $X_4^+(w)X_2^-(z) = X_2^-(z)X_4^+(w)$. □

Lemma 6.5. *In the algebra $\mathcal{U}(\widehat{R})$, we have*

$$e_{36}^\pm(z) = f_{63}^\pm(z) = 0, \quad e_{45}^\pm(z) = f_{54}^\pm(z) = 0.$$

Proof. We only verify a case of the first relation. By (5-9) and (5-11), we have $M_{13} = M'_{13}$, and so get the relations

$$(6-31) \quad \sum_{i=1}^8 c_{i6} \left(\frac{z}{w}\right) l_{1i}^\pm(w) l_{1i'}^\pm(z) = l_{13}^\pm(z) l_{16}^\pm(w),$$

$$(6-32) \quad \sum_{i=1}^8 c_{i6} \left(\frac{z}{w}\right) l_{2i}^\pm(w) l_{1i'}^\pm(z) = a_{12} \left(\frac{z}{w}\right) l_{13}^\pm(z) l_{26}^\pm(w) + b_{12} \left(\frac{z}{w}\right) l_{23}^\pm(z) l_{16}^\pm(w),$$

$$(6-33) \quad \sum_{i=1}^8 c_{i6} \left(\frac{z}{w}\right) l_{3i}^\pm(w) l_{1i'}^\pm(z) = a_{13} \left(\frac{z}{w}\right) l_{13}^\pm(z) l_{36}^\pm(w) + b_{13} \left(\frac{z}{w}\right) l_{33}^\pm(z) l_{16}^\pm(w).$$

Let $(f_{32}^\pm(w) f_{21}^\pm(w) - f_{31}^\pm(w)) \cdot (6-31) - f_{32}^\pm(w) \cdot (6-32) + (6-33)$. Through a lot of calculations, we obtain

$$(6-34) \quad \sum_{i=3}^8 c_{i6} \left(\frac{z}{w}\right) k_3^\pm(w) e_{3i}^\pm(w) l_{1i'}^\pm(z) \\ = a_{13} \left(\frac{z}{w}\right) l_{13}^\pm(z) k_3^\pm(w) e_{36}^\pm(w) + b_{13} \left(\frac{z}{w}\right) k_3^\pm(w) k_1^\pm(z) \\ \cdot \{(e_{12}^\pm(w) - e_{12}^\pm(z)) e_{26}^\pm(w) - e_{16}^\pm(w)\}.$$

And from $M_{23} = M'_{23}$, we obtain

$$(6-35) \quad \sum_{i=1}^8 c_{i6} \left(\frac{z}{w}\right) l_{1i}^\pm(w) l_{2i'}^\pm(z) = a_{21} \left(\frac{z}{w}\right) l_{23}^\pm(z) l_{16}^\pm(w) + b_{21} \left(\frac{z}{w}\right) l_{13}^\pm(z) l_{26}^\pm(w),$$

$$(6-36) \quad \sum_{i=1}^8 c_{i6} \left(\frac{z}{w}\right) l_{2i}^\pm(w) l_{2i'}^\pm(z) = l_{23}^\pm(z) l_{26}^\pm(w),$$

$$(6-37) \quad \sum_{i=1}^8 c_{i6} \left(\frac{z}{w}\right) l_{3i}^\pm(w) l_{2i'}^\pm(z) = a_{23} \left(\frac{z}{w}\right) l_{23}^\pm(z) l_{36}^\pm(w) + b_{23} \left(\frac{z}{w}\right) l_{33}^\pm(z) l_{26}^\pm(w).$$

Calculating $(f_{32}^\pm(w) f_{21}^\pm(w) - f_{31}^\pm(w)) \cdot (6-35) - f_{32}^\pm(w) \cdot (6-36) + (6-37) - f_{21}^\pm(z) \cdot (6-34)$, we get

$$(6-38) \quad \sum_{i=3}^8 c_{i6} \left(\frac{z}{w}\right) k_3^\pm(w) e_{3i}^\pm(w) k_2^\pm(z) e_{2i'}^\pm(z) \\ = a_{23} \left(\frac{z}{w}\right) k_2^\pm(z) e_{23}^\pm(z) k_3^\pm(w) e_{36}^\pm(w) + b_{23} \left(\frac{z}{w}\right) k_3^\pm(w) k_2^\pm(z) e_{26}^\pm(w).$$

Using $M_{23} = M'_{23}$, we come by the relations

$$(6-39) \quad \sum_{i=1}^8 c_{i6} \left(\frac{z}{w}\right) l_{1i}^{\pm}(w) l_{3i'}^{\pm}(z) = a_{31} \left(\frac{z}{w}\right) l_{33}^{\pm}(z) l_{16}^{\pm}(w) + b_{31} \left(\frac{z}{w}\right) l_{13}^{\pm}(z) l_{36}^{\pm}(w),$$

$$(6-40) \quad \sum_{i=1}^8 c_{i6} \left(\frac{z}{w}\right) l_{2i}^{\pm}(w) l_{3i'}^{\pm}(z) = a_{32} \left(\frac{z}{w}\right) l_{33}^{\pm}(z) l_{26}^{\pm}(w) + b_{32} \left(\frac{z}{w}\right) l_{23}^{\pm}(z) l_{36}^{\pm}(w),$$

$$(6-41) \quad \sum_{i=1}^8 c_{i6} \left(\frac{z}{w}\right) l_{3i}^{\pm}(w) l_{3i'}^{\pm}(z) = l_{33}^{\pm}(z) l_{36}^{\pm}(w).$$

Calculating

$$(f_{32}^{\pm}(w) f_{21}^{\pm}(w) - f_{31}^{\pm}(w)) \cdot (6-39) - f_{32}^{\pm}(w) \cdot (6-40) + (6-41) - f_{32}^{\pm}(z) \cdot (6-38) - f_{31}^{\pm}(z) \cdot (6-34),$$

we arrive at the relation

$$(6-42) \quad \sum_{i=3}^8 c_{i6} \left(\frac{z}{w}\right) k_3^{\pm}(w) e_{3i}^{\pm}(w) l_{3i'}^{\pm}(z) = k_3^{\pm}(z) k_3^{\pm}(w) e_{36}^{\pm}(w).$$

Setting $z = w$, we get $e_{36}^{\pm}(z) = 0$. The remaining relations can be proved in a similar way. \square

Lemma 6.6. *The following equations hold in $\mathcal{U}(\widehat{R})$:*

$$(6-43) \quad X_4^{\pm}(w) X_3^{\pm}(z) = (rs)^{\pm 1} X_3^{\pm}(z) X_4^{\pm}(w),$$

$$(6-44) \quad X_3^{\pm}(w) X_4^{\mp}(z) = X_4^{\mp}(z) X_3^{\pm}(w).$$

Proof. The arguments are similar for all relations so we only prove the relation $X_4^+(w) X_3^+(z) = rs X_3^+(z) X_4^+(w)$. By (5-9)–(5-12), we have $M_{34} = M'_{34}$, and then get the relation

$$\sum_{i=1}^8 c_{i5} \left(\frac{z}{w}\right) l_{3i}^{\pm}(w) l_{3i'}^{\pm}(z) = l_{34}^{\pm}(z) l_{35}^{\pm}(w)$$

through a similar calculation process as in Lemma 6.5, which yields

$$(6-45) \quad \sum_{i=3}^8 c_{i5} \left(\frac{z}{w}\right) k_3^{\pm}(w) e_{3i}^{\pm}(w) k_3^{\pm}(z) e_{3i'}^{\pm}(z) = k_3^{\pm}(z) e_{34}^{\pm}(z) k_3^{\pm}(w) e_{35}^{\pm}(w),$$

and from $M_{35} = M'_{35}$, we obtain

$$(6-46) \quad \sum_{i=3}^8 c_{i4} \left(\frac{z}{w}\right) k_3^{\pm}(w) e_{3i}^{\pm}(w) k_3^{\pm}(z) e_{3i'}^{\pm}(z) = k_3^{\pm}(z) e_{35}^{\pm}(z) k_3^{\pm}(w) e_{34}^{\pm}(w).$$

Furthermore, $M_{33} = M'_{33}$ and $M_{36} = M'_{36}$ give that

$$(6-47) \quad \sum_{i=3}^8 c_{i6} \left(\frac{z}{w} \right) k_3^\pm(w) e_{3i}^\pm(w) k_3^\pm(z) e_{3i'}^\pm(z) = k_3^\pm(z) k_3^\pm(w) e_{36}^\pm(w),$$

$$(6-48) \quad \sum_{i=3}^8 c_{i3} \left(\frac{z}{w} \right) k_3^\pm(w) e_{3i}^\pm(w) k_3^\pm(z) e_{3i'}^\pm(z) = k_3^\pm(z) e_{36}^\pm(z) k_3^\pm(w).$$

Combining (6-45) with (6-46), we get that

$$(6-49) \quad (rs)^{-1} X_4^+(w) X_3^+(z) - X_3^+(w) X_4^+(z) = X_3^+(z) X_4^+(w) - (rs)^{-1} X_4^+(z) X_3^+(w).$$

Taking (6-45) and (6-47), owing to Lemma 6.5 and the fact that $e_{36}^\pm(z) = 0$, we have

$$(6-50) \quad \sum_{i=4}^5 k_3^\pm(w) e_{3i}^\pm(w) k_3^\pm(z) e_{3i'}^\pm(z) \left(c_{i5} \left(\frac{z}{w} \right) - (r^{-1}s)^{\frac{1}{2}} c_{i6} \left(\frac{z}{w} \right) \right) \\ = k_3^\pm(z) e_{34}^\pm(z) k_3^\pm(w) e_{35}^\pm(w).$$

Using the relations (6-46) and (6-48), we get

$$(6-51) \quad \sum_{i=4}^5 k_3^\pm(w) e_{3i}^\pm(w) k_3^\pm(z) e_{3i'}^\pm(z) \left(c_{i4} \left(\frac{z}{w} \right) - (rs^{-1})^{\frac{1}{2}} c_{i3} \left(\frac{z}{w} \right) \right) \\ = k_3^\pm(z) e_{35}^\pm(z) k_3^\pm(w) e_{34}^\pm(w).$$

Exchanging z and w in the relation (6-51), and combining into (6-50), we obtain

$$(6-52) \quad z(rs)^{-1} X_4^+(w) X_3^+(z) - w X_3^+(w) X_4^+(z) \\ = z X_3^+(z) X_4^+(w) - w(rs)^{-1} X_4^+(z) X_3^+(w)$$

By (6-49) and (6-52), we get $X_4^+(w) X_3^+(z) = rs X_3^+(z) X_4^+(w)$. \square

Lemma 6.7. *The following equations hold in $\mathcal{U}(\widehat{R})$:*

$$(6-53) \quad [X_4^+(z), X_4^-(w)] \\ = (s^{-1} - r^{-1}) \left\{ \delta \left(\frac{z_-}{w_+} \right) k_5^-(w_+) k_3^-(w_+)^{-1} - \delta \left(\frac{z_+}{w_-} \right) k_5^+(z_+) k_3^+(z_+)^{-1} \right\}.$$

Proof. By (5-10) and (5-12), we have $M_{35} = M'_{35}$ and then get the relations

$$a_{53} \left(\frac{z}{w} \right) l_{13}^\pm(w) l_{35}^\pm(z) + b_{35} \left(\frac{z}{w} \right) l_{15}^\pm(w) l_{33}^\pm(z) = b_{31} \left(\frac{z}{w} \right) l_{15}^\pm(z) l_{33}^\pm(w) + a_{31} \left(\frac{z}{w} \right) l_{35}^\pm(z) l_{13}^\pm(w), \\ a_{53} \left(\frac{z}{w} \right) l_{23}^\pm(w) l_{35}^\pm(z) + b_{35} \left(\frac{z}{w} \right) l_{25}^\pm(w) l_{33}^\pm(z) = b_{32} \left(\frac{z}{w} \right) l_{25}^\pm(z) l_{33}^\pm(w) + a_{32} \left(\frac{z}{w} \right) l_{35}^\pm(z) l_{23}^\pm(w), \\ a_{53} \left(\frac{z}{w} \right) l_{33}^\pm(w) l_{35}^\pm(z) + b_{35} \left(\frac{z}{w} \right) l_{35}^\pm(w) l_{33}^\pm(z) = l_{35}^\pm(z) l_{33}^\pm(w), \\ a_{53} \left(\frac{z}{w} \right) l_{53}^\pm(w) l_{35}^\pm(z) + b_{35} \left(\frac{z}{w} \right) l_{55}^\pm(w) l_{33}^\pm(z) = b_{35} \left(\frac{z}{w} \right) l_{55}^\pm(z) l_{33}^\pm(w) + a_{35} \left(\frac{z}{w} \right) l_{35}^\pm(z) l_{53}^\pm(w).$$

By straightforward calculations, one can check that

$$\begin{aligned}
 & [X_4^+(z), X_4^-(w)] \\
 &= (s^{-1} - r^{-1}) \left\{ \delta \left(\frac{z_-}{w_+} \right) k_5^-(w_+) k_3^-(w_+)^{-1} - \delta \left(\frac{z_+}{w_-} \right) k_5^+(z_+) k_3^+(z_+)^{-1} \right\}.
 \end{aligned}$$

This completes the proof. □

Lemma 6.8. *The following equations hold in $\mathcal{U}(\widehat{\mathcal{R}})$:*

$$(6-54) \quad k_5^\pm(w) X_3^+(z) = \frac{rs(z_\pm - w)}{z_\pm s - rw} X_3^+(z) k_5^\pm(w),$$

$$(6-55) \quad X_3^-(z) k_5^\pm(w) = \frac{rs(z_\mp - w)}{z_\mp s - rw} k_5^\pm(w) X_3^-(z),$$

$$(6-56) \quad k_4^\pm(w) X_4^+(z) = \frac{wr - sz_\pm}{w - z_\pm} X_4^+(z) k_4^\pm(w),$$

$$(6-57) \quad X_4^-(z) k_4^\pm(w) = \frac{wr - sz_\mp}{w - z_\mp} k_4^\pm(w) X_4^-(z).$$

Proof. We only give a proof of one case of (6-54). Similarly, we give the other identities. Using again $M_{35} = M'_{35}$ and $M_{34} = M'_{34}$, we get

$$\sum_{i=1}^8 c_{i5} \left(\frac{z}{w} \right) l_{5i}^\pm(w) l_{3i'}^\pm(z) = b_{35} \left(\frac{z}{w} \right) l_{54}^\pm(z) l_{35}^\pm(w) + a_{35} \left(\frac{z}{w} \right) l_{34}^\pm(z) l_{55}^\pm(w)$$

through the same calculating process as in Lemma 6.5, which yields

$$(6-58) \quad \sum_{i=5}^8 c_{i5} \left(\frac{z}{w} \right) k_5^\pm(w) e_{5i}^\pm(w) k_3^\pm(z) e_{3i'}^\pm(z) = a_{35} \left(\frac{z}{w} \right) k_3^\pm(z) e_{34}^\pm(z) k_5^\pm(w).$$

From $M_{35} = M'_{35}$, we obtain

$$\sum_{i=1}^8 c_{i4} \left(\frac{z}{w} \right) l_{5i}^\pm(w) l_{3i'}^\pm(z) = b_{35} \left(\frac{z}{w} \right) l_{55}^\pm(z) l_{34}^\pm(w) + a_{35} \left(\frac{z}{w} \right) l_{35}^\pm(z) l_{54}^\pm(w),$$

we arrive at the relation

$$(6-59) \quad \sum_{i=5}^8 c_{i4} \left(\frac{z}{w} \right) e_{5i}^\pm(w) k_3^\pm(z) e_{3i'}^\pm(z) = b_{35} \left(\frac{z}{w} \right) k_5^\pm(w) k_3^\pm(z) e_{34}^\pm(w).$$

Using (6-58) and (6-59), due to the relation $e_{45}^\pm(z) = f_{54}^\pm(z) = 0$, we obtain

$$\begin{aligned}
 & k_5^\pm(w) k_3^\pm(z) e_{34}^\pm(z) \left(c_{55} \left(\frac{z}{w} \right) - c_{54} \left(\frac{z}{w} \right) \right) \\
 &= a_{35} \left(\frac{z}{w} \right) k_3^\pm(z) e_{34}^\pm(z) k_5^\pm(w) - b_{35} \left(\frac{z}{w} \right) k_5^\pm(w) k_3^\pm(z) e_{34}^\pm(w).
 \end{aligned}$$

Finally, we get

$$k_5^+(w)X_3^+(z) = \frac{rs(z_+-w)}{z_+s-rw}X_3^+(z)k_5^+(w).$$

This completes the proof. \square

Lemma 6.9.

$$(6-60) \quad k_4^\pm(z)k_5^\pm(w) = k_5^\pm(w)k_4^\pm(z),$$

$$(6-61) \quad k_4^\pm(z)k_5^\mp(w) \frac{z_\pm-w_\mp}{rz_\pm-sw_\mp} = \frac{z_\mp-w_\pm}{rz_\mp-sw_\pm}k_5^\mp(w)k_4^\pm(z).$$

Proof. The proof is similar to that of Lemma 6.5. \square

Lemma 6.10. *The following equations hold in $\mathcal{U}(\widehat{R})$:*

$$(6-62) \quad k_5^\pm(z)k_5^\pm(w) = k_5^\pm(w)k_5^\pm(z),$$

$$(6-63) \quad k_5^\pm(z)k_5^\mp(w) = k_5^\mp(w)k_5^\pm(z),$$

$$(6-64) \quad k_5^\pm(w)X_4^+(z) = \frac{rs(w-z_\pm)}{sw-rz_\pm}X_4^+(z)k_5^\pm(w),$$

$$(6-65) \quad X_4^-(z)k_5^\pm(w) = \frac{rs(w-z_\mp)}{sw-rz_\mp}k_5^\pm(w)X_4^-(z).$$

Proof. Here we only prove (6-64) as the other relations can be shown similarly. By (5-10) and (5-12), we have $M_{35} = M'_{35}$. Then we can get the relation

$$l_{55}^\pm(w)l_{35}^\pm(z) = b_{35}\left(\frac{z}{w}\right)l_{55}^\pm(z)l_{35}^\pm(w) + a_{35}\left(\frac{z}{w}\right)l_{35}^\pm(z)l_{55}^\pm(w).$$

By straightforward calculations, one checks that

$$k_5^+(w)X_4^+(z) = \frac{rs(z_+-w)}{rz_+-sw}X_4^+(z)k_5^+(w).$$

This completes the proof. \square

Proposition 6.11. *The following equations hold in $\mathcal{U}(\widehat{R})$*

$$(6-66) \quad \{X_2^-(z_1)X_2^-(z_2)X_4^-(w) - (r+s)X_2^-(z_1)X_4^-(w)X_2^-(z_2) \\ + rsX_4^-(w)X_2^-(z_1)X_2^-(z_2)\} + \{z_1 \leftrightarrow z_2\} = 0,$$

$$(6-67) \quad \{X_4^+(z_1)X_4^+(z_2)X_2^+(w) - (r+s)X_4^+(z_1)X_2^+(w)X_4^+(z_2) \\ + rsX_2^+(w)X_4^+(z_1)X_4^+(z_2)\} + \{z_1 \leftrightarrow z_2\} = 0,$$

$$(6-68) \quad \{rsX_2^+(z_1)X_2^+(z_2)X_4^+(w) - (r+s)X_2^+(z_1)X_4^+(w)X_2^+(z_2) \\ + X_4^+(w)X_2^+(z_1)X_2^+(z_2)\} + \{z_1 \leftrightarrow z_2\} = 0,$$

$$(6-69) \quad \{rsX_4^-(z_1)X_4^-(z_2)X_2^-(w) - (r+s)X_4^-(z_1)X_2^-(w)X_4^-(z_2) \\ + X_2^-(w)X_4^-(z_1)X_4^-(z_2)\} + \{z_1 \leftrightarrow z_2\} = 0.$$

Proof. The equations can be proved similarly as in [16]. □

Now we proceed to the case of general n . We first restrict the relation to $E_{ij} \otimes E_{kl}$, $2 \leq i, j, k, l \leq 2n - 1$. By induction, we get all the commutation relations we need except those between $X_1^\pm(z)$, $k_1^\pm(z)$, and $X_n^\pm(z)$, $k_{n+1}^\pm(z)$.

Lemma 6.12. *The following equations hold in $\mathcal{U}(\widehat{R})$:*

$$(6-70) \quad k_1^\pm(z)X_n^+(w) = rsX_n^+(w)k_1^\pm(z),$$

$$(6-71) \quad rsk_1^\pm(z)X_n^-(w) = X_n^-(w)k_1^\pm(z),$$

$$(6-72) \quad X_n^\pm(w)X_1^\pm(z) = X_1^\pm(z)X_n^\pm(w),$$

$$(6-73) \quad X_n^\pm(w)X_1^\mp(z) = X_1^\mp(z)X_n^\pm(w),$$

$$(6-74) \quad k_1^\pm(z)k_{n+1}^\pm(w) = k_{n+1}^\pm(w)k_1^\pm(z),$$

$$(6-75) \quad k_{n+1}^\pm(z)k_{n+1}^\pm(w) = k_{n+1}^\pm(w)k_{n+1}^\pm(z),$$

$$(6-76) \quad k_{n+1}^\pm(z)k_{n+1}^\mp(w) = k_{n+1}^\mp(w)k_{n+1}^\pm(z),$$

$$(6-77) \quad k_{n+1}^\pm(z)X_n^+(w) = \frac{rs(w-z_\pm)}{sw-rz_\pm}X_n^+(w)k_{n+1}^\pm(z),$$

$$(6-78) \quad X_n^-(w)k_{n+1}^\pm(z) = \frac{rs(w-z_\mp)}{sw-rz_\mp}k_{n+1}^\pm(z)X_n^-(w),$$

$$(6-79) \quad \frac{w_\pm - z_\mp}{w_\pm r - z_\mp s}k_{n+1}^\mp(w)k_1^\pm(z) = \frac{w_\mp - z_\pm}{w_\mp r - z_\pm s}k_1^\pm(z)k_{n+1}^\mp(w),$$

$$(6-80) \quad [X_n^+(z), X_n^-(w)] = (s^{-1} - r^{-1}) \left\{ \delta\left(\frac{z_-}{w_+}\right)k_{n+1}^-(w_+)k_{n-1}^-(w_+)^{-1} - \delta\left(\frac{z_+}{w_-}\right)k_{n+1}^+(z_+)k_{n-1}^+(z_+)^{-1} \right\}.$$

Proof. By straightforward calculations one checks that the preceding formulas are correct. □

Finally, we define the map $\tau : U_{r,s}(\widehat{\mathfrak{so}}_{2n}) \rightarrow \mathcal{U}(\widehat{R})$ as follows:

$$\begin{aligned} x_i^\pm(z) &\mapsto (r-s)^{-1}X_i^\pm(z(rs^{-1})^{\frac{i}{2}}), \\ x_n^\pm(z) &\mapsto (r-s)^{-1}X_n^\pm(z(rs^{-1})^{\frac{n-1}{2}}), \\ \varphi_i(z) &\mapsto k_{i+1}^+(z(rs^{-1})^{\frac{i}{2}})k_i^+(z(rs^{-1})^{\frac{i}{2}})^{-1}, \\ \psi_i(z) &\mapsto k_{i+1}^-(z(rs^{-1})^{\frac{i}{2}})k_i^-(z(rs^{-1})^{\frac{i}{2}})^{-1}, \\ \varphi_n(z) &\mapsto k_{n+1}^+(z(rs^{-1})^{\frac{n-1}{2}})k_{n-1}^+(z(rs^{-1})^{\frac{n-1}{2}})^{-1}, \\ \psi_n(z) &\mapsto k_{n+1}^-(z(rs^{-1})^{\frac{n-1}{2}})k_{n-1}^-(z(rs^{-1})^{\frac{n-1}{2}})^{-1}, \end{aligned}$$

where $1 \leq i \leq n - 1$, and satisfy all the relations of Proposition 6.13.

Proposition 6.13. *In $U_{r,s}(\widehat{\mathfrak{so}}_{2n})$, the generating series $x_i^\pm(z)$, $\varphi_i(z)$, $\psi_i(z)$, $x_n^\pm(z)$, $\varphi_n(z)$, and $\psi_n(z)$, and the relations between $x_i^\pm(z)$, $\varphi_i(z)$, and $\psi_i(z)$ are the same as in $U_{r,s}(\widehat{\mathfrak{sl}}_n)$; the other relations as follows:*

$$(6-81) \quad [\varphi_j(z), \varphi_n(w)] = 0, \quad [\psi_j(z), \psi_n(w)] = 0,$$

$$(6-82) \quad \varphi_j(z)\psi_n(w) = \frac{g_{jn}\left(\frac{z_-}{w_+}\right)}{g_{jn}\left(\frac{z_+}{w_-}\right)}\psi_n(w)\varphi_j(z), \quad 1 \leq j \leq n,$$

$$(6-83) \quad \begin{aligned} \varphi_n(z)x_{n-2}^\pm(w) &= (rs)^{\pm 1}g_{n,n-2}\left(\frac{z}{w_\pm}\right)^{\pm 1}x_{n-2}^\pm(w)\varphi_n(z), \\ \psi_n(z)x_{n-2}^\pm(w) &= (rs)^{\mp 1}g_{n,n-2}\left(\frac{w_\mp}{z}\right)^{\mp 1}x_{n-2}^\pm(w)\varphi_n(z), \end{aligned}$$

$$(6-84) \quad \begin{aligned} \varphi_n(z)x_n^\pm(w) &= g_{nn}\left(\frac{z}{w_\pm}\right)^{\pm 1}x_n^\pm(w)\varphi_n(z), \\ \psi_n(z)x_n^\pm(w) &= g_{nn}\left(\frac{w_\mp}{z}\right)^{\mp 1}x_n^\pm(w)\varphi_n(z), \end{aligned}$$

$$(6-85) \quad \begin{aligned} \varphi_n(z)x_{n-1}^\pm(w) &= (rs)^{\pm 1}x_{n-1}^\pm(w)\varphi_n(z), \\ \psi_n(z)x_{n-1}^\pm(w) &= (rs)^{\mp 1}x_{n-1}^\pm(w)\psi_n(z), \\ \varphi_n(z)x_l^\pm(w) &= x_l^\pm(w)\varphi_n(z), \end{aligned}$$

$$(6-86) \quad \begin{aligned} \psi_n(z)x_l^\pm(w) &= x_l^\pm(w)\psi_n(z), \quad 1 \leq l \leq n-3, \\ \varphi_t(z)x_n^\pm(w) &= x_n^\pm(w)\varphi_t(z), \\ \psi_t(z)x_n^\pm(w) &= x_n^\pm(w)\psi_t(z), \quad 1 \leq t \leq n-3, \end{aligned}$$

$$(6-87) \quad \begin{aligned} \varphi_{n-2}(z)x_n^\pm(w) &= g_{n,n-2}\left(\frac{z}{w_\pm}\right)^{\pm 1}x_n^\pm(w)\varphi_{n-2}(z), \\ \psi_{n-2}(z)x_n^\pm(w) &= g_{n,n-2}\left(\frac{w_\mp}{z}\right)^{\mp 1}x_n^\pm(w)\varphi_{n-2}(z), \end{aligned}$$

$$(6-88) \quad \begin{aligned} x_{n-2}^\pm(z)x_n^\pm(w) &= g_{n,n-2}\left(\frac{z}{w}\right)^{\pm 1}x_n^\pm(w)x_{n-2}^\pm(z), \\ x_i^\pm(z)x_n^\pm(w) &= \langle w'_n, w_i \rangle^{\pm 1}x_n^\pm(w)x_i^\pm(z), \quad a_{in} = 0 \end{aligned}$$

$$(6-89) \quad [x_n^+(z), x_j^-(w)] = (s^{-1} - r^{-1})\delta_{jn} \left\{ \delta\left(\frac{z_-}{w_+}\right)\psi_n(w_+) - \delta\left(\frac{z_+}{w_-}\right)\varphi_n(z_+) \right\}, \quad j \leq n,$$

$$(6-90) \quad \begin{aligned} \{x_{n-2}^\pm(z_1)x_{n-2}^\pm(z_2)x_n^\pm(w) - (r^{\pm 1} + s^{\pm 1})x_{n-2}^\pm(z_1)x_n^\pm(w)x_{n-2}^\pm(z_2) \\ + (rs)^{\pm 1}x_n^\pm(w)x_{n-2}^\pm(z_1)x_{n-2}^\pm(z_2)\} + \{z_1 \leftrightarrow z_2\} = 0, \end{aligned}$$

$$(6-91) \quad \begin{aligned} \{x_n^\pm(z_1)x_n^\pm(z_2)x_{n-2}^\pm(w) - (r^{\mp 1} + s^{\mp 1})x_n^\pm(z_1)x_{n-2}^\pm(w)x_n^\pm(z_2) \\ + (rs)^{\mp 1}x_{n-2}^\pm(w)x_n^\pm(z_1)x_n^\pm(z_2)\} + \{z_1 \leftrightarrow z_2\} = 0, \end{aligned}$$

where $z_+ = zr^{\frac{\epsilon}{2}}$ and $z_- = zs^{\frac{\epsilon}{2}}$. We set $g_{ij}^{\pm}(z) = \sum_{n \in \mathbb{Z}_+} c_{ijn}^{\pm} z^n$, a formal power series in z , and it can be expressed as follows:

$$g_{ij}^{\pm}(z) = \frac{\langle w'_j, w_i \rangle^{\pm 1} z - (\langle w'_j, w_i \rangle \langle w'_i, w_j \rangle^{-1})^{\pm \frac{1}{2}}}{z - (\langle w'_i, w_j \rangle \langle w'_j, w_i \rangle)^{\pm \frac{1}{2}}}.$$

Acknowledgements

We thank Professors Zhengwei Liu, Jinsong Wu, and Yilong Wang for inviting Hu to report the current results in the international workshop on Advances in Quantum Algebras during Jan. 23–27, 2024 in BIMSA (Beijing Institute of Mathematical Sciences and Applications). We also thank ECNU for partial support of Xu’s master thesis where the basic R-matrices were announced in March 2023.

References

- [1] G. Benkart and S. Witherspoon, “Two-parameter quantum groups and Drinfel’d doubles”, *Algebr. Represent. Theory* **7**:3 (2004), 261–286. MR Zbl
- [2] N. Bergeron, Y. Gao, and N. Hu, “Drinfel’d doubles and Lusztig’s symmetries of two-parameter quantum groups”, *J. Algebra* **301**:1 (2006), 378–405. MR Zbl
- [3] N. Bergeron, Y. Gao, and N. Hu, “Representations of two-parameter quantum orthogonal and symplectic groups”, pp. 1–21 in *Proceedings of the International Conference on Complex Geometry and Related Fields*, edited by Z. Chen et al., AMS/IP Stud. Adv. Math. **39**, Amer. Math. Soc., Providence, RI, 2007. MR Zbl
- [4] X. Y. Chen, N. H. Hu, and X. L. Wang, “Convex PBW-type Lyndon bases and restricted two-parameter quantum group of type F_4 ”, *Acta Math. Sin. (Engl. Ser.)* **39**:6 (2023), 1053–1084. MR Zbl
- [5] J. T. Ding and I. B. Frenkel, “Isomorphism of two realizations of quantum affine algebra $U_q(\widehat{\mathfrak{gl}(n)})$ ”, *Comm. Math. Phys.* **156**:2 (1993), 277–300. MR Zbl
- [6] V. G. Drinfeld, “Hopf algebras and the quantum Yang–Baxter equation”, *Dokl. Akad. Nauk SSSR* **283**:5 (1985), 1060–1064. In Russian; translated in *Sov. Math. Dokl.* **32** (1985), 245–258. MR Zbl
- [7] V. G. Drinfeld, “A new realization of Yangians and of quantum affine algebras”, *Dokl. Akad. Nauk SSSR* **296**:1 (1987), 13–17. In Russian; translated in *Sov. Math. Dokl.* **36** (1988), 212–216. MR Zbl
- [8] V. G. Drinfeld, “Quantum groups”, pp. 798–820 in *Proceedings of the International Congress of Mathematicians, Vol. 1, 2* (Berkeley, CA, 1986), Amer. Math. Soc., Providence, RI, 1987. MR
- [9] L. D. Faddeev, N. Yu. Reshetikhin, and L. A. Takhtajan, “Quantization of Lie groups and Lie algebras”, pp. 299–309 in *Yang–Baxter equations in integrable systems*.
- [10] H. Garland, “The arithmetic theory of loop groups”, *Inst. Hautes Études Sci. Publ. Math.* **52** (1980), 5–136. MR Zbl
- [11] N. Hu and Y. Pei, “Notes on two-parameter quantum groups, II”, *Comm. Algebra* **40**:9 (2012), 3202–3220. MR Zbl
- [12] N. Hu and H. Zhang, “Generating functions with τ -invariance and vertex representations of quantum affine algebras $U_{r,s}(\hat{\mathfrak{g}})$, I: simply-laced cases”, 2014. arXiv 1401.4925

- [13] N. Hu, M. Rosso, and H. Zhang, “Two-parameter quantum affine algebra $U_{r,s}(\widehat{\mathfrak{sl}}_n)$, Drinfeld realization and quantum affine Lyndon basis”, *Comm. Math. Phys.* **278**:2 (2008), 453–486. MR Zbl
- [14] N. Hu, X. Xu, and R. Zhuang, “*RLL*-realization of two-parameter quantum affine algebra of type $B_n^{(1)}$ ”, 2024. arXiv 2405.06587
- [15] M. Jimbo, “A q -difference analogue of $U(\mathfrak{g})$ and the Yang–Baxter equation”, *Lett. Math. Phys.* **10**:1 (1985), 63–69. MR Zbl
- [16] N. Jing and M. Liu, “ R -matrix realization of two-parameter quantum affine algebra $U_{r,s}(\widehat{\mathfrak{gl}}_n)$ ”, *J. Algebra* **488** (2017), 1–28. MR Zbl
- [17] N. Jing, M. Liu, and A. Molev, “Isomorphism between the R -matrix and Drinfeld presentations of quantum affine algebra: type C ”, *J. Math. Phys.* **61**:3 (2020), art. id. 031701. MR
- [18] N. Jing, M. Liu, and A. Molev, “Isomorphism between the R -matrix and Drinfeld presentations of quantum affine algebra: types B and D ”, *SIGMA Symmetry Integrability Geom. Methods Appl.* **16** (2020), art. id. 043. MR Zbl
- [19] A. Klimyk and K. Schmüdgen, *Quantum groups and their representations*, Springer, 1997. MR Zbl
- [20] N. Y. Reshetikhin and M. A. Semenov-Tian-Shansky, “Central extensions of quantum current groups”, *Lett. Math. Phys.* **19**:2 (1990), 133–142. MR Zbl
- [21] N. Y. Reshetikhin, L. A. Takhtajan, and L. D. Faddeev, “Quantization of Lie groups and Lie algebras”, *Algebra i Analiz* **1**:1 (1989), 178–206. In Russian; translated in *Leningrad Math. J.* **1** (1990), 193–225. MR Zbl
- [22] M. Takeuchi, “A two-parameter quantization of $GL(n)$ (summary)”, *Proc. Japan Acad. Ser. A Math. Sci.* **66**:5 (1990), 112–114. MR Zbl
- [23] X. Zhong, N. Hu, and N. Jing, “*RLL*-realization of two-parameter quantum affine algebra in type $C_n^{(1)}$ ”, 2024. arXiv 2405.06597

Received February 8, 2024. Revised May 17, 2024.

RUSHU ZHUANG

52205500005@stu.ecnu.edu.cn

NAIHONG HU

nhhu@math.ecnu.edu.cn

XIAO XU

52275500004@stu.ecnu.edu.cn

(all authors)

SCHOOL OF MATHEMATICAL SCIENCES

MOE-KLMEA & SH-KLPMMP

EAST CHINA NORMAL UNIVERSITY

SHANGHAI

CHINA

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Fakultät für Mathematik
Universität Wien
Vienna, Austria
matthias.aschenbrenner@univie.ac.at

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Atsushi Ichino
Department of Mathematics
Kyoto University
Kyoto 606-8502, Japan
atsushi.ichino@gmail.com

Robert Lipshitz
Department of Mathematics
University of Oregon
Eugene, OR 97403
lipshitz@uoregon.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Dimitri Shlyakhtenko
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
shlyakht@ipam.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Ruixiang Zhang
Department of Mathematics
University of California
Berkeley, CA 94720-3840
ruixiang@berkeley.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2024 is US \$645/year for the electronic version, and \$875/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2024 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 329 No. 2 April 2024

Concentration inequalities for Paley–Wiener spaces SYED HUSAIN and FRIEDRICH LITTMANN	201
Characterizing the Fourier transform by its properties MATEUSZ KRUKOWSKI	217
Reduction types of CM curves MENTZELOS MELISTAS	233
The local character expansion as branching rules: nilpotent cones and the case of $SL(2)$ MONICA NEVINS	259
Extremely closed subgroups and a variant on Glauberman’s Z^* -theorem HUNG P. TONG-VIET	303
Vishik equivalence and similarity of quasilinear p -forms and totally singular quadratic forms KRISTÝNA ZEMKOVÁ	327
RLL -realization of two-parameter quantum affine algebra in type $D_n^{(1)}$ RUSHU ZHUANG, NAIHONG HU and XIAO XU	357