

*Pacific
Journal of
Mathematics*

HIGHER-GENUS QUANTUM K -THEORY

YOU-CHENG CHOU, LEO HERR AND YUAN-PIN LEE

Volume 330 No. 1

May 2024

HIGHER-GENUS QUANTUM K -THEORY

YOU-CHENG CHOU, LEO HERR AND YUAN-PIN LEE

We prove genus g invariants in quantum K -theory are determined by genus zero invariants of a smooth stack in the spirit of K. Costello's result in Gromov–Witten theory.

0. Introduction

Let X be a smooth quasiprojective variety over \mathbb{C} . Let $\overline{M}_{g,R}(X, \beta)$ be the space of genus g , R -pointed stable maps to X with degree β . The perfect obstruction theory on $\overline{M}_{g,R}(X, \beta)$ [Behrend and Fantechi 1997] endows the moduli stack with a “virtual structure sheaf” $\mathcal{O}_{\overline{M}_{g,R}(X, \beta)}^{\text{vir}}$ [Lee 2004].

Let $\alpha_i \in K^\circ(X)$ and L_i be the universal cotangent line bundles. When the insertion

$$\Omega := \sum_I a_I \prod_{i=1}^R L_i^{k_i} \otimes ev_i^* \alpha_i$$

has an action by S_R , the permutation-equivariant pushforward

$$(1) \quad \sum_j (-1)^j H^j(\overline{M}_{g,R}(X, \beta), \mathcal{O}^{\text{vir}} \otimes \Omega)$$

is an element in the Grothendieck group of S_R -representations with \oplus , i.e., virtual representations. We can also take a subgroup of S_R instead. These are by definition the *permutation-equivariant quantum K -invariants*.

The main theorem of this paper is the following.

Theorem 0.1 (see Theorem 3.1). *Genus g quantum K -invariants on X can be computed from permutation-equivariant genus zero quantum K -invariants on*

$$[\text{Sym}^{g+1} X] = [X^{g+1}/S_{g+1}].$$

A similar statement holds for X a smooth DM stack with projective coarse moduli space.

MSC2020: 14N35, 19E08.

Keywords: quantum K -theory, Costello's pushforward formula.

We believe the higher-genus *permutation-equivariant* quantum K -theory of X can also be computed from genus zero theory of $[\mathrm{Sym}^{g+1}X]$ by extending our methods.

When the target, X itself or $[\mathrm{Sym}^{g+1}X]$, is a Deligne–Mumford stack, the definition of $\overline{M}_{g,R}(X, \beta)$ involves twisted/orbifold curves and twisted stable maps. The domain curves are families of pointed nodal curves with cyclic gerbe structures at the marked points and nodes, such that the gerbe structures at the nodes are *balanced*. This means they are locally stack quotients of the node $R[x, y]/(xy)$ by the antidiagonal action

$$\zeta \cdot (x, y) := (\zeta x, \zeta^{-1} y), \quad \zeta \in \mu_r.$$

Note that we do *not* require the μ_r -gerbe structures at the marked points to be trivial in families as in [Costello 2006]. *Twisted stable maps* are *representable* morphisms from twisted curves to the target with finite automorphisms. $\overline{M}_{g,R}(X, \beta)$ are the moduli stacks of (twisted) stable maps with the discrete data g, R, β .

Hence, the marked points are no longer literal “points”, but gerbes. Due to the nontrivial gerbes at the marked points, the evaluation maps have the natural codomain a partially rigidified inertia stack $\overline{I}(X)$, instead of the inertia stack IX . This has been done in [Chen and Ruan 2002; 2004] and [Abramovich et al. 2008] in the context of cohomology and Chow groups.

We only need one class pulled back from $\overline{I}(-)$ as opposed to ordinary K -theory, which comes from $[\mathrm{Sym}^k X]$ for some $k \leq g + 1$ in Section 3B. We do not need the full K -theory of $\overline{I}([\mathrm{Sym}^{g+1}X])$.

The appearance of permutation-equivariant K -theory is quite natural, not simply a “technical clutch”. In cohomological Gromov–Witten theory, we often rely on the fact that the substacks (“strata”) appearing in “common operations” (e.g., fixed-point loci in torus localizations or the components of inertia stacks of the moduli) are *variants* of known quantities in the sense of induction. These variants can be identified with the actual known quantities in Gromov–Witten theory by simple modifications. For example, for the purpose of computing Gromov–Witten invariants, we have

$$\int_{[\overline{M}/S_n]} \cdots = \frac{1}{n!} \int_{[\overline{M}]} \pi^*(\cdots).$$

These equalities are no longer true in quantum K -theory. In fact, we have

$$\chi([\overline{M}/S_n], \cdots) = \chi_{S_n}(\overline{M}, \pi^*(\cdots))^{S_n}.$$

We note that K -theory on $[\overline{M}/S_n]$ can be identified with the S_n -equivariant K -theory on \overline{M} , and $\chi_{S_n}(-)^{S_n}$ is the pushforward in the S_n -equivariant theory, i.e., the S_n -invariant part $(-)^{S_n}$ of alternating sum of sheaf cohomologies viewed as S_n representations $\chi_{S_n}(-)$. This necessitates permutation-equivariant quantum K -theory.

Quantum K -theory has already been defined for stacks; see, e.g., [Tonita and Tseng 2013] and [Zhang, Section 2.4]. A comparison of the quantum K -theories with trivial and nontrivial gerbes at marked points can be found in [Zhang, Remark 2.8]. See also [Abramovich et al. 2002, Sections 4.4, 4.5] in the cohomological context. We allow nontrivial gerbes and recall the basic definitions in Section 1E.

Quantum K -invariants are roughly Gromov–Witten invariants computed in K -theory instead of cohomology or Chow groups. The idea of computing genus g Gromov–Witten invariants of any smooth projective variety X in terms of genus zero quantum K -invariants of quotient stack $[\mathrm{Sym}^{g+1} X]$ goes back to M. Kontsevich and was independently obtained in K. Costello’s thesis [Costello 2006]. This paper can be considered as a K -theoretic version of this circle of ideas.

The calculation of genus zero quantum K -theoretic invariants is simpler and self-contained, while the higher-genus invariants necessarily involve invariants of lower genus. Genus-0 quantum K -theory is much better understood, with additional *finite difference* structure in addition to the usual D -module structure.

Quantum K -theory has connections with modern enumerative geometry, integrable systems, representation theory, geometric combinatorics and theoretical physics. Its influence on theoretical physics is largely its relation to 3-dimensional topological field theory. See the pioneering works of N. Nekrasov, H. Jockers, P. Mayr etc. [Jockers and Mayr 2019; 2020]. For its connection to representation theory, see [Okounkov 2017]. At the very onset of the quantum K -theory, it was intimately connected to integrable systems. See, for example, [Givental and Lee 2003]. It has also inspired much progress in geometric combinatorics through works like [Buch and Mihalcea 2011; Buch et al. 2013; 2020]. Most of these works are in genus zero. We hope that our algorithm will prove useful in the further development of higher-genus quantum K -theory.

We work exclusively with schemes, stacks, etc. locally of finite-type over the complex numbers \mathbb{C} . In particular, they are locally noetherian.

1. Higher-genus quantum K -invariants

Let C' be a general genus g smooth curve with a general divisor B of degree $d = g + 1$. There is exactly one ramified cover $f : C' \rightarrow \mathbb{P}^1$ of degree d with ramification divisor $B = f^*\infty$ over infinity by Riemann–Roch (see [Costello 2006, Lemma 6.0.1] and [Herr and Wise 2023, Theorem 3.12]). The following facts come from Costello [2006].

- This entails a birational map between moduli spaces (Lemma 1.16).
- By adding stack structure \sim , we can make $f : \widetilde{C}' \rightarrow \widetilde{\mathbb{P}}^1$ a finite étale cover. This is pulled back along a map $\widetilde{\mathbb{P}}^1 \rightarrow BS_d$ from a stacky genus zero curve to the moduli space BS_d of finite étale degree d covers.

We can similarly interpolate between genus g and genus zero maps to a fixed smooth, quasiprojective target X .

Remark 1.1. Write $\langle d \rangle = \{1, 2, \dots, d\}$ for the ordered set of d elements. An S_d -torsor $P \rightarrow X$ is equivalent to the data of a finite étale degree d cover

$$T' := (P \times \langle d \rangle) / S_d \rightarrow X.$$

The universal d -sheeted cover

$$(\text{pt} \times \langle d \rangle) / S_d \rightarrow BS_d$$

can be noncanonically identified with the map $BS_{d-1} \rightarrow BS_d$ induced by any of the d inclusions $S_{d-1} \subseteq S_d$.

Consider a twisted, representable stable map $\tilde{C}' \rightarrow X$ together with a finite étale degree d cover $\tilde{C}' \rightarrow \tilde{C}$ of a stacky curve \tilde{C} of genus zero. To promote \tilde{C}' to a marked curve, we need only order the fibers of the marked points of \tilde{C} .

Our data is pulled back from the universal finite étale degree d cover mapping to X :

$$\begin{array}{ccc} \tilde{C}' & \xrightarrow{\quad \Gamma \quad} & [(X^d \times \langle d \rangle) / S_d] \\ \downarrow & & \downarrow \\ \tilde{C} & \longrightarrow & [\text{Sym}^d X] = [X^d / S_d] \end{array}$$

and the whole diagram has finitely many automorphisms over the right arrow if and only if the map $\tilde{C} \rightarrow [\text{Sym}^d X]$ is stable.

Definition 1.2. The stack $\tilde{\mathcal{K}}_{0,n}([\text{Sym}^d X])$ parameterizes families $\tilde{C} \rightarrow S$ of twisted curves of genus zero with n marked points and a representable map $\tilde{C} \rightarrow [\text{Sym}^d X]$ together with an ordering of the fibers over the marked points. The marked points of \tilde{C} may be nontrivial gerbes over S .

The stack $\tilde{\mathcal{K}}_{0,n}([\text{Sym}^d X])$ equivalently parametrizes families of ramified d -sheeted covers $C' \rightarrow C$ together with maps $C' \rightarrow X$ that have finitely many automorphisms. All ramification points are marked and the fibers above the marked points of C are all the marked points of C' . By “ordering of the fibers”, we mean that the fibers of $C' \rightarrow C$ over each marked point of C must be ordered, a torsor for a product of symmetric groups. We later consider variants where less of the marked points of $C' \rightarrow C$ are ordered; see Figure 2.

Our twisted/stacky stable maps and curves are different from [Costello 2006]. For families of curves $\tilde{C} \rightarrow S$ over a base scheme S , the i -th marked point of \tilde{C} may be a *nontrivial* μ_{r_i} gerbe for $r_i \in \mathbb{Z}_{\geq 1}$. We fix the orders r_i later.

We want to apply the K -theoretic version of Costello’s pushforward formula [Chou et al. 2023, Theorem 2.7] to a square from [Herr and Wise 2023, Section 3.2] introduced in Section 1A:

$$(2) \quad \begin{array}{ccc} \tilde{\mathcal{K}}_{\Xi}(S^d X) & \xrightarrow{q} & \overline{M}_{g,R}(X) \\ \pi' \downarrow & \lrcorner & \downarrow \pi \\ \tilde{\mathfrak{K}}_{\Xi}(BS_d) & \xrightarrow{p} & \mathfrak{M}_{g,R} \end{array}$$

The stacks $\overline{M}_{g,R}(X)$ of stable maps and $\mathfrak{M}_{g,R}$ of prestable curves are standard. We do not fix a curve class β , so this space is a disjoint union over choices of β .

We denote $\tilde{\mathcal{K}}_{\Xi}(S^d X) \subseteq \tilde{\mathcal{K}}_{0,n}([\mathrm{Sym}^d X])$ a substack with appropriate discrete invariants fixed in Section 1B. The stack $\tilde{\mathfrak{K}}_{\Xi}(BS_d)$, denoted $\tilde{\mathfrak{M}}_{0,n}(BS_d)$ in [Herr and Wise 2023, before Lemma 3.6], is approximately the stack of prestable maps $\tilde{C} \rightarrow BS_d$ from the genus zero twisted base curves parameterized in $\tilde{\mathcal{K}}_{\Xi}(S^d X)$.

The obstruction theory for π' is pulled back from π . The problem is that p is of degree

$$e = k!(g!)^{\#J} (g!)^k,$$

while the K -theoretic virtual pushforward formula so far only applies to birational maps. We decompose p as a finite étale torsor of degree e composed with a birational map to which the pushforward formula applies.

1A. Costello's square (2). We describe (2). Write $\langle d \rangle = \{1, 2, \dots, d\}$. A subset $A \subseteq \langle \ell \rangle$ will be fixed later; the symbols $\overline{M}_{g,R}(X)$, $\mathfrak{M}_{g,R}$ refer to moduli stacks of ordinary stable maps and prestable curves with $R = \ell - \#A$ marked points. We do not fix the curve class β for simplicity. We assume $R \geq 1$.

The substack $\tilde{\mathcal{K}}_{\Xi}(S^d X) \subseteq \tilde{\mathcal{K}}_{0,n}([\mathrm{Sym}^d X])$ parametrizes stable maps of genus zero curves to $[\mathrm{Sym}^d X]$, identified with triples $C \leftarrow C' \rightarrow X$ above. The Ξ refers to fixed discrete invariants (see Section 1B): ramification profiles of $C' \rightarrow C$, the numbers n and ℓ of marked points for C and C' , the genus of C' , and the degree d of $C' \rightarrow C$. The number $\ell \leq dn$ is the sum of the degrees of the fibers in the ramification profiles. These invariants satisfy Riemann–Hurwitz to ensure that the space is nonempty:

$$(3) \quad 2g - 2 = -2d + \sum_{P \in C'} (e_P - 1).$$

The functor q forgets the marked points $A \subseteq \langle \ell \rangle$ of C' and then takes the *stabilization* $\overline{C}' \rightarrow X$ of the resulting map $C' \rightarrow X$.

The map π' forgets the stable map to X . To make the diagram commute, π' must remember the stabilization \overline{C}' of $C' \rightarrow X$. Define the stack $\tilde{\mathfrak{K}}_{\Xi}(BS_d)$ of triples $C \leftarrow C' \rightarrow D$, where $C' \rightarrow C$ is a ramified cover of type Ξ and $C' \rightarrow D$ a partial stabilization after forgetting $A \subseteq \langle \ell \rangle$. The map p sends this triple to D . The square (2) is cartesian and p is proper by Lemma 3.9 and Corollary 3.7 of [Herr and Wise 2023], respectively.

Remark 1.3. The degree $e = k!(g!)^{\#J} (g!)^k$ differs from both [Herr and Wise 2023, Theorem 3.12] and [Costello 2006, Lemma 6.0.1]. Our use of nontrivial gerbes instead of trivialized gerbes accounts for the difference from [Herr and Wise 2023]. Costello’s version is reconciled in Remark 3.15 of loc. cit. Our degree can be computed using the proof of Theorem 3.12 of loc. cit. or by taking into account the degrees of the universal gerbes.

The degree e is the order of a group $\Gamma = (S_g)^J \times S_g \wr S_k$ that reorders marked points of $C' \rightarrow C$ discussed in Section 1D.

The stabilization $\bar{C}' \rightarrow X$ was omitted in [Costello 2006], leading to nonproper moduli stacks or noncommutative diagrams. This could be rectified using Costello’s technology of weighted graphs instead of our partial stabilizations.

1B. Specifying Ξ . We unpack our discrete data:

$$\Xi = \left\{ \begin{array}{l} g(C') = g, g(C) = 0, d = g + 1, \\ \langle \ell \rangle \rightarrow \langle n \rangle \text{ is } \langle k \rangle \times \langle d \rangle \xrightarrow{pr_1} \langle k \rangle, J \times \langle d \rangle \xrightarrow{pr_1} J, I \mapsto \infty, \\ \forall j \in J, r_j = 1, \forall j \in \langle k \rangle, r_j = 2, \gamma : I \rightarrow \mathbb{Z}_{\geq 1}, r_\infty = \text{lcm}(\gamma(i)), \\ \bigsqcup_J B\mu_1 = * \rightarrow BS_d, \bigsqcup_{i \in \langle k \rangle} B\mu_2 \xrightarrow{\phi} BS_d. \end{array} \right\}$$

See Figure 1 for an example which is Figure 2 in [Herr and Wise 2023].

Ramification profiles are specified by an action of μ_r on an unordered set of size d . Take a small loop around $p \in C$, and its lifts to C' identify which of the d sheets come together over p . Encode this action in a map $B\mu_r \rightarrow BS_d$ up to isomorphism.

Remark 1.4. The category of maps $BG \rightarrow BH$ has:

- Objects: homomorphisms $f : G \rightarrow H$.
- Morphisms $f_1 \rightarrow f_2$: elements $h \in H$ which conjugate one morphism to another $f_1 = hf_2h^{-1}$. They are all isomorphisms.




Figure 1. A cover in Ξ , $g = 3$, $d = 4$. Marked points are black if forgotten $A \subseteq \langle \ell \rangle$ and white if remembered under the map to $\mathfrak{M}_{g,R}$. The space $\widetilde{\mathfrak{R}}_{\Xi^*}^*(BS_d)$ forgets the ordering on the black marked points.

Objects of the category $\text{Hom}(B\mu_r, BS_d)$ can be identified with actions $\mu_r \circ \langle d \rangle$. Isomorphisms between two such actions are relabelings of the set $\langle d \rangle$ of d elements. So an isomorphism class of functors $B\mu_r \rightarrow BS_d$ is an action of μ_r on an unlabeled set with d elements.

The action $\mu_r \circ \langle d \rangle$ contains the information of a ramification point $C' \rightarrow C$ of a map of curves. The stacky quotient $[\langle d \rangle / \mu_r]$ is the fiber of $\tilde{C}' \rightarrow \tilde{C}$ over the point $B\mu_r \in C$. To extract the set-theoretic fiber, we take the coarse moduli space $\langle d \rangle / \mu_r$. This gives an unlabeled set with some number of elements between 1 and d . In families, $B\mu_r$ is allowed to be a nontrivial gerbe.

A point $p \in C$ is *simply ramified* if its fiber consists of $d - 1$ points, where exactly two of the d sheets come together and the other points in the fiber are unramified. This corresponds to the action $\mu_r \circ \langle d \rangle$ with one 2-cycle and the rest of the points fixed, up to reordering $\langle d \rangle$.

Let $k \geq 0$ be an integer, $g = g(C')$ be the genus of C' , and fix the degree $d = g + 1$. Divide the n marked points of C into three sets:

- $\{\infty\}$: write $I \subseteq C'$ for the fiber over this point $\infty \in C$. This point has ramification described by a map $B\mu_{r_\infty} \rightarrow BS_d$ or function $\gamma : I \rightarrow \mathbb{Z}_{\geq 1}$. That is, $\gamma(i)$ is the size of the stabilizer of i in the corresponding action $\mu_{r_\infty} \circ \langle d \rangle$.
- J : these points $J \subseteq C$ have no ramification.
- $\langle k \rangle$: these points have simple ramification.

This gives partitions

$$\langle n \rangle = J \sqcup \langle k \rangle \sqcup \{\infty\}, \quad \langle \ell \rangle = J \times \langle g + 1 \rangle \sqcup \langle kg \rangle \sqcup I.$$

The map $\langle \ell \rangle \rightarrow \langle n \rangle$ on marked points is compatible with these partitions.

The j -th marked point of C is a μ_{r_j} -gerbe, where $r_j = 1$ at unramified points $j \in J$, $r_j = 2$ for the k simple ramification points and ∞ has $r_\infty = \text{lcm}(\gamma(i))$ the least common multiple of the ramification function γ on I . These data are subject to a constraint easier seen with trivialized gerbes: the sum $B\mathbb{Z} \rightarrow BS_d$ of the classifying space maps from all the composites $\mathbb{Z} \rightarrow \mu_r \rightarrow S_d$ be zero, lest the space be empty. This corresponds to the presentation of $\pi_1(\mathbb{P}^1 \setminus \langle n \rangle)$ via generators whose product is trivial.

Let $A \subseteq \langle \ell \rangle$ consist of all of $\langle kg \rangle$, none of I , and a subset of $J \times \langle g + 1 \rangle$ such that $J \times \langle g + 1 \rangle \setminus A \rightarrow J$ is a bijection. Note that the set $\langle \ell \rangle \setminus A = I \sqcup J$ has R elements.

Take $k = \#I + 3g - 1$ so that all the dimensions agree [Herr and Wise 2023, Theorem 3.12]:

$$\dim \tilde{\mathfrak{K}}_{\Xi}(BS_d) = \dim \mathfrak{M}_{g,R}.$$

We can now prove our main equality between virtual fundamental classes. We first recall their definition in K -theory.

1C. K -theory. Let Y be a finite-type noetherian algebraic stack. The K -theory of Y is the K -theory of a category of lis-ét sheaves on Y , for which there are two main options:

- $K_\circ(Y)$: coherent sheaves, otherwise known as G -theory $G(Y)$.
- $K^\circ(Y)$: locally free sheaves of finite rank.

We work with \mathbb{Q} -coefficients, tensoring these groups up to \mathbb{Q} -vector spaces

$$K_\circ(Y) = K_\circ(Y) \otimes \mathbb{Q}, \quad K^\circ(Y) = K^\circ(Y) \otimes \mathbb{Q}.$$

These groups are generated by classes $[F]$ of coherent/locally free lis-ét sheaves F on Y , modulo relations $[F'] + [F''] = [F]$ for each exact sequence

$$0 \rightarrow F' \rightarrow F \rightarrow F'' \rightarrow 0.$$

See [Chou et al. 2023, Section 1] for discussion.

The groups K° , K_\circ coincide on Y whenever every coherent sheaf F admits a finite resolution by locally free sheaves. Under certain hypotheses on Y , this is equivalent to Y being a quotient stack [Edidin et al. 2001, Remark 2.15].

Let $f : X \rightarrow Y$ be a map between finite-type noetherian algebraic stacks. Pullback and pushforward of sheaves sometimes induce maps on K° and K_\circ .

- K° : pullback f^* always exists and pushforward f_* makes sense when $X \rightarrow Y$ is finite étale.
- K_\circ : pullback f^* exists when f is flat. Armed with a perfect obstruction theory, we can also define a pullback $f^!$ even if f is not flat.

If f is proper and of DM-type, define the pushforward f_* on K_\circ -theory as the alternating sum

$$(4) \quad f_* F := \sum_i (-1)^i R^i f_* F.$$

We must check that this sum is finite.

When the map $f : X \rightarrow Y$ is clear from context, we write $\beta|_X = f^* \beta$ for classes $\beta \in K^\circ(Y)$ or $\beta \in K_\circ(Y)$ without risk of confusion.

Lemma 1.5. *Let $p : X \rightarrow Y$ be a proper, DM-type morphism between finite-type noetherian algebraic stacks. The pushforward*

$$p_* : K_\circ(X) \rightarrow K_\circ(Y).$$

of (4) is well defined on K_\circ -theory.

Proof. We argue that the sum (4) is finite. Write \bar{X} for the relative coarse moduli space of the map p and N for a number larger than the dimensions of the fibers of $\bar{X} \rightarrow Y$. This is possible using quasicompactness of Y .

We claim $R^i p_* F$ vanishes for $i > N$ for any coherent sheaf F . The claim is étale local in Y and so is the formation of the relative coarse moduli space \bar{X} , so we can assume Y is an affine scheme. The map $t : X \rightarrow \bar{X}$ is finite flat, so pushforward is exact $Rt_* = t_*$. This reduces to the representable case $\bar{X} \rightarrow Y$. Because Y is affine, it results from dimensional vanishing [Stacks 2005–, 0A4R]. \square

Pushforward from a proper DM stack to a point is denoted χ .

Example 1.6. If the morphism $p : X \rightarrow Y$ is not of DM-type, the pushforward need not be well defined. Take $B\mathbb{G}_m \rightarrow \text{pt}$. The cohomology of $B\mathbb{G}_m$ is freely generated as a ring by the first Chern class of the universal line bundle and does not vanish in any degree.

Example 1.7. Let G be a finite group. Sheaf pushforward along $p : BG \rightarrow \text{pt}$ sends a complex G -representation V to the invariant subspace V^G . The pushforward is then the alternating sum of group cohomology

$$\chi(V) = \sum (-1)^i [H^i(G, V)].$$

Because we work over \mathbb{C} , the structure sheaf of pt is $\mathcal{O}_{\text{pt}} = \mathbb{C}$. Likewise G -representations V on $BG = [\text{Spec } \mathbb{C}/G]$ are complex representations, and the order of the group $\#G$ is invertible in V . The group cohomology therefore vanishes:

$$H^i(G, V) = 0, \quad i \neq 0.$$

The alternating sum is just the first term $\chi(V) = [V^G]$.

The projection formula holds in both K° and K_\circ , where defined

$$(5) \quad f_*(\alpha \otimes f^*\beta) = f_*\alpha \otimes \beta.$$

This results from the formula on the level of sheaves [Stacks 2005–, 08EU].

The main K -theory classes we are interested in are the fundamental class $[\mathcal{O}_X] \in K_\circ(X)$ and the *virtual fundamental class* (also known as *virtual structure sheaf*) [Lee 2004, Section 2.3; Qu 2018, Definition 2.2; Chou et al. 2023, Definition 1.2]. Consider a map $f : X \rightarrow M$ from a DM stack X to a smooth stack M endowed with a perfect obstruction theory $C_{X/M} \subseteq E$. The virtual fundamental class $[\mathcal{O}_X^{\text{vir}}]$ is the image of the structure sheaf of the normal cone $[\mathcal{O}_{C_{X/M}}]$ under the isomorphism [Chou et al. 2023, Remark 1.6]

$$[\mathcal{O}_X^{\text{vir}}] = \sigma^*[\mathcal{O}_{C_{X/M}}], \quad \sigma^* : K_\circ(E) \simeq K_\circ(X).$$

Example 1.8. Let $\pi : Y = BG \times X \rightarrow X$ be a trivial gerbe for a finite group G . Suppose X has a perfect obstruction theory over some M and Y is given the induced perfect obstruction theory. Then the virtual fundamental class pulls back $\pi^*[\mathcal{O}_X^{\text{vir}}] = [\mathcal{O}_Y^{\text{vir}}]$.

Example 1.7 describes π_* as taking G -invariants of a representation. Then we have $\pi_* \pi^*[\mathcal{O}_X] = [\mathcal{O}_X]$. Using the projection formula, this implies that the virtual fundamental class also pushes forward:

$$\pi_*[\mathcal{O}_Y^{\text{vir}}] = \pi_*(\pi^*[\mathcal{O}_X^{\text{vir}}] \otimes [\mathcal{O}_Y]) = [\mathcal{O}_X^{\text{vir}}] \otimes \pi_* \pi^*[\mathcal{O}_X] = [\mathcal{O}_X^{\text{vir}}].$$

Proposition 1.9. *Let $\pi : \mathcal{G} \rightarrow X$ be a gerbe banded by a finite group G . The base X is a scheme or algebraic stack which we emphasize lies over \mathbb{C} . Then the structure sheaf pushes forward to the structure sheaf, both as sheaves and in K -theory:*

$$R\pi_* \mathcal{O}_{\mathcal{G}} = \pi_* \mathcal{O}_{\mathcal{G}} = \mathcal{O}_X, \quad \pi_*[\mathcal{O}_{\mathcal{G}}] = [\mathcal{O}_X] \in K_0(X).$$

The same holds for virtual fundamental classes if \mathcal{G} is given the induced perfect obstruction theory from X :

$$\pi_*[\mathcal{G}]^{\text{vir}} = [X]^{\text{vir}} \quad \text{in } K_0(X).$$

Proof. The statement on sheaves implies that on K -theoretic classes and is local in X . We can then assume that \mathcal{G} is trivial, fitting in a pullback square:

$$\begin{array}{ccc} \mathcal{G} & \longrightarrow & BG \\ \downarrow & \lrcorner & \downarrow \\ X & \longrightarrow & \text{pt} \end{array}$$

Example 1.7 covers the case of $BG \rightarrow \text{pt}$, and the general case results from cohomology and base change applied to this square.

The statement on virtual fundamental classes results from Example 1.8. \square

Remark 1.10. The proof of Proposition 1.9 does not work for schemes over \mathbb{Z} . The groups $H^i(G, V)$ for $i \neq 0$ are torsion, and so are the sheaves $R^i \pi_* V$ for any coherent sheaf on \mathcal{G} . But this does not mean they vanish in $K_0 \otimes \mathbb{Q}$. Tensoring $-\otimes \mathbb{Q}$ kills K -theoretic classes that are torsion in the group law on K -theory, not the classes of sheaves that themselves are torsion.

The trivial gerbe $[\text{Spec } \mathbb{Z}/G] \rightarrow \text{Spec } \mathbb{Z}$ satisfies Example 1.7 and Proposition 1.9, because the classes of torsion groups vanish in the K -theory of the integers. But this statement does not localize.

We need two related theorems on the behavior of (virtual) fundamental classes under pushforward. These extend Hironaka's theorem and Costello's theorem, respectively.

Theorem 1.11 (Hironaka's pushforward theorem [Chou et al. 2023, Proposition 2.3]). *Let $p : X \rightarrow Y$ be a proper birational map of smooth DM stacks. The pushforward of the fundamental class of X is that of Y in K -theory:*

$$p_*[\mathcal{O}_X] = [\mathcal{O}_Y] \quad \text{in } K_0(Y).$$

Our Costello-type pushforward theorem was originally in the more general context of log geometry. We remove log structures in our citation for simplicity.

Theorem 1.12 (Costello’s pushforward theorem [Chou et al. 2023, Theorem 2.7]). *Consider a pullback square of algebraic stacks*

$$\begin{array}{ccc} X & \xrightarrow{p} & Y \\ \downarrow & \lrcorner & \downarrow \\ M & \xrightarrow{q} & N \end{array}$$

with X, Y DM stacks and M, N smooth. Suppose $Y \rightarrow N$ is equipped with a perfect obstruction theory and $X \rightarrow M$ is given the pullback perfect obstruction theory. If the map q is proper birational, the pushforward of the virtual class of X is that of Y

$$p_*[\mathcal{O}_X^{\text{vir}}] = [\mathcal{O}_Y^{\text{vir}}] \quad \text{in } K_0(Y).$$

To use these theorems, it is important that the relevant maps are proper and birational. Birational maps $f : X \rightarrow Y$ of stacks must have an open dense subset of each X and Y that are isomorphic.

Remark 1.13. For stacks, *pure degree one* [Herr and Wise 2023, Definition 2.3] and birational are not the same. The map $p : B\mathbb{Z}/2 \sqcup B\mathbb{Z}/2 \rightarrow \text{pt}$ is pure degree one but not birational. The pushforward of the fundamental class is not the fundamental class:

$$p_*[\mathcal{O}_{B\mathbb{Z}/2 \sqcup B\mathbb{Z}/2}] = 2 \cdot [\mathcal{O}_{\text{pt}}] \quad \text{in } K_0(\text{pt}).$$

If a morphism of schemes is of pure degree one, it is birational. More generally, if $X \rightarrow Y$ is a morphism of stacks of pure degree one inducing a *representable* morphism $U \rightarrow V$ on open dense substacks $U \subseteq X, V \subseteq Y$, it is birational.

1D. An intermediary stack. The moduli stack $\tilde{\mathfrak{K}}_{\Xi}(BS_d)$ parametrizes a triple $C \leftarrow C' \rightarrow D$ of curves over any base S . We introduce a variant $\tilde{\mathfrak{K}}_{\Xi}^*(BS_d)$ to describe how virtual classes push forward in Proposition 1.17.

The functor $p : \tilde{\mathfrak{K}}_{\Xi}(BS_d) \rightarrow \mathfrak{M}_{g,n}$ sends such a triple to D . The map p is proper of degree $e = k!(g!)^{\#J}(g!)^k$ [Herr and Wise 2023, Section 3]. This map forgets everything about $C' \rightarrow C$, including the ordering of the forgotten marked points under $C' \rightarrow D$.

Let $A \subseteq \langle \ell \rangle$ be the points forgotten under p as in Figure 1. Let $\tilde{\mathfrak{K}}_{\Xi}^*(BS_d)$ be the space similar to $\tilde{\mathfrak{K}}_{\Xi}(BS_d)$, but where the marked points $A \subseteq \langle \ell \rangle$ are *unordered*. The forgetful map $\tilde{\mathfrak{K}}_{\Xi}(BS_d) \rightarrow \tilde{\mathfrak{K}}_{\Xi}^*(BS_d)$ is a torsor under

$$\Gamma := (S_g)^J \times S_g \wr S_k.$$

The group $S_g \wr S_k := S_k \ltimes (S_g)^k$ is the wreath product. There is a short exact sequence

$$1 \rightarrow S_g^k \rightarrow S_g \wr S_k \rightarrow S_k \rightarrow 1$$

and a section $S_k \dashrightarrow S_g \wr S_k$ of the quotient. We may view $S_g \wr S_k$ as a subgroup of S_{gk} by choosing an identification of $\langle gk \rangle$ with $\langle g \rangle \times \langle k \rangle$.

The k copies of S_g reorder the unramified points in the fibers with simple ramification points, while S_k reorders the fibers themselves and their images $\langle k \rangle \subseteq C$.

Example 1.14. Isomorphisms of curves in $\tilde{\mathcal{R}}_{\Xi^*}^*(BS_d)$ need not stabilize the unordered marked points. For example, \mathbb{P}^1 with three unordered points has automorphism group S_3 by interchanging the points $0, 1, \infty$. The moduli space of genus zero curves with three unordered points is then BS_3 .

These choices of ordering certain marked points can also be made on the moduli of stable maps to the stack $[\mathrm{Sym}^d X]$.

Definition 1.15. Let $\tilde{\mathcal{K}}_{\Xi}(S^d X) \subseteq \tilde{\mathcal{K}}_{0,n}([\mathrm{Sym}^d X])$ be the moduli space of representable stable maps to $[\mathrm{Sym}^d X]$ with discrete invariants Ξ . This parameterizes étale, d -sheeted covers $\tilde{C}' \rightarrow \tilde{C}$ with minimal stack structure together with stable maps $\tilde{C}' \rightarrow X$. The curves may have nontrivial gerbes at marked points. All the marked points of C' and C are ordered.

Define $\mathcal{K}_{\Xi}^*(S^d X)$ analogously to $\tilde{\mathcal{R}}_{\Xi^*}^*(BS_d)$ by forgetting the ordering on the marked points of C' corresponding to $A \subseteq \langle \ell \rangle$ (see Figure 2).

$$\begin{array}{ccccc}
 & & \tilde{\mathcal{K}}_{\Xi}(S^d X) & & \\
 & \swarrow / & \downarrow & \searrow \Gamma & \\
 \tilde{\mathcal{M}}_{\Xi}(S^d X) & & & & \mathcal{K}_{\Xi}^*(S^d X) \longrightarrow \bar{\mathcal{M}}_{g,R}(X) \\
 & \searrow \psi & \downarrow / \tilde{\Gamma} & \swarrow \phi & \\
 & & \bar{\mathcal{K}}_{\Xi}(S^d X) & & \\
 & \swarrow / S_k & & \nwarrow (d:1)^J \times / S_{\#I} &
 \end{array}$$

Figure 2. The stacks of stable maps to a fixed target X . $\bar{\mathcal{M}}_{g,R}(X)$ is the ordinary space of stable maps to X . The rest are spaces of stacky genus zero maps $\tilde{C} \rightarrow [\mathrm{Sym}^d X]$. These can be interpreted as ramified finite covers $C' \rightarrow C$ of nonstacky curves together with a map $C' \rightarrow X$ satisfying a stability condition. The difference between $\tilde{\mathcal{M}}_{\Xi}(S^d X)$, $\tilde{\mathcal{K}}_{\Xi}(S^d X)$, $\mathcal{K}_{\Xi}^*(S^d X)$, $\bar{\mathcal{K}}_{\Xi}(S^d X)$ lies in which points of C' , C are ordered. The maps “ $/G$ ” between them are quotients by various groups G reordering the marked points. The one exception is $\mathcal{K}_{\Xi}^*(S^d X) \rightarrow \bar{\mathcal{K}}_{\Xi}(S^d X)$, which is a quotient followed by a $d^{\#J}$ -sheeted cover (denoted $(d:1)^J$).

Extend (2) to the cartesian diagram

$$(6) \quad \begin{array}{ccccc} & & q & & \\ & & \curvearrowright & & \\ \tilde{\mathcal{K}}_{\Xi}(S^d X) & \xrightarrow{v} & \mathcal{K}_{\Xi}^*(S^d X) & \xrightarrow{w} & \bar{M}_{g,R}(X) \\ \downarrow \pi' & \lrcorner & \downarrow & \lrcorner & \downarrow \pi \\ \tilde{\mathfrak{K}}_{\Xi}(BS_d) & \xrightarrow{v} & \tilde{\mathfrak{K}}_{\Xi}^*(BS_d) & \xrightarrow{\omega} & \mathfrak{M}_{g,R} \\ & & p & & \end{array}$$

Lemma 1.16. *The map $\omega : \tilde{\mathfrak{K}}_{\Xi}^*(BS_d) \rightarrow \mathfrak{M}_{g,R}$ is proper and birational.*

Proof. Fix a generic smooth D and prescribed ramification divisor $B = \sum_{i \in I} d(i)[i]$ over ∞ specified by Ξ . The proof of [Herr and Wise 2023, Theorem 3.12] shows that there is exactly one cover $C' \rightarrow C$ with ramification in B and $C' \rightarrow D$ a partial stabilization. The map is thus proper and of pure degree one, but this is not yet sufficient by Remark 1.13.

We argue that ω is generically representable, hence birational. The proof of [Herr and Wise 2023, Theorem 3.12] shows that if $D \in \mathfrak{M}_{g,R}(X)$ is general, the preimage under ω is exactly one cover $C' \rightarrow C$ with $C' = D$. Consider automorphisms

$$\begin{array}{ccc} C' & \xrightarrow{\sim} & C' \\ \downarrow & & \downarrow \\ C & \xrightarrow{\sim} & C \end{array}$$

of the map $C' \rightarrow C$. These form a subgroup of automorphisms of C' because $C' \rightarrow C$ is an epimorphism. Since the map $\underline{\text{Aut}}(C' \rightarrow C) \rightarrow \underline{\text{Aut}}(D)$ is injective, the map is generically representable and hence birational. \square

Hironaka's pushforward theorem (see Theorem 1.11) equates their fundamental classes:

$$w_*[\mathcal{O}_{\tilde{\mathfrak{K}}_{\Xi}^*(BS_d)}] = [\mathcal{O}_{\mathfrak{M}_{g,n}}] \quad \text{in } K_{\circ}(\mathfrak{M}_{g,n}).$$

Costello's pushforward theorem (see Theorem 1.12) likewise equates the virtual fundamental classes:

Proposition 1.17. *The fundamental class pushes forward along the map w in (6):*

$$w_*[\mathcal{O}_{\tilde{\mathfrak{K}}_{\Xi}^*(BS_d)}] = [\mathcal{O}_{\mathfrak{M}_{g,n}}] \quad \text{in } K_{\circ}(\mathfrak{M}_{g,n}).$$

As a result, the virtual fundamental class pushes forward the same way:

$$w_*[\mathcal{O}_{\tilde{\mathcal{K}}_{\Xi}(S^d X)}^{\text{vir}}] = [\mathcal{O}_{\bar{M}_{g,R}(X)}^{\text{vir}}] \quad \text{in } K_{\circ}(\bar{M}_{g,R}(X)).$$

The wreath product $S_g \wr S_k$ arises naturally as the automorphism group of the projection $\langle g \rangle \times \langle k \rangle \rightarrow \langle k \rangle$ of marked points of $C' \rightarrow C$:

Remark 1.18 (thanks to J. Rufus Lawrence). The iterated stack-theoretic symmetric product $[\mathrm{Sym}^k[\mathrm{Sym}^g X]]$ is isomorphic to the global quotient

$$[\mathrm{Sym}^k[\mathrm{Sym}^g X]] \simeq [X^{gk}/(S_g \wr S_k)].$$

To make sense of $[\mathrm{Sym}^k-]$ applied to a stack \mathcal{S} , use its functor of points

$$[\mathrm{Sym}^k \mathcal{S}](T) = \left\{ \begin{array}{ccc} T' & \longrightarrow & \mathcal{S} \\ k:1 \downarrow & & \\ T & & \end{array} \right\},$$

where $T' \rightarrow T$ is a k -sheeted cover that is part of the moduli.

Consider the evaluation map corresponding to a point $C' \rightarrow C$ of $\mathcal{K}_{\Xi}^*(S^d X)$ over a base T . Upon ordering the k -marked points i of C , we get a map to $[\mathrm{Sym}^g X]$ corresponding to each i . Ordering the k -marked points of C entails a k -sheeted cover of T with a map to $[\mathrm{Sym}^g X]$. These evaluation maps are precisely

$$\mathcal{K}_{\Xi}^*(S^d X) \rightarrow [\mathrm{Sym}^k[\mathrm{Sym}^g X]] = [X^{gk}/(S_g \wr S_k)].$$

Two more stacks $\tilde{M}_{\Xi}(S^d X)$, $\bar{\mathcal{K}}_{\Xi}(S^d X)$. The stack $\tilde{\mathcal{K}}_{\Xi}(S^d X)$ is the simplest because all the marked points of C' and C are ordered, but we will not actually use it for our theorem. The variant $\mathcal{K}_{\Xi}^*(S^d X)$ above is virtually birational to $\bar{M}_{g,R}(X)$. We need two more variants, completing Figure 2.

Definition 1.19. Let $\tilde{M}_{\Xi}(S^d X)$ be the moduli space of representable twisted stable maps $C \rightarrow [\mathrm{Sym}^d X]$. It is the same as $\tilde{\mathcal{K}}_{\Xi}(S^d X)$, except the marked points of C' are not ordered. The only difference from twisted stable maps $C \rightarrow [\mathrm{Sym}^d X]$ in the literature is the nontrivial gerbes.

The quotient maps $\tilde{\mathcal{K}}_{\Xi}(S^d X) \rightarrow \tilde{M}_{\Xi}(S^d X)$, $\tilde{\mathcal{K}}_{\Xi}(S^d X) \rightarrow \mathcal{K}_{\Xi}^*(S^d X)$ forget different marked points, so there is not a map between them. Define $\bar{\mathcal{K}}_{\Xi}(S^d X)$ to forget all the marked points of both, so only the points of C that are not in $\langle k \rangle \subseteq C$ are ordered.

As shown in Figure 2, there are quotient maps from $\tilde{\mathcal{K}}_{\Xi}(S^d X)$ to all the others $\tilde{M}_{\Xi}(S^d X)$, $\bar{\mathcal{K}}_{\Xi}(S^d X)$, $\mathcal{K}_{\Xi}^*(S^d X)$ by various groups reordering marked points. The map $\tilde{M}_{\Xi}(S^d X) \rightarrow \bar{\mathcal{K}}_{\Xi}(S^d X)$ quotients by S_k , while $\mathcal{K}_{\Xi}^*(S^d X) \rightarrow \bar{\mathcal{K}}_{\Xi}(S^d X)$ is a composite of many d -sheeted covers indexed by J and a quotient by $S_{\#I}$.

Remark 1.20. Remark that $R = 1$ when $\#I = 1$ and $J = \emptyset$. In that case, $k = 3g$. The map $\mathcal{K}_{\Xi}^*(S^d X) \rightarrow \bar{\mathcal{K}}$ is an isomorphism precisely when one of the conditions holds:

- $R = 1$.
- $\#I = 1$ and $g = 0$.

See Figure 2.

Remark 1.21. The finite étale maps

$$\begin{array}{ccc} \tilde{\mathcal{K}}_{\Xi}(S^d X) & \xrightarrow{v} & \mathcal{K}_{\Xi}^*(S^d X) \\ \downarrow & & \downarrow \phi \\ \tilde{M}_{\Xi}(S^d X) & \xrightarrow{\psi} & \bar{\mathcal{K}}_{\Xi}(S^d X) \end{array}$$

in Figure 2 all equate virtual fundamental classes under pullback:

$$v^*[\mathcal{O}_{\mathcal{K}_{\Xi}^*(S^d X)}^{\text{vir}}] = [\mathcal{O}_{\tilde{\mathcal{K}}_{\Xi}(S^d X)}^{\text{vir}}], \text{ etc.}$$

1E. Gromov–Witten invariants in the K_{\circ} -theory of stacks. Quantum K_{\circ} -theoretic invariants have been defined variously in the literature [Tonita and Tseng 2013] [Zhang, Section 2.4]. Our definition parallels [Abramovich et al. 2008], adding in ψ classes. Our invariants differ by a scaling factor due to conventions over whether gerbes at marked points are trivialized; see [Tonita and Tseng 2013; Zhang, Remark 2.8] or the original [Abramovich et al. 2002, Sections 4.4, 4.5] for comparison. We allow nontrivial gerbes.

For any moduli space \mathcal{K} of stable maps $C \rightarrow Y$ from n -pointed curves, there is an evaluation map

$$\text{ev} : \mathcal{K} \rightarrow Y^n.$$

If Y is a stack and we take representable twisted stable maps $\tilde{C} \rightarrow Y$ in say $\mathcal{K} = \bar{M}_{g,n}(Y)$, the “points” of \tilde{C} are not quite points, but μ_r -banded gerbes. The evaluation map doesn’t produce points of Y , but cyclic gerbes mapping representably to Y . Cyclic gerbes representably embedded in Y form the *rigidified cyclotomic inertia* stack $\bar{I}(Y)$ [Abramovich et al. 2008, Section 3], so the evaluation map is

$$\text{ev} : \mathcal{K} \rightarrow (\bar{I}(Y))^n.$$

The stack $\bar{I}(Y)$ is closely related to the inertia stack IY . The universal gerbe over $\bar{I}(Y)$ can be identified with representable maps from the trivial gerbe

$$\text{Hom}^{\text{rep}}(B\mu_r, Y).$$

Fixing an isomorphism $\mu_r \simeq \mathbb{Z}/r$ over \mathbb{C} , we get a map to the inertia stack

$$\begin{array}{ccc} \text{Hom}^{\text{rep}}(B\mu_r, Y) & \longrightarrow & IY \\ \downarrow & & \\ \bar{I}(Y) & & \end{array}$$

by composing $B\mathbb{Z} \rightarrow B\mu_r \rightarrow Y$. See Section 3C for a worked example.

Instead of pulling back K_{\circ} -theoretic classes from Y , we pull back from $K_{\circ}(\bar{I}(Y))$. In our case, $Y = [\text{Sym}^d X]$ and our evaluation map is

$$\text{ev}_{C,\infty} : \tilde{\mathcal{K}}_{\Xi}(S^d X) \rightarrow \bar{I}([\text{Sym}^d X]).$$

Now we add in ψ classes. Write $\mathcal{U} \rightarrow \mathfrak{M}_{g,n}$ for the universal curve and L_i for the conormal bundle at the i -th marked section σ_i :

$$L_i := N_{\sigma_i} \simeq \sigma_i^* T_{\mathcal{U}/\mathfrak{M}_{g,n}}.$$

Use the same notation for their pullback to any moduli space with prestable curves, for example, $\mathcal{K}_{g,n}(Y)$, $\overline{\mathcal{M}}_{g,n}$. The classes of $L_i \in K^\circ(\mathfrak{M}_{g,n})$ and their pullbacks to various moduli stacks are referred to as ψ classes.

Beware that maps between moduli spaces involving stabilization do not have the same ψ classes. We compare the ψ class of a map $C \rightarrow X$ with the stabilization C^{st} in Section 2.

The stack $\mathcal{K}_{g,n}(Y)$ supports an obstruction theory relative to $\mathfrak{M}_{g,n}$ that lets us define virtual fundamental classes $\mathcal{O}_{\mathcal{K}_{g,n}(Y)}^{\text{vir}}$ in $K_\circ(\mathcal{K}_{g,n}(Y))$. Classes

$$\alpha_1, \dots, \alpha_n \quad \text{in } K_\circ(\overline{\mathcal{I}}(Y))$$

and exponents e_1, \dots, e_n give rise to a (descendent) *Gromov–Witten invariant*

$$\langle \alpha_1 L_1^{e_1}, \dots, \alpha_n L_n^{e_n} \rangle = \chi \left(\mathcal{O}_{\mathcal{K}_{g,n}(Y)}^{\text{vir}} \otimes \prod ev_i^* \alpha_i \otimes L_i^{e_i} \right) \quad \text{in } \mathbb{Q}.$$

We are equally interested in power series of these invariants.

When $Y = [\text{Sym}^d X]$, there are two twisted curves $\tilde{\mathcal{C}}' \rightarrow \tilde{\mathcal{C}}$ and hence two evaluation maps and two sets of ψ classes. The ψ classes of $\tilde{\mathcal{C}}'$ are the same as the marked points of $\tilde{\mathcal{C}}$ below. The main technical problem in Theorem 3.1 will be converting between classes pulled back along the evaluation map of $\tilde{\mathcal{C}}'$ and that of $\tilde{\mathcal{C}}$. Genuine Gromov–Witten invariants have classes $ev^* \alpha$ pulled back from the evaluation map of $\tilde{\mathcal{C}}$, not that of $\tilde{\mathcal{C}}'$. This convention parrots [Abramovich et al. 2008].

1F. Permutation-equivariant K -theory. The K -theory of $\overline{\mathcal{K}}_\Xi(S^d X)$ is equivalent to permutation-equivariant K -theory of $\tilde{\mathcal{M}}_\Xi(S^d X)$, as in [Givental 2017]. Ordinary quantum K -theory entails “correlators” defined as the integrals:

$$\langle \alpha_1 L_1^{m_1}, \dots, \alpha_n L_n^{m_n} \rangle := \chi \left(\mathcal{O}_{\overline{\mathcal{K}}_\Xi(S^d X)}^{\text{vir}} \otimes \prod ev_i^* (\alpha_i) L_i^{m_i} \right) \quad \text{in } K_\circ(\text{pt}) = \mathbb{Q}.$$

How can we compute Euler characteristics of $\overline{\mathcal{K}}_\Xi(S^d X)$ by working on $\tilde{\mathcal{M}}_\Xi(S^d X)$? Take the pullback square

$$\begin{array}{ccc} \tilde{\mathcal{M}}_\Xi(S^d X) & \longrightarrow & \text{pt} \\ \psi \downarrow & \lrcorner & \downarrow \\ \overline{\mathcal{K}}_\Xi(S^d X) & \longrightarrow & BS_k \xrightarrow{-S_k} \text{pt} \end{array}$$

Denote pushforward along $\overline{\mathcal{K}}_\Xi(S^d X) \rightarrow BS_k$ by $\chi_{S_k}(-)$. This remembers the S_k -representation on the virtual vector space $\chi(\psi^*(-))$. To get the ordinary Euler

characteristic $\chi(-)$ on $\bar{\mathcal{K}}_{\Xi}(S^d X)$, we have to take the virtual S_k -invariants of the virtual representation $\chi_{S_k}(-)$:

$$\chi(-) = (\chi_{S_k}(-))^{S_k}.$$

These should be derived invariants, but we work over \mathbb{Q} . The higher-group cohomologies of S_k valued in a representation all vanish, so the distinction is moot.

The class $\psi^*(-)$ on $\tilde{M}_{\Xi}(S^d X)$ will be a Gromov–Witten invariant where the insertions at the marked points forgotten under ψ are identical. In our case, we only care about insertions away from those forgotten marked points. We will only integrate classes at ordered marked points of both $\bar{\mathcal{K}}_{\Xi}(S^d X)$ and $\tilde{M}_{\Xi}(S^d X)$.

Even if we insert away from the permuted points $A \subseteq \langle \ell \rangle$, the action of Γ still nontrivially permutes the sections:

Example 1.22. Consider S_3 acting on $\bar{M}_{0,4} = \mathbb{P}^1$ by permuting the last three marked points. The generating function of the S_3 -invariant quantum K -invariants on $\bar{M}_{0,4}$ with only L_1 can be calculated

$$\chi\left([\bar{M}_{0,4}/S_3], \frac{1}{1 - q_1 L_1}\right) = \frac{1}{(1 - q_1^2)(1 - q_1^3)}.$$

Indeed, $L_1 = \mathcal{O}(1)$, $\bar{M}_{0,4} = \mathbb{P}^1$, and $H^{\geq 1}(\bar{M}_{0,4}, L_1^d) = 0$ for all $d \geq 0$. The case $d = 1$ has the sections the linear functions on \mathbb{P}^1 , which are never S_3 -invariant (up to Möbius transformations). It is easy to see that $d = 2$ and $d = 3$ have invariant sections. In fact, $[\bar{M}_{0,4}/S_3] = \mathbb{P}(2, 3)$ and the formula follows [Lee and Qu 2014].

1G. Grothendieck–Riemann–Roch (GRR). This expository section explains why Grothendieck–Riemann–Roch (GRR) does not reduce equivariant Euler characteristics to ordinary ones on stacks, the way it would for schemes.

Using Grothendieck–Riemann–Roch for schemes, one would expect an equality

$$[\mathcal{O}_{\bar{\mathcal{K}}_{\Xi}(S^d X)}^{\text{vir}}] \stackrel{?}{=} [\mathcal{O}_{\mathcal{K}_{\Xi}^*(S^d X)}^{\text{vir}}]^{\oplus \#\Gamma} \quad \text{in } K_o(\mathcal{K}_{\Xi}^*(S^d X))_{\mathbb{Q}}.$$

Coupled with the projection formula, this would reduce permutation equivariant integrals to ordinary ones.

For schemes, this holds. Let $\pi : P \rightarrow X$ be a G -torsor with P, X schemes for some finite group G . GRR gives a commutative square

$$\begin{array}{ccc} K_o(P) & \longrightarrow & A_*(P)_{\mathbb{Q}} \\ \pi_* \downarrow & & \downarrow \pi_* \\ K_o(X) & \longrightarrow & A_*(X)_{\mathbb{Q}}. \end{array}$$

The Todd classes cancel out since $P \rightarrow X$ is étale and $T_X|_P = T_P$, so one can take the horizontal arrows as the Chern character isomorphisms. Since the pushforward

in Chow groups gives $\pi_*[P] = \#G \cdot [X]$ and the horizontal isomorphisms send 1 to 1, we have

$$(7) \quad \pi_*[\mathcal{O}_P] = [\mathcal{O}_X^{\oplus \#G}].$$

These formulas do not hold for stacks!

Take $P = \text{pt}$, $X = BG$. The analogous GRR square

$$(8) \quad \begin{array}{ccc} K_\circ(\text{pt}) & \longrightarrow & A_*(\text{pt})_{\mathbb{C}} \\ \downarrow & & \downarrow \\ K_\circ(BG) & \longrightarrow & A_*(BG)_{\mathbb{C}} \end{array}$$

still commutes [Edidin 2013, Section 5], and the Todd class terms vanish. But the lower horizontal arrow is not multiplicative and does not send 1 to 1! One cannot identify $\pi_*\mathcal{O}_{\text{pt}}$ and $\mathcal{O}_{BG}^{\oplus \#G}$.

Example 1.23. Let $G = \mathbb{Z}/2$ and consider the quotient map $\text{pt} \rightarrow BG$. The inertia stack is

$$IBG = BG \sqcup \text{pt},$$

so its rational Chow groups are $A_*(IBG) = \mathbb{C}^{\oplus 2}$. The GRR square for the quotient $\pi : \text{pt} \rightarrow BG$ is then

$$\begin{array}{ccc} K_\circ(\text{pt}) & \xrightarrow{ch(-)} & A_*(\text{pt}) \\ \downarrow \pi_* & & \downarrow \pi_* \\ K_\circ(BG) & \xrightarrow{ch(-)} & A_*(IBG) \simeq \mathbb{C}^{\oplus 2}. \end{array}$$

The Todd classes are trivial here.

The Chern character of the trivial representation \mathcal{O}_{BG} is $(1, 1)$. By GRR [Edidin 2013, Theorem 5.4], the Chern character of $\pi_*\mathcal{O}_{\text{pt}}$ is

$$ch(\pi_*\mathcal{O}_{\text{pt}}) = (2, 0) \quad \text{in } A_*(IBG) \simeq \mathbb{C}^{\oplus 2}.$$

We can see that $\pi_*\mathcal{O}_{\text{pt}} \neq \mathcal{O}_{BG}^2$.

Remark 1.24. One can define a Borel equivariant K_\circ theory for stacks in which formula (7) holds using [Noohi 2012]. One can equip them with virtual fundamental classes and study Borel equivariant quantum K -theory.

The problem with (8) is that $\text{pt} \rightarrow BG$ introduces stack structure. It is representable, but points in BG have more automorphisms than pt does. We want the opposite of representable, that the automorphism groups of points surject. We can prove a version of (7) in this setting [Herr and Lee].

2. Stabilization and ψ classes

2A. Costello's lemma for stabilizing ψ classes. We recall the following three categories defined in [Costello 2006, Section 3]: Γ^u , which contains the label of nodal curves; Γ^t , which contains the label of twisted nodal curves; and Γ^c , which contains labels of twisted marked curves \mathcal{C} and \mathcal{C}' with an étale morphism $\mathcal{C}' \rightarrow \mathcal{C}$. They are related by the diagram

$$\Gamma^c \xrightarrow{s,t} \Gamma^t \xrightarrow{r} \Gamma^u,$$

where r, t stand for source and target of the étale map $\mathcal{C}' \rightarrow \mathcal{C}$ and r maps \mathcal{C} to its coarse moduli space. These categories will depend on a semigroup A . Furthermore, one can relate the functors between graphs into morphisms of stacks of moduli of curves by applying the functor \mathfrak{M} .

Given $\eta \in \Gamma^c$ consider the diagram

$$\mathfrak{M}_\eta \xrightarrow{s} \mathfrak{M}_{s(\eta)} \xrightarrow{r} \mathfrak{M}_{r(s(\eta))} \xrightarrow{\pi} \mathfrak{M}_{v(I)},$$

where $I \subsetneq T(s(\eta))$ is a chosen finite subset such that after removing tails in I , $s(\eta)$ remains stable. Let $v(I)$ be obtained from $r(s(\eta))$ by removing the tails in I and denote the resulting contraction map by π .

To compare the pullback of ψ classes via s, r and π , we define the notation $S(e, t, I)$ as follows: Consider the diagram

$$\mathfrak{M}_\gamma \rightarrow \mathfrak{M}_{r(s(\eta))} \xrightarrow{\pi} \mathfrak{M}_{v(I)},$$

where $\gamma \rightarrow r(s(\eta))$ is a contraction of Γ^u . Now let $t \in T(\gamma) \setminus I$ and $e \in E(\gamma)$. Let

$$S(e, t, I) := \begin{cases} 1 & \text{if } t \text{ is in a vertex of } \gamma_e \text{ contracted after forgetting the tails } I, \\ 0, & \text{otherwise.} \end{cases}$$

Here γ_e is obtained by contracting all edges of γ except e .

Costello described the pullback of ψ classes in Chow groups.

Lemma 2.1 [Costello 2006, Section 4.1]. *For each $t \in T(v(I))$, we have*

$$s^* r^* \pi^*(\psi_t) = m(t)\psi_t - \sum_{f:\gamma \rightarrow \eta} S(f, t, I)[\mathfrak{M}_f],$$

where the sum is over $f : \gamma \rightarrow \eta$ in Γ^c with $\#E(t(\gamma)) = \#E(f) = 1$, $\mathfrak{M}_f \hookrightarrow \mathfrak{M}_\eta$ is the closed substack supported on the image of f , and

$$S(f, t, I) := \sum_{e \in E(s(\gamma))} m(e) S(e, t, I)$$

is the corresponding multiplicity.

A similar result holds in K -theory:

Lemma 2.2. *For each $t \in T(v(I))$, we have*

$$s^*r^*\pi^*(L_t) = L_t^{\otimes m(t)} \otimes \mathcal{O}\left(-\sum_{f:\gamma \rightarrow \eta} S(f, t, I) \mathfrak{M}_f\right),$$

with $f : \gamma \rightarrow \eta$ and $S(f, t, I)$ defined in previous lemma.

We would like to write $\mathcal{O}(k\mathfrak{M}_f)$ in terms of a torsion sheaf.

2B. Explicit formulas for stabilizing ψ classes in K-theory. Before addressing stable maps to a target V , we work on the moduli of curves. Let $\pi : \bar{M}_{g,m+n} \rightarrow \bar{M}_{g,m}$. We introduce some notations below.

Decoration. Decorations index trees of rational curves to be contracted under forgetting points and stabilizing (see Figure 3).

For the special case $m = 1$, we denote a decoration of degree r as

$$\underline{a} = (a_{1,1}, \dots, a_{1,n_1}) \dots (a_{r,1}, \dots, a_{r,n_r}).$$

We further assume that

$$\{a_{1,1}, \dots, a_{1,n_1}, \dots, a_{r,1}, \dots, a_{r,n_r}\} \subset \{2, 3, \dots, n+1\} = [2, n+1]$$

and

$$a_{i,1} < a_{i,2} < \dots < a_{i,n_i} \quad \text{for all } i.$$

For the general case, a corresponding decoration is denoted by

$$\underline{a} = (\underline{a}_1, \underline{a}_2, \dots, \underline{a}_m),$$

where each \underline{a}_i is a decoration in the special case $m = 1$ and their disjoint set union forms a subset of $[m+1, m+n]$. We also denote it by \underline{a} if no confusion may occur.

Degeneration strata. Given $m = 1$ and a decoration \underline{a} , we define the corresponding degeneration strata of codimension r on $\bar{M}_{g,1+n}$ as

$$\bar{M}_{g,1+n} \supset D_{1,\underline{a}} := \left(\bigcap_{1 \leq i \leq r} D_{1a_{1,1} \dots a_{1,n_1} a_{2,1} \dots a_{i,n_i}} \right),$$

where $D_{abcde\dots}$ is the divisor with markings $abcde\dots$ lying on the rational tail. This is the closure of the locus where the curves have rational tails indexed by \underline{a} .

For the general case, given a decoration \underline{a} , we define the corresponding stratum

$$\bar{M}_{g,m+n} \supset D_{(m),\underline{a}} := \bigcap_{1 \leq i \leq m} D_{i,\underline{a}_i}.$$

Normal bundle. Given $m = 1$ and a strata $D_{1,\underline{a}}$ as above, we have

$$\mathcal{O}_{D_{1,\underline{a}}}^2 = \lambda_{-1}(\oplus_{i=1}^r L_{1i}) \mathcal{O}_{D_{1,\underline{a}}}.$$

Here L_{1i} is the normal bundle of the i -th node. More precisely, it can be described as

$$\mathcal{O}_{D_{1,a_{1,1}\dots a_{i,n_i}}}^2 = \lambda_{-1}(\tilde{L}_i) \mathcal{O}_{D_{1,a_{1,1}\dots a_{i,n_i}}}.$$

Then we define

$$L_{1i} := \tilde{L}_i|_{D_{1,\underline{a}}}.$$

For the general case, we denote by L_{ji} the normal bundle of the i -th node of the tail containing the marking j for $1 \leq j \leq m$.

Type. Given $m = 1$ and a decoration \underline{a} or its corresponding strata $D_{A_1\underline{a}}$, we define its *type* to be 0 if

$$a_{1,1} < a_{j,k} \quad \text{for any } (j, k) \neq (1, 1),$$

we define its type to be $\underline{l} = (l_1 \dots l_s)$ if

$$a_{1,1} > a_{2,1} > \dots > a_{l_1-1,1},$$

$$a_{l_1,1} > a_{l_1+1,1} > \dots > a_{l_2-1,1},$$

$$\vdots$$

$$a_{l_s,1} > a_{l_s+1,1} > \dots > a_{r,1}$$

and

$$a_{1,1} > a_{l_1,1} > \dots > a_{l_s,1}.$$

Given a general decoration $\underline{a} = (a_1, \dots, a_m)$, we define its type on each a_i as in the special case and also denote it by \underline{l} if no confusion may occur. We also say it is of type 0 if a_i is of type 0 for all i .

Polynomial corresponding to decoration. Given $m = 1$ and a decoration \underline{a} of type (0), we define its corresponding polynomial with r variables to be

$$F_{1,\underline{a}}(x_1, \dots, x_r) = \left(1 - \prod_{i=1}^r x_i\right).$$

For a partition \underline{a} of type (l_1, \dots, l_s) , we define its corresponding polynomial to be

$$F_{1,\underline{a}}(x_1, \dots, x_r) = \sum_{j=0}^s \left(1 - \prod_{i=l_j}^{l_{j+1}-1} x_i\right).$$

Here we set $l_0 = 1$ and $l_{s+1} = r + 1$.

For the general case, given a decoration $\underline{a} = (a_1, \dots, a_m)$, we define F_{i,a_i} as in the special case for all $i = 1, \dots, m$.




Figure 3. Rational tails corresponding to the decoration $\underline{a} = (\underline{a}_1, \underline{a}_2, \underline{a}_3)$ of type 0, with $\underline{a}_1 = (a_{111})(a_{121})(a_{131}, a_{132}, a_{133}) = (4)(8)(5, 7, 14)$, $\underline{a}_2 = (a_{211})(a_{221}, a_{222}, a_{223})(a_{231}) = (6)(10, 12, 13)(11)$, $\underline{a}_3 = (a_{311}) = (9)$. The smallest numbered marked point on each rational tail (except 1, 2, and 3) must be on the \mathbb{P}^1 farthest from the main component to be type 0.

Difference operator. We define the difference operator δ on a (multi)variable polynomial $F(\underline{x}) = F(x_1, \dots, x_n)$ as

$$\delta(F(\underline{x})) := \frac{F(\underline{x}) - \sum_i F(\underline{x})|_{x_i=1} + \dots + (-1)^n F(\underline{x})|_{x_1=\dots=x_n=1}}{(1-x_1) \cdots (1-x_n)}.$$

Lemma 2.3. *Let $\pi : \bar{M}_{g,m+n} \rightarrow \bar{M}_{g,m}$. Then we have*

$$\pi^*(L_1) = L_1 + \sum_{\underline{a}: \text{type } 0} (-1)^{\text{codim } D_{\underline{a}}} \mathcal{O}_{D_{\underline{a}}} \in K^0(\bar{M}_{g,m+n}).$$

Proof. We decompose π as

$$\bar{M}_{g,m+n} \xrightarrow{\pi_{m+n}} \bar{M}_{g,m+n-1} \rightarrow \dots \rightarrow \bar{M}_{g,m+1} \xrightarrow{\pi_{m+1}} \bar{M}_{g,m}$$

and compute $\pi^*(L_1) = \pi_{m+n}^* \dots \pi_{m+1}^*(L_1)$ step by step.

Given a stratum $D \subset \bar{M}_{g,m+n}$ of type 0, we consider the inclusion-exclusion formula

$$\pi_{m+n}^* \mathcal{O}_{\pi_{m+n}(D)} = \sum_{i=1}^r \mathcal{O}_{D_i} - \sum_{i < j} \mathcal{O}_{D_i \cap D_j} + \dots,$$

where $\pi_{m+n}^*(\pi_{m+n}(D)) = \bigcup_{i=1}^r D_i$ with D_i irreducible strata.

Note that \mathcal{O}_D will show up in either the first or the second term of the right-hand side depending on whether $\text{ft}_{n+m}(D)$ is stable or not. Since $\pi_{n+m}(D)$ is still of type 0. The lemma follows by induction. \square

Remark 2.4. This expression is not symmetric with respect to indices $m + 1, \dots, m + n$ since we chose a special order of pull-backs. Different choices of pull-back order will result in different expressions. Nevertheless, any expression will give the same element in $K^0(\overline{M}_{g,n+m})$.

Lemma 2.5. *Let $\pi : \overline{M}_{g,1+n} \rightarrow \overline{M}_{g,1}$, \underline{a} be a decoration, and G be any power series. Then*

$$\text{Coeff}(\mathcal{O}_{D_{1,\underline{a}}}, \pi^*G(L_1)) = \delta(G(L_1 - F_{1,\underline{a}})),$$

where $F_{1,\underline{a}} = F_{1,\underline{a}}(L_{11}, \dots, L_{1,\text{deg } \underline{a}})$ with L_{1j} defined above. When applying δ -operator on the right-hand side, we view $G(L_1 - F_{1,\underline{a}})$ as power series with variables $\{L_{ij}\}$ and view L_1 as constant.

Proof. We start with a special case $m = 1$ and $n = 2$. By previous lemma, we have

$$\pi^*L_1 = L_1 - \mathcal{O}_{D_{12}} - \mathcal{O}_{D_{13}} - \mathcal{O}_{D_{123}} + \mathcal{O}_{D_{(12)(3)}}.$$

Let $G(x)$ be any power series. To compute $\text{Coeff}(\mathcal{O}_{D_{(12)(3)}}, G(\pi^*L_1))$, we introduce the following process:

Write $x = \mathcal{O}_{D_{12}}$ and $y = \mathcal{O}_{D_{123}}$ and hence $\mathcal{O}_{D_{(12)(3)}} = xy$. Now we have

$$\begin{aligned} & \text{Coeff}(\mathcal{O}_{D_{(12)(3)}}, G(\pi^*L_1)) \\ &= \frac{G(L_1 - x - y + xy) + G(L_1 - y) + G(L_1 - x) + G(L_1)}{xy} \Big|_{x=1-L_{11}, y=1-L_{12}} \\ &= \delta(G(L_1 - (1 - L_{11}L_{12}))). \end{aligned}$$

Some remarks are in order:

- L_{11} and L_{12} are characterized by $x^2 = (1 - L_{11})x$ and $y^2 = (1 - L_{12})y$.
- The second equality above follows from the definition of δ . Here we view $G(L_1 - (1 - L_{11}L_{12}))$ as power series in L_{11} and L_{12} .
- $1 - L_{11}L_{12}$ is exactly the polynomial $F_{1,(12)(3)}(L_{11}, L_{12})$, i.e., the polynomial corresponding to the strata $D_{(12)(3)}$.

For general case, we can compute the coefficient of $D_{1,\underline{a}}$ using the above computation process. It suffices to find the polynomial corresponding to $D_{1,\underline{a}}$. To find the polynomial, we compute

$$- \sum_{\substack{\underline{a}': \text{type } 0 \\ D_{1,\underline{a}'} \supset D_{1,\underline{a}}}} (-1)^{\text{codim } D_{1,\underline{a}'}} \mathcal{O}_{D_{1,\underline{a}'}} \cdot \mathcal{O}_{D_{1,\underline{a}'}} \Big|_{D_{1,\underline{a}}} = F_{1,\underline{a}}(L_{11}, \dots, L_{1 \text{ deg } \underline{a}}) \mathcal{O}_{D_{1,\underline{a}}}.$$

A direct computation shows that it is exactly the polynomial we defined above. \square

Theorem 2.6. *Let $\pi : \bar{M}_{g,m+n} \rightarrow \bar{M}_{g,m}$ and $G = G(x_1, \dots, x_m)$ be any power series of m variables. Then*

$$\pi^* G(L_1, \dots, L_m) = G(L_1, \dots, L_m) + \sum_{\underline{a}: \text{all decorations}} \mathcal{O}_{D_{(m),\underline{a}}} \delta(G(L_1 - F_{1,\underline{a}_1}, \dots, L_m - F_{m,\underline{a}_m})),$$

where $F_{i,\underline{a}_i} = F_{i,\underline{a}_i}(L_{i1}, \dots, L_{i \deg \underline{a}_i})$ with L_{ij} defined above. When applying the δ -operator on the right-hand side, we view

$$G(L_1 - F_{1,\underline{a}_1}, \dots, L_m - F_{m,\underline{a}_m})$$

as power series with variables $\{L_{ij}\}$ and view L_i as constant.

Proof. It follows from the definition that

$$D_{(m),\underline{a}} := \bigcap_{1 \leq i \leq m} D_{i,\underline{a}_i},$$

and the following observation that if $F(\underline{x}_1, \dots, \underline{x}_r) = \prod_{i=1}^r F_i(\underline{x}_i)$, then

$$\delta(F(\underline{x}_1, \dots, \underline{x}_r)) = \prod_{i=1}^r \delta(F_i(\underline{x}_i)).$$

Here \underline{x}_i could be multiindices. □

Corollary 2.7. *Let $\pi : \bar{M}_{g,m+n} \rightarrow \bar{M}_{g,m}$. Then we have*

$$\pi^* e^{\sum_{i=1}^m r_i L_i} = e^{\sum_{i=1}^m r_i L_i} + \sum_{\underline{a}: \text{all decorations}} \mathcal{O}_{D_{(m),\underline{a}}} \prod_{i=1}^m \delta(e^{r_i(L_i - F_{i,\underline{a}_i})}).$$

Use the same assumptions as in the previous theorem when applying δ -operator on the right-hand side.

Proof. Take G to be $e^{\sum_{i=1}^m r_i L_i}$ and apply the previous theorem. Notice that if $F(\underline{x}_1, \dots, \underline{x}_r) = \prod_{i=1}^r F_i(\underline{x}_i)$, then

$$\delta(F(\underline{x}_1, \dots, \underline{x}_r)) = \prod_{i=1}^r \delta(F_i(\underline{x}_i)).$$

Here \underline{x}_i could be multiindices. □

2C. Application to the map $w : \mathcal{K}_{\Xi}^*(S^d X) \rightarrow \bar{M}_{g,R}(X)$. We continue to write $R = \ell - \#A = I \sqcup J$ for the number of marked points of D . Consider the map $p : \tilde{\mathfrak{K}}_{\Xi}(BS_d) \rightarrow \mathfrak{M}_{g,R}$ sending a triple $C \leftarrow C' \rightarrow D$ to D as before.

Write M_i , L'_i , and L_i for the cotangent line bundles on D , C' and C respectively. Let $\iota : \langle R \rangle \subseteq \langle \ell \rangle$ be the inclusion of marked points such that $\iota(i) \in C'$ maps to $i \in D$. Write $\bar{\iota}(i)$ for the corresponding point of C .

Given $f : \gamma \rightarrow \eta$ lying over D , we define $F_{f, \bar{\iota}(i)}$ as

$$F_{f, \bar{\iota}(i)} = F_{i, \underline{a}_i}(L_{i_1}^{m(1)}, \dots, L_i^{m(\deg \underline{a}_i)}).$$

The power m is given by the ramification between nodes, F_{i, \underline{a}_i} is defined in the point target case, and \underline{a}_i is the combinatorial type of f . The definition of type is exactly the same as the point case.

Proposition 2.8. *Let $G = G(x_1, \dots, x_R)$ be any power series of R variables. Then*

$$\begin{aligned} & p^*(G(M_1, \dots, M_R)) \\ &= G(L_{\bar{\iota}(1)}^{m(t_1)}, \dots, L_{\bar{\iota}(R)}^{m(t_R)}) + \sum_{f: \gamma \rightarrow \eta} \mathcal{O}_{\mathfrak{M}_f} \delta(G(L_{\bar{\iota}(1)}^{m(t_R)} - F_{f, \bar{\iota}(1)}, \dots, L_{\bar{\iota}(R)}^{m(t_R)} - F_{f, \bar{\iota}(R)})), \end{aligned}$$

where δ only applies on variables $\{L_{ij}\}$.

Proof. The proof is similar to the point case. We only need to take care of the difference coming from ramification points.

For the power on marked points, note that $p^*M_i = (L'_i)^{m(t_i)} + \text{torsion part}$ and $L'_i = L_{\bar{\iota}(i)}$ since $\tilde{C}' \rightarrow \tilde{C}$ is étale.

For the definition of $F_{f, \bar{\iota}(i)}$, if D is a divisor, note that

$$\mathcal{O}_{mD} := \mathcal{O} - \mathcal{O}(-mD) = \mathcal{O} - (\mathcal{O} - \mathcal{O}_D)^m = \delta(L_e^m) \mathcal{O}_D,$$

where L_e is characterized by $\mathcal{O}_D^2 = (1 - L_e)\mathcal{O}_D$. This explains the power in the definition of $F_{f, \bar{\iota}(i)}$. \square

Remark 2.9. The cover $\tilde{C}' \rightarrow \tilde{C}$ is étale, so any marked point $i \in \tilde{C}'$ and its image $s \in \tilde{C}$ will have the same ψ classes $L'_i = L_s$. Proposition 2.8 writes the pullback $p^*G(M_1, \dots, M_R)$ as a power series in L'_i ; plugging in L_s for each L'_i in the fiber of $s \in \tilde{C}$ rewrites this pullback as a power series $H(L_1, \dots, L_n)$. This power series is invariant under $\underline{\text{Aut}}(\tilde{C}'|\tilde{C})$, giving an analogous formula on $\mathcal{K}_{\Xi}^*(S^d X)$:

$$\omega^*G(M_1, \dots, M_R) = H^*(L_1, \dots, L_n).$$

Remark 2.10. The map $\phi : \mathcal{K}_{\Xi}^*(S^d X) \rightarrow \bar{\mathcal{K}}_{\Xi}(S^d X)$ is a finite étale map. The ψ classes of $\mathcal{K}_{\Xi}^*(S^d X)$ are pulled back from those of $\bar{\mathcal{K}}_{\Xi}(S^d X)$. If $H(\vec{L})$ is a power series with coefficients $\alpha_I \in K^\circ(\mathcal{K}_{\Xi}^*(S^d X))$ in K -theory, write $H^\phi(\vec{L})$ for the power series on $\bar{\mathcal{K}}_{\Xi}(S^d X)$ with coefficients the pushforwards $\phi_*\alpha_I$ of the coefficients of $H(\vec{L})$. If $\beta \in K_\circ(\bar{\mathcal{K}}_{\Xi}(S^d X))$, $\alpha \in K^\circ(\mathcal{K}_{\Xi}^*(S^d X))$, the projection formula equates

$$\phi_*(\phi^*\beta \otimes H(\vec{L}) \otimes \alpha) = \beta \otimes H^\phi(\vec{L}) \otimes \phi_*\alpha.$$

Remark 2.11. We sketch how to compute the coefficients $\mathcal{O}_{\mathfrak{M}_f}$ in terms of divisors on the moduli space of curves, and then how to deal with those divisors in quantum K -theory. This makes the power series $H^\phi(\vec{L})$ computable.

A map of covers $f : \gamma \rightarrow \eta$ induces a map $f_* : \mathfrak{M}_\gamma \rightarrow \mathfrak{M}_\eta$ with image a closed substack \mathfrak{M}_f . The map $f_* : \mathfrak{M}_\gamma \rightarrow \mathfrak{M}_f$ is a combination of a finite étale torsor for the automorphisms $\underline{\text{Aut}}(f|\eta)$ of the source over the target and a gerbe part for the stacky points. By Proposition 1.9, the gerbe part does not affect the pushforward. The pushforward is then related to the regular representation of the sheaf $\underline{\text{Aut}}(f|\eta)$.

After the above reductions, it remains to explain that quantum K -invariants involving the torsion structure sheaf \mathcal{O}_S supported on boundary strata S can be written in terms of the “usual” quantum K -invariants. In the cohomological Gromov–Witten theory, this is achieved by the splitting axiom, as the boundary stratum consist of the substacks indexed by the “dual graphs” exhibiting the imposed nodes of the general curves. In quantum K -theory a parallel splitting axiom is also available, albeit in a more sophisticated form. See [Givental 2000] and the genus reduction and splitting axioms in [Lee 2004, Section 4.3] for details. One will then have to push these coefficients forward as outlined in Section 3A.

3. Main theorem

Let $\alpha_1, \dots, \alpha_R \in K^\circ(X)$ be classes and form

$$\alpha = \alpha_1 \boxtimes \cdots \boxtimes \alpha_R = \alpha_1|_{X^R} \otimes \cdots \otimes \alpha_R|_{X^R} \in K^\circ(X^R).$$

Let $G(\vec{M})$ a power series in the ψ classes of $\bar{M}_{g,R}(X)$ with coefficients arbitrary classes in $K^\circ(\bar{M}_{g,R}(X))$.

Form the Gromov–Witten invariant

$$\chi\left(\mathcal{O}_{\bar{M}_{g,R}(X)}^{\text{vir}} \otimes ev^* \prod \alpha_i G(\vec{M})\right) \quad \text{in } K_\circ(\text{pt}) = \mathbb{Q}.$$

Write $H(\vec{L})$ for the power series in the ψ -classes L_i which is equal to $\omega^* G(\vec{M})$ by Remark 2.9. Likewise write $H^\phi(\vec{L})$ with the power series with coefficients given by the pushforwards of those of $H(\vec{L})$ as in Remark 2.10.

Theorem 3.1. *Gromov–Witten invariants on $\bar{M}_{g,R}(X)$ are equal to S_k -invariant Euler characteristics on the space $\tilde{M}_\Xi(S^d X)$ of stable genus zero maps to $[\text{Sym}^d X]$:*

$$\begin{aligned} \chi(\mathcal{O}_{\bar{M}_{g,R}(X)}^{\text{vir}} \otimes ev^* \alpha G(\vec{M})) &= \chi(\mathcal{O}_{\mathcal{K}_\Xi^*(S^d X)}^{\text{vir}} \otimes ev^* \alpha H(\vec{L})) \\ &= \chi(\mathcal{O}_{\bar{\mathcal{K}}_\Xi(S^d X)}^{\text{vir}} \otimes \phi_*(ev^* \alpha) H^\phi(\vec{L})) \\ &= \chi_{S_k}(\mathcal{O}_{\tilde{M}_\Xi(S^d X)}^{\text{vir}} \otimes \psi^*(\phi_*(ev^* \alpha) H^\phi(\vec{L})))^{S_k}. \end{aligned}$$

Proof. For the first equality, apply the projection formula for $w : \mathcal{K}_\Xi^*(S^d X) \rightarrow \bar{M}_{g,R}(X)$ and the equality of virtual fundamental classes from Proposition 1.17. The evaluation maps are compatible and the power series $H(\vec{L})$ is designed to be the pullback $w^* G(\vec{M})$.

The second equality results from the projection formula and the pullback

$$\phi^*[\mathcal{O}_{\bar{\mathcal{K}}_{\Xi}(S^d X)}^{\text{vir}}] = [\mathcal{O}_{\mathcal{K}_{\Xi}^*(S^d X)}^{\text{vir}}].$$

The power series $H^\phi(\vec{L})$ applies ϕ_* to the coefficients, so we are using the projection formula for each monomial of $H(\vec{L})$ one at a time.

The third equality results from the S_k -quotient $\psi : \tilde{M}_{\Xi}(S^d X) \rightarrow \bar{\mathcal{K}}_{\Xi}(S^d X)$ as in Section 1F. The pullback

$$\begin{array}{ccc} \tilde{M}_{\Xi}(S^d X) & \longrightarrow & \text{pt} \\ \psi \downarrow & \lrcorner & \downarrow \\ \bar{\mathcal{K}}_{\Xi}(S^d X) & \longrightarrow & BS_k \end{array}$$

equates the underlying vector space of $\chi_{S_k}(V)$ with the pullback $\chi(\psi^*V)$. The pushforward map $BS_k \rightarrow \text{pt}$ then takes the quotient $(-)^{S_k}$ by S_k . \square

We have reduced Gromov–Witten invariants on $\bar{M}_{g,R}(X)$ to some equivariant Euler characteristics on $\tilde{M}_{\Xi}(S^d X)$. But are these Euler characteristics actually Gromov–Witten invariants?

Lemma 3.2. *The equivariant Euler characteristic*

$$\chi\left(\mathcal{O}_{\bar{\mathcal{K}}_{\Xi}(S^d X)}^{\text{vir}} \otimes \phi_*\left(ev^* \prod \alpha_i\right) H^\phi(\vec{L})\right)$$

is a “Gromov–Witten invariant” in genus zero. In other words, the class $\phi_* ev^* \prod \alpha_i$ can be described as the pullback of a class via the evaluation map of $\bar{\mathcal{K}}_{\Xi}(S^d X)$.

We spend the rest of the section proving Lemma 3.2. This lemma was essentially left to the reader in [Costello 2006], although it is simpler in Chow groups than in K -theory. Reducing the part $H^\phi(\vec{L})$ is left to the reader, following Remark 2.11 and the process we outline for the evaluation classes.

3A. Turning equivariant Euler characteristics on $\tilde{M}_{\Xi}(S^d X)$ into proper Gromov–Witten invariants. We need to show that $\phi_* ev^* \alpha$ is pulled back from the evaluation map on $\bar{\mathcal{K}}_{\Xi}(S^d X)$. We first show it is pulled back from a natural map

$$\bar{\mathcal{K}}_{\Xi}(S^d X) \rightarrow ([\text{Sym}^d X])^J \times [\text{Sym}^{\#I} X].$$

In the covers $C' \rightarrow C$ parameterized by $\bar{\mathcal{K}}_{\Xi}(S^d X)$, none of the marked points of C' are ordered. Write $\mathcal{Q}_j \rightarrow \bar{\mathcal{K}}_{\Xi}(S^d X)$ for $j \in J$ for the d -sheeted cover of preimages of $j \in C$ in C' . Likewise, let $\mathcal{P} \rightarrow \bar{\mathcal{K}}_{\Xi}(S^d X)$ be the $S_{\#I}$ -torsor ordering the preimages in C' of $\infty \in C$. The product over $\bar{\mathcal{K}}_{\Xi}(S^d X)$ of all these d -sheeted covers and the $S_{\#I}$ -torsor is \mathcal{K}^* :

$$\mathcal{K}_{\Xi}^*(S^d X) = \prod_{\bar{\mathcal{K}}_{\Xi}(S^d X)} \mathcal{Q}_j \times_{\bar{\mathcal{K}}_{\Xi}(S^d X)} \mathcal{P} \rightarrow \bar{\mathcal{K}}_{\Xi}(S^d X).$$

Considering the map $\mathcal{Q}_j \rightarrow X$ as a d -sheeted cover of $\bar{\mathcal{K}}$, it is parameterized by a map to $[\mathrm{Sym}^d X]$:

$$\begin{array}{ccc} \mathcal{Q}_j & \xrightarrow{\quad \Gamma \quad} & X \times [\mathrm{Sym}^{d-1} X] \\ \downarrow & & \downarrow w_j \\ \bar{\mathcal{K}}_{\Xi}(S^d X) & \longrightarrow & [\mathrm{Sym}^d X] \end{array}$$

The equivariant map $\mathcal{P} \rightarrow X^I$ is also parameterized by a map to a symmetric product stack, but with different total space:

$$\begin{array}{ccc} \mathcal{P} & \xrightarrow{\quad \Gamma \quad} & X^{\#I} \\ \downarrow & & \downarrow w_I \\ \bar{\mathcal{K}}_{\Xi}(S^d X) & \longrightarrow & [\mathrm{Sym}^{\#I} X] \end{array}$$

There is then a pullback square

$$(9) \quad \begin{array}{ccccccc} \mathcal{K}_{\Xi}^*(S^d X) & \longrightarrow & \prod \mathcal{Q}_j \times \mathcal{P} & \longrightarrow & X^J \times ([\mathrm{Sym}^{d-1} X])^J \times X^{\#I} & \longrightarrow & X^R \\ \downarrow \phi & \lrcorner & \downarrow & \lrcorner & \downarrow w & & \\ \bar{\mathcal{K}}_{\Xi}(S^d X) & \xrightarrow{\Delta} & \bar{\mathcal{K}}_{\Xi}(S^d X)^R & \longrightarrow & ([\mathrm{Sym}^d X])^J \times [\mathrm{Sym}^{\#I} X] & & \end{array}$$

Take classes $\alpha_1, \dots, \alpha_R \in K^{\circ}(X)$. Write

$$\alpha := \alpha_1 \boxtimes \dots \boxtimes \alpha_R$$

for the tensor product of the pullback of these classes to $X^J \times ([\mathrm{Sym}^{d-1} X])^J \times X^I$. Pullback and pushforward in K° theory along cartesian squares commute (5), so the resulting classes on $\bar{\mathcal{K}}_{\Xi}(S^d X)$ are the same:

$$(10) \quad \phi_*(\alpha|_{\mathcal{K}_{\Xi}^*(S^d X)}) = (w_* \alpha)|_{\bar{\mathcal{K}}_{\Xi}(S^d X)} \quad \text{in } K^{\circ}(\bar{\mathcal{K}}_{\Xi}(S^d X)).$$

3B. Evaluation maps. We have shown that $\phi_* ev^* \alpha$ is pulled back from the map $\bar{\mathcal{K}}_{\Xi}(S^d X) \rightarrow ([\mathrm{Sym}^d X])^J \times [\mathrm{Sym}^{\#I} X]$. We need to compare this with the natural evaluation map on $\bar{\mathcal{K}}_{\Xi}(S^d X)$.

Gromov–Witten invariants are certain integrals of K theoretic classes pulled back from the evaluation maps as defined in Section Section 1E. To make sense of this, we need to be pedantic about the correct evaluation map for each target stack.

For spaces parameterizing multiple curves C_1, C_2 such as Hurwitz stacks, there is more than one evaluation map. We default to the evaluation maps of the base curve of the cover to be correct.

Example 3.3. The space $\tilde{\mathcal{M}}_{\Xi}(S^d X)$ parameterizes ramified covers $C' \rightarrow C$ together with a map $C' \rightarrow X$. There are n ordered marked points of C and ℓ unordered

points of C' . The resulting evaluation map is

$$\text{ev} : \tilde{\mathcal{M}}_{\Xi}(S^d X) \rightarrow \bar{\mathbb{I}}([\text{Sym}^d X])^n.$$

Example 3.4. The space $\bar{\mathcal{K}}_{\Xi}(S^d X)$ is similar to $\tilde{\mathcal{M}}_{\Xi}(S^d X)$ but harder, because the k gerbey points on C are not even ordered. That part of the evaluation map lands in a symmetric stack of a symmetric stack

$$\text{ev} : \bar{\mathcal{K}}_{\Xi}(S^d X) \rightarrow \bar{\mathbb{I}}([\text{Sym}^d X])^J \times [\text{Sym}^k[\text{Sym}^d X]] \times [\text{Sym}^d X].$$

The evaluation maps fit in a commutative square:

$$\begin{array}{ccc} \tilde{\mathcal{M}}_{\Xi}(S^d X) & \xrightarrow{\text{ev}} & \bar{\mathbb{I}}([\text{Sym}^d X])^J \times ([\text{Sym}^d X])^k \times [\text{Sym}^d X] \\ \downarrow & & \downarrow \\ \bar{\mathcal{K}}_{\Xi}(S^d X) & \xrightarrow{\text{ev}} & \bar{\mathbb{I}}([\text{Sym}^d X])^J \times [\text{Sym}^k[\text{Sym}^d X]] \times [\text{Sym}^d X] \end{array}$$

We won't need classes on the middle factor $\bar{\mathbb{I}}([\text{Sym}^k[\text{Sym}^d X]])$. Write ev' for the projection away from this factor on the evaluation map of $\bar{\mathcal{K}}_{\Xi}(S^d X)$:

$$\text{ev}' : \bar{\mathcal{K}}_{\Xi}(S^d X) \rightarrow \bar{\mathbb{I}}([\text{Sym}^d X])^J \times [\text{Sym}^d X].$$

Remark 3.5. The 2-functor $\bar{\mathbb{I}}(-)$ does not distribute over products because of the representability requirement. For example, the identity map on $B\mathbb{Z}/2 \times B\mathbb{Z}/2$ is representable, but it doesn't factor through a representable map to either factor. It is more accurate to say the evaluation map lands in the product of $\bar{\mathbb{I}}(-)$ applied to each factor, so it is a product of the evaluation maps for each marked point.

Our ramification points are μ_r -banded gerbes mapping to BS_d . Given a map $B\mu_r \rightarrow BS_d$, we can extract the set theoretic fiber of the stacky point of $\tilde{C}' \rightarrow \tilde{C}$ by taking the set-theoretic quotient $\langle d \rangle / \mu_r$ of the corresponding action as in Remark 1.4.

More generally, we have a μ_r -gerbe $\mathcal{G} \rightarrow T$ with a map $\mathcal{G} \rightarrow [\text{Sym}^d X]$. This means a finite étale cover $\tilde{\mathcal{G}} \rightarrow \mathcal{G}$ of degree d . The coarse moduli space of \mathcal{G} is T , and that of $\tilde{\mathcal{G}}$ is a finite étale cover $T' \rightarrow T$. The degree k of this cover is some number less than d . The map $\tilde{\mathcal{G}} \rightarrow X$ factors through T' because X is a scheme. This procedure gives a map

$$(11) \quad c : \bar{\mathbb{I}}([\text{Sym}^d X]) \rightarrow \bigsqcup_{k \leq d} [\text{Sym}^k X],$$

sending $\mathcal{G} \rightarrow [\text{Sym}^d X]$ to $T' \rightarrow X$. We describe this map in detail in the next Section Section 3C.

Because we fixed the discrete data Ξ for $\bar{\mathcal{K}}_\Xi(S^d X)$, we know which component of each factor of $\bigsqcup_{k \leq d} [\text{Sym}^k X]$ it maps to

$$\begin{array}{ccc} \bar{\mathcal{K}}_\Xi(S^d X) & \longrightarrow & ([\text{Sym}^d X])^J \times [\text{Sym}^{\#I} X] \\ \text{ev}' \downarrow & & \downarrow \text{inc} \\ \bar{\mathcal{I}}([\text{Sym}^d X])^J \times [\text{Sym}^d X] & \xrightarrow{c} & (\bigsqcup_k [\text{Sym}^k X])^{J \cup \infty} \end{array}$$

Let β be a class in $K^\circ([\text{Sym}^d X]^J \times [\text{Sym}^{\#I} X])$. Because the map inc is an inclusion of components, we have

$$\text{inc}^* \text{inc}_* \beta = \beta.$$

Then the pullback $\beta|_{\bar{\mathcal{K}}_\Xi(S^d X)}$ is the same as the class

$$\text{ev}'^* c^* \text{inc}_* \beta \quad \text{in } K^\circ(\bar{\mathcal{K}}_\Xi(S^d X)).$$

Proof of Lemma 3.2. Take $\beta = w_* \alpha$ above. Then

$$\phi_*(\alpha|_{\mathcal{K}_\Xi^*(S^d X)}) = (w_* \alpha)|_{\bar{\mathcal{K}}_\Xi(S^d X)} = \text{ev}'^* c^* \text{inc}_* w_* \alpha,$$

using equation (10) and the discussion immediately above. This expresses the factor $\phi_*(\alpha|_{\mathcal{K}_\Xi^*(S^d X)})$ in the Gromov–Witten invariant as a class pulled back via the evaluation map on $\bar{\mathcal{K}}_\Xi(S^d X)$. □

3C. Describing the map c . We describe the map c in detail using the inertia stack $I([\text{Sym}^d X])$. This section is purely expository, and an example is given at the end.

The inertia stack of a quotient stack $[Y/G]$ is the disjoint union of the quotients

$$I([Y/G]) = \bigsqcup_{g \in G \text{ conj classes}} [Y^g/C_g]$$

of the fixed locus Y^g by the centralizer $C_g \subseteq G$.

For $[\text{Sym}^d X]$, we have $Y = X^d$ and $G = S_d$. Conjugacy classes of S_d are indexed by cycle types, the multiset of lengths of cycles. For example,

$$g_1 := (12)(34)(5) \in S_5 \mapsto \{2, 2, 1\}, \quad g_2 := (234)(761)(5)(89) \in S_9 \mapsto \{3, 3, 1, 2\}.$$

Let N_s be the number of cycles of length s , $N := \sum N_s$ the total number of cycles, and t the cardinality of the set of distinct lengths in the cycle type. We include all cycles of length one N_1 , so $\sum i N_i = d$. For g_1, g_2 above,

$$g_1 \mapsto N_1 = 1, N_2 = 2, t = 2, \quad g_2 \mapsto N_1 = 1, N_2 = 1, N_3 = 2, t = 3.$$

The centralizer of a cycle type g is

$$C_g = S_{N_1} \times S_{N_2} \rtimes (\mathbb{Z}/2)^{N_2} \times S_{N_3} \rtimes (\mathbb{Z}/3)^{N_3} \cdots \times S_{N_t} \rtimes (\mathbb{Z}/t)^{N_t}.$$

The fixed locus $(X^d)^g \subseteq X^d$ is the multidagonal

$$(X^d)^g = \{(x_1, \dots, x_d) \mid x_i = x_j \text{ if } i, j \text{ in the same cycle}\},$$

so $(X^d)^g \cong X^N$. For g_1, g_2 , we have

$$(X^d)^{g_1} = \{(x, x, y, y, z)\} \in X^5, \quad (X^d)^{g_2} = \{(x, x, x, y, y, y, z, w, w)\} \subseteq X^9.$$

Write $H_i = S_{N_i} \times (\mathbb{Z}/i)^{N_i} = \mathbb{Z}/i \wr S_{N_i}$. There is an exact sequence

$$0 \rightarrow (\mathbb{Z}/i)^{N_i} \rightarrow H_i \rightarrow S_{N_i} \rightarrow 1$$

and a splitting $S_{N_i} \subseteq H_i$. The subgroups \mathbb{Z}/i act trivially on the diagonal fixed locus $X \subseteq X^{N_i}$, so the stack quotient is a trivial gerbe

$$[X/(\mathbb{Z}/i)] = X \times B\mathbb{Z}/i.$$

The quotient $X^N/\prod H_i$ is the product of symmetric products

$$X^N/\prod H_i = \prod_i [\mathrm{Sym}^{N_i}[X/(\mathbb{Z}/i)]] = \prod_i [\mathrm{Sym}^{N_i}(X \times B\mathbb{Z}/i)].$$

On each component, there is a map to a single symmetric product

$$\prod_i [\mathrm{Sym}^{N_i}(X \times B\mathbb{Z}/i)] \rightarrow \prod_i [\mathrm{Sym}^{N_i} X] \xrightarrow{c'} [\mathrm{Sym}^N X].$$

The second map c' takes t covers $T'_i \rightarrow T$ of degrees N_i and assembles them into one cover $T' = \bigsqcup T'_i \rightarrow T$ of degree N . This gives a map

$$(12) \quad \bar{c} : I([\mathrm{Sym}^d X]) \rightarrow \bigsqcup_{N \leq d} [\mathrm{Sym}^N X].$$

The reader can check this coincides with the map c defined in (11).

Lemma 3.6. *There is a commutative diagram*

$$\begin{array}{ccc} I([\mathrm{Sym}^d X]) & & \\ \downarrow & \searrow \bar{c} & \\ \bar{I}([\mathrm{Sym}^d X]), & \xrightarrow{c} & \bigsqcup_{k \leq d} [\mathrm{Sym}^k X] \end{array}$$

where c is the map (11) and \bar{c} is (12).

We explain the above from the point of view of covers. A T -point of $I([\mathrm{Sym}^d X])$ is a map $T \times B\mu_r \rightarrow [\mathrm{Sym}^d X]$, which is a d -sheeted cover $P \rightarrow T \times B\mu_r$. Write P_0

for the pullback d -sheeted cover of T :

$$\begin{array}{ccc} P_0 & \longrightarrow & P \\ \downarrow & \lrcorner & \downarrow \\ T & \longrightarrow & T \times B\mu_r \end{array}$$

so $P = [P_0/\mu_r]$.

Fix a generator $\mathbb{Z}/r \simeq \mu_r$. If T is a geometric point, $P_0 \simeq \langle d \rangle$ and $\mu_r \curvearrowright P_0$ is an element $\sigma \in S_d$. Its order is the lcm of the cycle lengths, which divides r ; r is equal to the lcm when the map $T \times B\mu_r \rightarrow [\text{Sym}^d X]$ is representable. Reordering $P_0 \simeq \langle d \rangle$ conjugates σ , so σ is well defined as a conjugacy class.

Even if T is not a geometric point, this defines a locally constant function

$$T \mapsto S_d / \text{ad} S_d, \quad t \mapsto [\sigma]$$

from T to the conjugacy classes of S_d . This decomposes $I([\text{Sym}^d X])$ into components corresponding to cycle type, or partitions of d .

Given a partition $d = \sum i N_i$ corresponding to the cycle type of $\sigma \in S_d$, a point $T \rightarrow I([\text{Sym}^d X])$ factors through the corresponding component if its fibers at geometric points are isomorphic to $\langle d \rangle / \sigma$. The N_i different orbits of i points may be interchanged in families over T , and each such orbit may vary in a $B\mathbb{Z}/i$ family. The collection of orbits of i points is parameterized by the stack

$$[\text{Sym}^{N_i}(B\mathbb{Z}/i)].$$

We have $\prod [\text{Sym}^{N_i}(X \times B\mathbb{Z}/i)]$ instead because we also need a map to X .

The case $d = 5$. If $d = 5$, the seven cycle types/partitions are

$$5, \quad 4+1, \quad 3+2, \quad 3+1+1, \quad 2+2+1, \quad 2+1+1+1, \quad 1+1+1+1+1.$$

The corresponding components of $I([\text{Sym}^5 X])$ are

$$\begin{aligned} X \times B\mathbb{Z}/5, \quad X^2 \times B\mathbb{Z}/4, \quad X^2 \times B\mathbb{Z}/3 \times B\mathbb{Z}/2, \quad X/(\mathbb{Z}/3) \times X^2/S_2, \\ X \times X^2/(S_2 \times (\mathbb{Z}/2)^2), \quad X/\mathbb{Z}/2 \times X^3/S_3, \quad X^5/S_5. \end{aligned}$$

Project away from the cyclic gerbes $B\mathbb{Z}/i$:

$$X, \quad X^2, \quad X^2, \quad X \times [\text{Sym}^2 X], \quad [\text{Sym}^2 X] \times X, \quad X \times [\text{Sym}^3 X], \quad [\text{Sym}^5 X].$$

For example, consider a trivial gerbe $b : B\mu_4 \rightarrow [\text{Sym}^5 X]$ mapping to the symmetric product. This corresponds to a 5-sheeted cover $P \rightarrow B\mu_4$, counting stacky multiplicity.

The composite $\text{pt} \rightarrow B\mu_4 \rightarrow [\text{Sym}^5 X] \rightarrow BS_5$ parameterizes a 5-sheeted cover of the point, which we trivialize and view as $\langle 5 \rangle$. The action of μ_4 on $\langle 5 \rangle$ can be

viewed as a map $\mu_4 \rightarrow S_5$. Choose a generator to identify $\mu_4 \simeq \mathbb{Z}/4$ and let $\sigma \in S_5$ be the image of the generator.

The element σ has order dividing 4. Take, for example,

$$\sigma = g_1 = (12)(34)(5).$$

Its order 2 is not 4, so the classifying map b is not representable.

The stack quotient of $\langle 5 \rangle$ by $\mathbb{Z}/4 \cdot \sigma$ is a disjoint union

$$B\mu_2 \sqcup B\mu_2 \sqcup B\mu_4.$$

This is the 5-sheeted cover $P \rightarrow B\mu_4$. The corresponding point $\text{pt} \rightarrow I([\text{Sym}^5 X])$ factors through the component of the partition $2 + 2 + 1$, i.e., $X \times X^2 / (S_2 \times (\mathbb{Z}/2)^2)$ above.

The cover comes with a map $P \rightarrow X$. The map \bar{c} takes coarse moduli spaces of $P \rightarrow B\mu_4$, obtaining $\langle 3 \rangle \rightarrow \text{pt}$. The map $P \rightarrow X$ factors through $\langle 3 \rangle$ because X is a scheme with no stack structure. This sends the component $X \times X^2 / (S_2 \times (\mathbb{Z}/2)^2)$ to $[\text{Sym}^3 X]$.

4. Elliptic curves example

We apply our theorem in the case of elliptic curves, with $g = R = 1$, $d = 2$, and $X = \text{pt}$. The symmetric product is the classifying stack $[\text{Sym}^d X] = BS_d$. The results are reassuring but not surprising. See Figure 4.

We know the map w is proper and birational. For elliptic curves, more is true.

Lemma 4.1. *The map $w : \tilde{\mathcal{K}}_{\Xi^*}^*(BS_d) \rightarrow \bar{\mathcal{M}}_{1,1}$ is an isomorphism.*

Proof. The map on coarse moduli spaces is an isomorphism $\mathbb{P}^1 = \mathbb{P}^1$. It remains to show the stack structure is the same; i.e., the automorphism groups of the admissible covers $C' \rightarrow C$ are the same as that of the stabilization $D = C'^{\text{st}}$. We already checked this for a generic elliptic curve in the proof of Lemma 1.16.




Figure 4. An elliptic curve C' and its double cover of $C = \mathbb{P}^1$ simply ramified at four points: Ξ with $g = 1$, $d = 2$, $k = 0$. Marked points are white if remembered and black if forgotten under the map $p : \tilde{\mathcal{K}}_{\Xi}^*(BS_d) \rightarrow \mathfrak{M}_{1,1}$.




Figure 5. The admissible cover $C' \rightarrow C$ which stabilizes to the nodal cubic $C'^{\text{st}} = D$. The components of C' are labeled C'_0, C'_1 and they each doubly cover a component of C . The pair of nodes P is the intersection $C'_0 \cap C'_1$ of the components.

We want to show $\underline{\text{Aut}}(C' \rightarrow C) = \underline{\text{Aut}}(D)$. Again, it helps that automorphisms of the covering map are a subgroup of automorphisms of the source $\underline{\text{Aut}}(C' \rightarrow C) \subseteq \underline{\text{Aut}}(C')$. These automorphisms must send ramification points to ramification points, and it will be clear that they also send the preimage $I \in C'$ of infinity to itself.

For all the smooth elliptic curves D , the source is already stable $C' = D$ and the unique map $C' \rightarrow C = \mathbb{P}^1$ is the quotient by the elliptic involution. This identifies $\underline{\text{Aut}}(C' \rightarrow C) \simeq \underline{\text{Aut}}(D)$ for all smooth elliptic curves with automorphism group $\mathbb{Z}/2$. The smooth curves $j = 0, 1728$ remain, as does the singular cubic.

The curve $j = 1728$ has the equation

$$y^2 = x^3 - x.$$

The automorphism group $\mathbb{Z}/4$ is generated by $(x, y) \mapsto (-x, iy)$. This commutes with the automorphism $x \mapsto -x$ of \mathbb{P}^1 . Likewise $j = 0$ has the equation

$$y^2 = x^3 - 1.$$

Letting ζ be a sixth root of unity, the automorphism group is generated by the map $(x, y) \mapsto (\zeta^2 x, \zeta^3 y)$. This also commutes with an automorphism of \mathbb{P}^1 .

For the singular elliptic curve D , the preimage is an admissible cover $C' \rightarrow C$ with both source and target reducible. See Figure 5. Each consists of two components, labeled C'_i, C_i for $i = 0, 1$. Assume that the point in $I \subseteq C'$ lies on C'_0 .

The map restricts to two double covers $C'_i \rightarrow C_i$ of \mathbb{P}^1 's. The preimage $P \subseteq C'$ of the singular point of C is a pair of nodes joining C'_0, C'_1 .

Any automorphism of C' must restrict to an automorphism of each component because of the marked point. Any automorphism φ of C' restricts to an automorphism of the pair of nodes P which determines φ . This is because φ must preserve

at least three points on each component, the nodes and the ramification points. All such automorphisms lie over C , so $\underline{\text{Aut}}(C' \rightarrow C) = \underline{\text{Aut}}(P) = \mathbb{Z}/2$. This is the same automorphism group of the singular stabilization $D = C'^{\text{st}}$, the nodal cubic. \square

The map $\phi : \mathcal{K}_{\Xi}^*(BS_d) \rightarrow \bar{\mathcal{K}}_{\Xi}(BS_d)$ is also an isomorphism by Remark 1.20, so Theorem 3.1 merely says that quantum K invariants on $\bar{M}_{1,1} = \mathbb{P}(4, 6)$ are equivariant Euler characteristics on its natural S_3 -cover by $\tilde{M}_{\Xi}(BS_d)$.

This cover is pulled back from the quotient $\mathbb{P}^1 \rightarrow \mathbb{P}(2, 3)$ by S_3 acting on $\lambda \in \mathbb{P}^1$. This map classically parameterizes the Legendre form

$$y^2 = x(x-1)(x-\lambda)$$

of an elliptic curve. The map $\tilde{M}_{\Xi}(BS_d) \rightarrow \bar{\mathcal{K}}_{\Xi}(BS_d) = \bar{M}_{1,1}$ then fits in a pullback square

$$\begin{array}{ccc} \tilde{M}_{\Xi}(BS_d) & \longrightarrow & \bar{M}_{1,1} \\ \downarrow & \lrcorner & \downarrow \\ \mathbb{P}^1 & \xrightarrow{/S_3} & \mathbb{P}(2, 3) \end{array}$$

with vertical arrows μ_2 -gerbes.

Acknowledgements

The authors wish to thank Sarah Arpin, Peter Bakic, Sebastian Bozlee, Renzo Cavalieri, Kevin Costello, Dan Edidin, Adeel Khan, Rufus Lawrence, Sam Molcho, Hsian-Hua Tseng, Jonathan Wise, and the Mathoverflow community for their help and support. This project originated in the work of Herr with Wise [Herr and Wise 2023].

Chou and Lee wish to acknowledge the partial supports from Academia Sinica, National Science and Technology Council, Simons Foundation and the University of Utah. Herr thanks Leiden University, University of Utah and the National Science Foundation for its support through RTG grant #1840190.

References

- [Abramovich et al. 2002] D. Abramovich, T. Graber, and A. Vistoli, “Algebraic orbifold quantum products”, pp. 1–24 in *Orbifolds in mathematics and physics* (Madison, WI, 2001), Contemp. Math. **310**, Amer. Math. Soc., Providence, RI, 2002. MR Zbl
- [Abramovich et al. 2008] D. Abramovich, T. Graber, and A. Vistoli, “Gromov–Witten theory of Deligne–Mumford stacks”, *Amer. J. Math.* **130**:5 (2008), 1337–1398. MR Zbl
- [Behrend and Fantechi 1997] K. Behrend and B. Fantechi, “The intrinsic normal cone”, *Invent. Math.* **128**:1 (1997), 45–88. MR Zbl
- [Buch and Mihalcea 2011] A. S. Buch and L. C. Mihalcea, “Quantum K -theory of Grassmannians”, *Duke Math. J.* **156**:3 (2011), 501–538. MR Zbl

- [Buch et al. 2013] A. S. Buch, P.-E. Chaput, L. C. Mihălcea, and N. Perrin, “Finiteness of cominuscule quantum K -theory”, *Ann. Sci. Éc. Norm. Supér.* (4) **46**:3 (2013), 477–494. MR Zbl
- [Buch et al. 2020] A. S. Buch, S. Chung, C. Li, and L. C. Mihălcea, “Euler characteristics in the quantum K -theory of flag varieties”, *Selecta Math. (N.S.)* **26**:2 (2020), art. id. 29. MR Zbl
- [Chen and Ruan 2002] W. Chen and Y. Ruan, “Orbifold Gromov–Witten theory”, pp. 25–85 in *Orbifolds in mathematics and physics* (Madison, WI, 2001), Contemp. Math. **310**, Amer. Math. Soc., Providence, RI, 2002. MR Zbl
- [Chen and Ruan 2004] W. Chen and Y. Ruan, “A new cohomology theory of orbifold”, *Comm. Math. Phys.* **248**:1 (2004), 1–31. MR Zbl
- [Chou et al. 2023] Y.-C. Chou, L. Herr, and Y.-P. Lee, “The log product formula in quantum K -theory”, *Math. Proc. Cambridge Philos. Soc.* **175**:2 (2023), 225–252. MR Zbl
- [Costello 2006] K. Costello, “Higher genus Gromov–Witten invariants as genus zero invariants of symmetric products”, *Ann. of Math.* (2) **164**:2 (2006), 561–601. MR Zbl
- [Edidin 2013] D. Edidin, “Riemann–Roch for Deligne–Mumford stacks”, pp. 241–266 in *A celebration of algebraic geometry*, Clay Math. Proc. **18**, Amer. Math. Soc., Providence, RI, 2013. MR Zbl
- [Edidin et al. 2001] D. Edidin, B. Hassett, A. Kresch, and A. Vistoli, “Brauer groups and quotient stacks”, *Amer. J. Math.* **123**:4 (2001), 761–777. MR Zbl
- [Givental 2000] A. Givental, “On the WDVV equation in quantum K -theory”, *Michigan Math. J.* **48** (2000), 295–304. MR Zbl
- [Givental 2017] A. Givental, “Permutation-equivariant quantum K -theory, I: Definitions. Elementary K -theory of $\overline{\mathcal{M}}_{0,n}/S_n$ ”, *Mosc. Math. J.* **17**:4 (2017), 691–698. MR Zbl
- [Givental and Lee 2003] A. Givental and Y.-P. Lee, “Quantum K -theory on flag manifolds, finite-difference Toda lattices and quantum groups”, *Invent. Math.* **151**:1 (2003), 193–219. MR Zbl
- [Herr and Lee] L. Herr and Y. P. Lee, “Quantum K -theory and log geometry”, preprint. In preparation.
- [Herr and Wise 2023] L. Herr and J. Wise, “Costello’s pushforward formula: errata and generalization”, *Manuscripta Math.* **171**:3-4 (2023), 621–642. MR Zbl
- [Jockers and Mayr 2019] H. Jockers and P. Mayr, “Quantum K -theory of Calabi–Yau manifolds”, *J. High Energy Phys.* **2019** (2019), art. id. 11. MR Zbl
- [Jockers and Mayr 2020] H. Jockers and P. Mayr, “A 3d gauge theory/quantum K -theory correspondence”, *Adv. Theor. Math. Phys.* **24**:2 (2020), 327–457. MR Zbl
- [Lee 2004] Y.-P. Lee, “Quantum K -theory, I: Foundations”, *Duke Math. J.* **121**:3 (2004), 389–424. MR Zbl
- [Lee and Qu 2014] Y.-P. Lee and F. Qu, “Euler characteristics of universal cotangent line bundles on $\overline{\mathcal{M}}_{1,n}$ ”, *Proc. Amer. Math. Soc.* **142**:2 (2014), 429–440. MR Zbl
- [Noohi 2012] B. Noohi, “Homotopy types of topological stacks”, *Adv. Math.* **230**:4-6 (2012), 2014–2047. MR Zbl
- [Okounkov 2017] A. Okounkov, “Lectures on K -theoretic computations in enumerative geometry”, pp. 251–380 in *Geometry of moduli spaces and representation theory*, IAS/Park City Math. Ser. **24**, Amer. Math. Soc., Providence, RI, 2017. MR Zbl
- [Qu 2018] F. Qu, “Virtual pullbacks in K -theory”, *Ann. Inst. Fourier (Grenoble)* **68**:4 (2018), 1609–1641. MR Zbl
- [Stacks 2005–] “The Stacks project”, 2005–, available at <https://stacks.math.columbia.edu>.

[Tonita and Tseng 2013] V. Tonita and H.-H. Tseng, “Quantum orbifold Hirzebruch–Riemann–Roch theorem in genus zero”, preprint, 2013. arXiv 1307.0262

[Zhang] M. Zhang, “Quantum K -theory of toric stacks”, preprint, available at <https://tinyurl.com/ZhangWb>.

Received June 2, 2023. Revised April 14, 2024.

YOU-CHENG CHOU
INSTITUTE OF MATHEMATICS
ACADEMIA SINICA
TAIPEI
TAIWAN
bensochou@gate.sinica.edu.tw

LEO HERR
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF LEIDEN
LEIDEN
NETHERLANDS
herr@math.utah.edu

YUAN-PIN LEE
INSTITUTE OF MATHEMATICS
ACADEMIA SINICA
TAIPEI
TAIWAN
ypleemath@gate.sinica.edu.tw

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Fakultät für Mathematik
Universität Wien
Vienna, Austria
matthias.aschenbrenner@univie.ac.at

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Atsushi Ichino
Department of Mathematics
Kyoto University
Kyoto 606-8502, Japan
atsushi.ichino@gmail.com

Robert Lipshitz
Department of Mathematics
University of Oregon
Eugene, OR 97403
lipshitz@uoregon.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Dimitri Shlyakhtenko
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
shlyakht@ipam.ucla.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

Ruixiang Zhang
Department of Mathematics
University of California
Berkeley, CA 94720-3840
ruixiang@berkeley.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2024 is US \$645/year for the electronic version, and \$875/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2024 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 330 No. 1 May 2024

Monotone twist maps and Dowker-type theorems	1
PETER ALBERS and SERGE TABACHNIKOV	
Unknotting via null-homologous twists and multitwists	25
SAMANTHA ALLEN, KENAN İNCE, SEUNGWON KIM, BENJAMIN MATTHIAS RUPPIK and HANNAH TURNER	
\mathbb{R} -motivic v_1 -periodic homotopy	43
EVA BELMONT, DANIEL C. ISAKSEN and HANA JIA KONG	
Higher-genus quantum K -theory	85
YOU-CHENG CHOU, LEO HERR and YUAN-PIN LEE	
Unknotted curves on genus-one Seifert surfaces of Whitehead doubles	123
SUBHANKAR DEY, VERONICA KING, COLBY T. SHAW, BÜLENT TOSUN and BRUCE TRACE	
On the Gauss maps of complete minimal surfaces in \mathbb{R}^n	157
DINH TUAN HUYNH	
Explicit bounds on torsion of CM abelian varieties over p -adic fields with values in Lubin–Tate extensions	171
YOSHIYASU OZEKI	