

*Pacific
Journal of
Mathematics*

OBSTRUCTION COMPLEXES IN GRID HOMOLOGY

YAN TAO

Volume 331 No. 2

August 2024

OBSTRUCTION COMPLEXES IN GRID HOMOLOGY

YAN TAO

Recently, Manolescu–Sarkar constructed a stable homotopy-type for link Floer homology, which uses grid homology and accounts for all domains that do not pass through a specific square. In doing so, they produced an obstruction chain complex of the grid diagram with that square removed. We define the obstruction chain complex of the full grid, without the square removed, and compute its homology. Though this homology is too complicated to immediately extend the Manolescu–Sarkar construction, we give results about the existence of sign assignments in grid homology.

1. Introduction

Link Floer homology, developed by Ozsváth and Szabó [2004a; 2008], and Rasmussen [2003] is an invariant of oriented links in three-manifolds which comes from Heegaard Floer homology, from [Ozsváth and Szabó 2004c; 2004b]. Manolescu et al. [2007; 2009a] and Ozsváth et al. [2015] gave a combinatorial description of the link Floer chain complex for a link in S^3 using grid diagrams, known as grid homology. A toroidal grid diagram is a $n \times n$ grid of squares, with the left and right edges identified and the top and bottom edges identified, together with markings X and O , such that each row and column contains exactly one X and one O . Given a grid diagram \mathbb{G} , drawing vertical segments from the X to the O in each column and horizontal segments — going under the vertical segments whenever they cross — from the O to the X in each row gives the diagram of an oriented link L ; we say that \mathbb{G} is a grid diagram for L . Figure 1 shows a 5×5 grid diagram for the trefoil. The grid chain complex is generated by unordered n -tuples of intersection points between the horizontal and vertical circles — Figure 1 shows an example of such a generator.

Grid diagrams have been useful in a variety of applications in Heegaard Floer homology. Manolescu et al. [2009b] and Manolescu and Ozsváth [2010] obtain the Heegaard Floer invariants of 3- and 4-manifolds using grid diagrams, which give algorithmically computable descriptions. Sarkar [2011] uses grid homology to give another proof of Milnor’s conjecture on the slice genus of torus

MSC2020: 57K18.

Keywords: link Floer, stable homotopy-type, spectrum, sign assignment.




Figure 1. A 5×5 grid diagram for the trefoil, along with the generator [51243] drawn with \bullet . Note that the generator is independent of the X and O markings.

knots. Ozsváth et al. [2008], Ng et al. [2008], Chongchitmate and Ng [2013], and Khandhawit and Ng [2010] use a version of grid homology to prove results about Legendrian knots.

Manolescu and Sarkar [2021] constructed a stable homotopy refinement of knot Floer homology from the grid chain complex, using framed flow categories in the sense of [Cohen et al. 1995]. The Manolescu–Sarkar construction uses only those domains that do not pass through a particular square on the grid, and uses obstruction theory. Their obstruction chain complexes CD_* and CDP_* , which we will henceforth denote \widehat{CD}_* and \widehat{CDP}_* , respectively, have simple enough homology to construct a stable homotopy-type. We will extend them to complexes CD_* and CDP_* which contain all domains in the grid, first with $\mathbb{Z}/2$ coefficients, which can be extended to \mathbb{Z} coefficients using obstruction theory. We take the first step towards extending the Manolescu–Sarkar construction, by computing the homology of CD_* and partially computing the homology of CDP_* .

To state our main results, we fix the following convention throughout the paper. For a ring R , R^{2^n} will denote the chain complex given by $R^{\binom{n}{k}}$ in grading k with no differentials, and $R[U]$ the chain complex given by R in every nonnegative even grading and 0 in every odd grading (which by definition has no differentials). We begin by showing that:

Proposition 1.1. $H_*(CD_*; \mathbb{Z}/2)$ is isomorphic to $\mathbb{Z}/2[U]$.

In order to frame the moduli spaces in the Manolescu–Sarkar construction, we will need a sign assignment for the grid diagram. A sign assignment is a particular way of orienting the index 1 domains in Heegaard Floer homology; equivalently, it is a particular assignment of 0 or 1 to each rectangle in the grid. The existence and uniqueness (up to gauge equivalence) of sign assignments for toroidal grid diagrams was constructed in [Manolescu et al. 2007]; see also [Gallais 2008] for an explicit construction. In the course of our later computations, we will provide a different proof of this fact via obstruction theory:

Theorem 1.2. *Sign assignments for CD_* exist and are unique up to gauge equivalence (equivalently, up to 1-coboundaries of CD_* with $\mathbb{Z}/2$ coefficients).*

Given a sign assignment for CD_* , we obtain a definition of CD_* in \mathbb{Z} coefficients. Perhaps unsurprisingly, we then obtain the following analogue of Proposition 1.1.

Proposition 1.3. *$H_*(CD_*; \mathbb{Z})$ is isomorphic to $\mathbb{Z}[U]$.*

In the end, our eventual goal is to extend the Manolescu–Sarkar construction over the full grid. Since the moduli spaces presented in [Manolescu and Sarkar 2021] exhibit some bubbling, we will compute the first few homology groups of CDP_* . Unfortunately, CDP_* has too much homology to immediately construct a stable homotopy-type. So instead, we will work towards constructing a framed 1-flow category, which is a formulation by Lobb et al. [2020] that still contains all the information needed to define invariants such as the second Steenrod square. This requires only a sign assignment and a frame assignment, whose obstructions lie in the following lower homologies.

Theorem 1.4. *We have that:*

- (0) $H_0(CDP_*; \mathbb{Z}/2)$ is isomorphic to $\mathbb{Z}/2$.
- (1) $H_1(CDP_*; \mathbb{Z}/2)$ is isomorphic to $(\mathbb{Z}/2)^n$.
- (2) $H_2(CDP_*; \mathbb{Z}/2)$ is isomorphic to $(\mathbb{Z}/2)^{\binom{n}{2}+1}$.
- (3) $H_3(CDP_*; \mathbb{Z}/2)$ is isomorphic to $(\mathbb{Z}/2)^{\binom{n}{3}+n}$.

In this paper, we’ll show existence and uniqueness of sign assignments for CDP_* .

Theorem 1.5. *A sign assignment s on CDP_* exists, and is unique up to gauge transformations and the values of*

$$s_j := s(c_{x^{\text{Id}}}, \vec{e}_j, (1)).$$

(The elements $(c_{x^{\text{Id}}}, \vec{e}_j, (1)) \in CDP_*$ will be defined later in Section 4.)

Just like for CD_* , we can use Theorem 1.5 to define CDP_* with \mathbb{Z} coefficients. We have the following analogue of Theorem 1.4.

It remains to find a frame assignment for CDP_* using the above homology computation, and to complete the construction of the 1-flow category for the full grid, which we will carry out in a future paper. This present paper may be treated as a prelude thereof.

2. The obstruction complex


Definitions related to grid diagrams are summarized below. For details, see [Manolescu et al. 2007; 2009a; Ozsváth et al. 2015].

- An index n grid diagram \mathbb{G} is a torus together with n α -circles (drawn horizontally) and n β -circles (drawn vertically). The complements of the α (respectively, β) circles are called the horizontal (respectively, vertical) annuli — the complements of the α and β circles are called the square regions.
- Each vertical and horizontal annulus contains exactly one X and O marking, which are arbitrarily labeled X_1, \dots, X_n and O_1, \dots, O_n .
- The horizontal (respectively, vertical) annuli can be labeled by which O -marking they pass through — write H_i (respectively, V_i) for the horizontal (respectively, vertical) annulus passing through O_i .
- Given a fixed planar drawing of the grid, we can also label the α circles $\alpha_1, \dots, \alpha_n$ from bottom to top, and the β circles β_1, \dots, β_n from left to right. The annuli can also be labeled by which sets of α or β circles they lie between — write $H_{(i)}$ (respectively, $V_{(i)}$) for the horizontal annulus between α_i and α_{i+1} (respectively, vertical annulus between β_i and β_{i+1}). Note that $H_{(n)}$ and $V_{(n)}$ lie between α_n and α_1 , and β_n and β_1 , respectively.
- A generator is an unordered n -tuple of points such that each α and β circle contains exactly one. Generators can equivalently be viewed a \mathbb{Z} -linear combination of n points, or alternatively as permutations — for a permutation $\sigma \in S_n$ the generator x^σ is the unique generator with a point at each $\alpha_{\sigma(i)} \cap \beta_i$. In this paper we will use the convention that $[a_1 a_2 \dots a_n]$ denotes the permutation $\sigma \in S_n$ where $\sigma(j) = a_j$ for each j . For instance, Figure 1 shows the generator $x^{[51243]}$, which we will interchangeably denote as $[51243]$.
- A domain is a \mathbb{Z} -linear combination of square regions with the property that $\partial(\partial D \cap \alpha) = y - x$ for some generators x, y . We say that D is a domain from x to y , and write $D \in \mathcal{D}(x, y)$. D is said to be positive if none of its coefficients are negative, in which case we would write $D \in \mathcal{D}^+(x, y)$.
- Given $D \in \mathcal{D}(x, y)$, $E \in \mathcal{D}(y, z)$, we get a domain $D * E \in \mathcal{D}(x, z)$ by adding D and E as 2-chains.
- The constant domain from a generator x to itself is the domain $c_x \in \mathcal{D}(x, x)$ whose coefficients are zero in every square region.
- For every domain D , there is an associated integer $\mu(D)$ called its Maslov index, which satisfies:
 - $\mu(D * E) = \mu(D) + \mu(E)$.
 - For a positive domain D , $\mu(D) \geq 0$, with equality if and only if D is some constant domain.
 - For $D \in \mathcal{D}^+(x, y)$, $\mu(D) = 1$ if and only if D is a rectangle: that is, its bottom left and top right corners are coordinates of x , its bottom right and top left


corners are coordinates of y , and the other $n - 2$ coordinates of x and y agree and do not lie in D .

- $\mu(D) = k$ if and only if D can be decomposed (not necessarily uniquely) into k rectangles $D = R_1 * \dots * R_k$.


It will be particularly helpful to classify positive index 2 domains, which are exactly those that can be decomposed as two rectangles. Thus every positive index 2 domain $D \in \mathcal{D}^+(x, y)$ is a horizontal or vertical annulus



or two rectangles (overlapping or disjoint)



or a hexagon of the following shape



(Here the generator x is shown with \bullet while y is shown with \circ .) Note that while a horizontal or vertical annulus admits exactly one decomposition into rectangles, all the other positive index 2 domains admit exactly two.

Given a grid diagram \mathbb{G} , we define the complex of positive domains, on which our desired sign assignment can be constructed as a cochain.

Definition 2.1. The complex of positive domains $CD_* = CD_*(\mathbb{G}; \mathbb{Z}/2)$ is freely generated over $\mathbb{Z}/2$ by the positive domains, with the homological grading being the Maslov index

$$CD_k = \mathbb{Z}/2\{\{(x, y, D) \mid D \in \mathcal{D}^+(x, y), \mu(D) = k\}\}.$$

Sometimes the generators x, y will be omitted. The differential $\partial : CD_k \rightarrow CD_{k-1}$ of $D \in \mathcal{D}^+(x, y)$ is given by

$$\partial(D) = \sum_{R * E = D} E + \sum_{E * R = D} E,$$

where R is a rectangle, and E is a positive domain.

Note that CD_* is independent of the placement of the X 's and O 's.

Lemma 2.2. (CD_*, ∂) is a chain complex, that is, $\partial^2 = 0$.

Proof. Let R and S denote rectangles, then

$$\partial^2(D) = \sum_{R*S*E=D} E + \sum_{R*E*S=D} E + \sum_{S*E*R=D} E + \sum_{E*S*R=D} E.$$

The second and third terms cancel (modulo 2). If $R * S$ is a hexagon or two rectangles, then it has exactly one other decomposition $R * S = R' * S'$, so $R * S * E$ and $R' * S' * E$ cancel in the first term. Similarly, $E * R * S$ and $E * R' * S'$ cancel in the last term. Finally, if $R * S$ is not a hexagon or two rectangles, it must be a horizontal or vertical annulus, and then the terms $E * R * S$ and $R * S * E$ in the first and last term cancel, and so $\partial^2(D) = 0$. \square

We now compute the homology CD_* by constructing filtrations, for which we need the following fact. Given two generators x and y , we say that $x \leq y$ if there exists a positive domain from y to x that does not intersect the topmost row $H_{(n)}$ or rightmost column $V_{(n)}$ of the grid. It is clear that the set of generators with \leq is a partially ordered set (which actually coincides with the opposite of the Bruhat ordering on the symmetric group S_n , see [Manolescu and Sarkar 2021, Section 3.2]).

Proof of Proposition 1.1. The proof is nearly identical to the proof of [Manolescu and Sarkar 2021, Proposition 3.4], so we present the most relevant parts. To $D \in CD_*$ associate $A(D) \in \mathbb{N}^n$ by its coefficients in the rightmost vertical annulus. Note that here, unlike in [Manolescu and Sarkar 2021], $A(D)$ is an n -tuple, since there is no assumption that domains do not pass through the top right corner. By definition, the differential only preserves or lowers $A(D)$, so it is a filtration on CD_* . Now let CD_*^a be the associated graded complex in filtration grading $A(D) = a$.

Let $M(D) = \min\{\text{coordinates of } A(D)\}$ — by definition, a positive domain D contains exactly $M(D)$ copies of the rightmost vertical annulus $V_{(n)}$, so write $D = D' * M(D)V_{(n)}$. $A(D')$ contains a 0, so without loss of generality (since the differential of CD_*^a does not change $A(D)$ and thus does not change where the 0 is located) D' does not contain the top right corner. Now let $B(D) \in \mathbb{N}^{n-1}$ be the coordinates of D' in the top row (except the top right corner). Similarly, $B(D)$ is a filtration on the associated graded complex CD_*^a , so let $CD_*^{a,b}$ be the associated graded complex in grading $A(D) = a, B(D) = b$.

Now fix (a, b) and consider the differential ∂ on $CD_*^{a,b}$. Consider the following new filtration. For any domain $D \in \mathcal{D}^+(x, y)$ with $A(D) = a, B(D) = b$, let the generator y be its filtration grading. With respect to the aforementioned partial ordering of the generators, ∂ preserves or decreases y since we only consider removing domains that do not pass through the topmost row and rightmost column. Therefore y is a filtration grading, so let $CD_*^{a,b,y}$ be the associated graded complex

with respect to this filtration. Unless $a = (l, l, \dots, l)$, $b = 0$, and $y = x^{\text{Id}}$, the proof in [Manolescu and Sarkar 2021] shows that $\text{CD}_*^{a,b,y}$ is acyclic. When $a = (l, l, \dots, l)$, $b = 0$, and $y = x^{\text{Id}}$, the complex $\text{CD}_*^{a,b,y}$ has one generator (since x^{Id} is maximal), which is represented by the domain $lV_{(n)}$, lying in grading $2l$.

Finally, because the associated graded complex has homology only in even gradings, CD_* must have the same homology. \square

In order to later remove obstructions in grading 2, we now explicitly find the generator U of $H_2(\text{CD}_*)$. We define the following index 2 domains (see Figure 2):

- A_1, \dots, A_{n-1} where A_i is the vertical annulus in the $(n - i)$ -th column from the left from the generator $[n23 \dots (n - i) 1(n - i + 1) \dots (n - 1)]$ to itself, and A_0 is the rightmost vertical annulus from the identity generator x^{Id} to itself.
- B_1, \dots, B_{n-1} where B_i is the horizontal annulus in the $(n - i)$ -th row from the bottom from the generator $[(n - i + 1) 23 \dots (n - i)(n - i + 2) \dots n1]$ to itself, and B_0 is the topmost horizontal annulus from the identity generator x^{Id} to itself.
- C_1, \dots, C_{n-2} where C_i is a hexagon from the generator

$$[n23 \dots (n - i) 1(n - i + 1) \dots (n - 1)]$$

to the generator $[12 \dots (n - i - 1) n(n - i) \dots (n - 1)]$.




Figure 2. The domains $A_i, B_i, C_i, D_i, E_i, F_{i,j}$, and $G_{i,j}$ in the special case of a 6×6 grid, where each domain is drawn from a generator x (drawn with \bullet) to a generator y (drawn with \circ).

- D_1, \dots, D_{n-2} where D_i is a hexagon from the generator

$$[(n-i+1)23 \dots (n-i)(n-i+2) \dots n1]$$

to the generator $[12 \dots (n-i-1)(n-i+1) \dots n(n-i)]$.

- E_1, \dots, E_{n-2} where E_i is a hexagon from the generator

$$[12 \dots (n-i-1)n(n-i+1) \dots (n-1)(n-i)]$$

to the generator $[12 \dots (n-i-2)n(n-i) \dots (n-1)(n-i-1)]$.

- $F_{i,1}, \dots, F_{i,n-i-2}$ for each $i = 1, \dots, n-3$, where $F_{i,j}$ is a hexagon from the generator

$$[12 \dots (n-i-j-2)(n-i-j)(n-i)(n-i-j+1) \dots (n-i-1)(n-i+1) \dots n(n-i-j-1)]$$

to the generator

$$[12 \dots (n-i-j-2)(n-i+1)(n-i-j)(n-i-j+1) \dots (n-i)(n-i+2) \dots n(n-i-j-1)].$$

- $G_{i,1}, \dots, G_{i,n-i-2}$ for each $i = 1, \dots, n-3$, where $G_{i,j}$ is a hexagon from the generator

$$[12 \dots (n-i-j-2)n(n-i-j-1)(n-i-j+1) \dots (n-i-1)(n-i-j)(n-i) \dots (n-1)]$$

to the generator

$$[12 \dots (n-i-j-2)n(n-i-j) \dots (n-i)(n-i-j-1)(n-i+1) \dots (n-1)].$$

Let

$$U := \sum_{i=0}^{n-1} (A_i + B_i) + \sum_{i=1}^{n-2} (C_i + D_i) + \sum_{i=1}^{n-2} E_i + \sum_{i=1}^{n-3} \sum_{j=1}^{n-i-2} (F_{i,j} + G_{i,j})$$

Proposition 2.3. U is the generator of $H_2(\text{CD}_*)$

Proposition 2.3 will follow from the following computational lemmas.

Lemma 2.4. U is a cycle in CD_2 (that is, $\partial U = 0$).

Proof. We consider the possible rectangles that appear in ∂U , starting with the following rectangles that will be useful to name for the purposes of giving signs later.

- $R_{1,2}, \dots, R_{1,n-1}$ where $R_{1,i}$ is the $1 \times i$ rectangle from

$$[n23 \dots (n-i+1)1(n-i+2) \dots (n-1)]$$

to $[n23 \dots (n-i)1(n-i+1) \dots (n-1)]$, and $R_{1,1}$ is the 1×1 rectangle from x^{ld} to $[n23 \dots (n-1)1]$.

- $R_{2,2}, \dots, R_{2,n-1}$ where $R_{2,i}$ is the $1 \times i$ rectangle from

$$[12 \dots (n-i) n(n-i+1) \dots (n-1)]$$

to $[12 \dots (n-i-1) n(n-i) \dots (n-1)]$, and $R_{2,1}$ is the 1×1 rectangle from x^{ld} to $[12 \dots (n-2) n(n-1)]$

- $R_{3,2}, \dots, R_{3,n-1}$ where $R_{3,i}$ is the $i \times 1$ rectangle from

$$[(n-i+1) 23 \dots (n-i)(n-i+2) \dots n1]$$

to $[(n-i) 23 \dots (n-i-1)(n-i+1) \dots n1]$, and $R_{3,1} = R_{1,1}$.

- $R_{4,2}, \dots, R_{4,n-1}$ where $R_{4,i}$ is the $i \times 1$ rectangle from

$$[12 \dots (n-i)(n-i+2) \dots n(n-i+1)]$$

to $[12 \dots (n-i-1)(n-i+1) \dots n(n-i)]$, and $R_{4,1} = R_{2,1}$.

- $R_{5,1}, \dots, R_{5,n-2}$, where $R_{5,i}$ is the $1 \times i$ rectangle from

$$[12 \dots (n-i-2)(n-i) n(n-i+1) \dots (n-1)(n-i-1)]$$

to $[12 \dots (n-i-2) n(n-i) \dots (n-1)(n-i-1)]$.

- $R_{6,1}, \dots, R_{6,n-2}$, where $R_{6,i}$ is the $i \times 1$ rectangle from

$$[12 \dots (n-i-2) n(n-i-1)(n-i+1) \dots (n-1)(n-i)]$$

to $[12 \dots (n-i-2) n(n-i) \dots (n-1)(n-i-1)]$.

We cancel each of these rectangles in the boundary as follows (see Figure 3):

- $R_{1,1}$ occurs in ∂U twice, from ∂A_0 and ∂B_0 , so it cancels in ∂U . $R_{1,i}$ occurs in ∂A_{i-1} and ∂C_{i-1} for $i = 2, \dots, n-1$, so they also cancel in ∂U .




Figure 3. The rectangles $R_{1,i}, R_{2,i}, R_{3,i}, R_{4,i}, R_{5,i}, R_{6,i}$, where each rectangle is drawn from a generator x (drawn with \bullet) to a generator y (drawn with \circ).

- $R_{2,1}$ occurs in ∂U twice, from ∂C_1 and ∂D_1 , so it cancels in ∂U , $R_{2,n-1}$ occurs in ∂A_{n-1} and ∂E_{n-2} , and for $i = 2, \dots, n-2$, $R_{2,i}$ occurs in ∂C_i and ∂E_i , so they also cancel in ∂U .
- For $i = 2, \dots, n-1$, $R_{3,i}$ occurs in ∂B_{i-1} and ∂D_{i-1} .
- For $i = 2, \dots, n-2$, $R_{4,i}$ occurs in ∂D_i and ∂E_{i-1} .
- $R_{5,1}$ occurs in ∂E_1 and $\partial F_{1,1}$; for $i = 2, \dots, n-2$, $R_{5,i}$ occurs in ∂E_i and $\partial F_{1,i-1}$.
- $R_{6,1}$ occurs in ∂E_1 and $\partial G_{1,1}$; for $i = 2, \dots, n-2$, $R_{6,i}$ occurs in ∂E_i and $\partial G_{1,i-1}$.

Next, we consider the following rectangles:

- $P_{i,1} \dots P_{i,n-i-1}$ for each $i = 2 \dots n-2$, where $P_{i,j}$ is the $1 \times j$ rectangle from $[12 \dots (n-i-j-1)(n-i-j+1)(n-i+1)(n-i-j+2) \dots (n-i)(n-i+2) \dots n(n-i-j)]$

to $[12 \dots (n-i-j-1)(n-i+1)(n-i-j+1) \dots (n-i)(n-i+2) \dots n(n-i-j)]$, and $P_{1,j} = R_{5,j}$ for each $j = 1, \dots, n-2$.

- $Q_{i,1} \dots Q_{i,n-i-1}$ for each $i = 2 \dots n-2$, where $Q_{i,j}$ is the $j \times 1$ rectangle from $[12 \dots (n-i-j-1)n(n-i-j)(n-i-j+2) \dots (n-i)(n-i-j+1)(n-i+1) \dots (n-1)]$

to $[12 \dots (n-i-j-1)n(n-i-j+1) \dots (n-i)(n-i-j)(n-i+1) \dots (n-1)]$, and $Q_{1,j} = R_{6,j}$ for each $j = 1, \dots, n-2$.

We cancel each of these rectangles in the boundary as follows (see Figure 4):

- $P_{n-2,1}$ occurs in $\partial F_{n-3,1}$ and ∂B_{n-2} . For $i = 2, \dots, n-3$, $P_{i,j}$ occurs in $\partial F_{i-1,j}$ and either $\partial F_{i,j-1}$ if $j \geq 2$ or $\partial F_{i,1}$ if $j = 1$.
- $Q_{n-2,1}$ occurs in $\partial G_{n-3,1}$ and ∂A_{n-2} . For $i = 2, \dots, n-3$, $Q_{i,j}$ occurs in $\partial G_{i-1,j}$ and either $\partial G_{i,j-1}$ if $j \geq 2$ or $\partial G_{i,1}$ if $j = 1$.




Figure 4. The rectangles $P_{i,j}$ and $Q_{i,j}$ in the special case of a 6×6 grid, where each domain is drawn from a generator x (drawn with \bullet) to a generator y (drawn with \circ).

Finally, the remaining rectangles have the following form:

- $R'_{1,1} \dots R'_{1,n-1}$, where $R'_{1,i}$ is the $(n-i) \times 1$ rectangle from $[n23 \dots (n-i) 1(n-i+1) \dots (n-1)]$

to $[12 \dots (n-i) n(n-i+1) \dots (n-1)]$.

- $R'_{2,1} \dots R'_{2,n-1}$, where $R'_{2,i}$ is the $1 \times (n-i)$ rectangle from $[(n-i+1) 23 \dots (n-i)(n-i+2) \dots n1]$

to $[12 \dots (n-i)(n-i+2) \dots n(n-i+1)]$.

- $P'_{i,2} \dots P'_{i,n-i-1}$ for $i = 1 \dots n-3$, where $P'_{i,j}$ is the $j \times 1$ rectangle from $[12 \dots (n-i-j-1)(n-i)(n-i-j+1) \dots (n-i-1)(n-i+1) \dots n(n-i-j)]$ to $[12 \dots (n-i-j-1)(n-i+1)(n-i-j+1) \dots (n-i)(n-i+2) \dots n(n-i-j)]$, and $P'_{i,1} = P_{i,1}$.

- $Q'_{i,2} \dots Q'_{i,n-i-1}$ for $i = 1 \dots n-3$, where $Q'_{i,j}$ is the $1 \times j$ rectangle from $[12 \dots (n-i-j-1) n(n-i-j+1) \dots (n-i-1)(n-i-j)(n-i) \dots (n-1)]$ to $[12 \dots (n-i-j-1) n(n-i-j+1) \dots (n-i)(n-i-j)(n-i+1) \dots (n-1)]$, and $Q'_{i,1} = Q_{i,1}$.

We cancel each of these rectangles in the boundary as follows (see Figure 5):




Figure 5. The rectangles $R'_{1,i}$, $R'_{2,i}$, $P'_{i,j}$, $Q'_{i,j}$ in the special case of a 6×6 grid, where each domain is drawn from a generator x (drawn with \bullet) to a generator y (drawn with \circ).

- $R'_{1,1}$ occurs in ∂B_0 and ∂C_1 , $R'_{1,n-1}$ occurs in ∂A_{n-1} and ∂C_{n-2} . $R'_{1,i}$ occurs in ∂C_{i-1} and ∂C_i for $i = 2, \dots, n-2$.
- $R'_{2,1}$ occurs in ∂A_0 and ∂D_1 , $R'_{2,n-1}$ occurs in ∂B_{n-1} and ∂D_{n-2} . $R'_{2,i}$ occurs in ∂D_{i-1} and ∂D_i for $i = 2, \dots, n-2$.
- $P'_{i,n-i-1}$ occurs in $\partial F_{i,n-i-2}$ and ∂B_i . For $2 \leq j \leq n-i-2$, $P'_{i,j}$ occurs in $\partial F_{i,j-1}$ and $\partial F_{i,j}$.
- $Q'_{i,n-i-1}$ occurs in $\partial G_{i,n-i-2}$ and ∂A_i . For $2 \leq j \leq n-i-2$, $Q'_{i,j}$ occurs in $\partial G_{i,j-1}$ and $\partial G_{i,j}$.

Since these are the only rectangles produced by $\partial A_i, \partial B_i, \partial C_i, \partial D_i, \partial E_i, \partial F_{i,j}, \partial G_{i,j}$, we conclude that indeed $\partial U = 0$. \square

Lemma 2.5. *U is not homologous to zero in CD_* .*

Proof. Let r be the 2-cochain which is 1 on the rightmost vertical annulus from any generator to itself, and zero on all other domains; we will first show that r is a cocycle, at which point it suffices to show that $r(U) \neq 0$. Let E be an index 3 domain. If E does not contain the rightmost vertical annulus, then clearly $\delta r(E) = 0$. If E does contain the rightmost vertical annulus, then E can be written exactly two ways as the product of the rightmost vertical annulus $V_{(n)}$ with an index 1 domain: $E = D * V_{(n)} = V_{(n)} * D$. So $\delta r(E) = 0$ and therefore r is a cocycle, and $r(U) = 1$ since U contains exactly one copy of the rightmost vertical annulus. \square

Proof of Proposition 2.3. This immediately follows from Lemmas 2.4 and 2.5 and Proposition 1.1. \square

3. Sign assignments

In order to extend CD_* over \mathbb{Z} coefficients (and to frame some of the 0-dimensional moduli spaces in the Manolescu–Sarkar construction), we need a sign assignment for CD_* , which is a particular $\mathbb{Z}/2$ -valued 1-cochain on CD_* . The following conditions for a sign assignment ensures that 1-dimensional moduli spaces are frameable, since their boundaries must have opposite signs; see [Manolescu and Sarkar 2021] for more details, and note also that this agrees with the sign assignments defined in [Manolescu et al. 2007; Gallais 2008], though we are giving a new proof of their existence.

Definition 3.1. A sign assignment for \mathbb{G} is a $\mathbb{Z}/2$ -valued 1-cochain s on CD_* such that:

- (1) (square rule) If D_1, D_2, D_3, D_4 are distinct rectangles and $D_1 * D_2 = D_3 * D_4 = E$ which is not an annulus, then $s(D_1) + s(D_2) = s(D_3) + s(D_4) + 1$.

- (2) (annuli) If D_1, D_2 are rectangles and $D_1 * D_2$ is a vertical annulus, then $s(D_1) = s(D_2) + 1$. If D_1, D_2 are rectangles such that $D_1 * D_2$ is a horizontal annulus, then $s(D_1) = s(D_2)$.

In order to prove that such a sign assignment exists, we will show that the 2-cocycle that we hypothesize to be δs is indeed a 2-coboundary.

Lemma 3.2. *Let T be the 2-cochain with the following values:*

- (1) (square rule) *For any index 2 domain D that is not an annulus, $T(D) = 1$.*
 (2) (annuli) *$T(V) = 1$ for all vertical annuli V , and $T(H) = 0$ for all horizontal annuli H .*

Then T is a 2-coboundary.

Proof. First, we show that T is a cocycle. Let E be any index 3 domain; we must show that $\langle T, \partial E \rangle = 0$. For every decomposition $E = D * A$, where A is a vertical or horizontal annulus, there is a corresponding decomposition $E = A * D$, so that A occurs an even number of times in ∂E . It now suffices to show that ∂E contains an even number of every other type of index 2 domain.

To every index 3 domain E from a generator x to a generator y , consider a graph with vertices x at level 3, y at level 0, and edges down 1 level corresponding to each way to break off an index 1 domain (see Figure 6 for an example of such a graph). Then each level 2 vertex has an index 2 domain to y , which decomposes into rectangles exactly two ways, so each level 2 vertex has downward degree 2, and each level 1 vertex has an index 2 domain from x , which decomposes into rectangles exactly two ways, so each level 1 vertex has upward degree 2. Therefore there are the same number of level 2 and level 1 vertices, so since each index 2 domain that shows up in ∂E corresponds to a level 2 or 1 vertex, there are an even number of index 2 domains. Since an even number of these are annuli, we must therefore have an even number of hexagons. This shows that $\langle T, \partial E \rangle = 0$, as desired.

By Propositions 1.1 and 2.3 it now suffices to show that $T(U) = 0$ where U is the generator of $H_2(\text{CD}_*)$. By definition, U consists of n annuli A_i , n annuli B_i , $n - 2$ hexagons C_i , $n - 2$ hexagons D_i , $n - 2$ hexagons E_i , $\binom{n-2}{2}$ hexagons $F_{i,j}$, and $\binom{n-2}{2}$ hexagons $G_{i,j}$, so for any T satisfying the conditions of Lemma 3.2,

$$T(U) \equiv n + 3(n - 2) + 2\binom{n-2}{2} \equiv 0 \pmod{2}$$

so that T is indeed a coboundary. □

Lemma 3.3. *Let T be the 2-coboundary from Lemma 3.2. Then $T = \delta s$ if and only if s is a sign assignment.*

Proof. This is clear from the definitions. □




Figure 6. An example of a positive index 3 domain from a generator x (drawn with \bullet) to a generator y (drawn with \circ), along with the graph defined in the proof of Proposition 1.3. The generators z_i are given by \circ on the i -th rectangle from the left and \bullet on the other two, while the generators w_i are given by \bullet on the i -th rectangle from the left and \circ on the other two.

Proof of Theorem 1.2. Existence immediately follows from Lemmas 3.2 and 3.3. For uniqueness, suppose $T = \delta s = \delta s'$. Then $\delta(s - s') = 0$, so $s - s'$ is a 1-cocycle, which is cohomologous to zero by Proposition 1.1, so there is a 0-cochain g such that $s = s' + \delta g$. □

Given a sign assignment s , we can use it to redefine CD_* in \mathbb{Z} coefficients:

Definition 3.4. $CD_*(\mathbb{G}; \mathbb{Z})$ is freely generated over \mathbb{G} by the positive domains, with the homological grading being the Maslov index. The differential $\partial : CD_k \rightarrow CD_{k-1}$ of $D \in \mathcal{D}^+(x, y)$ is given by

$$\partial(D) = \sum_{R * E = D} (-1)^{s(R)} E + (-1)^k \sum_{E * R = D} (-1)^{s(R)} E,$$

where R is a domain of index 1 from x to some generator z and E is a positive domain from z to y .

We now have analogues of Lemma 2.2 and Proposition 1.1 in \mathbb{Z} coefficients, in the following lemma and Proposition 1.3, respectively.

Lemma 3.5. (CD_*, ∂) is a chain complex.

Proof. The proof is similar to the proof of Lemma 2.2, except we must keep track of signs. □

Proof of Proposition 1.3. The proof is similar to the proof of Proposition 1.1. Specifically, our proof of Proposition 1.1 over $\mathbb{Z}/2$ adapts the proof of [Manolescu and Sarkar 2021, Proposition 3.4]. This proof is over \mathbb{Z} , and a similar adaptation will prove Proposition 1.3. □

4. The obstruction complex with partitions

The moduli spaces in the construction of the 1-flow category require more than just positive domains. Since periodic domains (annuli) can bubble, Manolescu and Sarkar [2021] introduces a new complex to keep track of the bubbles — since there are n different types of bubbles (corresponding to bubbling of the j -th horizontal or vertical annulus) which can be at the same or different heights, these correspond to n -tuples of ordered partitions.

It is convenient to use both of the following equivalent definitions of an ordered partition of a positive integer N (and when $N = 0$, a partition of N is the empty set).

- An ordered partition λ is a tuple of nonnegative integers $\lambda = (\lambda_1, \dots, \lambda_m)$ such that $N = \sum \lambda_i$ (m is called the *length* of the partition, and is denoted $l(\lambda)$).
- An ordered partition λ is a tuple $\epsilon(\lambda) = (\epsilon_1(\lambda), \dots, \epsilon_{N-1}(\lambda)) \in \{0, 1\}^{N-1}$, where an ϵ_i equaling 1 indicates a split at that point. For instance, the ordered partitions $(1, 1, 1)$, $(1, 2)$, $(2, 1)$, and (3) of 3 are written $(1, 1)$, $(1, 0)$, $(0, 1)$, and $(0, 0)$.

Besides annuli bubbling off (the second type of terms that will be in the differential — the first being terms in the differential of CD_*), there are two other boundary degenerations that occur with existing bubbles. Bubbles of the same type may come to the same height (the third type of term), and bubbles may go to height $\pm\infty$ (the fourth and final type of term). The corresponding changes to the partitions are described below.

Definition 4.1. The following changes to an ordered partition will describe the differential terms (see [Manolescu and Sarkar 2021, Definitions 4.1, 4.2, 4.3] for more details):

- A unit enlargement (at position k) increases N by 1 and adds a 1 to the tuple λ (at position k). The set of unit enlargements of λ is denoted $UE(\lambda)$.
- An elementary coarsening (at position k) replaces both terms λ_k and λ_{k+1} with one term $\lambda_k + \lambda_{k+1}$. The set of elementary coarsenings of λ is denoted $EC(\lambda)$.
- An initial reduction removes λ_1 (and decreases N by λ_1), and a final reduction removes λ_m (and decreases N by λ_m). The set of initial reductions (respectively, final reductions) of λ is denoted $IR(\lambda)$ (respectively, $FR(\lambda)$), where we consider both sets empty if $N = 0$.

We are now ready to define the complex of domains with partitions, CDP_* .

Definition 4.2. The complex of positive domains with $CDP_* = CDP_*(\mathbb{G}; \mathbb{Z}/2)$ is freely generated by triples of the form $D, \vec{N}, \vec{\lambda}$, where:

- $D \in \mathcal{D}^+(x, y)$ is a positive domain.
- $\vec{N} \in \mathbb{N}^n$ is an n -tuple of nonnegative integers, $\vec{N} = (N_1, \dots, N_n)$.

- $\vec{\lambda} = (\lambda_1, \dots, \lambda_n)$ is an n -tuple of ordered partitions for $\lambda_j = (\lambda_{j,1}, \dots, \lambda_{j,m_j})$, an ordered partition of N_j .

We denote $|\vec{N}| := \sum_{j=1}^n N_j$, and define $|\vec{l}(\vec{\lambda})| := \sum_{j=1}^n l(\lambda_j)$ be the total length of $\vec{\lambda}$. The grading of $(D, \vec{N}, \vec{\lambda})$ is given by the Maslov index of D plus $|\vec{l}(\vec{\lambda})|$. The differential is given by the sum of the following four terms.

- Type I terms, given by taking out a rectangle from D , just like in the differential of CD_* .
- Type II terms, given by taking out a vertical or horizontal annulus passing through O_j from D and performing a unit enlargement to λ_j .
- Type III terms, given by an elementary coarsening of one of the partitions λ_j .
- Type IV terms, given by taking the initial or final reduction of one of the partitions λ_j .

Precisely, we can write $\partial = \partial_1 + \partial_2 + \partial_3 + \partial_4$ where

$$\partial_1(D, \vec{N}, \vec{\lambda}) = \sum_{R * E = D} (E, \vec{N}, \vec{\lambda}) + \sum_{E * R = D} (E, \vec{N}, \vec{\lambda}),$$

$$\partial_2(D, \vec{N}, \vec{\lambda}) = \sum_{j=1}^n \sum_{D = E * H_j \text{ or } E * V_j} \sum_{\lambda'_j \in \text{UE}(\lambda_j)} (E, \vec{N} + \vec{e}_j, \vec{\lambda}'),$$

$$\partial_3(D, \vec{N}, \vec{\lambda}) = \sum_{j=1}^n \sum_{\lambda'_j \in \text{EC}(\lambda_j)} (D, \vec{N}, \vec{\lambda}'),$$

$$\partial_4(D, \vec{N}, \vec{\lambda}) = \sum_{j=1}^n \sum_{\lambda'_j \in \text{IR}(\lambda_j)} (D, \vec{N} - \lambda_{j,1} \vec{e}_j, \vec{\lambda}') + \sum_{j=1}^n \sum_{\lambda'_j \in \text{FR}(\lambda_j)} (D, \vec{N} - \lambda_{j,m_j} \vec{e}_j, \vec{\lambda}').$$

As in Definition 2.1, R is a rectangle, and the annuli H_j, V_j are the ones passing through the j -th O marking. We also use $\vec{\lambda}' := (\lambda_1, \dots, \lambda_{j-1}, \lambda'_j, \lambda_{j+1}, \dots, \lambda_n)$, and $\vec{e}_j := (0, \dots, 0, 1, 0, \dots, 0)$ with the 1 in the j -th position.

It will help us to classify the lower grading generators, that is, generators of $CDP_0, CDP_1, CDP_2, CDP_3$ (see [Manolescu and Sarkar 2021, Section 4.2] for details).

- (0) CDP_0 is generated by the constant domains with no partitions $(c_x, 0, 0)$ for some generator x .
- (1) CDP_1 is generated by rectangles with no partitions $(R, 0, 0)$ as well as triples of the form $(c_x, N \vec{e}_j, (N))$ for a constraint domain c_x .
- (2) CDP_2 is generated by Maslov index 2 domains with no partition $(D, 0, 0)$ (for a classification of the kinds of domains D , see above or [Ozsváth et al. 2015]), triples of the form $(R, N \vec{e}_j, (N))$ for a rectangle R , or a constant domain with partitions of total length 2. Specifically, we can have triples of the form $(c_x, N \vec{e}_j + M \vec{e}_k, ((N), (M)))$ (where $j \neq k$), or $(c_x, (N + M) \vec{e}_j, (N, M))$.

(3) Finally, CDP_3 is generated by Maslov index 3 domains with no partition, Maslov index 2 domains with a partition of the form $(D, N\vec{e}_j, (N))$, rectangles with a partition of total length 2, and constant domains with partitions of total length 3, which has the following cases:

- $(c_x, N_j \vec{e}_j + N_k \vec{e}_k + N_l \vec{e}_l, ((N_j), (N_k), (N_l)))$ for j, k, l distinct.
- $(c_x, (N_j + M_j) \vec{e}_j + N_k \vec{e}_k, ((N_j, M_j), (N_k)))$ for j, k distinct.
- $(c_x, (N_j + M_j + P_j) \vec{e}_j, (N_j, M_j, P_j))$.

Lemma 4.3. (CDP_*, ∂) is a chain complex.

Proof. The proof follows a similar case analysis to [Manolescu and Sarkar 2021, Lemma 4.4]. Write $\partial = \partial_1 + \partial_2 + \partial_3 + \partial_4$, where ∂_k is the type k term in the differential. Since ∂_1 is just the differential from CD_* , we have by Lemma 2.2 that $\partial_1^2 = 0$. Now for ∂_2^2 , the terms will correspond to removing two annuli (and doing two unit enlargements). If the annuli pass through two different O_i and O_j , then the corresponding term shows up twice, once in each order. If the annuli pass through the same O_j , then the corresponding term also shows up twice — once for each order in doing the unit enlargements. So, $\partial_2^2 = 0$. We can similarly show that

$$\partial_3^2 = 0, \quad \partial_1 \partial_2 + \partial_2 \partial_1 = 0, \quad \partial_1 \partial_3 + \partial_3 \partial_1 = 0, \quad \partial_2 \partial_3 + \partial_3 \partial_2 = 0, \quad \partial_1 \partial_4 + \partial_4 \partial_1 = 0$$

by doing the respective operations in two different orders.

Now consider $\partial_2 \partial_4 + \partial_4 \partial_2$, the terms of which correspond to a unit enlargement and an initial or final reduction, in either order. If one is done to λ_i and another to λ_j where $i \neq j$, then the two commute and cancel just like before. If both are done to λ_i , then all terms follow one of these cases:

- A unit enlargement not at the beginning, followed by an initial reduction. This cancels with the initial reduction followed by doing the enlargement one place earlier.
- A unit enlargement not at the end, followed by a final reduction. This cancels with the final reduction followed by the same enlargement.
- A unit enlargement at the beginning, followed by an initial reduction; or a unit enlargement at the end, followed by a final reduction. These cancel with each other.

Finally, consider the last terms of ∂^2 , $\partial_4^2 + \partial_3 \partial_4 + \partial_4 \partial_3$. Again there are some special types of terms:

- The elementary coarsening of λ_i by combining the first two parts, followed by a initial reduction of λ_i , cancels with two initial reductions of λ_i .
- The elementary coarsening of λ_i by combining the last two parts, followed by a final reduction of λ_i , cancels with two final reductions of λ_i .

where all the other terms cancel by doing the operations in two different orders. \square

We would like to compute the homology of CDP_* using successive filtrations, as in the proof of Proposition 1.1.

Proposition 4.4. *There is a filtration on CDP_* such that the associated graded has homology $(\mathbb{Z}/2)^{2^n} \otimes (\mathbb{Z}/2)[U]$.*

Proof. We again follow the proof of [Manolescu and Sarkar 2021, Proposition 4.6]. We can filter the complex CDP'_* in several steps. First we filter CDP_* by the quantity $A(D) \in \mathbb{N}^n$ which are the coefficients of D in the rightmost column. As in the proof of Proposition 1.1, we can assume without loss of generality that the minimum of $A(D)$ occurs in the top right corner, and then filter the associated graded CDP_*^a by $B(D) \in \mathbb{N}^{n-1}$ which are the coefficients of D in the topmost row. In the associated graded $\text{CDP}_*^{a,b}$, there are no type II terms in the differential, since such terms must decrease either A or B . Since $|\vec{N}|$ is kept constant by type I and type III terms and decreased by type IV terms, it is a filtration on $\text{CDP}_*^{a,b}$, so filtering by $|\vec{N}|$ and (as in the proof of Proposition 1.1) the end generator y gives a direct sum of complexes $\text{CDP}_*^{a,b,y,\vec{N}}$.

When $a \neq (l, l, \dots, l)$ or $b \neq 0$ or $y \neq x^{\text{Id}}$, filtering by the total length of $\vec{\lambda}$ removes all type III terms and keeps all type I terms, so $\text{CDP}_*^{a,b,y,\vec{N}}$ is a direct sum of complexes $\text{CD}_*^{a,b,y}$ which were all shown to be acyclic in the proof of [Manolescu and Sarkar 2021, Proposition 3.4]. Additionally, when $a = (l, l, \dots, l)$, $b = 0$, $y = x^{\text{Id}}$, and at least one $N_j > 1$, every generator of $\text{CDP}_*^{a,b,y,\vec{N}}$ is represented by some $(D, \vec{N}, \vec{\lambda})$ where $D = kV_n$, so we only have type III terms. The partitions of N_j are given by $(\epsilon_1, \dots, \epsilon_{N_j-1})$, where the elementary coarsenings just change a 1 to a 0. This gives a hypercube complex, which is acyclic. Therefore, we are only left with the associated graded complexes $\text{CDP}_*^{a,b,y,\vec{N}}$ where $a = (l, l, \dots, l)$, $b = 0$, $y = x^{\text{Id}}$, and every N_j is 0 or 1. \square

Corollary. $H_k(\text{CDP}_*; \mathbb{Z}/2)$ has rank at most

$$\sum_{l=0}^{\lfloor k/2 \rfloor} \binom{n}{k-2l}.$$

In the proof of Proposition 2.3, we found a cocycle that detects the generator of $H_2(\text{CD}_*)$. We will use a similar procedure to compute $H_0(\text{CDP}_*)$ through $H_3(\text{CDP}_*)$.

Proof of Theorem 1.4. ($k = 0$). This case is clear.

($k = 1$). The n generators of $H_1(\text{AssGr}(\text{CDP}_*))$ are the triples

$$g_j := (c_{x^{\text{Id}}}, \vec{e}_j, (1)),$$

which are still cycles in CDP_1 (because their initial and final reductions cancel). It will suffice to show that there exist n 1-cocycles r_j such that $r_j(g_k) = 1$ if and only

if $j = k$. Let f_j be the 1-cochain in CD_* such that $\delta f_j(D) = 1$ if and only if D is the vertical annulus V_j or the horizontal annulus H_j where f_j exists by Proposition 2.3 because the 2-cocycle which is 1 on V_j and H_j and zero on every other index 2 domain is a coboundary, since it is zero on the generator U of $H_2(CD_*)$. We can extend f_j to CDP_* by setting it equal to zero on all triples $(c_x, N\vec{e}_j, (N))$. Let N_j be the 1-cocycle that is the value of N_j in the triple $(D, \vec{N}, \vec{\lambda})$, and

$$r_j := N_j + f_j.$$

To show that r_j is a cocycle, we consider all possible triples $(D, \vec{N}, \vec{\lambda})$ in grading 2. If $N_j = 0$ and D is not the annulus V_j or H_j , then by definition $\delta r_j(D, \vec{N}, \vec{\lambda}) = 0$. If $N_j = 0$ and $D = V_j$ or H_j , then $\vec{N} = 0$ and

$$\delta r_j(D, 0, 0) = N_j(c_x, \vec{e}_j, (1)) + (f_j(R_1) + f_j(R_2)) = 1 + 1 = 0 \pmod{2},$$

where $D = R_1 * R_2$ is the decomposition into rectangles. Finally, if $N_j = M > 0$, there are three cases:

- D is a rectangle. In this case $\vec{N} = M\vec{e}_j$ and $\lambda_i = (M)$, so the initial and final reduction of λ_i cancel, and the only other differential terms are removing D . If D is a rectangle from x to y , then

$$\begin{aligned} \delta r_j(D, M\vec{e}_j, (M)) &= N_j(c_x, M\vec{e}_j, (M)) + N_j(c_y, M\vec{e}_j, (M)) \\ &= M + M = 0 \pmod{2}. \end{aligned}$$

- Some $N_k > 0$, where $k \neq j$. Then D must be a constant domain, and both λ_j and λ_k are length 1 partitions, so their initial and final reductions all cancel.
- D is a constant domain and $\lambda_j = (M_1, M_2)$ is a length 2 partition. In this case we have all of the type III and type IV differentials, which gives

$$\begin{aligned} \delta r_j(c_x, (M + N)\vec{e}_j, (M, N)) \\ &= N_j(c_x, M\vec{e}_j, (M)) + N_j(c_x, N\vec{e}_j, (N)) + N_j(c_x, (M + N)\vec{e}_j, (M + N)) \\ &= M + N + (M + N) = 0 \pmod{2}. \end{aligned}$$

Therefore r_j is a cocycle for each j , and by definition $r_j(g_j) = 1$ if and only if $j = k$, so the g_j are in fact the generators of $H_1(CDP_*)$.

($k = 2$). Now consider 2-cocycles. $(\mathbb{Z}/2)^{\binom{n}{2}}$ of the generators of $H_2(\text{AssGr}(CDP_*))$ are the triples

$$g_{j,k} := (c_{x^{\text{id}}}, \vec{e}_j + \vec{e}_k, ((1), (1))),$$

which are similarly still cycles in CDP_2 . The final generator will be given by a slight modification U' of $(U, 0, 0)$, where U is the generator of $H_2(CD_*)$. The boundary of U in CDP_* contains only pairs of triples of the form

$$(c_{x^j}, \vec{e}_j, (1)) \quad \text{and} \quad (c_{y^j}, \vec{e}_j, (1))$$

corresponding to type II differentials on the annuli A_j and B_j . Here we have that x^j and y^j are, respectively, the generators $[n23 \dots (n-j) 1(n-i+1) \dots (n-1)]$ and $[(n-j+1) 23 \dots (n-j)(n-j+2) \dots n1]$, as depicted in Figure 2. For each j , the generators x^j and y^j each have a planar domain (that is, a domain that does not intersect the topmost row or rightmost column of the grid), $D_{j,1}$ and $D_{j,2}$ respectively, from the identity generator x^{1d} . From Figure 2, we see that $D_{j,2}$ is the reflection of $D_{j,1}$ about the diagonal from the bottom left to the top right of the grid, so that $(-D_{j,1}) * D_{j,2}$ is an even index planar domain from x^j to y^j . This domain decomposes into an even number of planar rectangles $\pm R_{jk}$ (where each R_{jk} is positive), so that adding each $(R_{jk}, \vec{e}_j, (1))$ to $(U, 0, 0)$ will cancel the rest of its boundary, making a cycle U' .

In the proof of Proposition 2.3, we used the 2-cocycle r which is 1 on the rightmost vertical annulus and zero on every other 2-chain. Extending r to CDP_* by setting it equal to zero on every 2-chain with $|\vec{N}| > 0$ still gives a cocycle, since CDP_* has no new ways to create an annulus in the boundary, and we still have that $r(U') = 1$, while all of the $r(g_{j,k}) = 0$. Now it suffices to find $r_{j,k}$ such that $r_{j,k}(U') = 0$ for all j, k , and $r_{j,k}(g_{l,m}) = 1$ if and only if $\{l, m\} = \{j, k\}$. Let f_j be the 1-cocycles defined in the proof of (1), and let

$$f_j^k(R, \vec{N}, \vec{\lambda}) = N_k f_j(R),$$

where R is a rectangle and $\vec{\lambda}$ has total length 1 (and $f_j^k = 0$ on all other 2-chains). Now let $N_j N_k$ be the 2-cocycle that is the product of the values of N_j and N_k for a triple $(D, \vec{N}, \vec{\lambda})$, and let

$$r_{j,k} := N_j N_k + f_j^k + f_k^j.$$

To show that $r_{j,k}$ is a cocycle, we consider all possible triples $(D, \vec{N}, \vec{\lambda})$ in grading 3. If $N_j = 0$ (respectively, $N_k = 0$) and D does not contain the annulus V_j or H_j (respectively, V_k or H_k), then by definition $\delta r_{j,k}(D, \vec{N}, \vec{\lambda}) = 0$. If $N_j = 0$, $N_k > 0$ (or vice versa), and D contains V_j or H_j , then $D = V_j$ or H_j and all $N_l = 0$ for $l \neq k$, so that $\delta r_{j,k}(D, \vec{N}, \vec{\lambda}) = 0$ similarly to the proof of (1). Finally, if $N_j = M_j > 0$ and $N_k = M_k > 0$, there are three cases:

- D is a rectangle. In this case $\vec{N} = M_j \vec{e}_j + M_k \vec{e}_k$, $\lambda_j = (M_j)$, and $\lambda_k = (M_k)$, so the initial and final reductions of λ_j and λ_k cancel, and the only other differential terms are removing D . If D is a rectangle from x to y , then

$$\begin{aligned} &\delta r_{j,k}(D, \vec{N}, \vec{\lambda}) \\ &= N_j N_k (c_x, M_j \vec{e}_j + M_k \vec{e}_k, ((M_j), (M_k))) + N_i N_j (c_y, M_j \vec{e}_j + M_k \vec{e}_k, ((M_j), (M_k))) \\ &= M_i M_j + M_i M_j = 0 \pmod{2}. \end{aligned}$$

- Some $N_l > 0$, where $l \neq j, k$. Then D must be a constant domain, and all of λ_j , λ_k , and λ_l are length 1 partitions, so their initial and final reductions all cancel.

- D is a constant domain and $\lambda_j = (M_{j,1}, M_{j,2})$ is a length 2 partition (or symmetrically, $\lambda_k = (M_{k,1}, M_{k,2})$). In this case the initial and final reductions of λ_k cancel, but we have all of the type III and type IV differentials of λ_j , which give

$$\begin{aligned} \delta r_{j,k}(c_x, (M_{j,1} + M_{j,2}) \vec{e}_j + M_k \vec{e}_k, ((M_{j,1}, M_{j,2}), (M_k))) \\ = N_j N_k(c_x, M_{j,1} \vec{e}_j + M_k \vec{e}_k, ((M_{j,1}), (M_k))) \\ + N_j N_k(c_x, M_{j,2} \vec{e}_j + M_k \vec{e}_k, ((M_{j,2}), (M_k))) \\ + N_j N_k(c_x, (M_{j,1} + M_{j,2}) \vec{e}_j + M_k \vec{e}_k, ((M_{j,1} + M_{j,2}), (M_k))) \\ = M_k(M_{j,1} + M_{j,2} + (M_{j,1} + M_{j,2})) = 0 \pmod{2}. \end{aligned}$$

Therefore $r_{j,k}$ is a cocycle for all j, k , and by definition $r_{j,k}(U) = 0$. U' only adds an even number of planar rectangles with partitions that can contribute to f_j^k or f_k^j , no two of which can ever form an annulus, so $r_{j,k}(U') = 0$. Finally, by definition $r_{j,k}(g_{l,m}) = 1$ if and only if $\{l, m\} = \{j, k\}$, so these (along with U') are in fact the generators of $H_2(\text{CDP}_*)$.

($k = 3$). $\binom{n}{3}$ of the generators of $H_3(\text{AssGr}(\text{CDP}_*))$ are the triples

$$g_{j,k,l} := (c_{x^{\text{ld}}}, \vec{e}_j + \vec{e}_k + \vec{e}_l, ((1), (1), (1))),$$

which are similarly still cycles in CDP_3 . The other n generators are the triples U' obtained from U' by performing unit enlargements on N_j and adding the triples $(R_{jk}, 2\vec{e}_j, (2))$ defined previously (for this fixed j). Let $V_{(n)}$ be the rightmost vertical annulus and define the cochain

$$rr_j(D, \vec{N}, \vec{\lambda}) := \begin{cases} 0, & D \text{ does not contain } V_{(n)}, \\ r_j(D * -V_{(n)}, \vec{N}, \vec{\lambda}), & D \text{ contains } V_{(n)}, \end{cases}$$

where r_j is the 1-cocycle from the proof of (1). To show that rr_j are cocycles, we consider all $\delta rr_j(D, \vec{N}, \vec{\lambda})$ for triples $(D, \vec{N}, \vec{\lambda}) \in \text{CDP}_4$. If D does not contain $V_{(n)}$, then this quantity is zero by definition. If D contains $V_{(n)}$, then its Maslov index is at least 2, so that we have the following cases:

- D is an index 4 domain. In this case, $\vec{N} = 0$, so let $E = D * (-V_{(n)})$. If E is also an annulus V_k , then

$$\begin{aligned} \delta rr_j(D, \vec{N}, \vec{\lambda}) \\ = rr_j(A_1 * V_{(n)}, 0, 0) + rr_j(A_2 * V_{(n)}, 0, 0) + rr_j(E * B_1, 0, 0) + rr_j(E * B_2, 0, 0) \\ + rr_j(V_{(n)}, \vec{e}_k, (1)) + rr_j(E, \vec{e}_n, (1)) \quad (\text{where } A_1 * A_2 = E, B_1 * B_2 = V_{(n)}) \\ = r_j(A_1, 0, 0) + r_j(A_2, 0, 0) + r_j(c_x, \vec{e}_k, (1)) \\ + (r_j(B_1, 0, 0) + r_j(B_2, 0, 0) + r_j(c_x, \vec{e}_n, (1))) \quad (\text{added if and only if } k = n) \\ = 0 \end{aligned}$$

by definition of r_j (since this is just $\delta r_j(V_k)$ with possibly $\delta r_j(V_{(n)})$ added if $k = n$). If E is not an annulus, we similarly have

$$\begin{aligned} \delta rr_j(D, \vec{N}, \vec{\lambda}) &= rr_j(A_1 * V_{(n)}, 0, 0) + rr_j(A_2 * V_{(n)}, 0, 0) + rr_j(E * B_1, 0, 0) \\ &\quad + rr_j(E * B_2, 0, 0) + rr_j(E, \vec{e}_n, (1)) \quad (\text{where } A_1 * A_2 = E, B_1 * B_2 = V_{(n)}) \\ &= r_j(A_1, 0, 0) + r_j(A_2, 0, 0) + r_j(c_x, \vec{e}_k, (1)) = 0. \end{aligned}$$

• $D = R * V_{(n)}$ is an index 3 domain, where R is a rectangle from a generator x to a generator y . In this case, $\vec{N} = N\vec{e}_k$ and $\vec{\lambda} = \lambda_k = (N)$. Suppose that in the domain D , $V_{(n)} = A * B$, and $R = B$ (or symmetrically, $R = A$). In this case,

$$\begin{aligned} \delta rr_j(D, \vec{N}, \vec{\lambda}) &= rr_j(R * A, N\vec{e}_k, (N)) + rr_j(A * R, N\vec{e}_k, (N)) + rr_j(R, N\vec{e}_k + e_n, ((N), (1))) \\ &= r_j(c_x, N\vec{e}_k, (N)) + r_j(c_y, N\vec{e}_k, (N)) = 0. \end{aligned}$$

by definition of r_j . If R is not A or B , we similarly have

$$\begin{aligned} \delta rr_j(D, \vec{N}, \vec{\lambda}) &= rr_j(R * A, N\vec{e}_k, (N)) + rr_j(R * B, N\vec{e}_k, (N)) \\ &\quad + rr_j(V_{(n)}, N\vec{e}_k, (N)) \quad (\text{at the generator } x) \\ &\quad + rr_j(V_{(n)}, N\vec{e}_k, (N)) \quad (\text{at } y) \\ &\quad + rr_j(R, N\vec{e}_k + \vec{e}_n, ((N), (1))) \\ &= r_j(c_x, N\vec{e}_k, (N)) + r_j(c_y, N\vec{e}_k, (N)) = 0. \end{aligned}$$

• D is an index 2 domain. In this case, we have that $D = V_{(n)}$, $|\vec{\lambda}| = 2$, and $\delta rr_j(D, \vec{N}, \vec{\lambda}) = \delta r_j(c_x, \vec{N}, \vec{\lambda})$ since the type I and type II terms cannot possibly have an annulus; note that we previously showed this expression to be zero in showing that r_j is a cocycle.

Therefore rr_j is a cocycle, and by construction $rr_j(U'_k) = 1$ if and only if $k = j$, and all of the $rr_j(g_{k,l,m}) = 0$. It remains to find cocycles $r_{j,k,l}$ such that $r_{j,k,l}(U'_m) = 0$ for all j, k, l, m , and $r_{j,k,l}(g_{m,p,q}) = 1$ if and only if $\{m, p, q\} = \{j, k, l\}$. Let f_j be the 1-cocycles defined in the proof of (1), and let

$$f_j^{k,l}(R, \vec{N}, \vec{\lambda}) = N_k N_l f_j(R),$$

where R is a rectangle and λ_k, λ_l are both length 1 partitions (and $f_j^{k,l} = 0$ on all other 3-chains). Now let $N_j N_k N_l$ be the 3-cocycle that is the product of the values of N_j, N_k , and N_l for a triple $(D, \vec{N}, \vec{\lambda})$, and let

$$r_{j,k,l} = N_j N_k N_l + f_j^{k,l} + f_k^{j,l} + f_l^{j,k}.$$

To show that $r_{j,k,l}$ is a cocycle, we consider all possible triples $(D, \vec{N}, \vec{\lambda})$ in grading 4. If $N_j = 0$ (respectively, $N_k = 0$ or $N_l = 0$) and D does not contain the annulus V_j

or H_j (respectively, V_k or H_k , or V_l or H_l), then by definition $\delta r_{j,k,l}(D, \vec{N}, \vec{\lambda}) = 0$. If $N_j = 0$, $N_k > 0$, and $N_l > 0$ (or symmetrically, any other case where exactly one is zero), and D contains V_j or H_j , then $D = V_j$ or H_j and all $N_m = 0$ for $m \neq k, l$, so that $\delta r_{j,k,l}(D, \vec{N}, \vec{\lambda}) = 0$ similarly to the proof of (1). Finally, if $N_j = M_j > 0$, $N_k = M_k > 0$, and $N_l = M_l > 0$, there are three cases:

- D is a rectangle. In this case $\vec{N} = M_j \vec{e}_j + M_k \vec{e}_k + M_l \vec{e}_l$, $\lambda_j = (M_j)$, $\lambda_k = (M_k)$, and $\lambda_l = (M_l)$, so the initial and final reductions of λ_j , λ_k and λ_l cancel, and the only other differential terms are removing D . If D is a rectangle from x to y , then

$$\begin{aligned} \delta r_{j,k,l}(D, \vec{N}, \vec{\lambda}) &= N_j N_k N_l (c_x, M_j \vec{e}_j + M_k \vec{e}_k + M_l \vec{e}_l, ((M_j), (M_k), (M_l))) \\ &\quad + N_j N_k N_l (c_y, M_j \vec{e}_j + M_k \vec{e}_k + M_l \vec{e}_l, ((M_j), (M_k), (M_l))) \\ &= M_j M_k M_l + M_j M_k M_l = 0 \pmod{2}. \end{aligned}$$

- Some $N_m > 0$, where $m \neq j, k, l$. Then D must be a constant domain, and all of $\lambda_j, \lambda_k, \lambda_l$, and λ_m are length 1 partitions, so their initial and final reductions all cancel.
- D is a constant domain and $\lambda_j = (M_{j,1}, M_{j,2})$ is a length 2 partition (or symmetrically, $\lambda_k = (M_{k,1}, M_{k,2})$ or $\lambda_l = (M_{l,1}, M_{l,2})$). In this case the initial and final reductions of λ_k and λ_l cancel, but we have all of the type III and type IV differentials of λ_j , which give

$$\begin{aligned} &\delta r_{j,k,l}(c_x, (M_{j,1} + M_{j,2}) \vec{e}_j + M_k \vec{e}_k + M_l \vec{e}_l, ((M_{j,1}, M_{j,2}), (M_k), (M_l))) \\ &= N_j N_k N_l (c_x, M_{j,1} \vec{e}_j + M_k \vec{e}_k + M_l \vec{e}_l, ((M_{j,1}), (M_k), (M_l))) \\ &\quad + N_j N_k N_l (c_x, M_{j,2} \vec{e}_j + M_k \vec{e}_k + M_l \vec{e}_l, ((M_{j,2}), (M_k), (M_l))) \\ &\quad + N_j N_k N_l (c_x, (M_{j,1} + M_{j,2}) \vec{e}_j + M_k \vec{e}_k + M_l \vec{e}_l, ((M_{j,1} + M_{j,2}), (M_k), (M_l))) \\ &= M_k M_l (M_{j,1} + M_{j,2} + (M_{j,1} + M_{j,2})) = 0 \pmod{2} \end{aligned}$$

Therefore $r_{j,k,l}$ are cocycles, satisfying $r_{j,k,l}(F_m'') = 0$ for all j, k, l, m , and $r_{j,k,l}(c_{x^{\text{id}}}, \vec{e}_m + \vec{e}_p + \vec{e}_q, ((1), (1))) = 1$ if and only if $\{m, p, q\} = \{j, k, l\}$. \square

5. Sign assignments for domains with partitions

Similarly to Section 3, we find the criteria that the coboundary of a sign assignment for CDP_* must satisfy:

Definition 5.1. A sign assignment for CDP_* is a 1-cochain s on CDP_* such that:

- (1) $\delta s(D, 0, 0) = 1$ for any index 2 domain D that is not an annulus.
- (2) $\delta s(V, 0, 0) = 1$ for any vertical annulus V , and $\delta s(H, 0, 0) = 0$ for any horizontal annulus H .
- (3) $\delta s(R, (0, N\vec{e}_j, (N))) = 0$ for any rectangle R , any $N > 0$, and any j .

- (4) $\delta s(c_x, N\vec{e}_j + M\vec{e}_k, ((N), (M))) = 0$ for any constant domain c_x , any $N, M > 0$, and any j, k .
- (5) $\delta s(c_x, (N + M)\vec{e}_j, ((N), (M))) = 0$ for any constant domain c_x , any $N, M > 0$, and any j .

Proof of Theorem 1.5. Let T be the 2-cochain with values given by (1)–(5) of Definition 5.1. To see that T is a cocycle, we evaluate δT on all triples $(D, \vec{N}, \vec{\lambda})$ in grading 3. These are given by:

- $(D, 0, 0)$ where D is an index 3 domain. The proof of Lemma 3.3 shows that the contributions to δT by Type I differential terms all cancel. If D does not contain an annulus, these are all the differential terms. If D does contain an annulus $A = H_j$ or V_j , we can write $D = R * A$ for a rectangle R , so that the type II differential term gives $(R, (0, \vec{e}_j), (1))$, which does not contribute to δT by (3).
- $(D, N\vec{e}_j, (N))$ where D is an index 2 domain. Here the initial and final reduction of the partition both give $(D, 0, 0)$ so their contributions to δT cancel. The decompositions of D into rectangles do not contribute to δT by condition (3), and again if D is not an annulus, then these are the only other boundary terms. If D is an annulus, then either $D = H_j$, $D = V_j$, or D is some other annulus H_k or V_k . In the latter case, the type II differential term gives $(c_x, N\vec{e}_j + \vec{e}_k, ((N), (1)))$ which does not contribute to δT by (4). In the former case, the type II differential gives two terms, $(c_x, (N + 1)\vec{e}_j), (1, N))$ and $(c_x, (N + 1)\vec{e}_j, (N, 1))$, which do not contribute to δT by (5).
- $(R, \vec{N}, \vec{\lambda})$ where $R \in \mathcal{D}^+(x, y)$ is a rectangle and $\vec{\lambda}$ has total length 2. Here the type I differential removes R two ways, leaving either $(c_x, M\vec{e}_j + N\vec{e}_k, ((M), (N)))$ (and the corresponding term for c_y , which do not contribute to δT by (4)), or $(c_x, (M + N)\vec{e}_j), (M, N))$ (and the corresponding term for c_y , which do not contribute by (5)). All type III and type IV terms do not contribute by (3). Since R cannot possibly contain an annulus, there are no further terms so $\delta T(R, \vec{N}, \vec{\lambda}) = 0$.
- $(c_x, \vec{N}, \vec{\lambda})$ where c_x is a constant domain and $\vec{\lambda}$ has total length 3. None of these terms contribute to δT by (4) and (5).

Hence T is a cocycle, so it remains to show T is zero on every generator of $H_2(\text{CDP}_*)$ listed in the proof of Theorem 1.4. By definition, we have that every $T(c_x, \vec{e}_j + \vec{e}_k, ((1), (1))) = 0$. Also, $T(U, 0, 0) = 0$ by Lemma 3.2, so $T(U') = 0$ by condition (3), since these are the only types of triples added to $(U, 0, 0)$. Therefore T must be the zero cocycle by Theorem 1.4, so $T = \delta s$ for some s . The values s_j uniquely determine the $H^1(\text{CDP}_*)$ class of s by Theorem 1.4, so at that point s is unique up to gauge equivalence (like sign assignments for CD_*). □

There are two types of triples in grading 1 — rectangles with no partitions and constant domains with a length 1 partition. By uniqueness, the sign of a rectangle

with no partition in CDP_* agrees with the sign of that rectangle in CD_* , so it remains to compute the signs of constant domains with a length 1 partition.

Proposition 5.2. *For any constant domain c_x and any $N > 0$,*

$$s(c_x, N\vec{e}_j, (N)) = Ns_j \pmod{2}.$$

Proof. We first show that the sign is independent of the generator x . Let $R \in \mathcal{D}^+(x, y)$ be a rectangle. By (3) of Definition 5.1, we have

$$0 = \delta s(R, N\vec{e}_j, (N)) = s(c_x, N\vec{e}_j, (N)) + s(R, 0, 0) + s(R, 0, 0) + s(c_y, N\vec{e}_j, (N))$$

so that $s(c_x, N\vec{e}_j, (N)) = s(c_y, N\vec{e}_j, (N))$, and given any domain from x to y , we find a decomposition into rectangles and repeatedly apply this equation. Therefore we can now assume without loss of generality that $x = x^{\text{Id}}$. We will use the uniqueness of s up to the values s_j to proceed by induction on N . The base case is clear, and by (5) of Definition 5.1 we must have that

$$\begin{aligned} 0 &= \delta s(c_x, N\vec{e}_j, (1, N - 1)) \\ &= s(c_x, \vec{e}_j, (1)) + s(c_x, (N - 1)\vec{e}_j, (N - 1)) + s(c_x, N\vec{e}_j, (N)) \\ &= s_j + (N - 1)s_j \pmod{2} \end{aligned}$$

by the inductive hypothesis, so that $s(c_x, N\vec{e}_j, (N)) = Ns_j \pmod{2}$, which completes the induction. □

Remark 5.3. It would suffice by uniqueness to define a sign assignment on CDP_* by defining a sign assignment on CD_* and extending it by Proposition 5.2. Doing so would give another proof of Theorem 1.5.

Again, now that we have a sign assignment s , we can extend CDP_* to \mathbb{Z} coefficients. As in CD_* , the sign associated to breaking off a rectangle is the sign of the rectangle $s(R)$ given by the sign assignment. We now describe the sign of the other differential terms.

Definition 5.4. Let s be a sign assignment for CDP_* .

- Given an ordered partition λ and the unit enlargement

$$\lambda' = (\lambda_1, \dots, \lambda_{k-1}, 1, \lambda_k, \dots, \lambda_m),$$

the sign of the unit enlargement is

$$s(\lambda, \lambda') = k + 1 \pmod{2}.$$

- Given an ordered partition λ and the elementary coarsening

$$\lambda' = (\lambda_1, \dots, \lambda_{k-1}, \lambda_k + \lambda_{k+1}, \lambda_{k+2}, \dots, \lambda_m),$$

the sign of the elementary coarsening is

$$s(\lambda, \lambda') = k \pmod{2}.$$

- Given an ordered partition $\lambda = (\lambda_1, \dots, \lambda_m)$ and its initial reduction λ' , the sign of the reduction is given by

$$s(\lambda, \lambda') = \lambda_1 s_j \pmod{2}.$$

The sign of its final reduction is given by the same expression, with λ_m replacing λ_1 .

Definition 5.5. The complex of positive domains that have the partitions $\text{CDP}_* = \text{CDP}_*(\mathbb{G}; \mathbb{Z})$ is freely generated by triples of the form $D, \vec{N}, \vec{\lambda}$, where:

- $D \in \mathcal{D}^+(x, y)$ is a positive domain.
- $\vec{N} \in \mathbb{N}^n$ is an n -tuple of nonnegative integers, $\vec{N} = (N_1, \dots, N_n)$.
- $\vec{\lambda} = (\lambda_1, \dots, \lambda_n)$ is an n -tuple of ordered partitions, where $\lambda_j = (\lambda_{j,1}, \dots, \lambda_{j,m_j})$ is an ordered partition of N_j .

The grading of $(D, \vec{N}, \vec{\lambda})$ is given by the Maslov index of D plus the sum of the lengths of the λ_j (which is referred to as the total length of $\vec{\lambda}$). The differential is given by four terms, $\partial = \partial_1 + \partial_2 + \partial_3 + \partial_4$, where

$$\begin{aligned} \partial_1(D, \vec{N}, \vec{\lambda}) &= \sum_{R * \vec{E} = D} (-1)^{s(R)}(E, \vec{N}, \vec{\lambda}) + (-1)^{\mu(D)} \sum_{E * R = D} (-1)^{s(R)}(E, \vec{N}, \vec{\lambda}), \\ \partial_2(D, \vec{N}, \vec{\lambda}) &= (-1)^{\mu(D)} \sum_{j=1}^n (-1)^{l(\lambda_1) + \dots + l(\lambda_{j-1})} \sum_{\substack{D = E * H_j \text{ (horizontal)} \\ \lambda'_j \in \text{UE}(\lambda_j)}} (-1)^{1+s(\lambda_j, \lambda'_j)} (E, \vec{N} + \vec{e}_j, \vec{\lambda}') \\ &\quad + (-1)^{\mu(D)} \sum_{j=1}^n (-1)^{l(\lambda_1) + \dots + l(\lambda_{j-1})} \sum_{\substack{D = E * V_j \text{ (vertical)} \\ \lambda'_j \in \text{UE}(\lambda_j)}} (-1)^{s(\lambda_j, \lambda'_j)} (E, \vec{N} + \vec{e}_j, \vec{\lambda}'), \\ \partial_3(D, \vec{N}, \vec{\lambda}) &= (-1)^{\mu(D)} \sum_{j=1}^n (-1)^{l(\lambda_1) + \dots + l(\lambda_{j-1})} \sum_{\lambda'_j \in \text{EC}(\lambda_j)} (-1)^{s(\lambda_j, \lambda'_j)}(D, \vec{N}, \vec{\lambda}'), \\ \partial_4(D, \vec{N}, \vec{\lambda}) &= (-1)^{\mu(D)} \sum_{j=1}^n (-1)^{l(\lambda_1) + \dots + l(\lambda_{j-1})} \sum_{\lambda'_j \in \text{IR}(\lambda_j)} (-1)^{s(\lambda_j, \lambda'_j)}(D, \vec{N} - \lambda_{j,1} \vec{e}_j, \vec{\lambda}') \\ &\quad + (-1)^{\mu(D)} \sum_{j=1}^n (-1)^{l(\lambda_1) + \dots + l(\lambda_j)} \sum_{\lambda'_j \in \text{FR}(\lambda_j)} (-1)^{s(\lambda_j, \lambda'_j)}(D, \vec{N} - \lambda_{j,m_j} \vec{e}_j, \vec{\lambda}'). \end{aligned}$$

Remark 5.6. In the case that all $s_j = 0$, these signs agree with the signs of [Manolescu and Sarkar 2021, Definitions 4.1–4.3], with the exception of the type II differential.

Lemma 5.7. (CDP_*, ∂) is a chain complex.

Proof. The proof is similar to that of [Manolescu and Sarkar 2021, Lemma 4.4], which is the same case analysis of Lemma 4.3, except where we keep track of signs. In the case of

$$(\partial_4)^2 + \partial_3\partial_4 + \partial_4\partial_3 = 0,$$

we still have all but two cases canceling in pairs by reversing the order of the two operations. These two cases are:

- Two initial reductions and an elementary coarsening at the beginning, followed by an initial reduction. The former has sign

$$\lambda_1 s_j + \lambda_2 s_j \pmod{2}$$

and the latter has sign

$$1 + (\lambda_1 + \lambda_2) s_j \pmod{2},$$

which is the opposite sign.

- Two final reductions and an elementary coarsening at the end, followed by a final reduction. Note that final reductions have an extra sign of $l(\lambda_j)$ compared to initial reductions, so that including this extra sign, the former has sign

$$l(\lambda_j) + (l(\lambda_j) - 1) + \lambda_m s_j + \lambda_{m-1} s_j \pmod{2}$$

and the latter has sign

$$(l(\lambda_j) - 1) + (l(\lambda_j) - 1) + (\lambda_{m-1} + \lambda_m) s_j \pmod{2},$$

which is the opposite sign.

Finally, although we still have $\partial_2\partial_4 + \partial_4\partial_2 = 0$, the change to the sign of the type II differential gives a new set of cancellations

$$\partial_1^2 + \partial_2\partial_4 + \partial_4\partial_2 = 0.$$

For this case, suppose $D = A * E = E * A$ is the domain where $A = R * S$ is an annulus.

- If A is a vertical annulus V_j , then $s(R) + s(S) = 1$, so that removing R then S from the front has sign 1, while the type II differential that produces a unit enlargement at the front of λ_j followed by the initial reduction of λ_j has sign 0. Also, removing S then R from the back has sign 0 (since the Maslov index of the domain decreases once), while the type II differential that produces a unit enlargement at the end of λ_j followed by the final reduction of λ_j has sign $l(\lambda_j) + 1 + l(\lambda_j) = 1 \pmod{2}$.
- If A is a horizontal annulus H_j , then $s(R) + s(S) = 0$, so that removing R then S from the front has sign 0, while the type II differential that produces a unit enlargement at the front of λ_j followed by the initial reduction of λ_j has

sign 1. Also, removing S then R from the back has sign 1 (since the Maslov index of the domain decreases once), while the type II differential that produces a unit enlargement at the end of λ_j followed by the final reduction of λ_j has sign $l(\lambda_j) + l(\lambda_j) = 0 \pmod{2}$. \square

The analogue of Proposition 4.4 also holds over \mathbb{Z} .

Proposition 5.8. *There is a filtration on CDP_* such that the associated graded has homology $\mathbb{Z}^{2^n} \otimes \mathbb{Z}[U]$. In particular, $H_k(\text{CDP}_*)$ has rank at most*

$$\sum_{l=0}^{\lfloor k/2 \rfloor} \binom{n}{k-2l}.$$

Proof. The proof is identical to the proof of Proposition 4.4. \square

Acknowledgements

The author would like to thank Sucharit Sarkar for many helpful conversations, as well as the referee for many helpful suggestions and corrections. This work was supported by an NSF grant DMS-2136090.

References

- [Chongchitmate and Ng 2013] W. Chongchitmate and L. Ng, “An atlas of Legendrian knots”, *Exp. Math.* **22**:1 (2013), 26–37. MR Zbl
- [Cohen et al. 1995] R. L. Cohen, J. D. S. Jones, and G. B. Segal, “Floer’s infinite-dimensional Morse theory and homotopy theory”, pp. 297–325 in *The Floer memorial volume*, edited by H. Hofer et al., Progr. Math. **133**, Birkhäuser, Basel, 1995. MR Zbl
- [Gallais 2008] É. Gallais, “Sign refinement for combinatorial link Floer homology”, *Algebr. Geom. Topol.* **8**:3 (2008), 1581–1592. MR Zbl
- [Khandhawit and Ng 2010] T. Khandhawit and L. Ng, “A family of transversely nonsimple knots”, *Algebr. Geom. Topol.* **10**:1 (2010), 293–314. MR Zbl
- [Lobb et al. 2020] A. Lobb, P. Orson, and D. Schütz, “Khovanov homotopy calculations using flow category calculus”, *Exp. Math.* **29**:4 (2020), 475–500. MR Zbl
- [Manolescu and Ozsváth 2010] C. Manolescu and P. Ozsváth, “Heegaard Floer homology and integer surgeries on links”, preprint, 2010. To appear in *Geom. Topol.* arXiv 1011.1317
- [Manolescu and Sarkar 2021] C. Manolescu and S. Sarkar, “A knot Floer stable homotopy type”, preprint, 2021. arXiv 2108.13566
- [Manolescu et al. 2007] C. Manolescu, P. Ozsváth, Z. Szabó, and D. Thurston, “On combinatorial link Floer homology”, *Geom. Topol.* **11** (2007), 2339–2412. MR Zbl
- [Manolescu et al. 2009a] C. Manolescu, P. Ozsváth, and S. Sarkar, “A combinatorial description of knot Floer homology”, *Ann. of Math. (2)* **169**:2 (2009), 633–660. MR Zbl
- [Manolescu et al. 2009b] C. Manolescu, P. Ozsváth, and D. Thurston, “Grid diagrams and Heegaard Floer invariants”, preprint, 2009. To appear in *Ann. Math.* arXiv 0910.0078
- [Ng et al. 2008] L. Ng, P. Ozsváth, and D. Thurston, “Transverse knots distinguished by knot Floer homology”, *J. Symplectic Geom.* **6**:4 (2008), 461–490. MR Zbl

- [Ozsváth and Szabó 2004a] P. Ozsváth and Z. Szabó, “Holomorphic disks and knot invariants”, *Adv. Math.* **186**:1 (2004), 58–116. MR Zbl
- [Ozsváth and Szabó 2004b] P. Ozsváth and Z. Szabó, “Holomorphic disks and three-manifold invariants: properties and applications”, *Ann. of Math. (2)* **159**:3 (2004), 1159–1245. MR Zbl
- [Ozsváth and Szabó 2004c] P. Ozsváth and Z. Szabó, “Holomorphic disks and topological invariants for closed three-manifolds”, *Ann. of Math. (2)* **159**:3 (2004), 1027–1158. MR Zbl
- [Ozsváth and Szabó 2008] P. Ozsváth and Z. Szabó, “Holomorphic disks, link invariants and the multi-variable Alexander polynomial”, *Algebr. Geom. Topol.* **8**:2 (2008), 615–692. MR Zbl
- [Ozsváth et al. 2008] P. Ozsváth, Z. Szabó, and D. Thurston, “Legendrian knots, transverse knots and combinatorial Floer homology”, *Geom. Topol.* **12**:2 (2008), 941–980. MR Zbl
- [Ozsváth et al. 2015] P. S. Ozsváth, A. I. Stipsicz, and Z. Szabó, *Grid homology for knots and links*, Math. Surv. Monogr. **208**, Amer. Math. Soc., Providence, RI, 2015. MR Zbl
- [Rasmussen 2003] J. A. Rasmussen, *Floer homology and knot complements*, Ph.D. thesis, Harvard University, 2003, available at <https://www.proquest.com/docview/305332635>.
- [Sarkar 2011] S. Sarkar, “Grid diagrams and the Ozsváth–Szabó tau-invariant”, *Math. Res. Lett.* **18**:6 (2011), 1239–1257. MR Zbl

Received May 13, 2024. Revised August 1, 2024.

YAN TAO
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF CALIFORNIA, LOS ANGELES
LOS ANGELES, CA
UNITED STATES
ytao@math.ucla.edu

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Fakultät für Mathematik
Universität Wien
Vienna, Austria
matthias.aschenbrenner@univie.ac.at

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Atsushi Ichino
Department of Mathematics
Kyoto University
Kyoto 606-8502, Japan
atsushi.ichino@gmail.com

Robert Lipshitz
Department of Mathematics
University of Oregon
Eugene, OR 97403
lipshitz@uoregon.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Dimitri Shlyakhtenko
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
shlyakht@ipam.ucla.edu

Ruixiang Zhang
Department of Mathematics
University of California
Berkeley, CA 94720-3840
ruixiang@berkeley.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2024 is US \$645/year for the electronic version, and \$875/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2024 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 331 No. 2 August 2024

Long-time behavior of awesome homogeneous Ricci flows	187
ROBERTO ARAUJO	
Sobolev norms of L^2 -solutions to the nonlinear Schrödinger equation	217
ROMAN V. BESSONOV and SERGEY A. DENISOV	
Extrinsic polyharmonic maps into the sphere	259
ALI FARDOUN, STEFANO MONTALDO, CEZAR ONICIUC and ANDREA RATTO	
Balanced homogeneous harmonic maps between cones	283
BRIAN FREIDIN	
Rank growth of elliptic curves in S_4 - and A_4 -quartic extensions of the rationals	331
DANIEL KELIHER	
Obstruction complexes in grid homology	353
YAN TAO	