

*Pacific
Journal of
Mathematics*

ON DISJOINT STATIONARY SEQUENCES

MAXWELL LEVINE

Volume 332 No. 1

September 2024

ON DISJOINT STATIONARY SEQUENCES

MAXWELL LEVINE

We answer a question of Krueger by obtaining disjoint stationary sequences on successive cardinals. The main idea is an alternative presentation of a mixed support iteration, using it more explicitly as a variant of Mitchell forcing. We also use a Mahlo cardinal to obtain a model in which $\aleph_2 \notin I[\aleph_2]$ and there is no disjoint stationary sequence on \aleph_2 , answering a question of Gilton.

1. Introduction and background

In order to develop the theory of infinite cardinals, set theorists study a variety of objects that can potentially exist on these cardinals. The objects of interest for us are called *disjoint stationary sequences*. These were introduced by Krueger to answer a question of Abraham and Shelah about forcing clubs through stationary sets [2]. Beginning in joint work with Friedman, Krueger wrote a series of papers in this area, connecting a wide range of concepts and answering seemingly unrelated questions of Foreman and Todorćević [8; 17; 18; 19; 20; 21]. Our purpose is to further develop this area.

Krueger's new arguments generally hinged on the behavior of two-step iterations of the form $\text{Add}(\tau) * \mathbb{P}$. In order to extend the application of these arguments as widely as possible, Krueger developed the notion of mixed support forcing [18; 21], which had apparently been part of the folklore for some time. These forcings are to some extent an analog of the forcing that Mitchell used to obtain the tree property at double successors of regular cardinals. Their most notable feature is the appearance of quotients insofar as the forcings took the form $\mathbb{M} \simeq \overline{\mathbb{M}} * \text{Add}(\tau) * \mathbb{E}$ where $\overline{\mathbb{M}}$ is a partial mixed support iteration. The appearance of $\text{Add}(\tau)$ after the initial component, together with the preservation properties of the quotient \mathbb{E} , allowed Krueger's new arguments to go through various complicated constructions. Mixed support iterations have found several applications since [10], particularly in regard to guessing models [22].

Our main idea is to use a version of Mitchell forcing to accomplish the task of a mixed support iteration. Specifically, we prove that this version of Mitchell

MSC2020: 03E35, 03E55.

Keywords: set theory, forcing, independence, large cardinals.

forcing takes the form $\mathbb{M} \simeq \overline{\mathbb{M}} * \text{Add}(\tau) * \mathbb{E}$.¹ The trick used to obtain this structural property goes back to Mitchell's thesis and is also reminiscent of the one used by Cummings, S. D. Friedman, Magidor, Rinot, and Sinapova [5] to demonstrate that subtle variations in the definitions of Mitchell forcing — up to merely shifting a Lévy collapse by a single coordinate — can substantially alter the properties of the forcing extension. The benefit of the forcing used here is that it comes with a projection analysis of the sort that Abraham used for Mitchell forcing [1]. Both the forcing itself and its quotients are projections of products of the form $\mathbb{A} \times \mathbb{T}$ where \mathbb{A} has a good chain condition and \mathbb{T} has a good closure property. This allows us to obtain preservation properties conveniently, without having to delve into too many technical details. Abraham in fact used this projection analysis to extend Mitchell's result to successive cardinals. This is exactly what we do here for disjoint stationary sequences, answering the first component of a question of Krueger [21, Question 12.8]:

Theorem 1. *Suppose $\lambda_1 < \lambda_2$ are two Mahlo cardinals in V . Then there is a forcing extension in which there are disjoint stationary sequences on \aleph_2 and \aleph_3 .*

We lay out the basic definition and concepts in the following subsections and then develop the proof in Section 2. We also achieve one of Krueger's separations for successive cardinals, which answers a component of another one of his questions [21, Question 12.9]:

Theorem 2. *Suppose $\lambda_1 < \lambda_2$ are two Mahlo cardinals in V . Then there is a forcing extension in which for $\mu \in \{\aleph_1, \aleph_2\}$, there are stationarily many $N \in [H(\mu^+)]^\mu$ that are internally stationary but not internally club.*

The last main result is motivated by work of Gilton and Krueger, who answered a question from [5] by obtaining stationary reflection for subsets of $\aleph_2 \cap \text{cof}(\omega)$ together with failure of approachability at \aleph_2 (i.e., $\aleph_2 \notin I[\aleph_2]$) using disjoint stationary sequences [10]. This result used the fact that the existence of a disjoint stationary sequence implies failure of approachability. Gilton asked for the exact consistency strength of the failure of approachability at \aleph_2 together with the nonexistence of a disjoint stationary sequence on \aleph_2 [9, Question 9.0.15]. (He pointed out that Cox found this separation using PFA [3].) It is known that the failure of approachability requires the consistency strength of a Mahlo cardinal since \square_τ holds if τ^+ is not Mahlo in L [16] and \square_τ implies the approachability property $\tau^+ \in I[\tau^+]$ [6]. In Section 3 we show that a Mahlo cardinal is sufficient for the separation:

Theorem 3. *Suppose that λ is Mahlo in V . Then there is a forcing extension in which $\aleph_2 \notin I[\aleph_2]$ and there is no disjoint stationary sequence on \aleph_2 .*

¹The extent to which all variations of these forcings are equivalent or not is left as a loose end. Here we only deal with the case where the two-step iteration $\text{Add}(\tau) * \mathbb{P}$ takes the form $\text{Add}(\tau) * \text{Col}(\mu, \delta)$.

Disjoint stationary sequences are known to be interpretable in terms of canonical structure (see Fact 6 below), and the main idea for Theorem 3 is a simple master condition argument that exploits this connection.

1.1. Basic definitions. We assume familiarity with the basics of forcing and large cardinals. We use the following conventions: If \mathbb{P} is a forcing poset, then $p \leq q$ for $p, q \in \mathbb{P}$ means that p is stronger than q . We say that \mathbb{P} is κ -closed if for all $\leq_{\mathbb{P}}$ -decreasing sequences $\langle p_\xi : \xi < \tau \rangle$ with $\tau < \kappa$, there is a lower bound p , i.e., $p \leq p_\xi$ for all $\xi < \tau$. (Not all authors use this formulation of κ -closedness.) We say that \mathbb{P} has the κ -chain condition if all antichains $A \subseteq \mathbb{P}$ have cardinality strictly less than κ . All posets considered will be separative. Now we give our main definitions:

Definition 4. Given a regular cardinal μ , a *disjoint stationary sequence* on μ^+ is a sequence $\langle \mathcal{S}_\alpha : \alpha \in S \rangle$ such that

- $S \subseteq \mu^+ \cap \text{cof}(\mu)$ is stationary,
- \mathcal{S}_α is a stationary subset of $P_\mu(\alpha)$ for all $\alpha \in S$,
- $\mathcal{S}_\alpha \cap \mathcal{S}_\beta = \emptyset$ if $\alpha \neq \beta$.

We write $\text{DSS}(\mu^+)$ to say that there is a disjoint stationary sequence on μ^+ .

Definition 5. Given an uncountable regular κ and a set $N \in [H(\Theta)]^\kappa$,² we say:

- N is *internally unbounded* if for all $x \in P_\kappa(N)$, there is an $M \in N$ such that $x \subseteq M$.
- N is *internally stationary* if $P_\kappa(N) \cap N$ is stationary in $P_\kappa(N)$.
- N is *internally club* if $P_\kappa(N) \cap N$ is club in $P_\kappa(N)$.
- N is *internally approachable* if there is an increasing and continuous chain $\langle M_\xi : \xi < \kappa \rangle$ such that $|M_\xi| < \kappa$ and $\langle M_\eta : \eta < \xi \rangle \in M_{\xi+1}$ for all $\xi < \kappa$ such that $N = \bigcup_{\xi < \kappa} M_\xi$.

Although disjoint stationary sequences may seem unrelated to the separation of variants of internal approachability, there are deep connections here, for example:

Fact 6 (Krueger [21]). If μ is regular and $2^\mu = \mu^+$, then $\text{DSS}(\mu^+)$ is equivalent to the existence of a stationary set $U \subseteq [H(\mu^+)]^\mu$ such that every $N \in U$ is internally unbounded but not internally club.

1.2. Projections and preservation lemmas. Technically speaking, our main goal is to show that certain forcing quotients behave nicely. We will make an effort to demonstrate the preservation properties of these quotients directly. These quotients will be defined in terms of projections:

²See Jech's book [15] for details on stationary sets.

Definition 7. If \mathbb{P}_1 and \mathbb{P}_2 are posets, a *projection* is an onto map $\pi : \mathbb{P}_1 \rightarrow \mathbb{P}_2$ such that

- $p \leq q$ implies that $\pi(p) \leq \pi(q)$,
- if $r \leq \pi(p)$, then there is some $q \leq p$ such that $\pi(q) = r$.

A projection is *trivial* if for all $p, q \in \mathbb{P}_1$, if $\pi(p)$ and $\pi(q)$ are compatible, then p and q are compatible.

Trivial projections are basically isomorphisms:

Fact 8. If $\pi : \mathbb{P}_1 \rightarrow \mathbb{P}_2$ is a trivial projection, then $\mathbb{P}_1 \simeq \mathbb{P}_2$, that is, \mathbb{P}_1 and \mathbb{P}_2 are forcing-equivalent.

For our purposes, we are interested in the preservation of stationary sets. The chain condition gives us preservation fairly straightforwardly. The following fact is implicit in parts of the literature, and a version of it can be found in the form of Proposition 26.

Fact 9. If \mathbb{P} has the μ -chain condition and $S \subset P_\mu(X)$ is stationary, then \mathbb{P} forces that S is stationary in $P_\mu(X)$.

However, we must place demands on our stationary sets in order for them to be preserved by closed forcings.

Definition 10. Consider a regular uncountable cardinal μ and a stationary set $S \subset P_\mu(X)$. We say that S is *internally approachable of length τ* if for all $N \in S$ with $N \prec H(X)$, there is a continuous chain of elementary submodels $\langle M_i : i < \tau \rangle$ such that: $N = \bigcup_{i < \tau} M_i$ and for all $i < \tau$, $\langle M_i : i < j \rangle \in N$. In this case we write $S \subseteq \mathcal{JA}(\tau)$.

Here we are following Krueger's convention [21], which withholds the requirement that $|M_i| < \mu$ for $i < \tau$.

Fact 11. If $S \subset P_\mu(X) \cap \mathcal{JA}(\tau)$ is an internally approachable stationary set, $\tau < \mu$, and \mathbb{P} is μ -closed, then \mathbb{P} forces that S is stationary in $P_\mu(X)^V$.³

1.3. Costationarity of the ground model. The notion of ground model costationarity is a key ingredient in arguments pertaining to disjoint stationary sequences. It will specifically give us the disjointness, since we will be picking stationary sets that are not added by initial segments of these forcings.

Gitik obtained the classical result:

Fact 12 (Gitik [12]). If $V \subset W$ are models of ZFC with the same ordinals, $W \setminus V$ contains a real, and κ is a regular cardinal in W such that $(\kappa^+)^W \leq \lambda$, then $P_\kappa^W(\lambda) \setminus V$ is stationary as a subset of $P_\kappa(\lambda)$ in the model W .

³See [7] for discussion of related facts.

Because we will need Fact 11, we will actually use Krueger's refinement of Gitik's theorem:

Fact 13 (Krueger [21]). Suppose $V \subset W$ are models of ZFC with the same ordinals, $W \setminus V$ contains a real, μ is a regular cardinal in W , and $X \in V$ is such that $(\mu^+)^W \subseteq X$, and that in W , Θ is a regular cardinal such that $X \subset H(\Theta)$. Then in W the set $\{N \in P_\mu(H(\Theta)) \cap \mathcal{J}\mathcal{A}(\omega) : N \cap X \notin V\}$ is stationary.

2. The new Mitchell forcing

2.1. Defining the forcing. Here we will illustrate the basic idea of this paper by using our new take on Mitchell forcing to prove a known result:

Fact 14 (Krueger [21]). If λ is a Mahlo cardinal and $\mu < \lambda$ is regular, there is a forcing extension in which $2^\omega = \mu^+ = \lambda$ and there is a disjoint stationary sequence on λ .

Specifically, we will define a forcing $\mathbb{M}^+(\tau, \mu, \lambda)$ such that the model W in Fact 14 can be realized as an extension by $\mathbb{M}^+(\omega, \mu, \lambda)$.

For standard technical reasons, we define a poset isomorphic to $\text{Add}(\tau, \lambda)$:

Definition 15. Given a regular τ and a set of ordinals Y , we let $\text{Add}^*(\tau, Y)$ be the poset consisting of partial functions $p : \{\delta \in Y : \delta \text{ is inaccessible}\} \times \tau \rightarrow \{0, 1\}$ where $|\text{dom } p| < \tau$. We let $p \leq_{\text{Add}^*(\tau, Y)} q$ if and only if $p \supseteq q$.

In later subsections we will conflate $\text{Add}(\tau, \lambda)$ and $\text{Add}^*(\tau, \lambda)$ to simplify notation.

Definition 16. Let λ be inaccessible and let $\tau < \mu < \lambda$ be regular cardinals such that $\tau^{<\tau} = \tau$. We define a forcing $\mathbb{M}^+(\tau, \mu, \lambda)$ that consists of pairs (p, q) such that

- (1) $p \in \text{Add}^*(\tau, \lambda)$;
- (2) q is a function such that
 - (a) $\text{dom } q \subset \{\delta < \lambda : \delta \text{ is inaccessible}\}$,
 - (b) $|\text{dom } q| < \mu$,
 - (c) for all $\delta \in \text{dom } q$, $q(\delta)$ is an $\text{Add}^*(\tau, \delta + 1)$ -name such that

$$p \restriction ((\delta + 1) \times \tau) \Vdash_{\text{Add}^*(\tau, \delta + 1)} \text{“} q(\delta) \in \text{Col}(\mu, \delta)\text{”}.$$

We let $(p, q) \leq (p', q')$ if and only if

- (i) $p \leq_{\text{Add}^*(\tau, \lambda)} p'$,
- (ii) $\text{dom } q \supseteq \text{dom } q'$,
- (iii) for all $\delta \in \text{dom } q'$, $p \restriction ((\delta + 1) \times \tau) \Vdash_{\text{Add}^*(\tau, \delta + 1)} \text{“} q(\delta) \leq_{\text{Col}(\mu, \delta)} q'(\delta)\text{”}$.

First we go through the more routine properties that one would expect of this forcing.

Proposition 17. $\mathbb{M}^+(\tau, \mu, \lambda)$ is τ -closed and λ -Knaster.

Proof. Closure uses the facts that $\text{Add}^*(\tau, \lambda)$ is τ -closed and $\Vdash_{\text{Add}^*(\tau, \delta+1)}$ “ $\dot{\text{Col}}(\mu, \delta)$ is μ -closed” for all δ . For Knasterness: consider $\{(p_i, q_i) : i < \lambda\} \subseteq \mathbb{M}^+(\tau, \mu, \lambda)$, then fix a regular $\rho \in (\mu, \lambda)$ and find a stationary subset of $\lambda \cap \text{cof}(\rho)$ on which $\text{dom } p_i, \text{dom } q_i$ are fixed, and then proceed with a standard delta system lemma argument. \square

Crucially, we get a nice termspace:

Definition 18. Let $\mathbb{T} = \mathbb{T}(\mathbb{M}^+(\tau, \mu, \lambda))$ be the poset consisting of conditions q such that

- (1) $\text{dom } q \subset \{\delta < \lambda : \delta \text{ is inaccessible}\}$,
- (2) $|\text{dom } q| < \mu$,
- (3) for all $\delta \in \text{dom } q$, $\Vdash_{\text{Add}^*(\tau, \delta+1)}$ “ $q(\delta) \in \dot{\text{Col}}(\mu, \delta)$ ”.

Most importantly, we let $q \leq q'$ if and only if

- (i) $\text{dom } q \supseteq \text{dom } q'$,
- (ii) for all $\delta \in \text{dom } q$, $\Vdash_{\text{Add}^*(\tau, \delta+1)}$ “ $q(\delta) \leq q'(\delta)$ ”.

Proposition 19. There is a projection $\text{Add}^*(\tau, \lambda) \times \mathbb{T}(\mathbb{M}^+(\tau, \mu, \lambda)) \rightarrow \mathbb{M}^+(\tau, \mu, \lambda)$.

Proof. We let π be the projection with the definition $\pi(p, q) = (p, q)$. This is automatically order-preserving because the ordering $\leq_{\text{Add}^*(\tau, \lambda) \times \mathbb{T}}$ is coarser than the ordering $\leq_{\mathbb{M}^+(\tau, \mu, \lambda)}$. For obtaining the density condition, suppose $(r, s) \leq_{\mathbb{M}^+(\tau, \mu, \lambda)} (p_0, q_0)$. We want to find some (p_1, q_1) such that $(p_1, q_1) \leq_{\text{Add}^*(\tau, \lambda) \times \mathbb{T}} (p_0, q_0)$ and $(p_1, q_1) \leq_{\mathbb{M}^+(\tau, \mu, \lambda)} (r, s)$. To do this, we first let $p_1 = r$, and then we define q_1 with $\text{dom } q_1 = \text{dom } r$ such that at each coordinate $\delta \in \text{dom } q_1$, we use standard arguments on names to show that we can get both $p_0 \upharpoonright ((\delta + 1) \times \tau) \Vdash_{\text{Add}^*(\tau, \lambda)}$ “ $q_1(\delta) \leq s(\delta)$ ” as well as $1_{\text{Add}^*(\tau, \lambda)} \Vdash_{\text{Add}^*(\tau, \lambda)}$ “ $q_1(\delta) \leq q_0(\delta)$ ”. \square

Proposition 20. $\mathbb{T} = \mathbb{T}(\mathbb{M}^+(\tau, \mu, \lambda))$ is μ -closed.

Proof. This is an application of the mixing principle. Given a $\leq_{\mathbb{T}}$ -decreasing sequence $\langle q_i : i < \tau \rangle$ with $\tau < \mu$ we let $d = \bigcup_{i < \tau} \text{dom } q_i$. Then we define a lower bound \bar{q} with domain d such that for all $\delta \in d$, $\bar{q}(\delta)$ is a canonically defined name for a lower bound of the $q_i(\delta)$'s (where i is large enough that $\delta \in \text{dom } q_i$). \square

Then we get the standard consequences of the termspace analysis:

Proposition 21. The following are true in any extension by $\mathbb{M}^+(\tau, \mu, \lambda)$:

- (1) V -cardinals up to and including μ are cardinals.
- (2) For all $\alpha < \lambda$, $|\alpha| = \mu$.

(3) $\lambda = \mu^+$.

(4) $2^\tau = \lambda$.

Proof. (1) follows from the projection analysis and the fact that \mathbb{T} is μ -closed and $\text{Add}^*(\tau, \lambda)$ is τ^+ -cc, and from τ -closure of $\mathbb{M}^+(\tau, \mu, \lambda)$. (2) follows from the fact that for all inaccessible $\delta < \lambda$, $\mathbb{M}^+(\tau, \mu, \lambda)$ projects onto $\text{Add}^*(\tau, \delta) * \text{Col}(\mu, \delta)$. (3) follows from (1) and (2) plus λ -Knasterness. (4) follows from the fact that $\mathbb{M}^+(\mu, \lambda)$ projects onto $\text{Add}^*(\tau, \lambda)$, so it forces that $2^\tau \geq \lambda$. Since the poset has size λ and λ is inaccessible, it also forces that $2^\tau \leq \lambda$. \square

The following lemma is the crux of the new idea.

Lemma 22. *If $\delta_0 < \lambda$ is inaccessible, then there is a forcing equivalence*

$$\mathbb{M}^+(\tau, \mu, \lambda) \simeq \mathbb{M}^+(\tau, \mu, \delta_0) * \text{Add}(\tau) * \mathbb{E},$$

where $\mathbb{M}^+(\tau, \mu, \delta_0) * \text{Add}(\tau)$ forces that \mathbb{E} is a projection of a product of a μ -closed forcing and a τ^+ -cc forcing.

Proof. In particular, we will show that there is a forcing equivalence $\mathbb{M}^+(\tau, \mu, \lambda) \simeq \mathbb{M}^+(\tau, \mu, \delta_0) * \text{Add}(\tau) * (\mathbb{F} \times \mathbb{G})$ where, in the extension by $\mathbb{M}^+(\tau, \mu, \delta_0) * \text{Add}(\tau)$,

- \mathbb{G} is a projection of a product of a μ -closed forcing and $\text{Add}^*(\tau, \lambda)$, and
- \mathbb{F} is μ -closed.

The statement of the lemma can then be obtained by merging \mathbb{F} with the closed component of the product that projects onto \mathbb{G} .

First we describe \mathbb{F} and \mathbb{G} . To do this, we fix some notation. Given $Y \subseteq \lambda$, we let π_{Add}^Y denote the projection $(p, q) \mapsto p \upharpoonright (Y \times \tau)$ from $\mathbb{M}^+(\tau, \mu, \lambda)$ onto $\text{Add}^*(\tau, Y)$. For any poset \mathbb{P} , we employ the convention that $\Gamma(\mathbb{P})$ denotes a canonical name for a \mathbb{P} -generic. If $X \subset \mathbb{P}$, then we use the notation $\uparrow X := \{q \in \mathbb{P} : \exists p \in X, p \leq q\}$.

We will let

$$\mathbb{F} := \text{Col}(\mu, \delta_0)^V [(\uparrow(\pi_{\text{Add}}^{\delta_0} \text{''} \Gamma(\mathbb{M}^+(\tau, \mu, \delta_0)))) \times \Gamma(\text{Add}(\tau))]$$

if we are working in an extension by $\mathbb{M}^+(\tau, \mu, \delta_0) * \text{Add}(\tau)$. (In other words, the poset \mathbb{F} will be the version of $\text{Col}(\mu, \delta_0)$ as interpreted in the extension of V by $\text{Add}^*(\tau, \delta_0 + 1)$ where the initial coordinates come from $\mathbb{M}^+(\tau, \mu, \delta_0)$ and the last coordinate comes from the additional copy of $\text{Add}(\tau)$ that occupies the coordinate δ_0 in $\text{Add}^*(\tau, \delta_0 + 1)$.)

Still working in an extension by $\mathbb{M}^+(\tau, \mu, \delta_0) * \text{Add}(\tau)$, the poset \mathbb{G} consists of pairs (p, q) such that

- (1) $p \in \text{Add}^*(\tau, (\delta_0, \lambda))$;
- (2) q is a function such that
 - (a) $\text{dom } q \subset \{\delta \in (\delta_0, \lambda) : \delta \text{ is inaccessible}\}$,

(b) $|\text{dom } q| < \mu$,

(c) for all $\delta \in \text{dom } q$, $p \upharpoonright ((\delta_0, (\delta + 1)) \times \tau) \Vdash_{\text{Add}^*(\tau, (\delta_0, \delta + 1))} \text{“}q(\delta) \in \dot{\text{Col}}(\mu, \delta)\text{”}$.

The ordering is the one analogous to that of $\mathbb{M}^+(\tau, \mu, \lambda)$. An easy adaptation of the arguments for the projection analysis for $\mathbb{M}^+(\tau, \mu, \lambda)$ will then give a projection analysis for \mathbb{G} .

The rest of the proof of the lemma consists of verifying the more substantial claims.

Claim 23. $\mathbb{M}^+(\tau, \mu, \lambda) \simeq \mathbb{M}^+(\tau, \mu, \delta_0) * \text{Add}(\tau, 1) * (\mathbb{F} \times \mathbb{G})$.

Proof. We identify $\mathbb{M}^+(\tau, \mu, \delta_0) * \text{Add}(\tau, 1) * (\mathbb{F} \times \mathbb{G})$ with the dense subset of conditions $((r, s), t, u, (\dot{r}', \dot{s}'))$ such that \dot{s}' is forced to have a specific domain in V . The fact that this subset is dense follows from the fact that $\mathbb{M}^+(\tau, \mu, \lambda) * \text{Add}(\tau, 1)$ has the μ -covering property.

We will argue that there is a trivial projection defined by

$$\pi : (p, q) \mapsto \left(\underbrace{(p \upharpoonright (\delta_0 \times \tau), q \upharpoonright \delta_0)}_{\mathbb{M}^+(\mu, \delta_0)}, \underbrace{p \upharpoonright (\{\delta_0\} \times \tau)}_{\text{Add}(\tau)}, \underbrace{q^*(\delta_0)}_{\mathbb{F}}, \underbrace{(\bar{p}, \bar{q})}_{\mathbb{G}} \right)$$

such that

- $\bar{p} := p \upharpoonright ((\delta_0, \lambda) \times \tau)$;
- $q^*(\delta_0)$ is obtained by changing $q(\delta_0)$ from an $\text{Add}^*(\tau, \delta_0 + 1)$ -name to an $\text{Add}(\tau)$ -name as interpreted in the extension by the relevant generic, namely $(\uparrow(\pi_{\text{Add}}^{\delta_0} \text{''} \Gamma(\mathbb{M}^+(\tau, \mu, \delta_0))))$;
- \bar{q} has domain (δ_0, λ) , and for each $\delta \in (\delta_0, \lambda)$, $\bar{q}(\delta)$ has changes analogous to the changes made to $q^*(\delta_0)$.

It is clear that π is order-preserving. We also want to show that if

$$((r, s), t, u, (\dot{r}', \dot{s}')) \leq_{\mathbb{M}^+(\tau, \mu, \delta_0) * \text{Add}(\tau) * (\mathbb{F} \times \mathbb{G})} \pi(p_0, q_0)$$

then there is some $(p_1, q_1) \leq_{\mathbb{M}^+(\mu, \lambda)} (p_0, q_0)$ such that we have $\pi(p_1, q_1) \leq ((r, s), t, u, (\dot{r}', \dot{s}'))$. This can be done by taking

- $p_1 = r^* \cup \tilde{t} \cup r'$ where $r^* \leq r$ decides t and \dot{r}' and \tilde{t} writes t as a partial function $\{\delta\} \times \tau \rightarrow \{0, 1\}$,
- $q_1 = s \cup \tilde{u} \cup \tilde{s}'$ where \tilde{u} reinterprets u as an $\text{Add}^*(\delta_0 + 1)$ -name and for each $\delta \in \text{dom } \dot{s}'$, \tilde{s}' reinterprets $\dot{s}'(\delta)$ as an $\text{Add}^*(\delta + 1)$ -name.

Last, we argue that $\pi(p_0, q_0) = \pi(p_1, q_1)$ implies that (p_0, q_0) and (p_1, q_1) are compatible. Suppose that (p_0, q_0) and (p_1, q_1) are incompatible. If p_0 and p_1 are incompatible as elements of $\text{Add}^*(\tau, \lambda)$, then one of $p_i \upharpoonright (\delta_0 \times \tau)$, $p_i \upharpoonright (\{\delta_0\} \times \tau)$, and $p_i \upharpoonright ((\delta_0, \lambda) \times \tau)$ must be distinct for $i = 0$ and $i = 1$. Otherwise, there is some $p' \leq p_0, p_1$ and some $\delta \in \text{dom } q_0 \cap \text{dom } q_1$ inaccessible such that $p' \Vdash \text{“}q_0(\delta) \perp q_1(\delta)\text{”}$,

which implies that $q_0(\delta) \neq q_1(\delta)$. Therefore, one of $q_i \restriction \delta_0$, $q_i(\delta_0)$, or $q_i \restriction (\delta_0, \lambda)$ is distinct for $i \in \{0, 1\}$. \square

Claim 24. $\Vdash_{\mathbb{M}^+(\tau, \mu, \delta_0) * \text{Add}(\tau, 1)} \text{“}\mathbb{F} \text{ is } \mu\text{-closed”}$.

Proof. In fact, our argument will also show that

$$\Vdash_{\mathbb{M}^+(\tau, \mu, \delta_0) * \text{Add}(\tau, 1)} \text{“}\mathbb{F} = \text{Col}(\mu, \delta_0)\text{”}.$$

We fix some arbitrary generics:

- G is $\mathbb{M}^+(\tau, \mu, \delta_0)$ -generic over V ,
- r is $\text{Add}(\tau, 1)$ -generic over $V[G]$,
- H is the $\text{Add}^*(\tau, \delta_0)$ -generic induced from G by $\pi_{\text{Add}}^{\delta_0}$,
- K is the generic for the quotient of $\mathbb{M}^+(\tau, \mu, \delta_0)$ by $\text{Add}^*(\tau, \delta_0)$, i.e., the generic such that $V[H][K] = V[G]$,
- T is the generic for the termspace forcing $\mathbb{T}(\mathbb{M}^+(\tau, \mu, \delta_0))$, so that $V[G] \subset V[T][H]$.

It is enough to argue that $V[G][r] \models \text{“}\mathbb{F} \text{ is } \mu\text{-closed”}$ knowing that $V[H][r] \models \text{“}\mathbb{F} \text{ is } \mu\text{-closed”}$. Because adjoining G does not change the definition of $\text{Add}(\tau, 1)$, and because K is defined in terms of the subsets of τ adjoined by the filter H , we have $V[G][r] = V[H][K][r] = V[H][r][K]$. Therefore, it is enough to show that K does not add $< \mu$ -sequences over $V[H][r]$, so that $V[H][r]$'s version of $\text{Col}(\mu, \delta_0)$ remains μ -closed in $V[G][r]$. We have

$$V[H][r] \subset V[H][r][K] = V[H][K][r] = V[G][r] \subset V[T][H][r] = V[H][r][T].$$

Recall Easton's lemma, which states in part that if \mathbb{A} is μ -cc and \mathbb{B} is μ -closed, then $\Vdash_{\mathbb{A}} \text{“}\mathbb{B} \text{ is } \mu\text{-distributive”}$. Easton's lemma implies that T does not add new $< \mu$ -sequences over $V[H][r]$ since the forcing adjoining r is μ -cc over $V[H]$ and the forcing adjoining T is μ -closed over $V[H]$. Therefore K does not add new $< \mu$ -sequences over $V[H][r]$ since it is an intermediate factor of the extension. \square

This completes the proof of the lemma. \square

Now we have an application for the case where $\tau = \omega$.

Proposition 25. *If λ is Mahlo, then $\Vdash_{\mathbb{M}^+(\omega, \mu, \lambda)} \text{DSS}(\lambda)$.*

This basically repeats Krueger's argument for [21, Theorem 9.1].

Proof. Let G be $\mathbb{M}^+(\omega, \mu, \lambda)$ -generic over V . The set of V -inaccessibles in λ will form the stationary set $S \subset \mu^+ \cap \text{cof}(\mu)$ carrying the disjoint stationary sequence in the extension by $\mathbb{M}^+(\omega, \mu, \lambda)$. For every such $\delta \in S$, let \bar{G} be the generic on $\mathbb{M}^+(\omega, \mu, \delta)$ induced by G and let r be the $\text{Add}(\omega)$ -generic induced by G via $\pi_{\text{Add}}^{\{\delta\}}$. We use Fact 13 to obtain a stationary set $S_\delta^* \subset P_\mu(H(\delta))^{V[\bar{G}][r]}$ such that

for all $N \in \mathcal{S}_\delta^*$, $N \cap \delta \notin V[\bar{G}]$ and such that N is also internally approachable by a ω -sequence. Therefore we can apply Lemma 22 with Fact 11 and then Fact 9 to find that \mathcal{S}_δ^* is stationary in $V[G]$. We then let $\mathcal{S}_\delta = \{N \cap \delta : N \in \mathcal{S}_\delta^*\}$, and we see that $\langle \mathcal{S}_\delta : \delta \in S \rangle$ is a disjoint stationary sequence. \square

2.2. Proving the main theorems. Now we will apply the new version of Mitchell forcing to answer Krueger’s questions. We can readily prove Theorem 1, which states that we can obtain $\text{DSS}(\aleph_2) \wedge \text{DSS}(\aleph_3)$:

Proof of Theorem 1. Begin with a ground model V in which $\lambda_1 < \lambda_2$ and the λ ’s are Mahlo. Let $\mathbb{M}_1 = \mathbb{M}^+(\omega, \aleph_1, \lambda_1)$. (Any λ_1 -sized forcing that turns λ_1 into \aleph_2 and adds a disjoint stationary sequence on \aleph_2 would work, so we could also use a more standard mixed support iteration.) Then let $\dot{\mathbb{M}}_2$ be an \mathbb{M}_1 -name for $\mathbb{M}^+(\omega, \lambda_1, \lambda_2)$. We argue that if G_1 is \mathbb{M}_1 -generic over V and G_2 is $\dot{\mathbb{M}}_2[G_1]$ -generic over $V[G_1]$, then $V[G_1][G_2] \models \text{“DSS}(\lambda_1) \wedge \text{DSS}(\lambda_2)\text{”}$. We get $\text{DSS}(\lambda_2)$ from the fact that λ_2 remains Mahlo in $V[G_1]$ together with Proposition 25, so we only need to argue that the disjoint stationary sequence $\vec{S} := \langle \mathcal{S}_\alpha : \alpha \in S \rangle \in V[G_1]$ remains a disjoint stationary sequence in $V[G_1][G_2]$.

Working in $V[G_1]$, preservation of \vec{S} follows from the projection analysis: Let H_1 and H_2 be chosen so that H_1 is $\mathbb{T} := \mathbb{T}(\mathbb{M}_2)$ -generic over $V[G_1]$, H_2 is $\text{Add}(\omega, \lambda_2)^{V[G_1]}$ -generic over $V[G_1][H_1]$, and $V[G_1][G_2] \subseteq V[G_1][H_1][H_2]$. Since \mathbb{T} is λ_1 -closed, it preserves stationarity of S and the \mathcal{S}_α ’s, and $\text{Add}(\omega, \lambda_2)^{V[G_1]}$ still has the countable chain condition in $V[G_1][H_1]$. It follows that the stationarity of S is preserved in $V[G_1][H_1][H_2]$, as well as the stationarity of the \mathcal{S}_α ’s (by Fact 9). Therefore \vec{S} is a disjoint stationary sequence on λ_1 in $V[G_1][G_2]$. \square

It will take a bit more work to show how to obtain Theorem 2 in the same model for Theorem 1. (Recall that Theorem 2 states that we can simultaneously separate internally stationary and internally club for $[H(\aleph_2)]^{\aleph_1}$ and $[H(\aleph_3)]^{\aleph_2}$.) Note that we cannot just apply Fact 6 because $2^\omega = \aleph_3$ in the model for Theorem 1, plus it is consistent that there can be a stationary set which is internally unbounded but not internally stationary [19].

We will give some facts on preservation of the distinction between stationary sets that are internally stationary but not internally club:

Proposition 26. *Suppose \mathbb{P} is ν -closed and $S \subseteq [X]^\delta$ is a stationary set such that $|[X]^\delta| \leq \nu$ and $|X| > 1$. Then $\Vdash_{\mathbb{P}} \text{“}S \text{ is stationary in } [X]^\delta\text{”}$.*

Proof. Let \dot{C} be a \mathbb{P} -name for a club in $[X]^\delta$ and let $\vec{x} = \langle x_\xi : \xi \leq \bar{\nu} \rangle$ enumerate $[X]^\delta$ (where $\bar{\nu} \leq \nu$). Note that we have $\delta < 2^\delta \leq |X|^\delta \leq \nu$, so conditions in \mathbb{P} can decide names for elements of \dot{C} . We construct a sequence $\vec{z} = \langle z_\xi : \xi < \bar{\nu} \rangle \subseteq [X]^\delta$ and a $\leq_{\mathbb{P}}$ -descending sequence $\langle p_\xi : \xi < \bar{\nu} \rangle$ using the closure of \mathbb{P} such that for all $\xi < \bar{\nu}$, $p_\xi \Vdash \text{“}x_\xi \subseteq z_\xi \in \dot{C}\text{”}$ and $p_\xi \parallel \text{“}x_\xi \in \dot{C}\text{”}$.

Then let D be the set $\{x_\xi : \exists \zeta < \bar{v}, p_\zeta \Vdash "x_\xi \in \dot{C}"\}$. We can argue that D is a club: It is unbounded because of the sets chosen for z_ξ . It is closed because if $\langle x_{\xi_i} : i < \bar{\delta} \rangle \subseteq D$ (for $\bar{\delta} \leq \delta$) is an \subseteq -increasing sequence such that we have $p_{\zeta_i} \Vdash "x_{\xi_i} \in \dot{C}"$, and $\zeta^* = \sup_{i < \bar{\delta}} \zeta_i$, then $p_{\zeta^*} \Vdash "\bigcup_{i < \bar{\delta}} x_{\xi_i} \in \dot{C}"$.

There is some $w \in D \cap S$. If p_ξ is such that $p_\xi \Vdash "w \in \dot{C}"$, then we have $p_\xi \Vdash "\dot{C} \cap S \neq \emptyset"$. □

Proposition 27. *Suppose $|[H(\theta)]^\delta| \leq \nu$. Let \mathbb{F} have the δ -chain condition and let \mathbb{G} be ν -closed. If there is a stationary set $S \subseteq [H(\theta)]^\delta$ consisting of sets that are internally stationary but not internally club, then $\mathbb{F} \times \mathbb{G}$ forces that there is a stationary set consisting of sets that are internally stationary but not internally club.*

Proof. Since \mathbb{G} preserves the chain condition of \mathbb{F} , we show that preservation of the distinction can be achieved by forcing with \mathbb{G} and then \mathbb{F} . The poset \mathbb{G} preserves the distinction by Proposition 26 and the fact that it does not change $H(\theta)$.

Now we argue that \mathbb{F} preserves the distinction. Let S be the witnessing stationary set in V and let $X = H(\theta)^V$. If G is \mathbb{F} -generic over V , let $Y = H(\theta)^{V[G]}$ and let $S^* = \{M \in [Y]^\delta : M \cap X \in S\}$. We will argue that S^* witnesses the relevant statement in $V[G]$. Let \dot{S}^* be a name for S^* .

To see that \dot{S}^* is forced to be stationary, let \dot{C} be a name for a club in $[Y]^\delta$. Given $p \in \mathbb{F}$, let $D = \{z : \exists \dot{w}, p \Vdash "\dot{w} \in \dot{C}", \dot{w} \cap X = z\}$. Then D is a club in $[X]^\delta$ as regarded in V , so there is some $z \in S$, and hence $p \Vdash "\dot{w} \in \dot{C} \cap \dot{S}^*"$.

Next we argue that members of \dot{S}^* are forced not to be internally club. Suppose for contradiction, then, that p forces $\dot{M} \in \dot{S}^*$ to be internally club as witnessed by \dot{c} , and also that $N = \dot{M} \cap X$ where $N \in S$. Let $d = \{z : \exists \dot{w}, p \Vdash "\dot{w} \in \dot{c}", \dot{w} \cap X = z\}$. Then d is a club in $P_\mu(N)$ since if $a \in z \subseteq N$ then $a \in N$ and if $p \Vdash "\dot{w} \cap X = z"$ then in particular $p \Vdash "\dot{w} \in \dot{M}"$, so $z \in N$. This contradicts the fact that N consists of sets that are not internally club.

Finally, we argue that \dot{S}^* is forced to be internally stationary. Let \dot{M} be forced by p to be in \dot{S}^* and that $N = \dot{M} \cap X$. Let G be generic with $p \in G$ and work in $V[G]$. Then $\{w \subseteq P_\delta(\dot{M}) : \exists z \in N, w = z[G]\}$ is a club as regarded in $V[G]$. As in the argument for stationarity, any name \dot{c} for a club in $P_\delta(\dot{M})$ can produce a corresponding club d in the ground model. Then we can find some $z \in N \cap d$ and if G is generic with $p \in G$ then $z[G] \in \dot{c} \cap \dot{M}$. □

We use a concept from Harrington and Shelah to handle Mahlo cardinals [13]:

Definition 28. Let λ be Mahlo and let \mathcal{N} be a model of some fragment of ZFC. We say that $\mathcal{M} \prec \mathcal{N}$ is *rich* if

- (1) $\lambda \in \mathcal{M}$;
- (2) $\bar{\lambda} := \mathcal{M} \cap \lambda \in \lambda$;
- (3) $\bar{\lambda}$ is an inaccessible cardinal in \mathcal{N} ;

- (4) the size of \mathcal{M} is $\bar{\lambda}$;
(5) \mathcal{M} is closed under $<\bar{\lambda}$ -sequences and $\bar{\lambda} < \lambda$.

Lemma 29. *If λ is Mahlo, then $\mathbb{M}^+(\omega, \mu, \lambda)$ forces that there are stationarily many $Z \in [\mu^+]^\mu$ which are internally stationary but not internally club.*

This follows Krueger's proof of [21, Theorem 10.1], making necessary changes for Mahlo cardinals, and including enough details to show that we can get the necessary preservation of stationarity simply from the projection analysis. We do not need guessing functions (which are used in Krueger's argument) because we are only obtaining one instance of separation per large cardinal.

Proof of Lemma 29. Define $\mathbb{M} := \mathbb{M}^+(\omega, \mu, \lambda)$ and let \dot{C} be an \mathbb{M} -name for a club in $([H(\mu^+)]^\mu)^{V[\mathbb{M}]}$. We want to find an \mathbb{M} -name \dot{Z} for an element of $([H(\mu^+)]^\mu)^{V[\mathbb{M}]} \cap \dot{C}$ that is internally stationary but not internally club. Let \dot{F} be an \mathbb{M} -name for a function $(H(\mu^+)^{V[\mathbb{M}]})^{<\omega} \rightarrow H(\mu^+)^{V[\mathbb{M}]}$ with the property that all of its closure points are in \dot{C} . Let Θ be as large as needed for the following discussion and let \mathcal{N} be the structure $(H(\Theta), \in, <_\Theta, \mathbb{M}, \dot{F}, \lambda, \mu)$ where $<_\Theta$ is a well-ordering of $H(\Theta)$.

Since λ is Mahlo, we can find some $\mathcal{K} \prec \mathcal{N}$ with $\mu \subset \mathcal{K}$ that is a rich submodel of cardinality $\bar{\lambda}$. Now set G to be \mathbb{M} -generic over V . Note that $H(\lambda)^{V[G]} = H(\lambda)[G]$ because \mathbb{M} has the λ -chain condition and $\mathbb{M} \subset H(\lambda)$. We will argue that $Z := \mathcal{K}[G] \cap H(\lambda)[G]$ is what we are looking for.

Claim 30. $Z \in C := \dot{C}[G]$.

Proof. We have $\bar{\lambda} \leq |Z| \leq |\mathcal{K}| \leq \bar{\lambda}$ and $\bar{\lambda}$ has cardinality μ in $\mathcal{N}[G]$, so $Z \in [H(\lambda)^{V[G]}]^\mu$. If $a_1, \dots, a_n \in Z$, there are \mathbb{M} -names $\dot{b}_1, \dots, \dot{b}_n \in \mathcal{K} \cap H(\lambda)$ such that $a_i = \dot{b}_i^G$ for all $1 \leq i \leq n$. By elementarity, \mathcal{K} contains the $<_\Theta$ -least maximal antichain $A \subset \mathbb{M}$ of conditions deciding $\dot{F}(\dot{b}_1, \dots, \dot{b}_n)$. Since $|A| < \lambda$, we have $|A| \in \mathcal{K} \cap \lambda = \bar{\lambda}$, so it will follow that $A \subset \mathcal{K}$. Therefore if $p \in G \cap A$, then $p \in M$ in particular, so $p \Vdash \dot{F}(\dot{b}_1, \dots, \dot{b}_n) = \dot{b}_*$ for some $\dot{b}_* \in \mathcal{K} \cap H(\lambda)$ where we automatically get $\dot{b}_* \in H(\bar{\lambda})$, and therefore

$$F(a_1, \dots, a_n) = a_* := \dot{b}_*^G \in \mathcal{K}[G] \cap H(\lambda)[G] = Z$$

(where of course $F := \dot{F}[G]$). □

For the rest of the proof let $\bar{G} := \pi_{\mathcal{K}}(G)$ where $\pi_{\mathcal{K}}$ is the Mostowski collapse relative to \mathcal{K} . Since $\pi_{\mathcal{K}}(\mathbb{M}) = \mathbb{M}^+(\omega, \mu, \bar{\lambda})$, there is an extension $\pi_{\mathcal{K}} : \mathcal{K}[G] \cong \pi_{\mathcal{K}}(\mathcal{K})[\bar{G}]$. We also define $h := \pi_{\mathcal{K}}(H(\lambda)[G] \cap \mathcal{K}[G])$.

Claim 31. Z is internally stationary.

Proof. First, we argue that $S := P_\mu(h)^{\mathcal{N}[\bar{G}]}$ is stationary as a subset of $P_\mu(h)^{\mathcal{N}[G]}$ in $\mathcal{N}[G]$. By Lemma 22, the quotient \mathbb{M}/\bar{G} is a projection of a forcing of the form

$\mathbb{A}_1 * (\dot{\mathbb{T}} \times \mathbb{A}_2)$ where \mathbb{A}_1 has the countable chain condition, $\dot{\mathbb{T}}$ is an \mathbb{A}_1 -name for a μ -closed forcing, and \mathbb{A}_2 also has the countable chain condition. Let K_1 , K_T , and K_2 be respective generics such that $V[G] \subseteq V[\bar{G}][K_1][K_T][K_2]$. Working in $\mathcal{N}[\bar{G}]$, note that $S' := S \cap \mathcal{J}\mathcal{A}(\omega)$ is stationary, and therefore has its stationarity preserved in $V[\bar{G}][K_1]$ by Fact 9.

We must also show that the stationarity of S' will be preserved by countably closed forcings over $\mathcal{N}[\bar{G}][K_1]$. Suppose $\langle M_n : n < \omega \rangle$ witnesses internal approachability of some $N \in S'$ in $V[\bar{G}]$ with respect to the structure $H(\lambda^+)^{V[\bar{G}]}$, and let $M_\omega := \bigcup_{n < \omega} M_n$. Then we can see that $\langle M_n[K_1] : n < \omega \rangle$ is a chain of elementary submodels of $H(\lambda)[\bar{G}][K_1] = H(\lambda)^{V[\bar{G}][K_1]}$. We also have $M_n[K_1] \cap V[\bar{G}] = M$ and $M_\omega[K_1] \cap V[\bar{G}] = M_\omega \in S'$ with $M_\omega[K_1] \prec H(\lambda)^{V[\bar{G}][K_1]}$. If we choose the M_n 's to be elementary substructures of $H(\lambda^+)^{V[\bar{G}]}$ ($\in, <^*, \dot{C}, \dots$) where $<^*$ is a well-ordering and \dot{C} is an $\mathbb{A}_1 * \dot{\mathbb{T}}$ -name for a club, then an argument almost exactly like the one showing that internal approachability is preserved (i.e., the proof of Fact 11) will show that S' is stationary in $\mathcal{N}[\bar{G}][K_1][K_T]$.

Then the extension of $\mathcal{N}[\bar{G}][K_1][K_T][K_2]$ over $\mathcal{N}[\bar{G}][K_1][K_T]$ preserves the stationarity of S' by another application of Fact 9, so we get stationarity in $\mathcal{N}[G]$.

Now that we have established preservation of stationarity of S' , we can finish the argument. Since $|h| = \mu$ in $\mathcal{N}[G]$, we can write $h = \bigcup_{i < \mu} x_i$ where $\langle x_i : i < \mu \rangle$ is a continuous and \subset -increasing chain of elements of $P_\mu(h)$. (This is *not* a chain through $P_\mu(h)^{\mathcal{N}[\bar{G}]}$.) The chain is a club in $P_\mu(h)^{\mathcal{N}[G]}$, in which S' is stationary, so there is a stationary $X \subseteq \mu$ such that $\{x_i : i \in X\} \subseteq S'$. Since $S' \subseteq S$, it follows that $i \in X$ implies that $x_i = \pi_{\mathcal{X}}(y_i)$ for some $y_i \in Z$. Therefore $\langle y_i : i < \mu \rangle$ is \subset -increasing with union Z , and in particular $\langle y_i : i \in X \rangle$ is stationary in Z . \square

Claim 32. Z is *not* internally club.

Proof. Suppose for contradiction that Z is internally club and hence that there is a \subset -increasing and continuous chain $\langle Z_i : i < \mu \rangle \in \mathcal{N}[G]$ with $|Z_i| < \mu$ for all $i < \mu$ and $\bigcup_{i < \mu} Z_i = Z$. So for all $i < \mu$, $Z_i \subset Z$, and so $\langle \pi_{\mathcal{X}}[Z_i] : i < \mu \rangle$ is an \subset -increasing and continuous chain with union h . If we let $W_i := \pi_{\mathcal{X}}[Z_i]$ for all $i < \mu$, then the fact that $|W_i| < \mu$ implies that $W_i = \pi_{\mathcal{X}}(Z_i) \in \mathcal{K}[\bar{G}]$. Therefore $\langle W_i : i < \mu \rangle$ is a continuous and \subset -increasing chain of sets in $P_\mu(h)$ with union h .

We define a set $U \in \mathcal{N}[\bar{G}][r]$ (where r is the generic induced by G from $\pi_{\text{Add}}^{\{\lambda\}}$) as

$$\{A \in P_\mu(H(\chi)) \cap \mathcal{J}\mathcal{A}(\omega) : A \cap h \notin \mathcal{N}[\bar{G}]\}.$$

We have a real in $\mathcal{N}[\bar{G}][r] \setminus \mathcal{N}[\bar{G}]$ and $(\mu^+)^{\mathcal{N}[\bar{G}][r]} = \lambda \subset H(\lambda)$. Hence we apply Fact 13 to see that U is stationary in $\mathcal{N}[\bar{G}][r]$, and it remains stationary in $\mathcal{N}[G]$ by the preservation properties of the quotient (i.e., Lemma 22 combined with Facts 11 and 9). Therefore in $\mathcal{N}[G]$, since $h \subseteq H(\chi)^{\mathcal{N}[\bar{G}][r]}$, $\{A \cap h : A \in U\}$ is stationary in $P_\mu(h)$. Since $\langle W_i : i < \mu \rangle$ is club in h , there is some $i < \mu$ such that $W_i = A \cap h$

for some $A \in U$. But by definition, $A \cap h \notin \mathcal{N}[\bar{G}]$, but $W_i \in \mathcal{K}[\bar{G}] \subset \mathcal{N}[\bar{G}]$, so this is a contradiction. \square

This completes the proof of the lemma. \square

Proof of Theorem 2. Let \mathbb{M}_1 be any λ_1 -sized forcing that turns λ_1 into \aleph_2 and adds stationarily many $N \in [H(\aleph_2)]^{\aleph_1}$ that are internally stationary but not internally club. Let $\dot{\mathbb{M}}_2$ be an \mathbb{M}_1 -name for $\mathbb{M}^+(\omega, \lambda_1, \lambda_2)$, let G_1 be \mathbb{M}_1 -generic over V , and let G_2 be $\dot{\mathbb{M}}_2[G_1]$ -generic over $V[G_1]$. Then we can see that the theorem holds in $V[G_1][G_2]$: the distinction between internally stationary and internally club on $[H(\aleph_2)]^{\aleph_1}$ is preserved in $V[G_1][G_2]$ by Proposition 27, and we get a distinction between internally stationary and internally club for $[H(\aleph_3)]^{\aleph_2}$ by Lemma 29. \square

3. A club forcing and a guessing sequence

3.1. A review of the tools. The main idea of the proof of Theorem 3 is to force a club through the complement of a canonical stationary set—that is, it is canonical in the sense that it is independent of a particular enumeration used to define it. This set is described as follows:

Fact 33 (Krueger [21]). Suppose μ is an uncountable regular cardinal and $\mu^{<\mu} \leq \mu^+$. Let $\underline{x} = \langle x_\alpha : \alpha < \mu^+ \rangle$ enumerate $[\mu^+]^{<\mu}$ and let

$$S(\underline{x}) := \{\alpha \in \mu^+ \cap \text{cof}(\mu) : P_\mu(\alpha) \setminus \langle x_\beta : \beta < \alpha \rangle \text{ is stationary}\}.$$

Then $\text{DSS}(\mu^+)$ holds if and only if $S(\underline{x})$ is stationary.

The natural thing to do is to define the following:

Definition 34. Let μ be an uncountable regular cardinal such that $\mu^{<\mu} = \mu^+$ and let \underline{x} and $S(\underline{x})$ be defined as in Fact 33. Then let $\mathcal{C}(\underline{x})$ be the set of closed bounded subsets p of μ^+ such that $p \cap S(\underline{x}) = \emptyset$. We let $p' \leq p$ if and only if $p' \cap (\max p + 1) = p$.

Proposition 35. Assuming $\mu^{<\mu} \leq \mu^+$, $\mathcal{C}(\underline{x})$ is μ^+ -distributive.

Sketch of proof. If $S(\underline{x})$ is nonstationary, then the result is trivial. If it is stationary, then $S(\underline{x})$ does not contain a stationary set of approachable points [17, Corollary 3.7]. Since $\mu^{<\mu} \leq \mu^+$ there is going to be a stationary set S^* of approachable points, which without loss of generality is disjoint from $S(\underline{x})$. Then a standard distributivity argument applies (see Cox's explanation [3]). \square

We will also crucially need a characterization of diamonds. The following appears in joint work with Gilton and Stejskalová [11, Lemma 3.12].

Fact 36. The following are equivalent:

(1) λ is Mahlo and $\diamond_\lambda(\text{Reg})$ (where of course $\text{Reg} = \{\tau < \lambda : \tau \text{ regular}\}$) holds.

(2) There is a function $\ell : \lambda \rightarrow V_\lambda$ such that for every transitive structure \mathcal{N} satisfying a rich fragment of ZFC that is closed under λ^+ -sequences in V , the following holds: for every $A \in \mathcal{N}$ with $A \in H(\lambda^+)$ and any $a \in \mathcal{N}$ with $|a| < \lambda$, there is a rich $\mathcal{M} \prec \mathcal{N}$ with $a \cup \{\ell\} \cup \{A\} \subset \mathcal{M}$ such that $\ell(\bar{\lambda}) = \pi_{\mathcal{M}}(A)$ (where $\bar{\lambda} = \mathcal{M} \cap \lambda$ and $\pi_{\mathcal{M}}$ is the Mostowski collapse).⁴

We can always use such an ℓ assuming the consistency of a Mahlo cardinal: If λ is Mahlo in a model V , then it is Mahlo in Gödel’s class L where $\diamond_\lambda(S)$ holds for all regular λ and stationary $S \subset \lambda$.

Two other forcings will be used, mostly for their black-boxed properties:

Definition 37. If T is a wide Aronszajn tree⁵ of cardinality \aleph_1 , let $\mathbb{B}(T)$ be Baumgartner’s forcing for specializing Aronszajn trees. It consists of finite functions $f : T \rightarrow \omega$ such that $f(x) \neq f(y)$ if $x \leq_T y$ or $y \leq_T x$. If $f, g \in \mathbb{B}(T)$, then $f \leq g$ if and only if $f \supseteq g$.

Definition 38. Let $S \subset [\aleph_2]^\omega$ be stationary. Then let $\mathbb{P}(S)$ be the forcing consisting of continuous, increasing, and countable chains $\langle M_\xi : \xi \leq \eta \rangle$ of elements of S . For $p, q \in \mathbb{P}(S)$, $p \leq q$ if and only if p end-extends q [8].

Fact 39. The following are true for these forcings:

- (1) For Aronszajn trees T of cardinality \aleph_1 , $\mathbb{B}(T)$ has the countable chain condition.
- (2) For $S \subset [\aleph_2]^\omega$ stationary, $\mathbb{P}(S)$ adds a closed unbounded set in $[\aleph_2^V]^\omega$ through S .
- (3) If $S \in V$, then $\text{Add}(\omega) * \dot{\mathbb{P}}(S)$ has the weak ω_1 -approximation property, that is, if \dot{f} is an $\text{Add}(\omega) * \dot{\mathbb{P}}(S)$ -name for a function $\omega_1 \rightarrow \text{ON}$ whose initial segments are in V , then \dot{f} is forced to be in V [17].
- (4) If $S \in V$, then $\text{Add}(\omega) * \dot{\mathbb{P}}(S)$ is proper [17].

3.2. The proof. Now we prove Theorem 3. Fix λ Mahlo. We can assume that $\diamond_\lambda(\text{Reg})$ holds, so let ℓ witness Fact 36.

Let $\mathbb{1} = \langle \mathbb{1}_\alpha, \dot{\mathbb{J}}_\alpha : \alpha < \lambda \rangle$ be a countable-support iteration of length λ such that if $\ell(\delta)$ is an $\mathbb{1}_\delta$ -name for a proper forcing then $\Vdash_{\mathbb{1}_\delta} \text{“}\dot{\mathbb{J}}_\delta = \ell(\delta)\text{”}$ and otherwise $\dot{\mathbb{J}}_\delta$ is forced to be the trivial forcing.⁶ We will argue momentarily that we have $\Vdash_{\mathbb{1}} \text{“}\aleph_1^{<\aleph_1} \leq \aleph_2 = \lambda\text{”}$, so we fix an $\mathbb{1}$ -name \dot{x} of $[\aleph_2]^{<\aleph_1}$ in the extension by $\mathbb{1}$ as well as a sequence of names $\langle \dot{x}_\alpha : \alpha < \aleph_2 \rangle$ that canonically represent the elements listed by \dot{x} . Then let $\dot{\mathcal{C}}$

⁴The original is stated with a different quantification — for all such \mathcal{N} , there exists a function, not the other way around. However, the proof works with the quantification used here.

⁵We say that T is a *wide Aronszajn tree* of cardinality \aleph_1 if it has no uncountable branches. This is meant to distinguish our situation from the case in which T has countable levels.

⁶See the work of Abraham [1] and Cummings and Foreman [4] for classical examples of forcings that use guessing functions in their definitions.

be an $\mathbb{1}$ -name for $\mathcal{C}(\dot{x})$. Let G be $\mathbb{1}$ -generic over V and let H be $\mathcal{C} := \dot{\mathcal{C}}[G]$ -generic over $V[G]$. Then the model in which the theorem is realized is $V[G][H]$.

Most of the desired properties of $V[G][H]$ follow easily. First of all, $V[G][H] \models \lambda = \aleph_2$: For all $\Theta < \lambda$ the forcing $\text{Col}(\aleph_1, \Theta)$ appears in the iteration, $\mathbb{1}$ has the λ -chain condition because the iterands have size less than $< \lambda$, and $\mathbb{1}$ preserves \aleph_1 because it is proper. Then adjoining H preserves \aleph_2 by the distributivity property noted above. The fact that $V[G][H] \models \neg \text{DSS}(\mu^+)$ follows from Fact 33 given that the generic object added by H is a club through the complement of the relevant stationary set. The main part of the work is to show that the approachability property fails.

If $\mathcal{M} \prec \mathcal{N}$ is rich and $\pi_{\mathcal{M}}$ is the Mostowski collapse relative to \mathcal{M} , we will typically denote $\pi_{\mathcal{M}}(a)$ as \bar{a} .

Lemma 40. $V[G][H] \models \neg \text{AP}(\aleph_2)$.

Proof. If $\text{AP}(\aleph_2)$ holds then this is forced by some condition $z \in \mathbb{1} * \mathcal{C}$. Assuming this is the case, we can derive a contradiction.

Claim 41. *Let $\mathcal{M} \prec \mathcal{N}$ be a rich model chosen to witness Fact 36 in the sense of having the properties that $\mathcal{M} \cap \lambda = \bar{\lambda}$, $z \in \mathcal{M}$, and $\ell(\bar{\lambda})$ is an $\mathbb{1} \upharpoonright \bar{\lambda}$ -name for*

$$\pi_{\mathcal{M}}(\dot{\mathcal{C}}(\dot{x}) * \text{Add}(\omega) * \mathbb{P}(Y) * \mathbb{B}(Y)),$$

where $Y = ([\lambda]^\omega)^{V[\mathbb{1} * \dot{\mathcal{C}}(\dot{x})]}$, and $\mathbb{P}(Y)$ is defined with respect to the interpretation of Y as a stationary set and $\mathbb{B}(Y)$ is defined with respect to the interpretation of Y as a tree ordered by end-extension.

Suppose $\bar{G}_0 * \bar{H}_0$ is $\bar{\mathbb{1}} * \bar{\mathcal{C}}$ -generic over V . Then there is a $G_0 * H_0$ which is $\mathbb{1} * \mathcal{C}(\dot{x})$ -generic over V such that if $j : \bar{\mathcal{M}} \rightarrow \mathcal{M} \subset \mathcal{N}$ is the inverse of the Mostowski collapse, then there is a lift $j : \bar{\mathcal{M}}[\bar{G}_0][\bar{H}_0] \rightarrow \mathcal{N}[G_0][H_0]$ with the property that $G_0 * H_0$ is an \aleph_1 -preserving extension over $V[\bar{G}_0][\bar{H}_0][\bar{K}_0][\bar{K}_1][\bar{K}_2]$ where $\bar{K}_0 * \bar{K}_1 * \bar{K}_2$ is $\text{Add}(\omega) * \bar{\mathbb{P}}(Y) * \bar{\mathbb{B}}(Y)$ -generic.

Proof of Claim 41. We will lift the elementary embedding $j : \bar{\mathcal{M}} \rightarrow \mathcal{N}$ to $j : \bar{\mathcal{M}}[\bar{G}_0][\bar{H}_0] \rightarrow \mathcal{N}[G_0][H_0]$. We therefore fix the notation $\bar{\lambda} = \mathcal{M} \cap \lambda$, and we have an $\bar{\mathbb{M}}$ -generic \bar{G}_0 , so we let $\mathcal{C} = \dot{\mathcal{C}}(\dot{x})[G_0]$.

To perform the lift, we need to show that we can absorb the generic \bar{H}_0 . The first stage is for handling G_0 . The forcing $\dot{\mathcal{C}}(\dot{x}) * \text{Add}(\omega) * \mathbb{P}(Y) * \mathbb{B}(Y)$ is an iteration of proper forcings and is therefore proper, and its image under $\pi_{\mathcal{M}}$ is proper for similar reasons. Hence, since it is also guessed, it is used in the iteration. Therefore G_0 takes the form $\bar{G}_0 * \bar{H}_0 * \bar{K}_0 * \bar{K}_1 * \bar{K}_2 * \bar{K}_3$ where \bar{K}_3 is just a remainder. The quotient preserves \aleph_1 since the whole forcing does.

To lift the embedding further, we use a master condition argument. Specifically, we want to show that $\cup \bar{H}_0 \cup \{\bar{\lambda}\}$ is a condition in \mathcal{C} . This follows because $\bar{\lambda} \notin S(\bar{x})$ as evaluated in $\mathcal{N}[G_0]$: Since $\bar{\mathcal{M}}^{< \bar{\lambda}} \subseteq \bar{\mathcal{M}}$ and $\mathbb{1} \upharpoonright \bar{\lambda}$ has the $\bar{\lambda}$ -chain condition, the evaluation $\langle x_\beta : \beta < \bar{\lambda} \rangle$ is equal to the countable subsets of $\bar{\lambda}$ in $\bar{\mathcal{M}}[\bar{G}_0]$. Therefore

$P_\mu(\bar{\lambda}) \setminus \langle x_\beta : \beta < \bar{\lambda} \rangle$ will be nonstationary because of the club added by $\mathbb{P}(Y)$. Hence we choose H_0 to be a generic containing $\cup \bar{H}_0 \cup \{\bar{\lambda}\}$. \square

Suppose then that $z \in \mathbb{M} * \dot{\mathcal{C}}(\underline{x})$ forces that approachability holds. By the claim, there is an embedding $\bar{\mathcal{M}}[\bar{G}][\bar{H}] \rightarrow \mathcal{N}[G][H]$ such that $V[G * H]$ is an extension over $V[\bar{G}][\bar{H}][\bar{K}_0][\bar{K}_1][\bar{K}_2]$ that preserves \aleph_1 where $\bar{K}_0 * \bar{K}_1 * \bar{K}_2$ is generic for $\pi_{\mathcal{M}}(\text{Add}(\omega) * \mathbb{P}(Y) * \mathbb{B}(Y))$. Since we are supposing that approachability holds, there is in $\mathcal{N}[G][H]$ a club $C \subseteq \aleph_2$ such that all of its points of cofinality \aleph_1 are approachable. By elementarity it follows that $\bar{\lambda} \in C$, so it is enough to show that $\bar{\lambda}$ cannot actually be an approachable point.

We need to show that \bar{Y} does not have a cofinal branch. By the weak ω_1 -approximation property of $\pi_{\mathcal{M}}(\text{Add}(\omega) * \dot{\mathbb{P}}(S))$ (Fact 39), \bar{Y} is a wide Aronszajn tree in $V[\bar{G}][\bar{H}][\bar{K}_0][\bar{K}_1]$ because no new cofinal branches are added. Moreover it has cardinality \aleph_1 in that model. If $D \subseteq \bar{\lambda}$ is a club of order-type ω_1 in $V[\bar{G}][\bar{H}][\bar{K}_0][\bar{K}_1]$, we can conflate \bar{Y} with $\{x \in ([\bar{\lambda}]^\omega)^{V[\bar{G}][\bar{H}]} : \sup x \in D\}$ so that it has height ω_1 . The forcing \bar{K}_2 adds a specializing function, therefore it remains a wide Aronszajn tree in any \aleph_1 -preserving extension, so in particular this is true for $V[G][H]$.

If $\bar{\lambda}$ were an approachable point as witnessed by (without loss of generality) a club E , then for all $\alpha \in E \cap D$, we have $E \cap \alpha \in ([\bar{\lambda}]^\omega)^{V[\bar{G}][\bar{H}]}$. Hence it would be implied that \bar{Y} has a cofinal branch, which is a contradiction. \square

Remark 42. This master condition argument can also be used to show that $\mathcal{C}(\underline{x})$ is distributive over $V[\mathbb{1}]$.

Now we are finished with the proof of Theorem 3.

4. Further directions

We propose some other considerations along the lines of the question: Why did we have to do more work to get Theorem 2 after obtaining Theorem 1? Or rather, is the assumption $2^\mu = \mu^+$ necessary for Fact 6?

Question 1. Is it consistent for μ regular that exactly one of DSS(μ^+) and “internally club and internally unbounded are distinct for $[H(\mu^+)]^\mu$ ” holds?⁷

On a similar note, the assumption that $2^\mu = |H(\mu^+)|$ is also used in a folklore result that assuming $2^\mu = \mu^+$, the distinction between internally unbounded and internally approachable for $[\mu^+]^\mu$ requires a Mahlo cardinal.

Question 2. What is the exact equiconsistency strength of the separation of internally approachable and internally unbounded for $[H(\mu^+)]^\mu$ for regular μ ?

⁷This question was answered by Jakob after the previous version of this paper was released [14].

Acknowledgements

Many thanks to Hannes Jakob for finding errors in the original version of this paper, and to the anonymous referees for further corrections and improvements.

References

- [1] U. Abraham, “Aronszajn trees on \aleph_2 and \aleph_3 ”, *Ann. Pure Appl. Logic* **24**:3 (1983), 213–230. MR Zbl
- [2] U. Abraham and S. Shelah, “Forcing closed unbounded sets”, *J. Symb. Logic* **48**:3 (1983), 643–657. MR Zbl
- [3] S. D. Cox, “Forcing axioms, approachability, and stationary set reflection”, *J. Symb. Log.* **86**:2 (2021), 499–530. MR Zbl
- [4] J. Cummings and M. Foreman, “The tree property”, *Adv. Math.* **133**:1 (1998), 1–32. MR Zbl
- [5] J. Cummings, S.-D. Friedman, M. Magidor, A. Rinot, and D. Sinapova, “The eightfold way”, *J. Symb. Log.* **83**:1 (2018), 349–371. MR
- [6] T. Eisworth, “Successors of singular cardinals”, pp. 1229–1350 in *Handbook of set theory*, edited by M. Foreman and A. Kanamori, Springer, 2010. MR Zbl
- [7] M. Foreman and M. Magidor, “Large cardinals and definable counterexamples to the continuum hypothesis”, *Ann. Pure Appl. Logic* **76**:1 (1995), 47–97. MR Zbl
- [8] S.-D. Friedman and J. Krueger, “Thin stationary sets and disjoint club sequences”, *Trans. Amer. Math. Soc.* **359**:5 (2007), 2407–2420. MR Zbl
- [9] T. D. Gilton, *On the infinitary combinatorics of small cardinals and the cardinality of the continuum*, Ph.D. thesis, University of California, Los Angeles, 2019, available at <https://www.proquest.com/docview/2327544356>. MR
- [10] T. Gilton and J. Krueger, “A note on the eightfold way”, *Proc. Amer. Math. Soc.* **148**:3 (2020), 1283–1293. MR Zbl
- [11] T. Gilton, M. Levine, and Š. Stejskalová, “Trees and stationary reflection at double successors of regular cardinals”, *J. Symb. Log.* **88**:2 (2023), 780–810. MR Zbl
- [12] M. Gitik, “Nonsplitting subset of $\mathcal{P}_\kappa(\kappa^+)$ ”, *J. Symb. Logic* **50**:4 (1985), 881–894. MR Zbl
- [13] L. Harrington and S. Shelah, “Some exact equiconsistency results in set theory”, *Notre Dame J. Formal Logic* **26**:2 (1985), 178–188. MR Zbl
- [14] H. Jakob, “Cascading variants of internal approachability”, preprint, 2024. arXiv 2404.18571
- [15] T. Jech, *Set theory*, 3rd ed., Springer, 2003. MR Zbl
- [16] R. B. Jensen, “The fine structure of the constructible hierarchy”, *Ann. Math. Logic* **4** (1972), 229–308. MR Zbl
- [17] J. Krueger, “Internally club and approachable”, *Adv. Math.* **213**:2 (2007), 734–740. MR Zbl
- [18] J. Krueger, “A general Mitchell style iteration”, *MLQ Math. Log. Q.* **54**:6 (2008), 641–651. MR Zbl
- [19] J. Krueger, “Internal approachability and reflection”, *J. Math. Log.* **8**:1 (2008), 23–39. MR Zbl
- [20] J. Krueger, “Internally club and approachable for larger structures”, *Fund. Math.* **201**:2 (2008), 115–129. MR Zbl
- [21] J. Krueger, “Some applications of mixed support iterations”, *Ann. Pure Appl. Logic* **158**:1-2 (2009), 40–57. MR Zbl

- [22] M. Viale, “Guessing models and generalized Laver diamond”, *Ann. Pure Appl. Logic* **163**:11 (2012), 1660–1678. MR Zbl

Received January 18, 2023. Revised October 19, 2024.

MAXWELL LEVINE
INSTITUTE OF MATHEMATICS
UNIVERSITY OF FREIBURG
FREIBURG IM BREISGAU
GERMANY

maxwell.levine@mathematik.uni-freiburg.de

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Fakultät für Mathematik
Universität Wien
Vienna, Austria
matthias.aschenbrenner@univie.ac.at

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Atsushi Ichino
Department of Mathematics
Kyoto University
Kyoto 606-8502, Japan
atsushi.ichino@gmail.com

Robert Lipshitz
Department of Mathematics
University of Oregon
Eugene, OR 97403
lipshitz@uoregon.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Dimitri Shlyakhtenko
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
shlyakht@ipam.ucla.edu

Ruixiang Zhang
Department of Mathematics
University of California
Berkeley, CA 94720-3840
ruixiang@berkeley.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2024 is US \$645/year for the electronic version, and \$875/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2024 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 332 No. 1 September 2024

Lagrangian cobordism of positroid links	1
JOHAN ASPLUND, YOUNGJIN BAE, ORSOLA CAPOVILLA-SEARLE, MARCO CASTRONOVO, CAITLIN LEVERSON and ANGELA WU	
Liouville equations on complete surfaces with nonnegative Gauss curvature	23
XIAOHAN CAI and MIJIA LAI	
On moduli and arguments of roots of complex trinomials	39
JAN ČERMÁK, LUCIE FEDORKOVÁ and JIŘÍ JÁNSKÝ	
On the transient number of a knot	69
MARIO EUDAVE-MUÑOZ and JOAN CARLOS SEGURA-AGUILAR	
Preservation of elementarity by tensor products of tracial von Neumann algebras	91
ILIJAS FARAH and SAEED GHASEMI	
Efficient cycles of hyperbolic manifolds	115
ROBERTO FRIGERIO, ENNIO GRAMMATICA and BRUNO MARTELLI	
On disjoint stationary sequences	147
MAXWELL LEVINE	
Product manifolds and the curvature operator of the second kind	167
XIAOLONG LI	