

*Pacific
Journal of
Mathematics*

**THETA-INDUCED DIFFUSION ON TATE ELLIPTIC CURVES
OVER NONARCHIMEDEAN LOCAL FIELDS**

PATRICK ERIK BRADLEY

Volume 334 No. 1

January 2025

THETA-INDUCED DIFFUSION ON TATE ELLIPTIC CURVES OVER NONARCHIMEDEAN LOCAL FIELDS

PATRICK ERIK BRADLEY

A diffusion operator on the K -rational points of a Tate elliptic curve E_q is constructed, where K is a nonarchimedean local field, as well as an operator on the Berkovich analytification E_q^{an} of E_q . These are integral operators for measures coming from a regular 1-form, and kernel functions constructed via theta functions. The second operator can be described via certain nonarchimedean curvature forms on E_q^{an} . The spectra of these self-adjoint bounded operators on the Hilbert spaces of L^2 -functions are identical and found to consist of finitely many eigenvalues. A study of the corresponding heat equations yields a positive answer to the Cauchy problem, and induced Markov processes on the curve. Finally, some geometric information about the K -rational points of E_q is retrieved from the spectrum.

1. Introduction

The uniformisation of a projective algebraic curve X addresses the problem of finding its universal cover Ω and the action of its fundamental group Γ such that $X \cong \Omega/\Gamma$. In the case of curves defined over the complex numbers, this problem is solved by showing that the only simply connected Riemann surfaces are up to conformal equivalence the open unit disc, the complex plane and the Riemann sphere. Hence, in the case of an elliptic curve E , the fundamental group $\pi_1(E, 0)$ can be represented as a lattice Λ acting on \mathbb{C} . In other words, $E \cong \mathbb{C}/\Lambda$. In the case of a nonarchimedean local field K , there are far more simply connected subdomains of $\mathbb{P}_1(K)$. However, the Tate elliptic curve $E_q(K)$ does have a uniformisation of the form $E_q(K) \cong K^\times/q^{\mathbb{Z}}$, where the multiplicative group K^\times is indeed simply connected, and the fundamental group is a multiplicative lattice depending on a parameter q in the ring of integers of K . Unlike in the complex case, not every elliptic curve has a uniformisation of this kind. It turns out that the ones for which this is possible are precisely the ones with split multiplicative reduction [17, Theorem V.5.3].

MSC2020: primary 14H52; secondary 58J35.

Keywords: Tate curve, p -adic diffusion, theta function, Laplacian, torsion points.

Brownian motion can be modelled by the heat equation, which describes a diffusion process on a given space. One of its mathematical meanings is to be a tool for extracting information about the space from the diffusion equation. Already the spectrum of the corresponding Laplacian operator reveals something about the space. In the p -adic context, the heat equation has been studied only on local fields [27] and on their subspaces [28]; see also [14] for Brownian motion on \mathbb{Q}_p^d . A multivariate version of the p -adic heat equation is studied in [15], where it is shown that its fundamental solution is the transition density of a Markov process. The author also studied heat equations on Mumford curves [4], inspired by W. Zúñiga-Galindo's approach to studying p -adic heat equations on finite graphs [28] and interpreting diffusion as local transitioning between the vertices of a reduction graph of the curve along edges. This means that only information about the skeleton of the Mumford curve, and these include Tate curves, should be retrievable through this approach. A retrieval of a reduction graph in the case of a Mumford curve or a Mumford-uniformisable abelian variety using Vladimirov–Taibleson-type operators as studied in [18; 24] was effected in [6] from their infinite spectra. So, a diffusion operator allowing to extract geometric information about p -adic spaces becomes a desired object.

Here we address the stated desideratum in the simplest higher-genus case of a Tate elliptic curve E_q by using the invariant measure $|\omega|$ on $E_q(K)$ coming from a regular differential form ω on $E_q(K)$. Notice that p -adic elliptic curves which are also 1-dimensional p -adic manifolds are necessarily Tate curves (possibly after a finite field extension), because elliptic curves with good reduction are not locally embeddable into the local field. So, this article studies the simplest example of a compact p -adic manifold which is also a projective algebraic variety.

The Berkovich analytification E_q^{an} captures all reduction graphs associated with the rational points of the curve for any (complete) extension field of K , it is natural to also construct an operator which acts on functions on E_q^{an} . In this way, the necessity to fix the field K becomes obsolete, as long as it is sufficiently large. This more unifying approach becomes possible through an integral operator which can be described as integrating over a signed Radon measure presented as the difference between two curvature forms of the type

$$c_1(\mathcal{L} \|\cdot\|),$$

where $\|\cdot\|$ is a continuous subharmonic metric on a line bundle \mathcal{L} which is ample. This so-called *Chambert-Loir measure* was developed in the context of non-archimedean Arakelov theory and is in the present case a finite linear combination of Dirac measures on the skeleton of E_q^{an} , and the metrics are obtained because the nonarchimedean Calabi–Yau problem for smooth strictly K -analytic curves has

been solved by Thuillier [23]. This is a case in which the Monge–Ampère equation

$$c_1(\mathcal{L}, \|\cdot\|)^n = \nu$$

for a given positive Radon measure on the analytification of an n -dimensional algebraic variety X and an invertible sheaf \mathcal{L} on X can be solved. In higher dimensions, this has also been solved under certain mild restrictions quite recently [3; 9]. The diffusion operator on E_q^{an} is obtained from the operator \mathcal{H}_θ on $E_q(K)$ described below essentially by pushing forward to the skeleton, and thus has an identical behaviour as \mathcal{H}_θ concerning the spectrum and the heat equation.

Before describing the operator \mathcal{H}_θ , let us view its first application (Theorem 4.1). It turns out that its spectrum can detect the presence or absence of two kinds of points:

- 2-torsion points in $E_q(K)$, and
- third roots of certain types of points in $E_q(K)$ depending on the valuation $\nu(q)$ of the curve parameter q .

In the case of their presence, also the absolute values of their representatives in the fundamental domain are detected. Furthermore:

- the parity of the integer number $\nu(q)$,

where $\nu(x) = -\log_{p^f}(x)$, is detectable from the spectrum. Here, f is the degree of the residue field extension. These results depend on knowing the field K , in particular its uniformiser π , and stands in contrast to [6], where the reduction graph structure is revealed from the spectrum. This result opens a new bridge between ultrametric diffusion and arithmetic. In particular the fact that the detectability of whether certain kinds of K -rational points exist can now be read off the spectrum of a p -adic operator begs the question of a deeper connection between p -adic diffusion on an elliptic curve and its arithmetic.

The new diffusion operator developed here is, like all p -adic diffusion operators known so far, an integral operator whose kernel function is defined via the absolute value of a meromorphic function on the Tate curve. This generalises the usual p -adic kernel functions, since those are given by functions depending on the absolute value of a coordinate function on the affine line, aka radial functions. However, unlike the Vladimirov–Taibleson operator, it is not expressed as a pseudodifferential operator, meaning that the Fourier transform is not used here. Its kernel function is constructed via theta functions on the multiplicative group K^\times which are invariant under the uniformising group $q^\mathbb{Z}$. It turns out that, in this way, a bounded linear operator is obtained which is self-adjoint on the Hilbert space of L^2 -functions with respect to integrating along the invariant measure $|\omega|$. More precisely:

Theorem 2.15. *The space $L^2(E_q(K), |\omega|)$ has an orthogonal decomposition*

$$L^2(E_q(K), |\omega|) = L^2(E_q(K))_\sigma \oplus L^2(E_q(K))_0$$

into \mathcal{H}_θ -invariant subspaces. The subspace $L^2(E_q(K))_\sigma$ is of finite dimension $v(q)$ and spanned by the indicator functions supported on the circles $S_\ell(0)$ of radius p^{-f_ℓ} centred in zero, with $\ell = 0, \dots, v(q) - 1$, and the subspace $L^2(E_q(K))_0$ is spanned by the wavelets on $E_q(K)$ obtained by pull-back of Kozyrev wavelets. The spectrum of \mathcal{H}_θ as a linear operator on $L^2(E_q(K), |\omega|)$ consists entirely of eigenvalues. These are the eigenvalues of a certain matrix acting on $L^2(E_q(K))_\sigma$, and the negative degree values corresponding to normalised wavelets on $E_q(K)$ supported in circles $S_k(x)$ with x inside the annulus $\{u \in K^\times \mid |q| < |u| \leq 1\}$, and $k = 0, \dots, v(q) - 1$.

This unsurprising result comes from viewing the Hilbert space $L^2(E_q(K), |\omega|)$ as a decomposition into a direct sum of L^2 -spaces of finitely many p -adic discs, each endowed with an individually scaled Haar measure coming from the differential form ω . What is new is that here the degree eigenvalues are grouped according to the circles contained inside the annulus forming a fundamental domain for $E_q(K)$. Hence, there is not only an infinite part of their ‘‘multiplicities’’ coming from shrinking the wavelets, but also an inherent finite part coming from a partitioning of circles into finitely many maximal subdiscs.

The next result is about very general p -adic operators on K -analytic manifolds of any finite dimension giving rise to Feller semigroups and solutions to corresponding heat equations under some technical assumptions. And the operators are more general of the form

$$\mathcal{J}f(x) = \int_X \{j(x, y)f(y) - j(y, x)f(x)\} |\omega(y)|,$$

as considered in [29; 30], where the kernel function $j(x, y)$ is such that the positive maximum principle is satisfied.

Theorem (cf. Theorem 3.3). *Under some technical assumptions, there exists a probability measure $p_t(x, \cdot)$ with $t \geq 0$, $x \in X$, on the Borel σ -algebra of X such that the Cauchy problem*

$$\begin{aligned} u(\cdot, t) &\in C^1([0, \tau], C_0(X, \mathbb{R})), \\ \frac{\partial}{\partial t} u(x, t) &= \int_X (j(x, y)u(y, t) - j(y, x)u(x, t)) |\omega(t)|, t \in [0, \tau], x \in X, \\ u(x, 0) &= u_0(x) \in C_0(X, \mathbb{R}), \end{aligned}$$

has a unique solution of the form

$$h(t, x) = \int_X h_0(y) p_t(x, |\omega(y)|).$$

In addition, $p_t(x, \cdot)$ is the transition function of a strong Markov process whose paths are right continuous and have no discontinuities other than jumps.

As a corollary, it follows that the new operator on $E_q(K)$ also describes a strong Markov process on $E_q(K)$ whose paths are right continuous and have no discontinuities other than jumps. The Cauchy problem for the corresponding heat equation has a positive answer depending uniquely on the initial condition in $C(E_q(K))$. The spectrum of the operator acting on $L^2(K)$ consists of finitely many eigenvalues, a part of which correspond to eigenfunctions of the Laplacian of a complete finite graph, and the other part comes from an infinite family of functions which restrict to the well-known p -adic Kozyrev wavelets supported inside a fundamental domain of $E_q(K)$, which is an annulus in K . Because of the transitions on a complete graph, one can view this new operator as nonlocal on a Tate curve, which is in contrast to the ones from [4] on Mumford curves.

The preceding general theorem, inspired by biology (cf. [29]) now raises the question, for further research, whether and how it is possible to extract information about special K -rational points on elliptic curves via the study of a directed diffusion process.

Coming back to self-adjoint operators, it is shown that \mathcal{H}_θ can be extended to an operator on E_q^{an} with corresponding properties on $L^2(E_q^{\text{an}})$; see Corollary 3.9. This extended operator $\mathcal{H}_{\theta, \sigma}$ can be written as integration against the Chambert-Loir measure $c_1(\mathcal{L}, \|\cdot\|_{\theta, x})$ for a suitably metrised ample line bundle \mathcal{L} :

Corollary 3.11. *The heat operator $\mathcal{H}_{\theta, \sigma}$ on $E_q^{\text{an}}(K)$ is obtained as an integral operator of the form*

$$\mathcal{H}_{\theta, \sigma} \psi(x) = \int_{E_q(K)^{\text{an}}} \psi c_1(\mathcal{L}, \|\cdot\|_{\theta, x}),$$

with metric $\|\cdot\|_{\theta, x} = e^{-g_{\theta, x}}$ and

$$g_{\theta, x} = \sum_{z \in \sigma(E_q^{\text{an}}(K))} \alpha_z g_z,$$

where $\alpha_z > 0$ and g_z is a continuous subharmonic function on E_q^{an} such that

$$c_1(\mathcal{L}, \|\cdot\|_z) = \delta_z$$

is a Dirac measure for $\|\cdot\|_z = e^{-g_z}$, $y \in \sigma(E_q^{\text{an}}(K)) \sqcup \sigma(x)$, and $x \in E_q^{\text{an}}(K)$.

Here, σ is the retraction map of E_q^{an} onto its skeleton.

A potential application outside of pure mathematics is seen in the analysis of topological data, in particular in the context of the ongoing DFG project *Distributed Simulation of Processes in Buildings and City Models*. In this context, an envisioned heat flow approach could lead to the possibility of verifying the topological correctness of CAD models obtained from point clouds on distributed computing systems.

The following section introduces the invariant measure $|\omega|$, explains how to construct a kernel function from theta functions, and studies the spectrum of the diffusion operator. The heat equation on $E_q(K)$, and the development of a diffusion process on E_q^{an} and its relationship to Chambert-Loir measures is treated in the third section. The short last section is devoted to extracting information about the Tate curve $E_q(K)$ from the spectrum.

Some words about notation are in order. The nonarchimedean local field used here is denoted by K . Its local ring is O_K and has a unique maximal ideal \mathfrak{m}_K . The uniformiser of K is denoted by π and the absolute value $|\cdot|$ on K is scaled so that

$$|\pi| = p^{-f},$$

where f is the degree of the extension of the residue field O_K/\mathfrak{m}_K over the finite field \mathbb{F}_p with p elements. The prime p is assumed not to be 2 or 3, as this is needed occasionally in order to not have to deal with the intricacies occurring with these primes. The Haar measure on K is denoted by μ_K , but also by $|dx|$ if x is the variable of integration. The reason for writing it in this way is because we are also working with measures $|\omega|$ coming from a differential 1-form, and these are locally written as

$$\omega|_U = f dx,$$

where f is a function defined on an open piece U of the space under consideration. In this case, the measure is locally written as

$$|\omega(x)| = |f(x)| |dx|$$

if $x \in U$ is the variable of integration. Indicator functions will be written as

$$\Omega(x \in B)$$

where B is a measurable set, or as

$$\Omega(|f(x)|) := \Omega(x \in B(f))$$

with

$$B(f) = \{x \in K \mid |f(x)| \leq 1\}$$

for some function $f : K \rightarrow K$.

The following function spaces are used:

$$\begin{aligned}\mathcal{D}(X) &= \{f : X \rightarrow \mathbb{C} \mid f \text{ is locally constant with compact support}\}, \\ C(X) &= \{f : X \rightarrow \mathbb{C} \mid f \text{ is continuous}\}, \\ L^2(X, \nu) &= \{f : X \rightarrow \mathbb{C} \mid f \text{ is square integrable with respect to } \nu\},\end{aligned}$$

where ν is a positive Borel measure on a topological space X . The elements of $\mathcal{D}(X)$ are also called *test functions* on X . In the situation of this article, the space of test functions is dense in the other two. Here, the space $C(X)$ is a Banach space with respect to the supremum norm $\|\cdot\|_\infty$, and the second space is a Hilbert space with respect to the pairing induced by ν .

An introduction to the theory of Berkovich analytic spaces can be found in [21] or, of course, [1]. The results of that theory needed here are the existence of a skeleton $I(X)$ of the Berkovich analytification X^{an} of a smooth projective variety X over K , and a map

$$\sigma : X^{\text{an}} \rightarrow I(X)$$

which is a deformation retraction. The topology of X^{an} is Hausdorff, and X^{an} is locally path-connected [2]. In the 1-dimensional case, the skeleton $I(X^{\text{an}})$ is a metrised graph which for the Tate curve can be identified with a circle.

2. Diffusion on the rational points of a Tate elliptic curve

From [20], [17] and [10, Chapter 5.1], one can take a comprehensive view on Tate's elliptic curves. Invariant functions on Tate curves are constructed in [20] and in [10, Chapter 5.1] via theta functions. This allows one to explicitly construct a measure and a kernel function.

Theorem 2.1 (Tate's uniformisation of Tate curves). *Let $q \in K^\times$, $|q| < 1$. Set*

$$s_k(q) = \sum_{n \geq 1} \frac{n^k q^n}{1 - q^n}, \quad a_4(q) = -5s_3(1), \quad a_6(q) = \frac{-5s_3(q) + 7s_5(q)}{12}.$$

Then $a_4(q)$ and $a_6(q)$ converge in K . The equation

$$E_q : y^2 + xy = x^3 + a_4(q)x + a_6(q)$$

defines an elliptic curve with discriminant

$$\Delta(E_q) = q \prod_{n \geq 1} (1 - q^n)^{24}$$

and j -invariant

$$j(E_q) = \frac{1}{q} + \sum_{n \geq 0} c(n)q^n,$$

with $c(n) \in \mathbb{Z}$.

A proof is given in [17, 1, Theorem V.3.1] (the coefficient 5 in $a_4(q)$ was inadvertently omitted there). The reader could also consult [16, §3 (37)] or [10, Theorem 5.1.10].

Lemma 2.2 [17, Chapter V.3]. *If*

$$|q| < |u| < |q|^{-1}$$

then the coordinates of E_q can be written as

$$X(u, q) = \frac{u}{(1-u)^2} + \sum_{d \geq 1} \left(\sum_{m|d} m(u^m + u^{-m} - 2) \right) q^d,$$

$$Y(u, q) = \frac{u^2}{(1-u)^3} + \sum_{d \geq 1} \left(\sum_{m|d} \left(\frac{(m-1)m}{2} u^m - \frac{m(m+1)}{2} u^{-m} + m \right) \right) q^2$$

Theorem 2.3 (Tate). *The Tate elliptic curve E_q can be written as*

$$E_q = \mathbb{C}_p^\times / q^{\mathbb{Z}}$$

with $q \in K^\times$ such that $|q| < 1$. The meromorphic functions on $E_q(K)$ are precisely the $q^{\mathbb{Z}}$ -invariant meromorphic functions on K^\times .

Proof. [20, Theorem 1], or [10, Theorem 5.1.4] together with the Weierstrass model for $\mathbb{C}_p/q^{\mathbb{Z}}$ [10, Theorem 5.1.10]. \square

2A. Measure on $E_q(K)$. Any nonsingular projective algebraic curve X of positive genus has regular differential 1-forms. If X is defined over a nonarchimedean local field K , a regular differential 1-form ω on X gives rise to a positive measure $|\omega|$ on the space $X(K)$ of K -rational points outside the vanishing locus of ω [11, Chapter 7.4]. A. Weil calls this a *gauge measure* [25, Chapter II.2.2]. Since the zero set of ω locally on each chart $U \rightarrow K$ has Haar measure zero [26, Lemma 3.1], one has a natural extension of $|\omega|$ to all of $X(K)$, where it locally on a chart U takes the form

$$\int_U |\omega| = \int_{U'} |f_U| d\mu_K$$

with μ_K the Haar measure on K , and $\omega|_U = f_U dx$ for some K -analytic function f_U with local coordinate x . Notice that in this description here, it is assumed that $X(K)$ is a 1-dimensional K -analytic manifold, which is the case only if X is a Mumford curve.

The Haar measure on K will also be written as

$$\mu_K = |dx|$$

in order to make visible the local coordinate x , and in order to distinguish it from the differential 1-form dx on \mathbb{A}_K^1 . In the case of the measure $|\omega|$, this will be effected by writing

$$|\omega(x)|$$

in particular when a clarification of the variable of integration is needed.

Returning to the Tate elliptic curve E_q , now assume that the parameter u from Lemma 2.2 is in the annulus

$$A_K(q) = \{u \in K^\times \mid |q| < |u| \leq 1\}$$

and look at the holomorphic differential

$$(1) \quad \omega = \frac{dx}{2y+x} = \frac{dX(u, q)}{2Y(u, q) + X(u, q)}$$

on E_q . By the Riemann–Roch Theorem [10, Proposition 5.1.2(2)], any holomorphic differential 1-form has no zeros on nonsingular projective curves of genus 1. Hence, the measure $|\omega|$ corresponding to ω , defined as in (1), is a gauge form in the sense of [25, Chapter II.2.2]. The following lemma has an explicit description of this measure:

Lemma 2.4. *On the annulus $A_K(q)$, the measure $|\omega|$ coming from ω , defined as in (1), takes the form*

$$|\omega(u)| = \frac{|du|}{|u|}$$

for $u \in A_K(q)$, where it is a positive measure which defines an $E_q(K)$ -invariant measure on the K -rational points $E_q(K)$ of the Tate curve E_q .

Proof. With

$$F(u, q) = 2Y(u, q) + X(u, q)$$

one obtains

$$\omega = \omega(u) = F(u, q)^{-1} \frac{dX(u, q)}{du} du$$

This yields the real-valued measure

$$|\omega(u)| = \left| F(u, q)^{-1} \frac{dX(u, q)}{du} \right| |du|$$

Now, Lemma 2.2 implies that

$$\begin{aligned} \left| F(u, q)^{-1} \frac{dX(u, q)}{du} \right| &= \left| \frac{(1-u)^3}{u(1+u)} \cdot \frac{1+u}{(1-u)^3} \right| + \text{higher order terms in } q \\ &= \frac{1}{|u|}, \end{aligned}$$

because all coefficients of q^d with $d \geq 1$ each have absolute value at most one, and because $|q| < |u| \leq 1$. It follows that

$$|\omega(u)| = \frac{|du|}{|u|}$$

as asserted. Since $A_K(q)$ is a fundamental domain of the action of the discrete group $q^{\mathbb{Z}}$, this proves that $|\omega(u)|$ with $u \in A_K(q)$ defines the positive measure on E_q associated with the holomorphic differential form ω .

To show how the measure $|\omega|$ pulls back under the action of the group $E_q(K)$, denote with $f_P : E_q \rightarrow E_q$ the translation with $P \in E_q(K)$. We claim that

$$f_P^* |\omega| = |\omega|$$

For this, w.l.o.g. take a representative v of P in the annulus $A_K(q)$, since $A(q)$ is a fundamental domain for $E_q(K)$. Then

$$f_P^* |\omega(u)| = |\omega(vu)| = \frac{|dvu|}{|vu|} \stackrel{(*)}{=} \frac{|v| |du|}{|v| |u|} = \frac{|du|}{|u|} = |\omega(u)|$$

as asserted, where $(*)$ holds true because of the transformation rule for the Haar measure $|du|$ on K . \square

2B. A Kernel function on a Tate curve. A theta function on K^\times for $E_q(K)$ is given by

$$\theta(z) = \prod_{n \geq 0} (1 - q^n z^{-1}) \prod_{n > 0} (1 - q^n z)$$

with $z \in K^\times$, [10, Definition 5.1.8]. For $x \in K^\times$, one defines

$$\theta_x(z) = \theta(x^{-1}z)$$

and obtains that any invertible meromorphic function on $E_q(K)$ can be written as

$$(2) \quad f(z) = \lambda \prod_{i=1}^N \theta_{x_i}(z)^{n_i}$$

where $\lambda \in K^\times$, and $x_i \in K^\times$ are the representatives of the finitely many zeros and poles of f with corresponding multiplicities $n_i \in \mathbb{Z}$, provided that

$$(3) \quad \sum_{i=1}^N n_i = 0 \quad \text{and} \quad \prod_{i=1}^N x_i^{n_i} \in q^{\mathbb{Z}};$$

see [10, p. 128]. Here the condition $\sum_{i=1}^N n_i = 0$ holds because $\text{div}(f)$ has degree zero according to Riemann–Roch [10, Proposition 5.1.2(2)].

Remark 2.5. The annulus $A_K(q)$ is a fundamental domain for the action of $q^{\mathbb{Z}}$ on K^\times giving rise to the universal covering map

$$\rho : K^\times \rightarrow E_q(K)$$

and Lemma 2.4 says that the measure $|\omega|$ pulls back to the Haar measure on K^\times , normalised such that circles centred in zero all have volume $1 - p^{-f}$, and so is itself a Haar measure on the Tate curve $E_q(K)$.

Let $x, y \in E_q(K)$, and define the function

$$(4) \quad g(x, y) = \frac{\theta(x^{-1}y)\theta(y^{-1}x)}{\theta(xy)^2} = \frac{\theta_x(y)}{\theta_{x^{-1}}(y)} \cdot \frac{\theta_y(x)}{\theta_{y^{-1}}(x)},$$

which is $q^{\mathbb{Z}}$ -invariant in both variables. As each factor on the right-hand side of (4) satisfies condition (3), it defines a meromorphic function on $E_q(K)$ for any fixed $x \in E_q(K)$, and also for any fixed $y \in E_q(K)$. Its divisor as a function on the surface $E_q(K)^2$ is

$$\operatorname{div}(g) = 2[y - x] - 2[y - x^{-1}] \in \operatorname{Div}(E_q(K)^2),$$

whose double zero is the diagonal $V(x - y)$, and whose double pole is the curve $V(xy - 1)$ in $E_1(K)^2$, where

$$V(F) = \{x \in X \mid F(x) = 0\}$$

is the vanishing set of a function $F \in K(X)$ for a variety X over K .

Lemma 2.6. *It holds true that*

$$|g(x, y)| = \frac{|\tilde{x}\tilde{y}||\tilde{x} - \tilde{y}|^2}{|1 - \tilde{x}\tilde{y}|^2}$$

for suitable representatives $(\tilde{x}, \tilde{y}) \in (K^\times)^2$ of $x, y \in E_q(K)^2$.

Proof. If $z \in A_q(K)$, then $|\theta(z)| = |1 - z^{-1}|$, which implies

$$|g(x, y)| = \frac{|1 - \tilde{y}^{-1}\tilde{x}||1 - \tilde{x}^{-1}\tilde{y}|}{|1 - \tilde{x}^{-1}\tilde{y}^{-1}|^2} = \frac{|\tilde{x} - \tilde{y}|^2}{|\tilde{x} - \tilde{y}^{-1}||\tilde{x}^{-1} - \tilde{y}|} = \frac{|\tilde{x}\tilde{y}||\tilde{x} - \tilde{y}|^2}{|1 - \tilde{x}\tilde{y}|^2}. \quad \square$$

A neighbourhood of the poles of the map $A_K(q)^2 \rightarrow \mathbb{R}$, $(x, y) \mapsto |g(x, y)|$ is the set

$$\mathcal{P}_g = \{(x, y) \in A_K(q)^2 \mid |xy| = 1 \text{ and } |x| \neq |y|\}$$

where $|x| \neq |y|$ is required, because if $|x| = |z|$ with $z \in E_q(K)[2] \cap A_K(q)$, then $x = uz$ with $|u| = 1$ and

$$\frac{|x - y|}{|1 - xy|} = \frac{|x||1 - x^{-1}y|}{|1 - xy|} = \frac{|x||1 - uz|}{|1 - uz|} = |x|,$$

for any $y \in A_K(q)$. This means that pairs (x, y) with $|x| = |y|$ are not poles of $|g(x, y)|$. The $|\omega|$ -measure of the set \mathcal{P}_g is nonzero, because it is a union of circles. Hence, it is reasonable to define the kernel function

$$H_\theta(x, y) = \begin{cases} |g(x, y)|, & \nexists \text{ representatives } (\tilde{x}, \tilde{y}) \in \mathcal{P}_g, \\ 1, & \exists \text{ representatives } (\tilde{x}, \tilde{y}) \in \mathcal{P}_g \end{cases}$$

for $x, y \in E_q(K)$. In this way, we obtain the linear operator

$$\mathcal{H}_\theta \psi(x) = \int_{E_q(K)} H_\theta(x, y) (\psi(y) - \psi(x)) |\omega(y)|$$

for $\psi \in \mathcal{D}(E_q(K))$ and its extension to a linear operator on $C(E_q(K))$ and on $L^2(E_q(K))$.

The significance of this kernel function is that it is controlled from the universal covering of the Tate curve via the function θ . This is an example of an operator invariant under the action of the fundamental group of $E_q(K)$, and generalises the concept of radial functions which is used for constructing kernel functions for integral operators on function spaces over K .

The corresponding degree function is

$$\text{deg}_{\mathcal{H}_\theta} : E_q(K) \rightarrow \mathbb{R}, \quad x \mapsto \int_{E_q(K)} H_\theta(x, y) |d\omega(y)|$$

and is defined wherever the integral converges. The adjacency operator is defined as

$$\mathcal{A}_\theta \psi(x) = \int_{E_q(K)} H_\theta(x, y) \psi(y) |\omega(y)|$$

for $x \in E_q(K)$.

Consider a partition of unity

$$1 = \sum_{k=0}^{v(q)-1} \eta_k,$$

with

$$\eta_\ell(x) = \Omega(x \in S_\ell) \in \mathcal{D}(E_q(K)),$$

where S_ℓ is the circle

$$S_\ell = S_\ell(0) = \{x \in K \mid |x| = |\pi^\ell|\}$$

for $\ell \in \mathbb{Z}$. Of interest are the matrix elements

$$A_\sigma(k, \ell) = \frac{\mathcal{A}_\theta \eta_\ell(\pi^k)}{1 - |\pi|}$$

for $k, \ell \in \mathbb{Z}/v(q)\mathbb{Z}$.

Now assume that $|q| < |\pi|^2$. All exponents m in expressions like

$$|w|^m$$

for $w \in A_K(q)$ representing a point in $E_q(K)$ are to be understood modulo $v(q)$, if not stated otherwise.

$$\mathbf{Lemma\ 2.7.} \quad A_\sigma(k, \ell) = \begin{cases} |\pi|^{k+\ell+2\min(k,\ell)} & \text{if } k + \ell \not\equiv 0, k \not\equiv \ell \pmod{v(q)}, \\ \frac{|\pi|^{4\ell}}{1 - |\pi|^3} & \text{if } k + \ell \not\equiv 0, k \equiv \ell \pmod{v(q)}, \\ 1 & \text{if } k + \ell \equiv 0 \pmod{v(q)}, \end{cases}$$

where in the expression $\min(k, \ell)$ it is assumed that $k, \ell \in \{0, \dots, v(q)\}$.

Proof. Case $k + \ell \equiv 0 \pmod{v(q)}$. This happens if and only if $|\pi^k \pi^\ell| \in |q|^\mathbb{Z}$. In this case,

$$A_\theta \eta_\ell(\pi^k) = \int_{|y|=|\pi|^\ell} |\omega(y)| = 1 - |\pi|,$$

which proves the assertion in this case.

Case $k + \ell \not\equiv 0 \pmod{v(q)}$ and $k \not\equiv \ell \pmod{v(q)}$. This happens if and only if $|\pi^k \pi^\ell| \notin |q|^\mathbb{Z}$ and $|\pi^k| \neq |\pi^\ell|$. If $0 \leq k < \ell < v(q)$, then

$$A_\theta \eta_\ell(\pi^k) = |\pi|^{k+\ell} \int_{|y|=|\pi|^\ell} |\pi|^{2k} \frac{dy}{|y|} = |\pi|^{3k+\ell} (1 - |\pi|),$$

while if that $0 \leq \ell < k < v(q)$, then

$$A_\theta \eta_\ell(\pi^k) = |\pi|^{k+\ell} \int_{|y|=|\pi|^\ell} |\pi|^{2\ell} \frac{|dy|}{|y|} = |\pi|^{k+3\ell} (1 - |\pi|).$$

This proves the assertion in this case.

Case $k + \ell \not\equiv 0 \pmod{v(q)}$ and $k \equiv \ell \pmod{v(q)}$. This is the case if and only if $|\pi^k| = |\pi^\ell|$. Then

$$\begin{aligned} A_\theta \eta_\ell(\pi^k) &= |\pi|^{2\ell} \int_{|y|=|\pi|^\ell} |\pi^\ell - y|^2 \frac{|dy|}{|y|} \\ &= |\pi|^\ell (1 - |\pi|) \sum_{v=\ell}^{\infty} |\pi|^{3v} = \frac{|\pi|^{4\ell} (1 - |\pi|)}{1 - |\pi|^3}. \end{aligned}$$

This proves the assertion in the remaining case. □

As an immediate consequence of Lemma 2.7, we have:

Corollary 2.8. *It holds true that*

$$\frac{\deg \mathcal{J}_{\mathcal{C}_\theta}(x)}{1 - |\pi|} = 1 + |\pi|^k \sum_{\substack{\ell=0 \\ v(q) \nmid \ell+k}}^{k-1} |\pi|^{3\ell} + |\pi|^{3k} \sum_{\substack{\ell=k+1 \\ v(q) \nmid \ell+k}}^{v(q)-1} |\pi|^\ell + \epsilon_q(k) |\pi|^{4k} > 0$$

where $|x| = |\pi|^k$ and

$$\epsilon_q(k) = \begin{cases} (1 - |\pi|^3)^{-1} & \text{if } v(q) \in 2\mathbb{Z} \text{ and } v(q) \nmid k, \\ 1 & \text{otherwise,} \end{cases}$$

with $k \in \{0, \dots, v(q) - 1\}$.

Lemma 2.9. *The following statements hold true:*

1. $H_\theta \in \mathcal{D}(E_q(K)^2)$.
2. The degree function $\deg_{\mathcal{H}_\theta}$ is defined everywhere on $E_q(K)$.
3. If $\sigma(x) = \sigma(x')$, then $\deg_{\mathcal{H}_\theta}(x) = \deg_{\mathcal{H}_\theta}(x')$. In particular, the degree function is locally constant on $E_q(K)$.

Proof. 1. Lemma 2.6 shows that H_θ is locally constant on the compact space $E_q(K)^2$, i.e., $H_\theta \in \mathcal{D}(E_q(K)^2)$.

2. This follows from 1.

3. This is an immediate consequence of Corollary 2.8. \square

2C. Spectrum of the diffusion operator on $E_q(K)$. We define a pairing $\langle \cdot, \cdot \rangle_\omega$ on $L^2(E_q(K), |\omega|)$ by

$$\langle \phi, \psi \rangle_\omega = \int_{E_q(K)} \phi(x) \overline{\psi(x)} |\omega(x)|$$

Lemma 2.10. *The operator \mathcal{H}_θ has the following properties:*

1. \mathcal{H}_θ is a bounded linear operator on $C(E_q(K))$ with respect to $\|\cdot\|_\infty$.
2. \mathcal{H}_θ is a self-adjoint bounded linear operator on $L^2(E_q(K), |\omega|)$.

Proof. 1. This is clear, since $H_\theta \in \mathcal{D}(E_q(K)^2)$ and $E_q(K)$ is a compact space.

2. Since the kernel function H_θ is symmetric, it follows that \mathcal{H}_θ is a symmetric operator on $L^2(E_q(K), |\omega|)$ which is also everywhere defined. By the Hellinger–Toeplitz Theorem [22, Theorem 2.10], it follows that \mathcal{H}_θ is also bounded on $L^2(E_q(K), |\omega|)$. \square

The following matrix is helpful:

$$(5) \quad L = (L_{k,\ell}) \in \mathbb{R}^{v(q) \times v(q)}, \quad \text{where } L_{k,\ell} = \mathcal{H}_\theta \eta_\ell(\pi^k).$$

Lemma 2.11. *The matrix L is diagonalisable. Its nonzero eigenvalues are all negative real numbers. The eigenvectors corresponding to eigenvalue zero are the constant vectors.*

Proof. All matrix elements

$$\mathcal{A}_\theta \eta_\ell(\pi^k) = (1 - |\pi|) A_\sigma(k, \ell)$$

are positive, by Lemma 2.7, and the degree function $\deg_{\mathfrak{g}_{\mathcal{C}_\theta}}$ is strictly positive, by Corollary 2.8. Also, the matrix L is symmetric. It follows that L is the Laplacian of a positively weighted, undirected complete graph G with $v(q)$ nodes. Hence, L is diagonalisable, and all eigenvalues are real, nonpositive. Since the graph G is connected, it follows that the only eigenvectors corresponding to eigenvalue zero are the constant vectors. \square

In [13], S.V. Kozyrev introduced p -adic wavelets and found they provide for an orthogonal decomposition of $L^2(\mathbb{Q}_p)$ into eigenspaces of the Vladimirov operator, meaning that its spectrum consists of eigenvalues corresponding to those wavelets, and that they are mutually orthogonal [13, Theorem 2]. They are defined as

$$\psi_{\gamma j n}(x) = p^{-\frac{\gamma}{2}} \chi(p^{\gamma-1} j x) \Omega(|p^\gamma x - n|_p)$$

with $\gamma \in \mathbb{Z}$, $n \in \mathbb{Q}_p/\mathbb{Z}_p$, and $j = 1, \dots, p-1$, and where $\chi : K \rightarrow S^1$ is an additive character into the complex unit circle. For a local field K , this extends readily to functions

$$\psi_{B,j}(x) = \mu_K(B)^{\frac{1}{2}} \chi_K(\pi^{d-1} \tau(j)) \Omega(x \in B)$$

where $\tau : O_K/\mathfrak{m}_K \rightarrow O_K$ is a lift of the residue field of K , $j \in (O_K/\mathfrak{m}_K)^\times$, and $B \subset K$ a disc of radius $|\pi|^{-d}$. These are the *Kozyrev wavelets* for K . It is well-known that the result [13, Theorem 2] readily extends to the local field K . However, not all of this theory is needed here. What is needed is this:

Lemma 2.12 (Kozyrev [12, Theorem 3.2.9]). *It holds true that*

$$\int_W \psi_{B,j}(y) |dy| = 0$$

for all measurable subsets $W \subseteq K$ containing B , and $j \in O_K/\mathfrak{m}_K$.

Definition 2.13. A *wavelet* on $E_q(K)$ is a function $\psi : E_q(K) \rightarrow \mathbb{C}$ supported inside an open $U \subset E_q(K)$ such that there is a local chart $\kappa : U \rightarrow K$ with $\kappa_* \psi$ a Kozyrev wavelet with support inside U .

It is immediately seen that a wavelet on $E_q(K)$ is defined by a Kozyrev wavelet $\psi_{B,j}$ supported inside $A_K(q)$. For simplicity, a wavelet on $E_q(K)$ will be written as such a Kozyrev wavelet $\psi_{B,j}$.

Lemma 2.14. *Let $\psi_{B,j}$ be a wavelet on $E_q(K)$. Then*

$$\int_W \psi_{B,j} |\omega| = 0$$

for all measurable subsets $W \subseteq E_q(K)$ containing the ball B , and $j \in O_K/\mathfrak{m}_K$.

Proof. View $\psi_{B,j}$ as a Kozyrev wavelet supported in $A_K(q)$. Since any ball in $A_K(q)$ is contained in some sphere

$$S_k(x) = \{z \in K \mid |z - x| = |\pi|^k\} \subset A_K(q)$$

with $k \in \{0, \dots, v(q) - 1\}$, obtain

$$\int_W \psi_{B,j} |\omega| = \int_{W'} \psi_{B,j}(y) \frac{|dy|}{|y|} = |x|^{-1} \int_{W'} \psi_{B,j}(y) |dy| = 0$$

where $W' \subset A_K(q)$ is a measurable set representing W , and the last equality follows from Lemma 2.12. \square

We can now prove the first theorem quoted in the Introduction; we repeat it for convenience.

Theorem 2.15. *The space $L^2(E_q(K), |\omega|)$ has an orthogonal decomposition*

$$L^2(E_q(K), |\omega|) = L^2(E_q(K))_\sigma \oplus L^2(E_q(K))_0$$

into \mathcal{H}_θ -invariant subspaces. The subspace $L^2(E_q(K))_\sigma$ is of finite dimension $v(q)$ and spanned by the indicator functions η_ℓ , with $\ell = 0, \dots, v(q) - 1$, and the subspace $L^2(E_q(K))_0$ is spanned by the wavelets on $E_q(K)$. The spectrum of \mathcal{H}_θ as a linear operator on $L^2(E_q(K), |\omega|)$ consists entirely of eigenvalues. These are the eigenvalues of the helpful matrix L of (5) acting on $L^2(E_q(K))_\sigma$, and the negative degree values $-\deg_{\mathcal{H}_\theta}(x)$ corresponding to normalised wavelets on $E_q(K)$ supported in $S_k(x)$ with $x \in A_K(q)$, and $k = 0, \dots, v(q) - 1$.

Proof. Since

$$\langle \psi_{B,j}, \eta_\ell \rangle_\omega = 0$$

for any pair (B, j) consisting of a ball $B \subset A_K(q)$ and $j \in O_K/\mathfrak{m}_K$ on the one hand, and of $\ell \in \{0, \dots, v(q) - 1\}$, it follows immediately that the given decomposition is an orthogonal decomposition of $L^2(E_q(K))$.

Clearly, the space $L^2(E_q(K))_\sigma$ is of finite dimension $v(q) \in \mathbb{N}$, spanned by the indicator functions η_ℓ with $\ell = 0, \dots, v(q) - 1$. Its orthogonal complement $L^2(E_q(K))_\sigma^\perp$ contains the space $L^2(E_q(K))_0$ spanned by the wavelets, and these form an orthonormal set in $L^2(E_q(K))_\sigma^\perp$. To see the converse inclusion, observe that $L^2(E_q(K))_0$ is isomorphic to a direct sum of spaces $L^2(B_r(a))_0$, where the $B_r(a)$ are p -adic balls forming a finite disjoint covering of $E_q(K)$, and

$$L^2(B_r(a)) = \mathbb{C} \Omega(x \in B_r(a)) \oplus L^2(B_r(a))_0,$$

with $L^2(B_r(a))$ having an orthonormal basis consisting of Kozyrev wavelets supported in $B_r(a)$; this is similar to [29, Proposition 2], except that in the actual decomposition of $L^2(E_q(K), |\omega|)$, each of the spaces $L^2(B_r(a))$ uses a rescaled

Haar measure, where the rescaling comes from the invariant differential form ω . This proves the asserted orthonormal basis of $L^2(E_q(K), |\omega|)$.

The eigenvalue of \mathcal{H}_θ corresponding to a wavelet $\psi_{B,j}$ with $B \subset S_k(x) \subset A_K(q)$ is now readily seen to be $-\deg_{\mathcal{G}_{\mathcal{H}_\theta}}(x)$, as asserted. This proves the theorem. \square

Remark 2.16. Since the support of a wavelet may be arbitrarily small, it follows that the operator \mathcal{H}_θ is not a compact linear operator on $L^2(E_q(K))$.

3. The heat equation

We now turn to the heat equation

$$(6) \quad \left(\frac{\partial}{\partial t} - \epsilon \mathcal{H}_\theta \right) u(x, t) = 0 \quad (\epsilon > 0)$$

for the Tate elliptic curve $E_q(K)$. The goal is to answer the corresponding Cauchy problem, verify that the heat operator $\epsilon \mathcal{H}_\theta$ generates a Markov process on $E_q(K)$ and write down the fundamental solution of (6).

The approach, suggested by the referee, will be to do this in a much more general setting in the following subsection, and then to specialise that result to the case of the heat operator $\epsilon \mathcal{H}_\theta$ on $E_q(K)$. Section 3B then will study a corresponding diffusion on the Berkovich analytification of E_q .

3A. Heat equation on K -analytic manifolds. Let X be a K -analytic manifold of dimension n , and let ω be a regular n -form on X . The gauge measure $|\omega|$ from [25, Chapter II.2.2] can be defined in any dimension $n \in \mathbb{N}$ outside of the vanishing set $V(\omega) \subset X$. Again, using [26, Lemma 3.1], it can be seen that $V(\omega)$ has measure zero, so $|\omega|$ again has a natural extension to all of X . The idea to follow here is to study operators of the form

$$(7) \quad \mathcal{J}f(x) = \int_X \{j(x, y)f(y) - j(y, x)f(x)\} |\omega(y)|$$

as considered in [29; 30]. In order to do this, write

$$X \times X = E_+ \cup E_-$$

with

$$E_+ \cap E_- = \Delta_X := \{(x, x) \mid x \in X\}$$

and a map

$$\iota : E_+ \rightarrow E_-, \quad (x, y) \mapsto (y, x)$$

which is bijective. Let

$$j_+ : E_+ \rightarrow \mathbb{R}_{\geq 0}, \quad j_- : E_- \rightarrow \mathbb{R}_{\geq 0}$$

be maps with

$$j_+|_{\Delta_X} = j_-|_{\Delta_X}$$

satisfying

$$(8) \quad j_-(x, y) \leq j_+(y, x) \quad \text{for } (x, y) \in E_-$$

and set

$$\begin{aligned} \mathcal{J}_- f(x) &= \int_{\{y:(x,y) \in E_-\}} \{j_-(x, y)f(y) - j_+(y, x)f(x)\} |\omega(y)|, \\ \mathcal{J}_+ f(x) &= \int_{\{y:(x,y) \in E_+\}} \{j_-(y, x)f(y) - j_+(x, y)f(x)\} |\omega(y)|, \end{aligned}$$

in order to obtain the operator

$$\mathcal{J} = \mathcal{J}_- + \mathcal{J}_+,$$

which acts on the space $C_0(X, \mathbb{R})$ of continuous functions vanishing at infinity via (7).

Lemma 3.1. *The operator \mathcal{J} satisfies the positive maximum principle.*

Proof. Let $f \in C_0(X, \mathbb{R})$, and let $x_0 \in X$ be a place where f takes its maximum, assumed positive. Then

$$\mathcal{J}f(x_0) \leq \int_X \{j(x_0, y) - j(y, x_0)\} |\omega(y)| f(x_0) \stackrel{(*)}{\leq} 0 \cdot f(x_0) \leq 0$$

where $(*)$ holds true because of (8). □

To be able to use the other conditions of the Hille–Yosida–Ray theorem below, observe that since X need not be compact, and \mathcal{J} may also be unbounded, a different approach than in the proof of [30, Theorem 3.1] is taken. The approach here is similar to that of the proof of [5, Lemma 5.1].

First, denote by $p_1, p_2 : X \times X \rightarrow X$ the projections onto the first and second coordinate, respectively. Let $\mathcal{P}(j) \subset X \times X$ be the set of poles of the kernel function $j(x, y)$, and let

$$P_z = p_2(p_1^{-1}(z) \cap \mathcal{P}(j))$$

for $z \in X$. Now, fix an atlas \mathcal{U} of X , and set

$$U_k(z) := \{y \in X \mid \text{locally with respect to } \mathcal{U} : d(y, P_z) \leq |\pi|^k\}$$

for $k \in \mathbb{N}$, where d is the distance in $U \in \mathcal{U}$, viewed as an analytic subdomain of K^n . Now, define

$$X_k(z) = X \setminus U_k(z)$$

and

$$\deg_k(z) = \int_{X_k(z)} j(y, z) |\omega(y)|$$

for $k \in \mathbb{N}$ and $z \in X$.

Assumption 1. The manifold X is assumed to be an open submanifold of a compact K -analytic manifold \bar{X} , and that $X \setminus \bar{X}$ is a zero set with respect to the measure $|\omega|$. The measure $|\omega|$ itself is assumed integrable on any chart $U \rightarrow K^n$, i.e., the differential form ω being on U of the form

$$\omega|_U = f_U dx_1 \wedge \cdots \wedge dx_n$$

with f_U a K -analytic function on U , satisfies

$$\int_U |f_U| |dx_1 \wedge \cdots \wedge dx_n| < \infty$$

where $|dx_1 \wedge \cdots \wedge dx_n|$ is the normalised Haar measure on K^n .

Assumption 2. It is assumed that $\mathcal{P}(j)$ is nowhere dense in $X \times X$, and that

$$\mu_{X^2}(\mathcal{P}(j)) = \int_{\mathcal{P}(j)} |\omega| \wedge |\omega| = 0.$$

Assumption 3. It is assumed that

$$\deg_k(z) < \infty$$

for all $z \in X$ and $k \in \mathbb{N}$.

Although the following result is not needed in the full generality of this subsection, it is nevertheless of independent interest.

Theorem 3.2. *Under Assumptions 1, 2, 3, and Hypothesis (8), the linear operator \mathcal{J} generates a Feller semigroup $e^{t\mathcal{J}}$ ($t \geq 0$) on $C_0(X, \mathbb{R})$.*

Proof. This is shown by checking the requirements for the Hille–Yosida–Ray theorem [8, Chapter 4, Lemma 2.1], which we do in three steps:

1. The domain of \mathcal{J} is dense in $C_0(X, \mathbb{R})$. This follows from Assumption 1.
2. The operator \mathcal{J} satisfies the positive maximum principle. This was proven in Lemma 3.1.
3. $\text{Ran}(\eta I - \mathcal{J})$ is dense in $C_0(X, \mathbb{R})$ for some $\eta > 0$. Since \mathcal{J} could be unbounded, a proof as in [30, Theorem 3.1] does not cover all cases here. Therefore, what follows is modelled after the proof of [5, Lemma 5.1]. The task is to find a solution of the equation

$$(9) \quad (\eta I - \mathcal{J})u = h$$

for some $\eta > 0$, and h in some dense subspace of $C_0(X, \mathbb{R})$. The equation can be formally rewritten as

$$(10) \quad u(z) - \frac{\int_X j(z, y)u(y) |\omega(y)|}{\eta - \deg(z)} = \frac{h(z)}{\eta - \deg(z)}$$

with

$$\deg(z) = \int_X j(y, z) |\omega(y)|,$$

which possibly does not converge, as \mathcal{J} might be unbounded. For this reason, study the operator

$$T_k u(z) = \frac{\int_{X_k(z)} j(y, z)u(y) |\omega(y)|}{\eta - \deg_k(z)}$$

for $k \gg 0$. Now,

$$|T_k u(z)| \leq \frac{\deg_k(z)}{|\eta - \deg_k(z)|} \|u\|_\infty < \infty,$$

by Assumption 3. This implies that

$$\|T_k\| \leq \frac{1}{1 - \eta/\deg_k(z)} < 1$$

for $k \gg 0$. Hence, $I - T_k$ has a bounded inverse as an operator on $C_0(X, \mathbb{R})$. Consequently, the range of $I - T_k$ is dense in $C_0(X, \mathbb{R})$ for $k \gg 0$. Now, let $h \in \mathcal{D}(X, \mathbb{R})$, the space of locally constant real-valued functions with compact support, and let $u_k, u_\ell \in C_0(X, \mathbb{R})$ be solutions of

$$(I - T_k)u_k = \frac{h}{\eta - \deg_k} \quad \text{and} \quad (I - T_\ell)u_\ell = \frac{h}{\eta - \deg_\ell}$$

for $k, \ell \gg 0$. Then

$$(11) \quad u_k - u_\ell = \frac{(I - T_\ell)(\eta - \deg_\ell) - (I - T_k)(\eta - \deg_k)}{(I - T_k)(I - T_\ell)(\eta - \deg_k)(\eta - \deg_\ell)} h$$

shows that (u_k) is a Cauchy sequence with respect to $\|\cdot\|_\infty$. Namely, first

$$(12) \quad \|T_k\| = \sup_{z \in X} \left| \frac{\deg_k(z)}{\eta - \deg_k(z)} \right| = \sup_{z \in X} \frac{1}{1 - \eta/\deg_k(z)}$$

is strictly increasing to 1 for $k \rightarrow \infty$. Hence, T_k converges to a bounded linear operator T on $C_0(X, \mathbb{R})$. Secondly, the numerator of the right hand side of (11) is

$$\eta(T_k - T_\ell) + (\deg_k - \deg_\ell) + (T_\ell \deg_\ell - T_k \deg_k)$$

whose first and second terms in norm become arbitrarily small as $\ell \geq k \rightarrow \infty$. The third term is

$$T_\ell \deg_\ell - T_k \deg_k = (T_\ell \deg_\ell - T_k \deg_\ell) + (T_k \deg_\ell - T_k \deg_k)$$

both of whose summands converge to 0 as $\ell \geq k \rightarrow \infty$. It follows that u_k converges to some $u \in C_0(X, \mathbb{R})$ which is seen to be a solution of (9) as follows: Namely, $(\eta + \deg_k)T_k$ converges to $(\eta + \deg)T$ for $k \rightarrow \infty$, where the limit operator coincides with the operator

$$Au(z) = \int_X j(z, y)u(y)|\omega(y)|,$$

which shows that the operator $T = \frac{A}{\eta - \deg}$ appearing in (10) is bounded. Now, u_k is a solution of

$$(\eta I - \mathcal{J}_k)u_k = h$$

with

$$\mathcal{J}_k = (\eta - \deg_k)T_k - \deg_k$$

which converges to \mathcal{J} for $k \rightarrow \infty$. As u_k converges to u , it follows that

$$(\eta I - \mathcal{J})u = (\eta I - \mathcal{J}_k)u + (\mathcal{J}_k - \mathcal{J})u$$

where

$$(\eta I - \mathcal{J}_k)u = (\eta I - \mathcal{J}_k)u_k + \mathcal{J}_k(u_k - u) = h + \mathcal{J}_k(u_k - u)$$

converges to h for $k \rightarrow \infty$, and

$$(\mathcal{J}_k - \mathcal{J})u \rightarrow 0$$

for $k \rightarrow \infty$. Hence, u is a solution of (9). This proves that $\text{Ran}(\eta I - \mathcal{J})$ contains $\mathcal{D}(X, \mathbb{R})$ which is dense in $C_0(X, \mathbb{R})$.

Since the limit operator of T_k does not depend on the choice of an atlas \mathcal{U} —cf. (12)—it follows that the existence of the solution u of (9) does also not depend on the choice of an atlas. This now proves the assertion. \square

We turn to the second theorem quoted in the Introduction.

Theorem 3.3. *There exists a probability measure $p_t(x, \cdot)$ with $t \geq 0$, $x \in X$, on the Borel σ -algebra of X such that the Cauchy problem*

$$\begin{aligned} u(\cdot, t) &\in C^1([0, \tau], C_0(X, \mathbb{R})), \\ \frac{\partial}{\partial t}u(x, t) &= \int_X (j(x, y)u(y, t) - j(y, x)u(x, t))|\omega(t)|, t \in [0, \tau], x \in X, \\ u(x, 0) &= u_0(x) \in C_0(X, \mathbb{R}), \end{aligned}$$

has a unique solution of the form

$$h(t, x) = \int_X h_0(y) p_t(x, |\omega(y)|).$$

In addition, $p_t(x, \cdot)$ is the transition function of a strong Markov process whose paths are right continuous and have no discontinuities other than jumps.

Proof. The proof will be the same as in the case of [28, Theorem 4.2], and is given here for the convenience of the reader.

Using the correspondence between Feller semigroups and transition functions, one sees from Theorem 3.2 that there is a uniformly stochastically continuous C_0 -transition function $p_t(x, |\omega|)$ satisfying condition (L) of [19, Theorem 2.10] such that

$$\exp(t\mathcal{J})h_0(x) = \int_X h_0(y) p_t(x, |\omega(y)|)$$

for $h_0 \in C_0(X, \mathbb{R})$; see, e.g., [19, Theorem 2.15]. Now, by using the correspondence between transition functions and Markov processes, there exists a strong Markov process whose paths are right continuous and have no discontinuities other than jumps; see [19, Theorem 2.12], for example. \square

We will now use Theorems 3.2 and 3.3 to address the following Cauchy problem: Find $h(t, x) \in C^1((0, \infty), E_q(K))$ such that

$$(13) \quad \left(\frac{\partial}{\partial t} - \epsilon \mathcal{H}_\theta \right) h(t, x) = 0, \quad h(0, x) = h_0(x),$$

for $t \geq 0, h_0 \in C(E_q(K))$.

Fact 3.4. *There exists a probability measure $p_t(x, \cdot)$ with $t \geq 0, x \in E_q(K)$, on the Borel σ -algebra of $E_q(K)$ such that the Cauchy problem (13) has a unique solution of the form*

$$h(t, x) = \int_{E_q(K)} h_0(y) p_t(x, |\omega(y)|).$$

In addition, $p_t(x, \cdot)$ is the transition function of a strong Markov process whose paths are right continuous and have no discontinuities other than jumps.

Proof. Hypothesis (8) is naturally satisfied by the kernel function. As Assumptions 1, 2, and 3 are clearly satisfied, Theorem 3.2 can be applied, and thus Theorem 3.3 yields the assertions. \square

As the corresponding semigroup is Feller, it describes a π -adic heat equation on $E_q(K)$. Consequently, there is a corresponding π -adic diffusion process in $E_q(K)$ attached to the heat equation (6).

Remark 3.5. As pointed out by the referee, it is possible to see Fact 3.4 as a special case of [29, Theorem 1], which itself finds an extension to general types of K -analytic manifolds in Theorem 3.3 here.

3B. Diffusion on the Berkovich analytification. In order to extend the diffusion theory to the Berkovich analytification E_q^{an} of the Tate curve E_q , it is necessary to solve the so-called Monge–Ampère equation

$$(14) \quad \mu = \lambda c_1(\mathcal{L}, \|\cdot\|)$$

where μ is a positive Radon measure on E_q^{an} . What this means is to find $\lambda > 0$ and a line bundle L on E_q^{an} such that the *Chambert-Loir measure* $c_1(\mathcal{L}, \|\cdot\|)$ on E_q^{an} coming from a subharmonic metric $\|\cdot\|$ on the line bundle L satisfies equation (14). Since E_q^{an} is a curve, this equation has been found to always have a solution by Thuillier in his dissertation [23, Corollary 3.4.18]. The solvability of the Monge–Ampère equation in a much more general nonarchimedean setting is settled in [9] in the mixed characteristic case.

Recall that a *harmonic function* on a weighted metrised graph G is a piecewise affine function $f : G \rightarrow \mathbb{R}$ such that for each point

$$\lambda_x(f) := \sum_{t \in T_x(G)} w_x(t) \lambda_{x,t} = 0$$

where $T_x(G)$ is the set of all tangent directions in each point of G , $w_x(t)$ the weight function in the point x along direction t , and $\lambda_{x,t}$ the linear part in the affine piece of f in the point x along t :

$$f = f(x) + \lambda_{x,t}(f)t$$

is the affine representation of f for this pair (x, t) . The corresponding map

$$\text{dd}^c : A^0(E_q^{\text{an}}) \rightarrow A^1(E_q^{\text{an}}), \quad f \mapsto \sum_{x \in G} \lambda_x(f) \delta_x,$$

is called *Laplacian*, and the kernel of dd^c consists of the harmonic functions in the space $A^0(E_q^{\text{an}})$ of global sections of piecewise affine functions on E_q^{an} , whereas $A^1(E_q^{\text{an}})$ are the global sections of the sheaf of locally finite signed measures on E_q^{an} .

A subharmonic function on an open $U \subseteq E_q^{\text{an}}$ is a function

$$u : U \rightarrow \mathbb{R} \cup \{-\infty\}$$

which is upper semicontinuous, not identically $-\infty$ on any connected component of U , and for any strictly K -affinoid domain Y of E_q^{an} and any harmonic function h on Y , it holds true that

$$u|_{\partial Y} \leq h|_{\partial Y} \quad \Rightarrow \quad u|_Y \leq h$$

A subharmonic metric on an invertible sheaf \mathcal{L} is a family of functions

$$\|\cdot\| : \Gamma(U, \mathcal{L}^\times) \rightarrow \mathbb{R}$$

with $U \subset E_q^{\text{an}}$ running through the open subsets, such that $-\log\|s\|$ is a subharmonic function on U , and for all open $U' \subset U \subset E_q^{\text{an}}$ and sections $s' \in \Gamma(U', \mathcal{L}^\times)$, $s \in \Gamma(U, \mathcal{L}^\times)$, it holds true that $s|_{U'} = fs'$ for some $f \in \Gamma(U', \mathcal{O}_{E_q^{\text{an}}}^\times)$ such that

$$-\log\|s\||_{U'} = -\log\|s'\| - \log|f|$$

Notice that the function $\log|f|$ is subharmonic [23, Proposition 3.1.6].

If $(\mathcal{L}, \|\cdot\|)$ is a metrised line bundle on E_q^{an} , then

$$\text{dd}^c \log\|s\|$$

for a section $s \in \Gamma(U, \mathcal{L}^\times)$ is in general only a distribution on $A^1(E_q^{\text{an}})$. Such is called a *current* of degree 1 on E_q^{an} . This current associated with $(\mathcal{L}, \|\cdot\|)$ is denoted by

$$c_1(\mathcal{L}, \|\cdot\|)$$

and is called its *curvature form*. It is positive if and only if $\|\cdot\|$ is a subharmonic metric on \mathcal{L} . And equation (14) asks for the existence of a curvature form representing a given positive Radon measure on E_q^{an} .

In order to relate the theory of metrised line bundles to the operator \mathcal{H}_θ on $E_q(K)$, observe that the measure

$$\mu_x = H_\theta(x, \cdot) |\omega(\cdot)|$$

is a positive Radon measure on $E_q(K)$. Hence, its pushforward

$$\mu_{\sigma(x)} := \sigma_* \mu_x$$

is a Radon measure on the skeleton $I(E_q^{\text{an}})$, and supported on $\sigma(E_q(K))$. Define also

$$\text{deg}_\theta(\sigma(x)) := \int_{I(E_q^{\text{an}})} \mu_{\sigma(x)}$$

for $x \in E_q(K)$, and solve the Monge–Ampère equation

$$\delta_z = c_1(\mathcal{L}, \|\cdot\|_z)$$

on the curve E_q^{an} , which is possible according to Thuillier with an ample line bundle \mathcal{L} on E_q^{an} . Notice that this is in fact the Calabi–Yau problem, where an ample line bundle \mathcal{L} on a projective K -variety X is given, and the question is whether for any given positive Radon measure ν on X^{an} with $\nu(X^{\text{an}}) = \text{deg}_\mathcal{L}(X)$, there exists a

continuous semipositive metric $\|\cdot\|$ on \mathcal{L} such that

$$\nu = c_1(\mathcal{L}, \|\cdot\|)^n$$

with $\dim(X^{\text{an}}) = n$. This has been recently solved in several important instances: if $n = 1$ and X is a smooth (analytic) curve [23]; when the residue characteristic is zero and X is smooth [3]; and in the mixed characteristic case for smooth X [9].

A *model* of the variety X is a normal scheme $\mathcal{X} \rightarrow \text{Spec } O_K$ which is flat and whose generic fibre is isomorphic to X . Given an ample line bundle \mathcal{L} on X , a *model metric* $\|\cdot\|_{\mathcal{L}}$ on \mathcal{L} is defined by an extension $\mathcal{L} \in \text{Pic}(\mathcal{X})_{\mathbb{Q}}$ of \mathcal{L} to some model \mathcal{X} of X . A model metric is *semipositive* if the line bundle \mathcal{L} is nef, i.e., if

$$\deg_{\mathcal{L}}(C) \geq 0$$

for all proper curves C on the special fibre \mathcal{X}_s of \mathcal{X} . A *semipositive continuous* metric $\|\cdot\|$ on \mathcal{L} is a uniform limit of semipositive metrics on \mathcal{L} . This gives rise to the *Chambert-Loir measure* $c_1(\mathcal{L}, \|\cdot\|)^n$, a positive Radon measure on X^{an} of mass $c_1(\mathcal{L})^n$, cf. [7]. So, the Calabi–Yau problem is to find a suitable continuous semipositive metric on \mathcal{L} such that the corresponding Chambert-Loir measure coincides with the given positive Radon measure.

In any case, we now obtain a measure

$$\nu_{\sigma(x)} = \mu_{\sigma(x)} - \deg_{\theta}(\sigma(x))\delta_{\sigma(x)}$$

on E_q^{an} for $x \in E_q(K)$ which is supported on the skeleton $\sigma(E_q(K)) \subset I(E_q^{\text{an}})$. This yields an operator

$$\mathcal{H}_{\theta, \sigma}\phi(z) = \int_{I(E_q^{\text{an}})} \phi \, d\nu_z$$

with $\phi \in \mathcal{D}(E_q^{\text{an}})$, where ν_z is the measure defined as

$$\nu_z = \begin{cases} \nu_{\sigma(x)} & \text{if } z = \sigma(x) \text{ for some } x \in E_q(K), \\ 0 & \text{otherwise,} \end{cases}$$

for $z \in E_q^{\text{an}}$.

Definition 3.6. A function $\phi : E_q^{\text{an}} \rightarrow \mathbb{C}$ is *radial* if $\phi(x) = \phi(\sigma(x))$ for all $x \in E_q^{\text{an}}$. The space of all radial test functions is denoted by $\mathcal{D}(E_q^{\text{an}})_{\sigma}$.

Helpful now is the averaging operator:

$$\mathcal{D}(E_q^{\text{an}}) \rightarrow \mathcal{D}(E_q^{\text{an}})_{\sigma}, \quad \phi \mapsto \text{av}(\phi)$$

with $\text{av}(\phi)$ defined as

$$x \mapsto \left(\int_{U(x)} |\sigma_*\omega| \right)^{-1} \int_{U(x)} \phi(y) |\sigma_*\omega(y)|$$

where

$$U(x) := \sigma^{-1}(\sigma(x))$$

is an open neighbourhood of $x \in E_q^{\text{an}}$. Clearly, functions of the form $av(\phi)$ are precisely the radial functions. Using the obviously defined pairing $\langle \cdot, \cdot \rangle_{\sigma_*\omega}$ on $L^2(E_q^{\text{an}}, |\sigma_*\omega|)$, obtain the orthogonal decomposition

$$L^2(E_q^{\text{an}}, |\sigma_*\omega|) = L^2(E_q^{\text{an}})_\sigma \oplus L^2(E_q^{\text{an}})_0$$

where $L^2(E_q^{\text{an}})_\sigma$ is spanned by the radial functions, and whose orthogonal complement is $L^2(E_q^{\text{an}})_\sigma^\perp = L^2(E_q^{\text{an}})_0$.

Let

$$U(\sigma(E_q(K))) := \sigma^{-1}(\sigma(E_q^{\text{an}}(K)))$$

The space $L^2(U(E_q(K))) \subset L^2(E_q^{\text{an}}, |\sigma_*\omega|)$ is a closed subspace and thus has the same property

$$(15) \quad L^2(U(E_q(K))) = L^2(U(E_q(K)))_\sigma \oplus L^2(U(E_q(K)))_0$$

and the part $L^2(U(E_q(K)))_0$ is spanned by the wavelets on $E_q(K)$. The reason is that that space coincides with the space of L^2 -functions on $E_q^{\text{an}}(K)$ supported outside the skeleton averaging to zero.

Lemma 3.7. *Assume that $\phi \in \mathcal{D}(E_q^{\text{an}})$ is a radial function. Then*

$$\mathcal{H}_{\theta,\sigma}\phi(z) = \begin{cases} \mathcal{H}_\theta\phi(x) & \text{if } \exists x \in E_q(K) \text{ with } \sigma(x) = \sigma(z) \\ 0 & \text{otherwise,} \end{cases}$$

where $z \in E_q^{\text{an}}$.

Proof. This is clear from the fact that $v_{\sigma(x)}$ is supported on $\sigma(E_q(K))$. □

Lemma 3.8. *Assume that $\phi \in L^2(U(E_q(K)))_0$. Then*

$$\mathcal{H}_{\theta,\sigma}\phi = \mathcal{H}_\theta\phi$$

holds true.

Proof. This follows immediately from the discussion before Lemma 3.7. □

Corollary 3.9. *The operator $\mathcal{H}_{\theta,\sigma}$ is a self-adjoint bounded linear operator on $L^2(E_q^{\text{an}}(K))$. Its spectrum coincides with that of \mathcal{H}_θ acting on $L^2(E_q^{\text{an}}, |\omega|)$, and there exists a probability measure $p_t(x, \cdot)$ with $t \geq 0$, $x \in E_q^{\text{an}}(K)$, on the Borel σ -algebra of $E_q^{\text{an}}(K)$ such that the Cauchy problem for the heat equation with $\mathcal{H}_{\theta,\sigma}$ has a unique solution of the form*

$$h(t, x) = \int_{E_q^{\text{an}}(K)} h_0(y) |\sigma_*\omega(y)|$$

where $h_0 \in C(E_q^{\text{an}}(K))$ is an initial condition for the heat equation. Furthermore, $p_t(x, \cdot)$ is the transition function of a Markov process whose paths are right continuous and have no discontinuities other than jumps.

Proof. This follows from the orthogonal decomposition (15), the isomorphism

$$L^2(U(E_q(K)))_0 \cong L^2(E_q(K))_0$$

observed before Lemma 3.7, the isomorphism

$$L^2(U(E_q(K)))_\sigma = L^2(E_q(K))_\sigma$$

between the radial parts, and Lemmas 3.7 and 3.8, and using Theorem 2.15 and Fact 3.4. Notice that the Feller semigroup property of the operator $\mathcal{H}_{\theta, \sigma}$ restricted to $C(U(E_q(K)))$ requires [4, Proposition 15]. \square

Apart from having a diffusion operator on the Berkovich analytification of a Tate curve, the added value of Corollary 3.9 is that the operator $\mathcal{H}_{\theta, \sigma}$ is controlled also by the Chambert-Loir measure $c_1(\mathcal{L}, \|\cdot\|_{\sigma(x)})$ equalling the Dirac measure on the point $\sigma(x)$ on the skeleton of the Tate curve. And also the push-forward of the positive Radon measure

$$\mu_x = H_\theta(x, \cdot) |\omega(\cdot)|$$

can be written as a Chambert-Loir measure

$$\mu_{\sigma(x)} = c_1(\mathcal{L}, \|\cdot\|_{\theta, x})$$

since we are dealing with the case of a curve.

Proposition 3.10. *The measure $\nu_{\sigma(x)}$ can be written as*

$$\nu_{\sigma(x)} = \sum_{z \in \sigma(E_q^{\text{an}}(K))} \alpha_z c_1(\mathcal{L}, \|\cdot\|_z) - \deg_\theta(\sigma(x)) c_1(\mathcal{L}, \|\cdot\|_{\sigma(x)})$$

with $\alpha_z > 0$, and $\|\cdot\|_y = e^{-g_y}$ with g_y a continuous subharmonic function on E_q^{an} such that

$$c_1(\mathcal{L}, \|\cdot\|_y) = \delta_y$$

for $y \in \sigma(E_q^{\text{an}}(K)) \sqcup \{\sigma(x)\}$ and $x \in E_q(K)$.

Proof. Clearly, $\nu_{\sigma(x)}$ is a linear combination of Dirac measures:

$$\nu_{\sigma(x)} = \sum_{z \in \sigma(E_q^{\text{an}}(K))} \alpha_z \delta_z - \deg_\theta(\sigma(x)) \delta_{\sigma(x)}$$

supported in $\sigma(E_q(K))$. Using the isomorphism

$$L^2(E(K), |\omega|) \cong L^2(U(E_q(K)))$$

obtained in the proof of Corollary 3.9, observe that the elements α_z are the corresponding entries in the helpful matrix L of (5). Using [23, Lemma 3.4.14], find that the Dirac measures are now seen to be of the form

$$\delta_y = \text{dd}^c(g_y)$$

with g_y subharmonic on $E_q^{\text{an}}(K)$ for $z \in \sigma(E_q^{\text{an}}(K)) \sqcup \{\sigma(x)\}$. Then $\|\cdot\|_y = e^{-g}$ are the desired corresponding metrics on \mathcal{L} proving the assertion. \square

The last result quoted in the Introduction is an immediate consequence of Proposition 3.10:

Corollary 3.11. *The heat operator $\mathcal{H}_{\theta,\sigma}$ on $E_q^{\text{an}}(K)$ is obtained as an integral operator of the form*

$$\mathcal{H}_{\theta,\sigma}\psi(x) = \int_{E_q^{\text{an}}} \psi c_1(\mathcal{L}, \|\cdot\|_{\theta,x})$$

with metric $\|\cdot\|_{\theta,x} = e^{-g_{\theta,x}}$ and

$$g_{\theta,x} = \sum_{z \in \sigma(E_q^{\text{an}}(K))} \alpha_z g_z$$

where α_z and g_z are as in Proposition 3.10, and $x \in \sigma(E_q(K))$.

4. On hearing the shape of a Tate curve

In [6, Corollary VI.1], it was shown how to reconstruct a reduction graph of a Mumford curve from the spectrum of a p -adic diffusion operator having an infinitely valued spectrum. The spectrum of the operator \mathcal{H}_{θ} has only finitely many values, and can also recover some information about the K -rational points of a Tate curve E_q themselves. This is shown in the following theorem:

Theorem 4.1. *Let E_q be a Tate elliptic curve defined over a local field K . Assume that all about K is known. Then the spectrum of the Laplacian operator \mathcal{H}_{θ} on $L^2(E_q(K), |\omega|)$, or of the Laplacian operator $\mathcal{H}_{\theta,\sigma}$ on $L^2(U(E_q(K)))$, respectively, can detect the following:*

1. *the presence of a 2-torsion point in $E_q(K)$.*
2. *the parity of $v(q)$.*
3. *the presence of $\pi^{\ell} \in E_q(K)$ which is a third root of π^{-k} , where k, ℓ are solutions for the congruence*

$$k + 3\ell \equiv 0 \pmod{v(q)}$$

in $\mathbb{Z}/v(q)\mathbb{Z}$.

Proof. From Lemmas 3.7 and 3.8, it follows that the spectra of both operators coincide. The assertions now follows from the explicit form of the negative degree eigenvalues given in Corollary 2.8. \square

Acknowledgements

Yassine El Maazouz, Stefan Kühnlein, Ángel Morán Ledezma and David Weisbart are warmly thanked for fruitful discussions. Frank Herrlich is thanked for guiding the author towards this beautiful area of mathematics when he was his student. The anonymous referee is thanked for helpful and clarifying suggestions towards substantially improving the paper. This work is partially supported by the Deutsche Forschungsgemeinschaft under project number 469999674.

References

- [1] V. G. Berkovich, *Spectral theory and analytic geometry over non-Archimedean fields*, Mathematical Surveys and Monographs **33**, American Mathematical Society, 1990. MR
- [2] V. G. Berkovich, “Smooth p -adic analytic spaces are locally contractible”, *Invent. Math.* **137**:1 (1999), 1–84. MR Zbl
- [3] S. Boucksom, C. Favre, and M. Jonsson, “Singular semipositive metrics in non-Archimedean geometry”, *J. Algebraic Geom.* **25**:1 (2016), 77–139. MR
- [4] P. E. Bradley, “Heat equations and wavelets on Mumford curves and their finite quotients”, *J. Fourier Anal. Appl.* **29**:5 (2023), art. id. 62, 26 pp. MR Zbl
- [5] P. E. Bradley, “Schottky-invariant p -adic diffusion operators”, preprint, 2024. To appear in *J. Fourier Anal. Appl.* Zbl arXiv 2405.17586
- [6] P. E. Bradley and A. Morán Ledezma, “Hearing shapes via p -adic Laplacians”, *J. Math. Phys.* **64**:11 (2023), art. id. 113502, 15 pp. MR Zbl
- [7] A. Chambert-Loir, “Mesures et équidistribution sur les espaces de Berkovich”, *J. Reine Angew. Math.* **595** (2006), 215–235. MR Zbl
- [8] S. N. Ethier and T. G. Kurtz, *Markov processes: characterization and convergence*, Wiley, New York, 1986. MR Zbl
- [9] Y. Fang, W. Gubler, , and K. Künnemann, “On the non-archimedean Monge–Ampère equation in mixed characteristic”, preprint, 2022. arXiv 2203.12282
- [10] J. Fresnel and M. van der Put, *Rigid analytic geometry and its applications*, Progress in Mathematics **218**, Birkhäuser, Boston, 2004. MR Zbl
- [11] J.-I. Igusa, *An introduction to the theory of local zeta functions*, AMS/IP Studies in Advanced Mathematics **14**, American Mathematical Society, Providence, RI, 2000. MR Zbl
- [12] A. Y. Khrennikov, S. V. Kozyrev, and W. A. Zúñiga-Galindo, *Ultrametric pseudodifferential equations and applications*, Encyclopedia of Mathematics and its Applications **168**, Cambridge University Press, 2018. MR Zbl
- [13] S. V. Kozyrev, “Wavelet theory as p -adic spectral analysis”, *Izv. Ross. Akad. Nauk Ser. Mat.* **66**:2 (2002), 149–158. In Russian; translated in *Izvestiya Math.* **66**:2 (2002), 367–376. MR Zbl
- [14] R. Rajkumar and D. Weisbart, “Components and exit times of Brownian motion in two or more p -adic dimensions”, *J. Fourier Anal. Appl.* **29**:6 (2023), art. id. 75, 28 pp. MR Zbl

- [15] J. J. Rodríguez-Vega and W. A. Zúñiga Galindo, “Taibleson operators, p -adic parabolic equations and ultrametric diffusion”, *Pacific J. Math.* **237**:2 (2008), 327–347. MR Zbl
- [16] P. Roquette, *Analytic theory of elliptic functions over local fields*, Hamburger Mathematische Einzelschriften (N.F.) **1**, Vandenhoeck & Ruprecht, Göttingen, 1970. MR Zbl
- [17] J. H. Silverman, *Advanced topics in the arithmetic of elliptic curves*, Graduate Texts in Mathematics **151**, Springer, 1994. MR Zbl
- [18] M. H. Taibleson, *Fourier analysis on local fields*, Princeton University Press, 1975. MR Zbl
- [19] K. Taira, *Boundary value problems and Markov processes*, 2nd ed., Lecture Notes in Mathematics **1499**, Springer, 2009. MR Zbl
- [20] J. Tate, “A review of non-Archimedean elliptic functions”, pp. 162–184 in *Elliptic curves, modular forms, and Fermat’s last theorem* (Hong Kong, 1993), Series in Number Theory **1**, International Press, Cambridge, MA, 1995. MR Zbl
- [21] M. Temkin, “Introduction to Berkovich analytic spaces”, pp. 3–66 in *Berkovich spaces and applications*, edited by A. Ducros et al., Lecture Notes in Math. **2119**, Springer, 2015. MR Zbl
- [22] G. Teschl, *Mathematical methods in quantum mechanics, with applications to Schrödinger operators*, 2nd ed., Graduate Studies in Mathematics **157**, American Mathematical Society, Providence, RI, 2014. MR Zbl
- [23] A. Thuillier, *Théorie du potentiel sur les courbes en géométrie analytique non archimédienne: applications à la théorie d’Arakelov*, PhD thesis, Université de Rennes 1, 2005. Zbl
- [24] V. S. Vladimirov, I. V. Volovich, and E. I. Zelenov, *p -adic analysis and mathematical physics*, Series on Soviet and East European Mathematics **1**, World Scientific, River Edge, NJ, 1994. MR Zbl
- [25] A. Weil, *Adeles and algebraic groups*, Progress in Mathematics **23**, Birkhäuser, Boston, 1982. MR Zbl
- [26] T. Yasuda, “The wild McKay correspondence and p -adic measures”, *J. Eur. Math. Soc. (JEMS)* **19**:12 (2017), 3709–3734. MR Zbl
- [27] W. A. Zúñiga-Galindo, “The non-Archimedean stochastic heat equation driven by Gaussian noise”, *J. Fourier Anal. Appl.* **21**:3 (2015), 600–627. MR Zbl
- [28] W. A. Zúñiga-Galindo, “Reaction-diffusion equations on complex networks and Turing patterns, via p -adic analysis”, *J. Math. Anal. Appl.* **491**:1 (2020), art. id. 124239, 39 pp. MR Zbl
- [29] W. A. Zúñiga-Galindo, “Eigen’s paradox and the quasispecies model in a non-Archimedean framework”, *Phys. A* **602** (2022), art. id. 127648, 18 pp. MR Zbl
- [30] W. A. Zúñiga-Galindo, “Ultrametric diffusion, rugged energy landscapes and transition networks”, *Phys. A* **597** (2022), art. id. 127221, 19 pp. MR Zbl

Received February 19, 2024. Revised October 15, 2024.

PATRICK ERIK BRADLEY
 INSTITUTE OF PHOTOGRAMMETRY AND REMOTE SENSING
 KARLSRUHE INSTITUTE OF TECHNOLOGY
 76131 KARLSRUHE
 GERMANY
 bradley@kit.edu