

*Pacific  
Journal of  
Mathematics*

**TRANPOSED POISSON SUPERALGEBRA STRUCTURES  
ON TWISTED  $N = 1$  BLOCK-LIE SUPERALGEBRA**

ANQI HUANG, YUN GAO AND JIANCAI SUN

# TRANPOSED POISSON SUPERALGEBRA STRUCTURES ON TWISTED $N = 1$ BLOCK-LIE SUPERALGEBRA

ANQI HUANG, YUN GAO AND JIANCAI SUN

**We investigate the transposed Poisson superalgebra structures on the twisted  $N = 1$  Block-Lie superalgebra  $\mathcal{S}(p, q)$ , where  $p$  and  $q$  are arbitrary complex numbers. We obtain that  $\mathcal{S}(p, q)$  admits only trivial transposed Poisson superalgebra structure for  $q \neq 0$  or  $p \notin \mathbb{Z}$ , while  $\mathcal{S}(p, q)$  has nontrivial transposed Poisson superalgebra structures for  $q = 0$  and  $p \in \mathbb{Z}$ , which are nonisomorphic with respect to  $p \in \mathbb{Z}$ .**

## 1. Introduction

Poisson algebras have their roots in Hamiltonian mechanics and have become a significant research topic in mathematics and physics. In 1809, D. Poisson introduced the operation of Poisson brackets on smooth functions while studying Lagrangian mechanics, thus giving smooth functions a Poisson structure. Thereafter, many researchers realized the importance of Poisson algebras and conducted extensive research from different perspectives. From string theory, quantum groups, and differential geometry, to integrable systems, algebraic geometry and representation theory, especially with the rise of noncommutative geometry, Poisson algebras have become an important branch of algebraic research (see [2; 4; 12]). More precisely, every Poisson algebra  $(\mathcal{L}, \cdot, [\cdot, \cdot])$  satisfies the *Leibniz rule*:

$$[x, y \cdot z] = [x, y] \cdot z + y \cdot [x, z] \quad \text{for } x, y, z \in \mathcal{L},$$

Transposed Poisson algebras (see [1]) are a generalization of Poisson algebras. In the definition of Poisson algebra, the Leibniz rule regards the Lie bracket operation as a derivation in an associative algebra. In the definition of transposed Poisson algebra, the transposed Leibniz rule treats the associative operation as a  $\frac{1}{2}$ -derivation in a Lie algebra. The transposed Leibniz rule, the compatibility condition of the transposed Poisson algebra  $(\mathcal{L}, \cdot, [\cdot, \cdot])$ , is articulated by

$$2z \cdot [x, y] = [z \cdot x, y] + [x, z \cdot y] \quad \text{for } x, y, z \in \mathcal{L}.$$

Y. Gao is the corresponding author.

MSC2020: 17B05, 17B40, 17B68.

Keywords: twisted  $N = 1$  Block-Lie superalgebra, transposed Poisson superalgebra,  $\frac{1}{2}$ -superderivation.

Therefore, the transposed Poisson algebra can be viewed as a dual structure of a Poisson algebra. In addition, the connection between the  $\frac{1}{2}$ -derivation of a Lie algebra and the transposed Poisson algebra is established in [6]. By applying this connection, all the transposed Poisson algebra structures can be obtained on the Witt algebra, Virasoro algebra, twisted Heisenberg–Virasoro algebra, Schrödinger–Virasoro algebra and extended Schrödinger–Virasoro algebra (see [6; 22]). The definition of the transposed Poisson superalgebra is also provided in [6]. A method for determining transposed Poisson algebra structures via the Kantor product is presented in [5]. The transposed Hom-Poisson algebras and the transposed BiHom-Poisson algebras are considered in [11; 13].

The study of antiderivation is generalized by Filippov (see [7]), in which the primary focus is on the  $\delta$ -derivations of prime Lie algebras with a nondegenerated symmetric invariant bilinear form. Specifically for a fixed element  $\delta$  in the ground field, a linear map  $\varphi$  on a Lie algebra  $(\mathcal{L}, [\cdot, \cdot])$  is called a  $\delta$ -derivation of  $\mathcal{L}$  if

$$\varphi([x, y]) = \delta([\varphi(x), y] + [x, \varphi(y)]) \quad \text{for } x, y \in \mathcal{L}.$$

The usual derivations can be viewed as 1-derivations. The centroid of a Lie algebra  $\mathcal{L}$ ,  $\text{Cent}(\mathcal{L})$ , is the space of linear maps  $\chi$  on  $\mathcal{L}$  satisfying  $\chi([x, y]) = [\chi(x), y] = [x, \chi(y)]$  for all  $x, y \in \mathcal{L}$ . It is easy to see that elements of the centroid are  $\frac{1}{2}$ -derivations. If  $\delta = 0$  and  $[\mathcal{L}, \mathcal{L}] \neq 0$ , then every nonzero linear map is called a 0-derivation. If  $[\mathcal{L}, \mathcal{L}] = \mathcal{L}$ , in particular,  $\mathcal{L}$  is a simple Lie algebra, then  $\mathcal{L}$  has no nonzero 0-derivations. If  $\delta = -1$ , then the linear map is called an *antiderivation*. The concept of the  $\delta$ -superderivation on nonassociative superalgebras is introduced by Kaygorodov, and it is proved that simple finite-dimensional Lie superalgebras over an algebraically closed field of characteristic 0 do not have nontrivial  $\delta$ -superderivations (see [8], [9]).

Block–Lie algebras are a class of infinite-dimensional simple Lie algebras introduced by Block in 1958 (see [3]). Since then, several generalizations of these algebras have been proposed (see [14; 15; 21]). The Block algebra  $B(q)$  for a fixed complex number  $q$  is defined in [18]. The notion of the Block superalgebra  $\mathfrak{K}(p)$  is introduced, and its finite-dimensional irreducible conformal modules are classified for any nonzero parameter  $p$  (see [17]). Based on the twisted rules from Ramond superalgebras to Neveu–Schwarz superalgebras, a twisted version of the  $\mathbb{Z}$ -graded conformal superalgebra  $\mathfrak{T}(p)$  is introduced in [16], where  $p$  is a nonzero parameter. Precisely speaking, the subscripts of the odd generators of the original  $\mathbb{Z}$ -graded algebra are shifted by  $\frac{1}{2}$ . Motivated by [16; 19], the parameters of twisted  $N = 1$  Block superalgebra are extended to include two nonzero parameters,  $p$  and  $q$ . Note that the special case of  $p \in \mathbb{C}$  and  $q = 1$  is considered in [16]. The nonweight modules over  $N = 1$  Lie superalgebras of Block type is studied in [19]. In particular, transposed Poisson algebra structures on Block–Lie algebras  $\mathcal{B}(q)$  and

Block-Lie superalgebras  $\mathcal{S}(q)$  are described in [10]. This work propels us to delve into transposed Poisson superalgebra structures of the twisted  $N = 1$  Block-Lie superalgebra  $\mathcal{S}(p, q)$  with two parameters.

This paper is organized as follows. In Section 2, we recall some basic definitions and the relation between transposed Poisson superalgebra structures and  $\frac{1}{2}$ -superderivations. In Section 3, we characterize all  $\frac{1}{2}$ -superderivations of the twisted  $N = 1$  Block-Lie superalgebra  $\mathcal{S}(p, q)$ . In Section 4, we present our main theorem about the transposed Poisson superalgebra structures on  $\mathcal{S}(p, q)$ ; see Theorem 4.1. More precisely, we prove in Theorem 4.1 that there is only nontrivial transposed Poisson superalgebra structure for  $q = 0$  and  $p \in \mathbb{Z}$ , otherwise, such structures are trivial. Throughout this paper, we denote by  $\mathbb{C}$ ,  $\mathbb{C}^*$ ,  $\mathbb{Q}$ ,  $\mathbb{Z}$  and  $\mathbb{Z}^*$  the complex numbers, nonzero complex numbers, rational numbers, integers and nonzero integers, respectively.

## 2. Preliminaries

In this section, we recall some definitions and notation for future convenience.

Motivated by the definitions of the Block-type algebra and the  $N = 1$  Lie superalgebra of Block type from [19; 20], we arrive at the subsequent definition.

**Definition 2.1.** Let  $p$  and  $q$  be fixed complex numbers. The twisted  $N = 1$  Block-Lie superalgebra  $\mathcal{S}(p, q)$  is defined as an infinite-dimensional Lie superalgebra over  $\mathbb{C}$  with the even part  $\mathcal{S}(p, q)_{\bar{0}} = \{L_{m,i} \mid m, i \in \mathbb{Z}\}$  and the odd part  $\mathcal{S}(p, q)_{\bar{1}} = \{G_{m+\frac{1}{2}, i+\frac{1}{2}} \mid m, i \in \mathbb{Z}\}$  together with the relations

$$(2-1) \quad [L_{m,i}, L_{n,j}] = ((n+p)(i+q) - (j+q)(m+p))L_{m+n, i+j} \\ = \begin{vmatrix} n+p & j+q \\ m+p & i+q \end{vmatrix} L_{m+n, i+j},$$

$$[L_{m,i}, G_{n+\frac{1}{2}, j+\frac{1}{2}}] = ((n+\frac{p}{2}+\frac{1}{2})(i+q) - (j+\frac{q}{2}+\frac{1}{2})(m+p))G_{m+n+\frac{1}{2}, i+j+\frac{1}{2}} \\ = \begin{vmatrix} n+\frac{p}{2}+\frac{1}{2} & j+\frac{q}{2}+\frac{1}{2} \\ m+p & i+q \end{vmatrix} G_{m+n+\frac{1}{2}, i+j+\frac{1}{2}},$$

$$[G_{m+\frac{1}{2}, i+\frac{1}{2}}, G_{n+\frac{1}{2}, j+\frac{1}{2}}] = 2qL_{m+n+1, i+j+1}$$

for  $m, n, i, j \in \mathbb{Z}$ .

Let  $\mathcal{L} = \mathcal{L}_{\bar{0}} \oplus \mathcal{L}_{\bar{1}}$  be a  $\mathbb{Z}_2$ -graded vector space. For  $i \in \mathbb{Z}_2 = \{\bar{0}, \bar{1}\}$ , if  $x \in \mathcal{L}_i$ , then  $|x|$  denotes the parity of  $x$ , that is,  $|x| = 0$  if  $x \in \mathcal{L}_{\bar{0}}$  or  $|x| = 1$  if  $x \in \mathcal{L}_{\bar{1}}$ .

We will briefly recall some definitions from [10].

**Definition 2.2.** Let  $\mathcal{L} = \mathcal{L}_{\bar{0}} \oplus \mathcal{L}_{\bar{1}}$  be a  $\mathbb{Z}_2$ -graded vector space equipped with two nonzero bilinear superoperations  $\cdot$  and  $[\cdot, \cdot]$ . The triple  $(\mathcal{L}, \cdot, [\cdot, \cdot])$  is called a

*transposed Poisson superalgebra* if  $(\mathcal{L}, \cdot)$  is a supercommutative associative superalgebra and  $(\mathcal{L}, [\cdot, \cdot])$  is a Lie superalgebra that satisfies the compatibility condition

$$(2-2) \quad 2z \cdot [x, y] = [z \cdot x, y] + (-1)^{|x||z|}[x, z \cdot y] \quad \text{for } x, z \in \mathcal{L}_i, y \in \mathcal{L}, i \in \mathbb{Z}_2.$$

**Definition 2.3.** Let  $(\mathcal{L}, [\cdot, \cdot])$  be a Lie superalgebra. A transposed Poisson superalgebra structure on  $(\mathcal{L}, [\cdot, \cdot])$  is a supercommutative associative multiplication  $\cdot$  on  $\mathcal{L}$  which makes  $(\mathcal{L}, \cdot, [\cdot, \cdot])$  a transposed Poisson superalgebra.

If  $\varphi$  is a homogeneous linear map  $\mathcal{L} \rightarrow \mathcal{L}$ , then  $|\varphi| = 0$  for  $\varphi(\mathcal{L}_i) \subseteq \mathcal{L}_i$  or  $|\varphi| = 1$  for  $\varphi(\mathcal{L}_i) \subseteq \mathcal{L}_{\bar{1}-i}$ ,  $i \in \mathbb{Z}_2$ .

**Definition 2.4.** Let  $(\mathcal{L}, [\cdot, \cdot])$  be a superalgebra and  $\varphi$  a homogeneous linear map  $\mathcal{L} \rightarrow \mathcal{L}$ . Then  $\varphi$  is called  $\frac{1}{2}$ -superderivation if it satisfies

$$(2-3) \quad \varphi([x, y]) = \frac{1}{2}([\varphi(x), y] + (-1)^{|\varphi||x|}[x, \varphi(y)]) \quad \text{for } x, y \in \mathcal{L}_i, i \in \mathbb{Z}_2.$$

We will use the notation  $\Delta(\mathcal{L})$  for the space of  $\frac{1}{2}$ -superderivations of Lie superalgebra  $\mathcal{L}$ .

Let  $L_z$  denote the operator of the left multiplication by an element  $z \in \mathcal{L}$ , that is,

$$L_z(x) = z \cdot x \quad \text{for } x \in \mathcal{L}.$$

Definitions 2.2 and 2.4 immediately imply the following key lemma.

**Lemma 2.5.** Let  $(\mathcal{L}, \cdot, [\cdot, \cdot])$  be a transposed Poisson superalgebra and  $z \in \mathcal{L}_i$ ,  $i \in \mathbb{Z}_2$ . Then the left multiplication  $L_z$  of  $(\mathcal{L}, \cdot)$  is a  $\frac{1}{2}$ -superderivation of  $(\mathcal{L}, [\cdot, \cdot])$  and  $|L_z| = |z|$ .

The basic example of a  $\frac{1}{2}$ -superderivation is the multiplication by a field element. Such  $\frac{1}{2}$ -superderivations will be called *trivial*.

**Theorem 2.6.** Let  $\mathcal{L}$  be a Lie superalgebra without nontrivial  $\frac{1}{2}$ -superderivations. Then all transposed Poisson superalgebra structures on  $\mathcal{L}$  are trivial.

Let  $\cdot$  be a transposed Poisson superalgebra structure on a Lie superalgebra  $(\mathcal{L}, [\cdot, \cdot])$ . Then any automorphism  $\phi$  of  $(\mathcal{L}, [\cdot, \cdot])$  induces another transposed Poisson superalgebra structure  $*$  on  $(\mathcal{L}, [\cdot, \cdot])$  given by

$$x * y = \phi(\phi^{-1}(x) \cdot \phi^{-1}(y)) \quad \text{for } x, y \in \mathcal{L}.$$

Clearly,  $\phi$  is an isomorphism of transposed Poisson superalgebras  $(\mathcal{L}, \cdot, [\cdot, \cdot])$  and  $(\mathcal{L}, *, [\cdot, \cdot])$ .

### 3. $\frac{1}{2}$ -superderivations of the twisted $N = 1$ Block–Lie superalgebra

In this section, we will investigate and describe all the  $\frac{1}{2}$ -superderivations of the twisted  $N = 1$  Block–Lie superalgebra  $\mathcal{S}(p, q)$ .

**3.1. Even  $\frac{1}{2}$ -superderivations of  $\mathcal{S}(p, q)$ .** In this subsection we consider only *even* linear maps  $\varphi : \mathcal{S}(p, q) \rightarrow \mathcal{S}(p, q)$ , i.e., those which satisfy  $\varphi(\mathcal{S}(p, q)_i) \subseteq \mathcal{S}(p, q)_i$  for  $i \in \mathbb{Z}_2$ . We thus have  $|\varphi| = 0$ , so  $\varphi$  is a  $\frac{1}{2}$ -superderivation of  $\mathcal{S}(p, q)$  if and only if  $\varphi$  is a usual  $\frac{1}{2}$ -derivation of  $\mathcal{S}(p, q)$ . We now write

$$\varphi = \sum_{r,s \in \mathbb{Z}} \varphi_{r,s},$$

where

$$(3-1) \quad \varphi_{r,s}(L_{m,i}) = d_{r,s}^{\bar{0}}(m, i)L_{m+r, i+s},$$

$$(3-2) \quad \varphi_{r,s}(G_{m+\frac{1}{2}, i+\frac{1}{2}}) = d_{r,s}^{\bar{1}}(m + \frac{1}{2}, i + \frac{1}{2})G_{m+r+\frac{1}{2}, i+s+\frac{1}{2}}$$

for some  $d_{r,s}^{\bar{0}}(m, i), d_{r,s}^{\bar{1}}(m + \frac{1}{2}, i + \frac{1}{2}) \in \mathbb{C}$ ,  $m, i, r, s \in \mathbb{Z}$ . Then we have  $\varphi \in \Delta^{\bar{0}}(\mathcal{S}(p, q))$  if and only if  $\varphi_{r,s} \in \Delta^{\bar{0}}(\mathcal{S}(p, q))$  for all  $r, s \in \mathbb{Z}$ .

**Lemma 3.1.** *Let  $\varphi_{r,s} : \mathcal{S}(p, q) \rightarrow \mathcal{S}(p, q)$ ,  $r, s \in \mathbb{Z}$ , be a linear map satisfying (3-1) and (3-2). Then  $\varphi_{r,s} \in \Delta^{\bar{0}}(\mathcal{S}(p, q))$  if and only if the these three conditions hold:*

$$(3-3) \quad 2 \begin{vmatrix} n+p & j+q \\ m+p & i+q \end{vmatrix} d_{r,s}^{\bar{0}}(m+n, i+j) \\ = \begin{vmatrix} n+p & j+q \\ m+r+p & i+s+q \end{vmatrix} d_{r,s}^{\bar{0}}(m, i) + \begin{vmatrix} n+r+p & j+s+q \\ m+p & i+q \end{vmatrix} d_{r,s}^{\bar{0}}(n, j),$$

$$(3-4) \quad 2 \begin{vmatrix} n+\frac{p}{2}+\frac{1}{2} & j+\frac{q}{2}+\frac{1}{2} \\ m+p & i+q \end{vmatrix} d_{r,s}^{\bar{1}}(m+n+\frac{1}{2}, i+j+\frac{1}{2}) \\ = \begin{vmatrix} n+\frac{p}{2}+\frac{1}{2} & j+\frac{q}{2}+\frac{1}{2} \\ m+r+p & i+s+q \end{vmatrix} d_{r,s}^{\bar{0}}(m, i) \\ + \begin{vmatrix} n+r+\frac{p}{2}+\frac{1}{2} & j+s+\frac{q}{2}+\frac{1}{2} \\ m+p & i+q \end{vmatrix} d_{r,s}^{\bar{1}}(n+\frac{1}{2}, j+\frac{1}{2}),$$

$$(3-5) \quad 2qd_{r,s}^{\bar{0}}(m+n+1, i+j+1) = q(d_{r,s}^{\bar{1}}(m+\frac{1}{2}, i+\frac{1}{2}) + d_{r,s}^{\bar{1}}(n+\frac{1}{2}, j+\frac{1}{2})).$$

*Proof.* By [Definition 2.1](#),  $\varphi_{r,s}$  is an even  $\frac{1}{2}$ -superderivation of  $\mathcal{S}(p, q)$  if and only if (2-3) hold on the basis  $\{L_{m,i}, G_{m+\frac{1}{2}, i+\frac{1}{2}} \mid m, i \in \mathbb{Z}\}$ . Namely, the subsequent three equations are satisfied:

$$(3-6) \quad 2 \begin{vmatrix} n+p & j+q \\ m+p & i+q \end{vmatrix} \varphi_{r,s}(L_{m+n, i+j}) = [\varphi_{r,s}(L_{m,i}), L_{n,j}] + [L_{m,i}, \varphi_{r,s}(L_{n,j})],$$

$$(3-7) \quad 2 \begin{vmatrix} n+\frac{p}{2}+\frac{1}{2} & j+\frac{q}{2}+\frac{1}{2} \\ m+p & i+q \end{vmatrix} \varphi_{r,s}(G_{m+n+\frac{1}{2}, i+j+\frac{1}{2}}) = [\varphi_{r,s}(L_{m,i}), G_{n+\frac{1}{2}, j+\frac{1}{2}}] \\ + [L_{m,i}, \varphi_{r,s}(G_{n+\frac{1}{2}, j+\frac{1}{2}})],$$

$$(3-8) \quad 4q\varphi_{r,s}(L_{m+n+1, i+j+1}) = [\varphi_{r,s}(G_{m+\frac{1}{2}, i+\frac{1}{2}}), G_{n+\frac{1}{2}, j+\frac{1}{2}}] \\ + [G_{m+\frac{1}{2}, i+\frac{1}{2}}, \varphi_{r,s}(G_{n+\frac{1}{2}, j+\frac{1}{2}})].$$

In view of (3-6), we can see that

$$\begin{aligned}
2 \begin{vmatrix} n+p & j+q \\ m+p & i+q \end{vmatrix} d_{r,s}^{\bar{0}}(m+n, i+j) L_{m+n+r, i+j+s} \\
&= 2 \begin{vmatrix} n+p & j+q \\ m+p & i+q \end{vmatrix} \varphi_{r,s}(L_{m+n, i+j}) \\
&= [\varphi_{r,s}(L_{m,i}), L_{n,j}] + [L_{m,i}, \varphi_{r,s}(L_{n,j})] \\
&= [d_{r,s}^{\bar{0}}(m, i) L_{m+r, i+s}, L_{n,j}] + [L_{m,i}, d_{r,s}^{\bar{0}}(n, j) L_{n+r, j+s}] \\
&= \begin{vmatrix} n+p & j+q \\ m+r+p & i+s+q \end{vmatrix} d_{r,s}^{\bar{0}}(m, i) L_{m+n+r, i+j+s} \\
&\quad + \begin{vmatrix} n+r+p & j+s+q \\ m+p & i+q \end{vmatrix} d_{r,s}^{\bar{0}}(n, j) L_{m+n+r, i+j+s}.
\end{aligned}$$

Thus, we come to (3-3). By (3-7), we observe

$$\begin{aligned}
2 \begin{vmatrix} n+\frac{p}{2}+\frac{1}{2} & j+\frac{q}{2}+\frac{1}{2} \\ m+p & i+q \end{vmatrix} d_{r,s}^{\bar{1}}(m+n+\frac{1}{2}, i+j+\frac{1}{2}) G_{m+n+r+\frac{1}{2}, i+j+s+\frac{1}{2}} \\
&= 2 \begin{vmatrix} n+\frac{p}{2}+\frac{1}{2} & j+\frac{q}{2}+\frac{1}{2} \\ m+p & i+q \end{vmatrix} \varphi_{r,s}(G_{m+n+\frac{1}{2}, i+j+\frac{1}{2}}) \\
&= [\varphi_{r,s}(L_{m,i}), G_{n+\frac{1}{2}, j+\frac{1}{2}}] + [L_{m,i}, \varphi_{r,s}(G_{n+\frac{1}{2}, j+\frac{1}{2}})] \\
&= [d_{r,s}^{\bar{0}}(m, i) L_{m+r, i+s}, G_{n+\frac{1}{2}, j+\frac{1}{2}}] + [L_{m,i}, d_{r,s}^{\bar{1}}(n+\frac{1}{2}, j+\frac{1}{2}) G_{n+r+\frac{1}{2}, j+s+\frac{1}{2}}] \\
&= \begin{vmatrix} n+\frac{p}{2}+\frac{1}{2} & j+\frac{q}{2}+\frac{1}{2} \\ m+r+p & i+s+q \end{vmatrix} d_{r,s}^{\bar{0}}(m, i) G_{m+n+r+\frac{1}{2}, i+j+s+\frac{1}{2}} \\
&\quad + \begin{vmatrix} n+r+\frac{p}{2}+\frac{1}{2} & j+s+\frac{q}{2}+\frac{1}{2} \\ m+p & i+q \end{vmatrix} d_{r,s}^{\bar{1}}(n+\frac{1}{2}, j+\frac{1}{2}) G_{m+n+r+\frac{1}{2}, i+j+s+\frac{1}{2}}.
\end{aligned}$$

Accordingly, we arrive at (3-4). With the use of (3-8), we have

$$\begin{aligned}
4q d_{r,s}^{\bar{0}}(m+n+1, i+j+1) L_{m+n+r+1, i+j+s+1} \\
&= 4q \varphi_{r,s}(L_{m+n+1, i+j+1}) \\
&= [\varphi_{r,s}(G_{m+\frac{1}{2}, i+\frac{1}{2}}), G_{n+\frac{1}{2}, j+\frac{1}{2}}] + [G_{m+\frac{1}{2}, i+\frac{1}{2}}, \varphi_{r,s}(G_{n+\frac{1}{2}, j+\frac{1}{2}})] \\
&= [d_{r,s}^{\bar{1}}(m+\frac{1}{2}, i+\frac{1}{2})(G_{m+r+\frac{1}{2}, i+s+\frac{1}{2}}), G_{n+\frac{1}{2}, j+\frac{1}{2}}] \\
&\quad + [G_{m+\frac{1}{2}, i+\frac{1}{2}}, d_{r,s}^{\bar{1}}(n+\frac{1}{2}, j+\frac{1}{2})(G_{n+r+\frac{1}{2}, j+s+\frac{1}{2}})] \\
&= 2q (d_{r,s}^{\bar{1}}(m+\frac{1}{2}, i+\frac{1}{2}) + d_{r,s}^{\bar{1}}(n+\frac{1}{2}, j+\frac{1}{2})) L_{m+n+r+1, i+j+s+1}.
\end{aligned}$$

So, we get (3-5).  $\square$

Inspired by [10], we proceed to analyze the  $\frac{1}{2}$ -superderivatives of  $\mathcal{S}(p, q)$ , considering whether  $p$  and  $q$  are zero or not.

**3.1.1. The case  $p, q \neq 0$ .**

**Lemma 3.2.** *Let  $\varphi = \sum_{r,s \in \mathbb{Z}} \varphi_{r,s}$  be an even  $\frac{1}{2}$ -superderivation of  $\mathcal{S}(p, q)$  and  $(r, s) \in \mathbb{Z} \times \mathbb{Z} \setminus \{(0, 0)\}$ . Then*

$$\varphi_{r,s} = 0.$$

*Proof.* Taking  $n = j = 0$  in (3-3), we obtain

$$(3-9) \quad (p(i-s) - q(m-r))d_{r,s}^{\bar{0}}(m, i) = ((r+p)(i+q) - (s+q)(m+p))d_{r,s}^{\bar{0}}(0, 0).$$

(1) For  $\frac{p}{q} \in \mathbb{Q}$ , there exists some  $k \in \mathbb{C}^*$  satisfying  $kp, kq \in \mathbb{Z}$ . Putting  $m = kp + r$  and  $i = kq + s$  in (3-9), it follows that

$$(qr - ps)d_{r,s}^{\bar{0}}(0, 0) = 0.$$

Clearly, we will discuss  $qr - ps$  being zero or nonzero.

Case 1:  $qr - ps \neq 0$ . We have  $d_{r,s}^{\bar{0}}(0, 0) = 0$ . And hence by (3-9), we can obtain

$$(3-10) \quad d_{r,s}^{\bar{0}}(m, i) = 0 \quad \text{if } i \neq \frac{q(m-r)}{p} + s.$$

If  $\frac{q(m-r)}{p} + s \in \mathbb{Z}$ , we set  $i = \frac{q(m-r)}{p} + s$ ,  $n = -m$ ,  $j = -i$ . Since

$$j = -i = -\frac{q(m-r)}{p} - s = \frac{q(n+r)}{p} - s \neq \frac{q(n-r)}{p} + s,$$

we can apply (3-10). It follows that

$$d_{r,s}^{\bar{0}}(n, j) = d_{r,s}^{\bar{0}}(-m, -i) = 0.$$

Substituting it into (3-3), we get

$$(3p + r - m)d_{r,s}^{\bar{0}}\left(m, \frac{q(m-r)}{p} + s\right) = 0.$$

Therefore, it leads to

$$d_{r,s}^{\bar{0}}\left(m, \frac{q(m-r)}{p} + s\right) = 0 \quad \text{if } m \neq 3p + r.$$

Taking  $m = 3p + r$ ,  $i = q(m-r)/p + s = 3q + s$  and  $n \notin \{0, 3p + r\}$  in (3-3), for  $d_{r,s}^{\bar{0}}(m+n, i+j) = 0$  and  $d_{r,s}^{\bar{0}}(n, j) = 0$ , we obtain

$$(3-11) \quad ((n+p)(2q+s) - (j+q)(2p+r))d_{r,s}^{\bar{0}}(3p+r, 3q+s) = 0.$$

If  $r + 2p \neq 0$ , then we take  $j \neq \frac{(n+p)(2q+s)}{r+2p} - q$ . So, we obtain

$$d_{r,s}^{\bar{0}}(3p+r, 3q+s) = 0.$$

If  $r + 2p = 0$ , it follows from (3-11) that

$$(n+p)(2q+s)d_{-2p,s}^{\bar{0}}(3p+r, 3q+s) = 0.$$

Due to  $qr - ps \neq 0$  and  $r = -2p$ , then we have  $2q + s \neq 0$ . It only needs to take  $n \neq -p$ , then we get

$$d_{-2p,s}^{\bar{0}}(p, 3q+s) = 0.$$

Case 2:  $qr - ps = 0$ . Since  $\frac{p}{q} = \frac{r}{s} \in \mathbb{Q}$ , there exists  $t \in \mathbb{C}^*$  satisfying  $tp, tq \in \mathbb{Z}$ . Taking  $r = tp, s = tq$  in (3-3), we have

$$(3-12) \quad 2 \begin{vmatrix} n+p & j+q \\ m+p & i+q \end{vmatrix} d_{tp,tq}^{\bar{0}}(m+n, i+j) = \begin{vmatrix} n+p & j+q \\ m+(t+1)p & i+(t+1)q \end{vmatrix} d_{tp,tq}^{\bar{0}}(m, i) \\ + \begin{vmatrix} n+(t+1)p & j+(t+1)q \\ m+p & i+q \end{vmatrix} d_{tp,tq}^{\bar{0}}(n, j).$$

Considering  $n = j = 0, m \neq 0$  in (3-12), we get

$$(pi - qm)d_{tp,tq}^{\bar{0}}(m, i) = (t+1)(pi - qm)d_{tp,tq}^{\bar{0}}(0, 0).$$

Clearly, our discussion will proceed based on whether  $pi - qm$  is zero or not.

Subcase 1:  $pi - qm \neq 0$ . We have

$$(3-13) \quad d_{tp,tq}^{\bar{0}}(m, i) = (t+1)d_{tp,tq}^{\bar{0}}(0, 0).$$

Taking  $m = -n \neq 0, j = -i$  and  $pi + qn \neq 0$  in (3-12), we obtain

$$(3-14) \quad 4d_{tp,tq}^{\bar{0}}(0, 0) = (t+2)d_{tp,tq}^{\bar{0}}(-n, i) + (t+2)d_{tp,tq}^{\bar{0}}(n, -i).$$

Then applying (3-13), we have  $d_{tp,tq}^{\bar{0}}(n, -i) = d_{tp,tq}^{\bar{0}}(-n, i) = (t+1)d_{tp,tq}^{\bar{0}}(0, 0)$ . As a result, (3-14) can be simplified as

$$(t+3)d_{tp,tq}^{\bar{0}}(0, 0) = 0.$$

If  $t \neq -3$ , then we arrive at  $d_{tp,tq}^{\bar{0}}(0, 0) = 0$ . And hence by (3-13), it follows that

$$d_{tp,tq}^{\bar{0}}(m, i) = 0.$$

If  $t = -3$ , by (3-13), we derive

$$(3-15) \quad d_{-3p,-3q}^{\bar{0}}(m, i) = -2d_{-3p,-3q}^{\bar{0}}(0, 0).$$

Letting  $i = j = 0$  and  $m, n, m \pm n \neq 0$  in (3-12), using (3-15), we are able to conclude that

$$2(n-m)d_{-3p,-3q}^{\bar{0}}(m+n, 0) = -(m+2n)d_{-3p,-3q}^{\bar{0}}(m, 0) + (2m+n)d_{-3p,-3q}^{\bar{0}}(-n, 0),$$

$$d_{-3p,-3q}^{\bar{0}}(0, 0) = 0.$$

Substituting it into (3-15), we see that

$$d_{-3p,-3q}^{\bar{0}}(m, i) = 0.$$

Hence, we can deduce that

$$(3-16) \quad d_{ip,tq}^{\bar{0}}(m, i) = 0 \quad \text{for } pi - qm \neq 0.$$

Subcase 2:  $pi - qm = 0$ . Since  $\frac{p}{q} = \frac{m}{i} \in \mathbb{Q}$ , there exists  $l \in \mathbb{C}^*$  satisfying  $lp, lq \in \mathbb{Z}$ . By substituting  $m = j = 0, n = lp$  and  $i = lq$  into (3-12), since  $pi - qm \neq 0, pj - qn \neq 0$  and (3-16), we arrive at  $d_{ip,tq}^{\bar{0}}(lp, 0) = d_{ip,tq}^{\bar{0}}(0, lq) = 0$ . So (3-12) can be written as

$$(l+2)d_{ip,tq}^{\bar{0}}(lp, lq) = 0.$$

Then, it yields

$$d_{ip,tq}^{\bar{0}}(lp, lq) = 0, \quad \text{if } l \neq -2.$$

If  $l = -2$ , then taking  $m = -2p, n = 2p, i = -2q$  and  $j = 0$  in (3-12) and considering (3-16), it follows that

$$(t-1)pqd_{ip,tq}^{\bar{0}}(-2p, -2q) = 0.$$

Since  $p, q \neq 0$ , then  $d_{ip,tq}^{\bar{0}}(-2p, -2q) = 0$  for  $t \neq 1$  is derived.

Therefore, we can conclude that  $d_{r,s}^{\bar{0}}(m, i) = 0$  for all  $(m, i) \in \mathbb{Z} \times \mathbb{Z}, (r, s) \notin \{(0, 0), (p, q)\}$  and  $d_{p,q}^{\bar{0}}(m, i) = 0$  unless  $(m, i) = (-2p, -2q)$ .

(2) For  $\frac{p}{q} \notin \mathbb{Q}$ , we can obtain, from (3-9),

$$(3-17) \quad d_{r,s}^{\bar{0}}(m, i) = \frac{(r+p)(i+q) - (s+q)(m+p)}{p(i-s) - q(m-r)} d_{r,s}^{\bar{0}}(0, 0) \quad \text{for } (m, i) \neq (r, s).$$

In the following analysis, we will proceed by the conditions whether  $r$  and  $s$  are zero or not.

Case 1:  $r, s \neq 0$ . Taking  $m = -r$  and  $j = -s$  into (3-9), we can see that

$$d_{r,s}^{\bar{0}}(-r, -s) = d_{r,s}^{\bar{0}}(0, 0).$$

Given the conditions  $m = 0, n = -r, i = -s$  and  $j = 0$  in (3-3), it follows that

$$2((p-r)(q-s) - pq)d_{r,s}^{\bar{0}}(-r, -s) = -2qrd_{r,s}^{\bar{0}}(0, -s) - 2psd_{r,s}^{\bar{0}}(-r, 0).$$

By combining

$$\begin{aligned}(qr - 2ps)d_{r,s}^{\bar{0}}(0, -s) &= (qr - 2ps - rs)d_{r,s}^{\bar{0}}(0, 0), \\ (ps - 2qr)d_{r,s}^{\bar{0}}(-r, 0) &= (ps - 2qr - rs)d_{r,s}^{\bar{0}}(0, 0),\end{aligned}$$

we can deduce that

$$d_{r,s}^{\bar{0}}(0, 0) = 0.$$

So it leads to

$$d_{r,s}^{\bar{0}}(m, i) = 0 \quad \text{for } (m, i) \neq (r, s).$$

Taking  $m = r$ ,  $i = s$ ,  $j \notin \{0, s\}$  and  $n \notin \{0, r, 2r\}$  into (3-3), we can get

$$((n + p)(2s + q) - (j + q)(2r + p))d_{r,s}^{\bar{0}}(r, s) = 0.$$

Since  $\frac{p}{q} \neq \mathbb{Q}$ , we have  $d_{r,s}^{\bar{0}}(r, s) = 0$ .

Case 2:  $r = 0$ ,  $s \neq 0$ . In this case, (3-17) becomes

$$(3-18) \quad d_{0,s}^{\bar{0}}(m, i) = \frac{p(i + q) - (s + q)(m + p)}{p(i - s) - qr} d_{0,s}^{\bar{0}}(0, 0) \quad \text{for } (m, i) \neq (0, s).$$

Substituting  $i = s$  and  $m \neq 0$  in (3-18), we have

$$d_{0,s}^{\bar{0}}(m, s) = \frac{s + q}{q} d_{0,s}^{\bar{0}}(0, 0) \quad \text{for } m \neq 0.$$

Putting  $m = 0$  and  $i = 2s$  into (3-18), we obtain

$$d_{0,s}^{\bar{0}}(0, 2s) = d_{0,s}^{\bar{0}}(0, 0).$$

Furthermore, considering  $m + n = 0$ ,  $m, n \neq 0$  and  $i = j = 0$  in (3-3), we can deduce that

$$\begin{aligned}4n(s + q)d_{0,s}^{\bar{0}}(0, 0) &= ((n + p)(2s + q) - (-n + p)(s + q))d_{0,s}^{\bar{0}}(m, s) \\ &\quad + ((n + p)(s + q) - (-n + p)(2s + q))d_{0,s}^{\bar{0}}(n, s).\end{aligned}$$

Therefore, it can be checked that

$$d_{0,s}^{\bar{0}}(m, i) = 0 \quad \text{for } (m, i) \neq (0, s).$$

Taking  $m = 0$ ,  $i = s$ ,  $j \notin \{0, s\}$  and  $n \neq 0$  into (3-3), we can get

$$((n + p)(2s + q) - p(j + q))d_{0,s}^{\bar{0}}(0, s) = 0.$$

Since  $\frac{p}{q} \neq \mathbb{Q}$ , we have  $d_{0,s}^{\bar{0}}(0, s) = 0$ .

Case 3:  $r \neq 0, s = 0$ . Similarly, it can be shown that  $d_{r,0}^{\bar{0}}(m, i) = 0$  for  $(m, i) \in \mathbb{Z} \times \mathbb{Z}$ .

Subsequently, we consider the value of  $d_{r,s}^{\bar{1}}(m + \frac{1}{2}, i + \frac{1}{2})$ . If  $(r, s) \notin \{(0, 0), (p, q)\}$ , (3-5) becomes

$$d_{r,s}^{\bar{1}}(m + \frac{1}{2}, i + \frac{1}{2}) + d_{r,s}^{\bar{1}}(n + \frac{1}{2}, j + \frac{1}{2}) = 0.$$

Taking  $m = n$  and  $i = j$ , we get

$$d_{r,s}^{\bar{1}}(m + \frac{1}{2}, i + \frac{1}{2}) = 0 \quad \text{for all } (m, i) \in \mathbb{Z} \times \mathbb{Z} \text{ and } (r, s) \notin \{(0, 0), (p, q)\}.$$

If  $(r, s) = (p, q)$ , it implies  $p, q \in \mathbb{Z}$ . Then taking  $m = n$  and  $i = j$  in (3-5), we get

$$d_{p,q}^{\bar{1}}(m + \frac{1}{2}, i + \frac{1}{2}) = d_{r,s}^{\bar{0}}(2m + 1, 2i + 1) = 0 \quad \text{for all } (m, i) \in \mathbb{Z} \times \mathbb{Z}.$$

Considering  $m = 0, i = 0, n = -2p - 1$  and  $j = -2q - 1$  in (3-5), we get

$$2d_{p,q}^{\bar{0}}(-2p, -2q) = d_{p,q}^{\bar{1}}(\frac{1}{2}, \frac{1}{2}) + d_{p,q}^{\bar{1}}(-2p - \frac{1}{2}, -2q - \frac{1}{2}) = 0.$$

So we obtain  $d_{r,s}^{\bar{0}}(m, i) = 0$  for all  $(m, i) \in \mathbb{Z} \times \mathbb{Z}$  and  $(r, s) \neq (0, 0)$ .  $\square$

**Lemma 3.3.** Let  $\varphi = \sum_{r,s \in \mathbb{Z}} \varphi_{r,s}$  be a  $\frac{1}{2}$ -derivation of  $\mathcal{S}(p, q)$  and  $(r, s) = (0, 0)$ . Then

$$d_{0,0}^{\bar{0}}(m, i) = d_{0,0}^{\bar{1}}(m' + \frac{1}{2}, i' + \frac{1}{2})$$

for all  $(m, i), (m', i') \in \mathbb{Z} \times \mathbb{Z}$ .

*Proof.* Writing (3-3) with  $(r, s) = (0, 0)$ , we have

$$(3-19) \quad ((n+p)(i+q) - (m+p)(j+q)) \\ \times (2d_{0,0}^{\bar{0}}(m+n, i+j) - d_{0,0}^{\bar{0}}(m, i) - d_{0,0}^{\bar{0}}(n, j)) = 0.$$

Taking  $n = j = 0$  in (3-19), we obtain

$$(pi - qm)(d_{0,0}^{\bar{0}}(m, i) - d_{0,0}^{\bar{0}}(0, 0)) = 0.$$

Therefore, we derive

$$(3-20) \quad d_{0,0}^{\bar{0}}(m, i) = d_{0,0}^{\bar{0}}(0, 0) \quad \text{if } pi - qm \neq 0.$$

Now, if  $pi - qm = 0$ , then we can assume  $m = kp$  and  $i = kq, k \in \mathbb{C}^*$ . We choose  $pj - qn \neq 0$  in (3-19), it becomes

$$(3-21) \quad (k+1)((n+p)q - (j+q)p) \\ \times (2d_{0,0}^{\bar{0}}(kp+n, kq+j) - d_{0,0}^{\bar{0}}(kp, kq) - d_{0,0}^{\bar{0}}(n, j)) = 0.$$

Since  $p(kq+j) - q(kp+n) \neq 0$ , by (3-20) we obtain

$$d_{0,0}^{\bar{0}}(kp+n, kq+j) = d_{0,0}^{\bar{0}}(n, j) = d_{0,0}^{\bar{0}}(0, 0).$$

Substituting it into (3-21), we have

$$(k+1)(d_{0,0}^{\bar{0}}(kp, kq) - d_{0,0}^{\bar{0}}(0, 0)) = 0.$$

Hence, it follows that

$$d_{0,0}^{\bar{0}}(kp, kq) = d_{0,0}^{\bar{0}}(0, 0) \quad \text{if } k \neq -1.$$

We substitute  $m = 0$ ,  $n = -p$ ,  $i = -q$  and  $j = 0$  into (3-19). By (3-20), it leads to

$$(3-22) \quad d_{0,0}^{\bar{0}}(-p, -q) = d_{0,0}^{\bar{0}}(0, 0).$$

Consequently, we can conclude that  $d_{0,0}^{\bar{0}}(m, i) = d_{0,0}^{\bar{0}}(0, 0)$  for all  $(m, i) \in \mathbb{Z} \times \mathbb{Z}$ .

Therefore, (3-5) becomes

$$d_{0,0}^{\bar{1}}(m + \frac{1}{2}, i + \frac{1}{2}) + d_{0,0}^{\bar{1}}(n + \frac{1}{2}, j + \frac{1}{2}) = 2d_{0,0}^{\bar{0}}(0, 0).$$

Taking  $m = n$  and  $i = j$ , we get

$$(3-23) \quad d_{0,0}^{\bar{1}}(m + \frac{1}{2}, i + \frac{1}{2}) = d_{0,0}^{\bar{0}}(0, 0) \quad \text{for all } (m, i) \in \mathbb{Z} \times \mathbb{Z}.$$

Combining with (3-22) and (3-23), we conclude that  $\varphi_{0,0} = d_{0,0}^{\bar{0}}(0, 0)id$ .  $\square$

From Lemmas 3.2 and 3.3, the subsequent proposition is directly derived.

**Proposition 3.4.** *Let  $p, q \in \mathbb{C}^*$ . Then*

$$\Delta^{\bar{0}}(\mathcal{S}(p, q)) = \langle id \rangle.$$

**3.1.2.** *The case  $p = 0, q \neq 0$ .*

**Lemma 3.5.** *Let  $\varphi = \sum_{r,s \in \mathbb{Z}} \varphi_{r,s}$  be a  $\frac{1}{2}$ -derivation of  $\mathcal{S}(0, q)$ ,  $r \in \mathbb{Z}^*$  and  $s \in \mathbb{Z}$ . Then*

$$\varphi_{r,s} = 0.$$

*Proof.* According to Lemma 2.3 from [10], we can obtain  $d_{r,s}^{\bar{0}}(m, i) = 0$  for all  $(m, i) \in \mathbb{Z} \times \mathbb{Z}$ . So, (3-5) becomes

$$d_{r,s}^{\bar{1}}(m + \frac{1}{2}, i + \frac{1}{2}) + d_{r,s}^{\bar{1}}(n + \frac{1}{2}, j + \frac{1}{2}) = 0.$$

Taking  $m = n$  and  $i = j$ , we get  $d_{r,s}^{\bar{1}}(m + \frac{1}{2}, i + \frac{1}{2}) = 0$  for all  $(m, i) \in \mathbb{Z} \times \mathbb{Z}$ .  $\square$

**Lemma 3.6.** *Let  $\varphi = \sum_{r,s \in \mathbb{Z}} \varphi_{r,s}$  be a  $\frac{1}{2}$ -derivation of  $\mathcal{S}(0, q)$ ,  $r = 0$  and  $s \in \mathbb{Z}^*$ . Then*

$$\varphi_{0,s} = 0.$$

*Proof.* If  $s \neq q$ , then we can obtain that  $d_{0,s}^{\bar{0}}(m, i) = 0$  for all  $(m, i) \in \mathbb{Z} \times \mathbb{Z}$  is established based on Lemma 2.4 in [10]. Hence, (3-5) becomes

$$d_{0,s}^{\bar{1}}\left(m + \frac{1}{2}, i + \frac{1}{2}\right) + d_{0,s}^{\bar{1}}\left(n + \frac{1}{2}, j + \frac{1}{2}\right) = 0.$$

Taking  $m = n$  and  $i = j$ , we get

$$d_{0,s}^{\bar{1}}\left(m + \frac{1}{2}, i + \frac{1}{2}\right) = 0 \quad \text{for all } (m, i) \in \mathbb{Z} \times \mathbb{Z}.$$

If  $s = q$ , then we arrive at  $d_{0,q}^{\bar{0}}(m, i) = 0$  unless  $(m, i) = (0, -2q)$ , which has been proved in Lemma 2.4 of [10]. Therefore (3-5) becomes

$$(3-24) \quad d_{0,q}^{\bar{1}}\left(m + \frac{1}{2}, i + \frac{1}{2}\right) + d_{0,q}^{\bar{1}}\left(n + \frac{1}{2}, j + \frac{1}{2}\right) = 0 \quad \text{for } (m+n+1, i+j+1) \neq (0, -2q),$$

$$(3-25) \quad d_{0,q}^{\bar{1}}\left(-n - \frac{1}{2}, -j - 2q - \frac{1}{2}\right) + d_{0,q}^{\bar{1}}\left(n + \frac{1}{2}, j + \frac{1}{2}\right) = 2d_{0,q}^{\bar{0}}(0, -2q).$$

By setting  $m = n = i = j$  into (3-24) and observing  $(m + n + 1, i + j + 1) = (2m + 1, 2i + 1) \neq (0, -2q)$ , we deduce, from (3-24),

$$d_{0,q}^{\bar{1}}\left(m + \frac{1}{2}, i + \frac{1}{2}\right) = 0 \quad \text{for all } (m, i) \in \mathbb{Z} \times \mathbb{Z}.$$

In the end, with the use of (3-25), we can obtain  $d_{0,q}^{\bar{0}}(0, -2q) = 0$ .  $\square$

**Lemma 3.7.** *Let  $\varphi = \sum_{r,s \in \mathbb{Z}} \varphi_{r,s}$  be a  $\frac{1}{2}$ -derivation of  $\mathcal{S}(0, q)$  and  $(r, s) = (0, 0)$ . Then*

$$d_{0,0}^{\bar{0}}(m, i) = d_{0,0}^{\bar{1}}\left(m' + \frac{1}{2}, i' + \frac{1}{2}\right)$$

for all  $(m, i), (m', i') \in \mathbb{Z} \times \mathbb{Z}$ .

*Proof.* As stated in Lemma 2.7 of [10], we can deduce  $d_{0,0}^{\bar{0}}(m, i) = d_{0,0}^{\bar{0}}(0, 0)$  for all  $(m, i) \in \mathbb{Z} \times \mathbb{Z}$ . Then, (3-5) becomes

$$d_{0,0}^{\bar{1}}\left(m + \frac{1}{2}, i + \frac{1}{2}\right) + d_{0,0}^{\bar{1}}\left(n + \frac{1}{2}, j + \frac{1}{2}\right) = 2d_{0,0}^{\bar{0}}(0, 0).$$

Taking  $m = n = i = j$ , we have

$$d_{0,0}^{\bar{1}}\left(m + \frac{1}{2}, i + \frac{1}{2}\right) = d_{0,0}^{\bar{0}}(0, 0)$$

for all  $(m, i) \in \mathbb{Z} \times \mathbb{Z}$ .  $\square$

Lemmas 3.5, 3.6 and 3.7 jointly yield the following proposition.

**Proposition 3.8.** *Let  $p = 0$  and  $q \in \mathbb{C}^*$ . Then*

$$\Delta^{\bar{0}}(\mathcal{S}(0, q)) = \langle id \rangle.$$

**3.1.3.** The case  $p \neq 0, q = 0$ . For this case, (3-3) and (3-4) are expressed as, respectively,

$$(3-26) \quad 2 \begin{vmatrix} n+p & j \\ m+p & i \end{vmatrix} d_{r,s}^{\bar{0}}(m+n, i+j) \\ = \begin{vmatrix} n+p & j \\ m+r+p & i+s \end{vmatrix} d_{r,s}^{\bar{0}}(m, i) + \begin{vmatrix} n+r+p & j+s \\ m+p & i \end{vmatrix} d_{r,s}^{\bar{0}}(n, j),$$

$$(3-27) \quad 2 \begin{vmatrix} n+\frac{p}{2}+\frac{1}{2} & j+\frac{1}{2} \\ m+p & i \end{vmatrix} d_{r,s}^{\bar{1}}(m+n+\frac{1}{2}, i+j+\frac{1}{2}) \\ = \begin{vmatrix} n+\frac{p}{2}+\frac{1}{2} & j+\frac{1}{2} \\ m+r+p & i+s \end{vmatrix} d_{r,s}^{\bar{0}}(m, i) + \begin{vmatrix} n+r+\frac{p}{2}+\frac{1}{2} & j+s+\frac{1}{2} \\ m+p & i \end{vmatrix} d_{r,s}^{\bar{1}}(n+\frac{1}{2}, j+\frac{1}{2}).$$

**Lemma 3.9.** Let  $\varphi = \sum_{r,s \in \mathbb{Z}} \varphi_{r,s}$  be a  $\frac{1}{2}$ -derivation of  $\mathcal{S}(p, 0)$ ,  $r \in \mathbb{Z}$  and  $s \in \mathbb{Z}^*$ . Then

$$\varphi_{r,s} = 0.$$

*Proof.* Taking  $n = j = 0$  in (3-26), we can see that

$$(3-28) \quad p(i-s)d_{r,s}^{\bar{0}}(m, i) = (i(r+p) - s(m+p))d_{r,s}^{\bar{0}}(0, 0).$$

Letting  $i = s$  and  $m \neq r$  in (3-28), we obtain

$$d_{r,s}^{\bar{0}}(0, 0) = 0.$$

Furthermore, substituting it into (3-28), it leads to

$$(i-s)d_{r,s}^{\bar{0}}(m, i) = 0 \quad \text{and} \quad d_{r,s}^{\bar{0}}(m, i) = 0 \quad \text{for } i \neq s.$$

Putting  $i = s, j \notin \{0, s\}$  and  $n \neq \frac{j(m+r+p)}{2s} - p$  in (3-26), then we have

$$d_{r,s}^{\bar{0}}(m+n, i+j) = 0 \quad \text{and} \quad d_{r,s}^{\bar{0}}(n, j) = 0.$$

Hence we obtain

$$(2s(n+p) - j(m+r+p))d_{r,s}^{\bar{0}}(m, s) = 0 \quad \text{and} \quad d_{r,s}^{\bar{0}}(m, s) = 0.$$

Therefore, we derive

$$(3-29) \quad d_{r,s}^{\bar{0}}(m, i) = 0 \quad \text{for all } (m, i) \in \mathbb{Z} \times \mathbb{Z}.$$

Taking into account both (3-27) and (3-29), then we infer

$$2 \begin{vmatrix} n+\frac{p}{2}+\frac{1}{2}j+\frac{1}{2} \\ m+pi \end{vmatrix} d_{r,s}^{\bar{1}}(m+n+\frac{1}{2}, i+j+\frac{1}{2}) = \begin{vmatrix} n+r+\frac{p}{2}+\frac{1}{2}j+s+\frac{1}{2} \\ m+pi \end{vmatrix} d_{r,s}^{\bar{1}}(n+\frac{1}{2}, j+\frac{1}{2}).$$

Substituting  $m = i = 0$ , we get

$$(j - s + \frac{1}{2})d_{r,s}^{\bar{1}}(n + \frac{1}{2}, j + \frac{1}{2}) = 0.$$

Since  $j, s \in \mathbb{Z}$ , then we have  $j - s + \frac{1}{2} \neq 0$ . Hence it follows that

$$d_{r,s}^{\bar{1}}(n + \frac{1}{2}, j + \frac{1}{2}) = 0$$

for all  $(n, j) \in \mathbb{Z} \times \mathbb{Z}$ . □

**Lemma 3.10.** Let  $\varphi = \sum_{r,s \in \mathbb{Z}} \varphi_{r,s}$  be a  $\frac{1}{2}$ -derivation of  $\mathcal{S}(p, 0)$ ,  $r \in \mathbb{Z}^*$  and  $s = 0$ .

(1) If  $r \neq p$ , then  $\varphi_{r,0} = 0$ .

(2) If  $r = p$ , then  $d_{p,0}^{\bar{0}}(m, i) = 0$  for all  $(m, i) \neq (-2p, 0)$  and  $d_{p,0}^{\bar{1}}(m + \frac{1}{2}, i + \frac{1}{2}) = 0$  for all  $(m, i) \in \mathbb{Z} \times \mathbb{Z}$ .

*Proof.* With  $s = 0$ , (3-26) can be expressed as

$$(3-30) \quad 2 \begin{vmatrix} n+p & j \\ m+p & i \end{vmatrix} d_{r,0}^{\bar{0}}(m+n, i+j) \\ = \begin{vmatrix} n+p & j \\ m+r+p & i \end{vmatrix} d_{r,0}^{\bar{0}}(m, i) + \begin{vmatrix} n+r+p & j \\ m+p & i \end{vmatrix} d_{r,0}^{\bar{0}}(n, j).$$

Taking  $n = j = 0$  in (3-30), we get

$$pid_{r,0}^{\bar{0}}(m, i) = (p+r)id_{r,0}^{\bar{0}}(0, 0).$$

Thus, we have

$$(3-31) \quad d_{r,0}^{\bar{0}}(m, i) = (1+rp^{-1})d_{r,0}^{\bar{0}}(0, 0) \text{ for } i \neq 0.$$

We proceed with a classification discussion based on whether or not  $p \in \mathbb{Z}$ .

Case 1:  $p \in \mathbb{Z}$ . Choosing  $m = -n$  and  $j = -i \neq 0$  in (3-30) and applying (3-31), then we obtain

$$4pd_{r,0}^{\bar{0}}(0, 0) = (r+2p)(d_{r,0}^{\bar{0}}(-n, i) + d_{r,0}^{\bar{0}}(n, -i)) \\ = 2(r+2p)(1+rp^{-1})d_{r,0}^{\bar{0}}(0, 0),$$

$$r(r+3p)d_{r,0}^{\bar{0}}(0, 0) = 0.$$

Thus, we get

$$d_{r,0}^{\bar{0}}(0, 0) = 0 \quad \text{for } r \neq -3p.$$

We take  $r = -3p$  and use (3-31). Hence it follows that

$$(3-32) \quad d_{r,0}^{\bar{0}}(m, i) = -2d_{r,0}^{\bar{0}}(0, 0) \quad \text{for } i \neq 0.$$

Putting  $m = n = 0$ ,  $i, j, i \pm j \neq 0$  in (3-30) and applying (3-32), then we can derive

$$\begin{aligned} 2p(i-j)d_{r,0}^{\bar{0}}(0, i+j) &= p(i+2j)d_{r,0}^{\bar{0}}(0, i) - p(2i+j)d_{r,0}^{\bar{0}}(0, j), \\ 2p(i-j)d_{r,0}^{\bar{0}}(0, 0) &= p(i+2j)d_{r,0}^{\bar{0}}(0, 0) - p(2i+j)d_{r,0}^{\bar{0}}(0, 0), \\ d_{r,0}^{\bar{0}}(0, 0) &= 0. \end{aligned}$$

Case 2:  $p \notin \mathbb{Z}$ . Observing that  $1+rp^{-1} \neq 0$  in this case, we take  $m = n = 0$  in (3-30), then we can obtain

$$2p(i-j)d_{r,0}^{\bar{0}}(0, i+j) = (p(i-j) - rj)d_{r,0}^{\bar{0}}(0, i) + (p(i-j) + ri)d_{r,0}^{\bar{0}}(0, j).$$

We choose  $i, j, i \pm j \neq 0$ . Then owing to (3-31) and  $1+rp^{-1} \neq 0$ , we have

$$d_{r,0}^{\bar{0}}(0, 0) = 0.$$

Hence by (3-31), we can deduce that

$$(3-33) \quad d_{r,0}^{\bar{0}}(m, i) = 0 \quad \text{for } i \neq 0.$$

It remains to analyze  $d_{r,0}^{\bar{0}}(m, 0)$ . To this end, putting  $m = 0$  and  $j = -i \neq 0$  in (3-30) and applying (3-33), hence we have

$$\begin{aligned} 2(2p+n)i d_{r,0}^{\bar{0}}(n, 0) &= (2p+n+r)i(d_{r,0}^{\bar{0}}(0, i) + d_{r,0}^{\bar{0}}(n, -i)), \\ (2p+n)d_{r,0}^{\bar{0}}(n, 0) &= 0. \end{aligned}$$

Therefore, we can conclude

$$d_{r,0}^{\bar{0}}(n, 0) = 0 \quad \text{for } n \neq -2p.$$

Taking  $m = -2p$ ,  $i = 0$  and  $j \neq 0$  in (3-30) and using (3-33), it follows that

$$\begin{aligned} 2pj d_{r,0}^{\bar{0}}(-2p+n, j) &= (p-r)j d_{r,0}^{\bar{0}}(-2p, 0) + pj d_{r,0}^{\bar{0}}(n, j), \\ (p-r)d_{r,0}^{\bar{0}}(-2p, 0) &= 0. \end{aligned}$$

So,  $d_{r,0}^{\bar{0}}(-2p, 0) = 0$  for  $r \neq p$  is derived.

If  $r \neq p$ , then (3-27) becomes

$$(3-34) \quad 2 \begin{vmatrix} n+\frac{p}{2}+\frac{1}{2} & j+\frac{1}{2} \\ m+p & i \end{vmatrix} d_{r,0}^{\bar{1}}(m+n+\frac{1}{2}, i+j+\frac{1}{2}) \\ = \begin{vmatrix} n+r+\frac{p}{2}+\frac{1}{2} & j+\frac{1}{2} \\ m+p & i \end{vmatrix} d_{r,0}^{\bar{1}}(n+\frac{1}{2}, j+\frac{1}{2}).$$

By taking  $m = i = 0$  into (3-34), we get

$$(j+\frac{1}{2})d_{r,0}^{\bar{1}}(n+\frac{1}{2}, j+\frac{1}{2}) = 0.$$

Since  $j \in \mathbb{Z}$ , then  $j + \frac{1}{2} \neq 0$  is inferred. Furthermore, we have

$$d_{r,0}^{\bar{1}}(n + \frac{1}{2}, j + \frac{1}{2}) = 0 \quad \text{for all } (n, j) \in \mathbb{Z} \times \mathbb{Z} \text{ and } r \neq p.$$

If  $r = p$ , the term in (3-27) that includes  $d_{p,0}^{\bar{0}}(m, i)$  is

$$\begin{aligned} ((n + \frac{p}{2} + \frac{1}{2})i - (j + \frac{1}{2})(m + 2p))d_{p,0}^{\bar{0}}(m, i) &= 0 \quad \text{for } (m, i) \neq (-2p, 0), \\ (j + \frac{1}{2})(-2p + p + p)d_{p,0}^{\bar{0}}(-2p, 0) &= 0 \quad \text{for } (m, i) = (-2p, 0). \end{aligned}$$

Therefore, the term with  $d_{p,0}^{\bar{0}}(m, i)$  in (3-27) vanishes. Equation (3-27) becomes

$$(3-35) \quad 2 \begin{vmatrix} n + \frac{p}{2} + \frac{1}{2} & j + \frac{1}{2} \\ m + p & i \end{vmatrix} d_{p,0}^{\bar{1}}(m + n + \frac{1}{2}, i + j + \frac{1}{2}) \\ = \begin{vmatrix} n + \frac{3p}{2} + \frac{1}{2} & j + \frac{1}{2} \\ m + p & i \end{vmatrix} d_{p,0}^{\bar{1}}(n + \frac{1}{2}, j + \frac{1}{2}).$$

Taking  $m = i = 0$  into (3-35), so we obtain

$$d_{p,0}^{\bar{1}}(n + \frac{1}{2}, j + \frac{1}{2}) = 0$$

for all  $(n, j) \in \mathbb{Z} \times \mathbb{Z}$ . □

According to this conclusion, we can get the following lemma.

**Lemma 3.11.** *Let  $p \in \mathbb{Z}^*$ . Then the linear map  $\alpha : \mathcal{S}(p, 0) \rightarrow \mathcal{S}(p, 0)$  is a  $\frac{1}{2}$ -derivation of  $\mathcal{S}(p, 0)$  such that*

$$\alpha(L_{m,i}) = \begin{cases} 0, & (m, i) \neq (-2p, 0), \\ L_{-p,0}, & (m, i) = (-2p, 0), \end{cases} \quad \alpha(G_{m+\frac{1}{2}, i+\frac{1}{2}}) = 0$$

for  $(m, i) \in \mathbb{Z} \times \mathbb{Z}$ .

*Proof.* We observe that  $\alpha = \sum_{r,s \in \mathbb{Z}} \alpha_{r,s} = \alpha_{p,0}$ . In view of Lemma 3.1 we need to check (3-3) and (3-4) for  $(r, s) = (p, 0)$  and

$$(3-36) \quad d_{p,0}^{\bar{0}}(m, i) = \begin{cases} 0, & (m, i) \neq (-2p, 0), \\ 1, & (m, i) = (-2p, 0), \end{cases}$$

$$(3-37) \quad d_{p,0}^{\bar{1}}(m + \frac{1}{2}, i + \frac{1}{2}) = 0, \quad (m, i) \in \mathbb{Z} \times \mathbb{Z}.$$

Firstly, we prove that (3-3) holds.

Case 1:  $(m, i), (n, j), (m + n, i + j) \neq (-2p, 0)$ . Then both sides of (3-3) are zero.

Case 2:  $(m, i) = (-2p, 0)$ . Then (3-3) becomes

$$2pjd_{p,0}^{\bar{0}}(-2p + n, j) = pj d_{p,0}^{\bar{0}}(n, j).$$

If  $j = 0$ , then it is trivially satisfied, otherwise both sides are zero by (3-36).

Case 3:  $(n, j) = (-2p, 0)$ . Then (3-3) becomes

$$2pid_{p,0}^{\bar{0}}(-2p + m, i) = pid_{p,0}^{\bar{0}}(m, i),$$

so this case is similar to Case 2.

Case 4:  $(m + n, i + j) = (-2p, 0)$ . Then (3-3) becomes

$$0 = pid_{p,0}^{\bar{0}}(-2p - n, i) + pid_{p,0}^{\bar{0}}(n, -i),$$

and again this holds by (3-36).

Next, we proceed to prove that (3-4) holds. Due to (3-37), (3-4) becomes

$$\left| \begin{array}{cc} n + \frac{p}{2} + \frac{1}{2} & j + \frac{1}{2} \\ m + 2p & i \end{array} \right| d_{p,0}^{\bar{0}}(m, i) = 0.$$

Given (3-36), it holds. □

The corollary is an immediate consequence of Lemmas 3.9 and 3.11.

**Corollary 3.12.** *Let  $\varphi$  be a  $\frac{1}{2}$ -derivation of  $\mathcal{S}(p, 0)$ ,  $r, s \in \mathbb{Z}$ .*

- (1) *If  $p \notin \mathbb{Z}$ , then  $\varphi_{r,s} = 0$  for  $(r, s) \neq (0, 0)$ .*
- (2) *If  $p \in \mathbb{Z}$ , then  $\varphi_{r,s} = 0$  for all  $(r, s) \notin \{(0, 0), (p, 0)\}$  and  $\varphi_{p,0} \in \langle \alpha \rangle$ .*

Finally we proceed to prove the case of  $r = s = 0$ .

**Lemma 3.13.** *Let  $\varphi$  be a  $\frac{1}{2}$ -derivation of  $\mathcal{S}(p, 0)$  and  $r = s = 0$ . Then*

$$d_{0,0}^{\bar{0}}(m, i) = d_{0,0}^{\bar{1}}(m' + \frac{1}{2}, i' + \frac{1}{2})$$

for all  $(m, i), (m', i') \in \mathbb{Z} \times \mathbb{Z}$ .

*Proof.* In this situation, (3-3) and (3-4) are presented as, respectively,

$$(3-38) \quad (i(n+p) - j(m+p))(2d_{0,0}^{\bar{0}}(m+n, i+j) - d_{0,0}^{\bar{0}}(m, i) - d_{0,0}^{\bar{0}}(n, j)) = 0,$$

$$(3-39) \quad (i(n + \frac{p}{2} + \frac{1}{2}) - (j + \frac{1}{2})(m+p)) \\ \times (2d_{0,0}^{\bar{1}}(m+n + \frac{1}{2}, i+j + \frac{1}{2}) - d_{0,0}^{\bar{0}}(m, i) - d_{0,0}^{\bar{1}}(n + \frac{1}{2}, j + \frac{1}{2})) = 0.$$

Taking  $n = j = 0$  and  $i \neq 0$  in (3-38), we obtain

$$(3-40) \quad d_{0,0}^{\bar{0}}(m, i) = d_{0,0}^{\bar{0}}(0, 0) \quad \text{for } i \neq 0.$$

We choose  $j \neq 0$  and  $n = -m$  in (3-38), then we arrive at

$$(m+p)(2d_{0,0}^{\bar{0}}(0, j) - d_{0,0}^{\bar{0}}(m, 0) - d_{0,0}^{\bar{0}}(-m, j)) = 0.$$

Due to (3-40), it follows that

$$d_{0,0}^{\bar{0}}(m, 0) = d_{0,0}^{\bar{0}}(0, 0) \quad \text{for } m \neq -p.$$

Substituting  $m = -p$ ,  $n = 0$  and  $j = -i \neq 0$  into (3-38) and using (3-40), we can derive that

$$\begin{aligned} 2d_{0,0}^{\bar{0}}(-p, 0) - d_{0,0}^{\bar{0}}(-p, i) - d_{0,0}^{\bar{0}}(0, -i) &= 0, \\ d_{0,0}^{\bar{0}}(-p, 0) &= d_{0,0}^{\bar{0}}(0, 0). \end{aligned}$$

Taking  $m = i = 0$  in (3-39), we can see that

$$p(j + \frac{1}{2})(2d_{0,0}^{\bar{1}}(n + \frac{1}{2}, j + \frac{1}{2}) - d_{0,0}^{\bar{0}}(0, 0) - d_{0,0}^{\bar{1}}(n + \frac{1}{2}, j + \frac{1}{2})) = 0.$$

Since  $p \neq 0$  and  $j \in \mathbb{Z}$ , then we have  $p(j + \frac{1}{2}) \neq 0$ . Furthermore, we are able to conclude that

$$d_{0,0}^{\bar{1}}(n + \frac{1}{2}, j + \frac{1}{2}) = d_{0,0}^{\bar{0}}(0, 0)$$

for all  $(n, j) \in \mathbb{Z} \times \mathbb{Z}$ . □

Directly from Lemmas 3.9, 3.13 and Corollary 3.12, we deduce the following proposition.

**Proposition 3.14.** *Let  $p \in \mathbb{C}^*$  and  $q = 0$ . Then*

$$\Delta^{\bar{0}}(\mathcal{S}(p, 0)) = \begin{cases} \langle id \rangle, & p \notin \mathbb{Z}, \\ \langle id, \alpha \rangle, & p \in \mathbb{Z}^*, \end{cases}$$

where  $\alpha$  is as in Lemma 3.11.

**3.1.4. The case  $p = q = 0$ .** Rewriting (3-4) with  $p = q = 0$ , we obtain

$$\begin{aligned} (3-41) \quad 2 \begin{vmatrix} n+\frac{1}{2} & j+\frac{1}{2} \\ m & i \end{vmatrix} d_{r,s}^{\bar{1}}(m+n+\frac{1}{2}, i+j+\frac{1}{2}) \\ = \begin{vmatrix} n+\frac{1}{2} & j+\frac{1}{2} \\ m+r & i+s \end{vmatrix} d_{r,s}^{\bar{0}}(m, i) + \begin{vmatrix} n+r+\frac{1}{2} & j+s+\frac{1}{2} \\ m & i \end{vmatrix} d_{r,s}^{\bar{1}}(n+\frac{1}{2}, j+\frac{1}{2}). \end{aligned}$$

**Lemma 3.15.** *Let  $\varphi = \sum_{r,s \in \mathbb{Z}} \varphi_{r,s}$  be a  $\frac{1}{2}$ -derivation of  $\mathcal{S}(0, 0)$  and let  $(r, s) \in \mathbb{Z} \times \mathbb{Z} \setminus \{(0, 0)\}$ . Then*

$$\varphi_{r,s} = 0.$$

*Proof.*  $d_{r,s}^{\bar{0}}(m, i) = 0$  for all  $(m, i) \in \mathbb{Z} \times \mathbb{Z}$  is derived from Lemma 2.9 in [10]. Then, (3-41) becomes

$$\begin{aligned} (3-42) \quad 2 \begin{vmatrix} n+\frac{1}{2} & j+\frac{1}{2} \\ m & i \end{vmatrix} d_{r,s}^{\bar{1}}(m+n+\frac{1}{2}, i+j+\frac{1}{2}) \\ = \begin{vmatrix} n+r+\frac{1}{2} & j+s+\frac{1}{2} \\ m & i \end{vmatrix} d_{r,s}^{\bar{1}}(n+\frac{1}{2}, j+\frac{1}{2}). \end{aligned}$$

Taking  $m = j = 0$  in (3-42), we obtain

$$(3-43) \quad \begin{aligned} 2i(n + \frac{1}{2})d_{r,s}^{\bar{1}}(n + \frac{1}{2}, i + \frac{1}{2}) &= i(n + r + \frac{1}{2})d_{r,s}^{\bar{1}}(n + \frac{1}{2}, \frac{1}{2}), \\ 2(n + \frac{1}{2})d_{r,s}^{\bar{1}}(n + \frac{1}{2}, i + \frac{1}{2}) &= (n + r + \frac{1}{2})d_{r,s}^{\bar{1}}(n + \frac{1}{2}, \frac{1}{2}) \quad \text{for } i \neq 0. \end{aligned}$$

It implies that  $d_{r,s}^{\bar{1}}(n + \frac{1}{2}, i + \frac{1}{2})$  is independent of  $i$  for  $i \neq 0$ . Substitution  $m = 0$ ,  $j = -i \neq 0$  into (3-42), it leads to

$$(3-44) \quad \begin{aligned} (2n + 1)d_{r,s}^{\bar{1}}(n + \frac{1}{2}, \frac{1}{2}) &= (n + r + \frac{1}{2})d_{r,s}^{\bar{1}}(n + \frac{1}{2}, -i + \frac{1}{2}) \\ &= (n + r + \frac{1}{2})d_{r,s}^{\bar{1}}(n + \frac{1}{2}, i + \frac{1}{2}). \end{aligned}$$

Multiplying (3-44) by  $2n + 1$  and using (3-43), we get

$$(2n + 1)^2 d_{r,s}^{\bar{1}}(n + \frac{1}{2}, \frac{1}{2}) = (n + r + \frac{1}{2})^2 d_{r,s}^{\bar{1}}(n + \frac{1}{2}, \frac{1}{2}).$$

Assuming  $d_{r,s}^{\bar{1}}(n + \frac{1}{2}, \frac{1}{2}) \neq 0$ , we obtain  $(2n + 1)^2 = (n + r + \frac{1}{2})^2$ , whence  $2n + 1 = \pm(n + r + \frac{1}{2})$ . It follows that  $n = r - \frac{1}{2}$  or  $n = -\frac{r}{3} - \frac{1}{2}$ , which contradicts  $n \in \mathbb{Z}$ .

Then utilizing (3-44) and  $n + r + \frac{1}{2} \neq 0$ , we have

$$d_{r,s}^{\bar{1}}(n + \frac{1}{2}, \frac{1}{2}) = \frac{2n + 1}{n + r + \frac{1}{2}} d_{r,s}^{\bar{1}}(n + \frac{1}{2}, \frac{1}{2}) = 0$$

for all  $(n, i) \in \mathbb{Z} \times \mathbb{Z}$ . □

**Lemma 3.16.** Let  $\varphi = \sum_{r,s \in \mathbb{Z}} \varphi_{r,s}$  be a  $\frac{1}{2}$ -derivation of  $\mathcal{S}(0, 0)$  and  $(r, s) = (0, 0)$ . Then

$$d_{0,0}^{\bar{1}}(m + \frac{1}{2}, i + \frac{1}{2}) = d_{0,0}^{\bar{0}}(m', i')$$

for  $(m, i) \in \mathbb{Z} \times \mathbb{Z}$ ,  $(m', i') \in \mathbb{Z} \times \mathbb{Z} \setminus \{(0, 0)\}$ .

*Proof.* According to Lemma 2.10 from [10], we can obtain

$$(3-45) \quad d_{0,0}^{\bar{0}}(m, i) = d_{0,0}^{\bar{0}}(m', i') \quad \text{for all } (m, i), (m', i') \in \mathbb{Z} \times \mathbb{Z} \setminus \{(0, 0)\}.$$

By substituting  $(r, s) = (0, 0)$  into (3-41), we get

$$(3-46) \quad \left| \begin{matrix} n + \frac{1}{2} & j + \frac{1}{2} \\ m & i \end{matrix} \right| (2d_{0,0}^{\bar{1}}(m + n + \frac{1}{2}, i + j + \frac{1}{2}) - d_{0,0}^{\bar{0}}(m, i) - d_{0,0}^{\bar{1}}(n + \frac{1}{2}, j + \frac{1}{2})) = 0.$$

We take  $m = j = 0$  and  $i \neq 0$  in (3-46). And since  $n + \frac{1}{2} \neq 0$ , we obtain

$$(3-47) \quad 2d_{0,0}^{\bar{1}}(n + \frac{1}{2}, i + \frac{1}{2}) = d_{0,0}^{\bar{0}}(0, i) + d_{0,0}^{\bar{1}}(n + \frac{1}{2}, \frac{1}{2}) \quad \text{for } i \neq 0.$$

Letting  $n = i = 0$  and  $m \neq 0$  in (3-46), and because of  $j + \frac{1}{2} \neq 0$ , we have

$$2d_{0,0}^{\bar{1}}(m + \frac{1}{2}, j + \frac{1}{2}) = d_{0,0}^{\bar{0}}(m, 0) + d_{0,0}^{\bar{1}}(12, j + \frac{1}{2}) \quad \text{for } m \neq 0.$$

Hence, due to (3-45), we arrive at

$$(3-48) \quad d_{0,0}^{\bar{1}}\left(n + \frac{1}{2}, \frac{1}{2}\right) = d_{0,0}^{\bar{1}}\left(\frac{1}{2}, i + \frac{1}{2}\right) \quad \text{for } n, i \neq 0.$$

Substituting  $m = -n \neq 0, i \neq 0$  and  $j = 0$  into (3-46), it leads to

$$(ni + \frac{1}{2}(n+i))\left(2d_{0,0}^{\bar{1}}\left(\frac{1}{2}, i + \frac{1}{2}\right) - d_{0,0}^{\bar{0}}(-n, i) - d_{0,0}^{\bar{1}}\left(n + \frac{1}{2}, \frac{1}{2}\right)\right) = 0.$$

If  $(n, i) \notin \{(0, 0), (-1, -1)\}$ , then we have  $ni + \frac{1}{2}(n+i) \neq 0$ . Therefore, it follows that

$$(3-49) \quad 2d_{0,0}^{\bar{1}}\left(\frac{1}{2}, i + \frac{1}{2}\right) = d_{0,0}^{\bar{0}}(-n, i) + d_{0,0}^{\bar{1}}\left(n + \frac{1}{2}, \frac{1}{2}\right) \quad \text{for } (n, i) \neq (0, 0), (-1, -1).$$

Now, combining (3-47), (3-48) and (3-49), we get

$$\begin{aligned} d_{0,0}^{\bar{1}}\left(n + \frac{1}{2}, \frac{1}{2}\right) &= d_{0,0}^{\bar{1}}\left(\frac{1}{2}, i + \frac{1}{2}\right) = d_{0,0}^{\bar{0}}(-n, i) \quad \text{for } n, i \neq 0 \text{ and } (n, i) \neq (-1, -1), \\ d_{0,0}^{\bar{1}}\left(n + \frac{1}{2}, i + \frac{1}{2}\right) &= d_{0,0}^{\bar{0}}(-n, i) = d_{0,0}^{\bar{0}}(n', i') \\ &\quad \text{for } (n, i) \neq (0, 0), (-1, -1) \text{ and } (n', i') \neq (0, 0). \end{aligned}$$

In particular, taking  $n = i = -1$  in (3-47), we have

$$2d_{0,0}^{\bar{1}}\left(-\frac{1}{2}, -\frac{1}{2}\right) = d_{0,0}^{\bar{0}}(0, -1) + d_{0,0}^{\bar{1}}\left(-\frac{1}{2}, \frac{1}{2}\right) = 2d_{0,0}^{\bar{0}}(0, -1) = 2d_{0,0}^{\bar{0}}(n', i')$$

for  $(n', i') \neq (0, 0)$ . Therefore, we can conclude

$$(3-50) \quad d_{0,0}^{\bar{1}}\left(m + \frac{1}{2}, i + \frac{1}{2}\right) = d_{0,0}^{\bar{0}}(m', i') \quad \text{for } (m, i), (m', i') \in \mathbb{Z} \times \mathbb{Z} \setminus \{(0, 0)\}$$

Taking  $n = j = 0$  in (3-46), we can deduce

$$(3-51) \quad (i - m)\left(2d_{0,0}^{\bar{1}}\left(m + \frac{1}{2}, i + \frac{1}{2}\right) - d_{0,0}^{\bar{0}}(m, i) - d_{0,0}^{\bar{1}}\left(\frac{1}{2}, \frac{1}{2}\right)\right) = 0.$$

If  $m \neq i$ , then we get  $(m, i) \neq (0, 0)$ . Consequently, it follows that  $d_{0,0}^{\bar{0}}(m, i) = d_{0,0}^{\bar{1}}\left(m + \frac{1}{2}, i + \frac{1}{2}\right)$ . Hence, from (3-51) we obtain that

$$d_{0,0}^{\bar{1}}\left(m + \frac{1}{2}, i + \frac{1}{2}\right) = d_{0,0}^{\bar{1}}\left(\frac{1}{2}, \frac{1}{2}\right) \quad \text{for } m \neq i.$$

Using (3-50), we have

$$d_{0,0}^{\bar{1}}\left(m + \frac{1}{2}, i + \frac{1}{2}\right) = d_{0,0}^{\bar{1}}\left(\frac{1}{2}, \frac{1}{2}\right) \quad \text{for } (m, i) \in \mathbb{Z} \times \mathbb{Z}. \quad \square$$

According to this conclusion, we are able to deduce the subsequent lemma.

**Lemma 3.17.** *Let  $p = q = 0$ . Then the linear map  $\beta : \mathcal{S}(0, 0) \rightarrow \mathcal{S}(0, 0)$  is a  $\frac{1}{2}$ -derivation of  $\mathcal{S}(0, 0)$  such that*

$$\beta(L_{m,i}) = \begin{cases} 0, & (m, i) \neq (0, 0), \\ L_{0,0}, & (m, i) = (0, 0), \end{cases} \quad \beta(G_{m+\frac{1}{2}, i+\frac{1}{2}}) = 0$$

for  $(m, i) \in \mathbb{Z} \times \mathbb{Z}$ .

*Proof.* We observe that  $\beta = \sum_{r,s \in \mathbb{Z}} \beta_{r,s} = \beta_{0,0}$ . In view of [Lemma 3.1](#), we need to check (3-3) and (3-4) for  $(r, s) = (0, 0)$  and

$$(3-52) \quad d_{0,0}^{\bar{0}}(m, i) = \begin{cases} 0, & (m, i) \neq (0, 0), \\ 1, & (m, i) = (0, 0), \end{cases}$$

$$(3-53) \quad d_{0,0}^{\bar{1}}\left(m + \frac{1}{2}, i + \frac{1}{2}\right) = 0$$

for  $(m, i) \in \mathbb{Z} \times \mathbb{Z}$ .

Firstly, we prove (3-3). In fact, we can split this proof into four cases:

- (1)  $(m, i), (n, j), (m+n, i+j) = (0, 0)$ ,
- (2)  $(m, i) = (0, 0)$ ,
- (3)  $(n, j) = (0, 0)$ ,
- (4)  $(m+n, i+j) \neq (0, 0)$ .

Obviously, under all these different cases, (3-3) is trivially satisfied.

Subsequently, we prove (3-4). Due to (3-53), (3-4) becomes

$$\left| \begin{array}{cc} n + \frac{1}{2} & j + \frac{1}{2} \\ m & i \end{array} \right| d_{0,0}^{\bar{0}}(m, i) = 0.$$

It clearly holds by (3-52). □

With [Lemmas 3.15](#) and [3.17](#), we can directly infer the subsequent proposition.

**Proposition 3.18.** *Let  $p = q = 0$ . Then*

$$\Delta^{\bar{0}}(\mathcal{S}(0, 0)) = \langle id, \beta \rangle,$$

where  $\beta$  is as in [Lemma 3.17](#).

By integrating [Propositions 3.4, 3.8, 3.14](#) and [3.18](#), we can deduce the following corollary.

**Corollary 3.19.** *Let  $p, q \in \mathbb{C}$ . Then*

$$\Delta^{\bar{0}}(\mathcal{S}(p, q)) = \begin{cases} \langle id \rangle, & q \neq 0, \\ \langle id, \alpha \rangle, & p \neq 0, q = 0, \\ \langle id, \beta \rangle, & p = q = 0. \end{cases}$$

where  $\alpha$  and  $\beta$  are as in [Lemmas 3.17](#) and [3.11](#).

**3.2. Odd  $\frac{1}{2}$ -derivations of  $\mathcal{S}(p, q)$ .** In this subsection we consider that a linear map  $\varphi : \mathcal{S}(p, q) \rightarrow \mathcal{S}(p, q)$  is *odd*, if  $\varphi(\mathcal{S}(p, q)_i) \subseteq \mathcal{S}(p, q)_{\bar{1}-i}$  for  $i \in \mathbb{Z}_2$ . In this case  $|\varphi| = 1$ , so  $\varphi$  is a  $\frac{1}{2}$ -superderivation of  $\mathcal{S}(p, q)$  if and only if

$$\begin{aligned} \varphi([x, y]) &= \frac{1}{2}([\varphi(x), y] + [x, \varphi(y)]), & x \in \mathcal{S}(p, q)_{\bar{0}}, \\ \varphi([x, y]) &= \frac{1}{2}([\varphi(x), y] - [x, \varphi(y)]), & x \in \mathcal{S}(p, q)_{\bar{1}}. \end{aligned}$$

Denote by  $\Delta^{\bar{1}}(\mathcal{S}(p, q))$  the space of odd  $\frac{1}{2}$ -superderivations of  $\mathcal{S}(p, q)$ . As usual, for any  $\varphi \in \Delta^{\bar{1}}(\mathcal{S}(p, q))$ , we write

$$\varphi = \sum_{r, s \in \mathbb{Z}} \varphi_{r+\frac{1}{2}, s+\frac{1}{2}},$$

where

$$(3-54) \quad \varphi_{r+\frac{1}{2}, s+\frac{1}{2}}(L_{m, i}) = d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(m, i)G_{m+r+\frac{1}{2}, i+s+\frac{1}{2}},$$

$$(3-55) \quad \varphi_{r, s}(G_{m+\frac{1}{2}, i+\frac{1}{2}}) = d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}\left(m + \frac{1}{2}, i + \frac{1}{2}\right)L_{m+r+1, i+s+1}$$

for some  $d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(m, i), d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}\left(m + \frac{1}{2}, i + \frac{1}{2}\right) \in \mathbb{C}, m, i, r, s \in \mathbb{Z}$ . We have

$$\varphi \in \Delta^{\bar{1}}(\mathcal{S}(p, q)) \quad \text{if and only if} \quad \varphi_{r+\frac{1}{2}, s+\frac{1}{2}} \in \Delta^{\bar{1}}(\mathcal{S}(p, q)) \quad \text{for all } r, s \in \mathbb{Z}.$$

**Lemma 3.20.** *Let*

$$\varphi_{r+\frac{1}{2}, s+\frac{1}{2}} : \mathcal{S}(p, q) \rightarrow \mathcal{S}(p, q)$$

*be a linear map satisfying (3-54) and (3-55). Then*

$$\varphi_{r+\frac{1}{2}, s+\frac{1}{2}} \in \Delta^{\bar{1}}(\mathcal{S}(p, q))$$

*if and only if the following three conditions hold:*

$$(3-56) \quad 2 \begin{vmatrix} n+p & j+q \\ m+p & i+q \end{vmatrix} d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(m+n, i+j) \\ = \begin{vmatrix} n+p & j+q \\ m+r+\frac{p}{2}+\frac{1}{2} & i+s+\frac{q}{2}+\frac{1}{2} \end{vmatrix} d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(m, i) \\ + \begin{vmatrix} n+r+\frac{p}{2}+\frac{1}{2} & j+s+\frac{q}{2}+\frac{1}{2} \\ m+p & i+q \end{vmatrix} d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(n, j),$$

$$(3-57) \quad 2 \begin{vmatrix} n+\frac{p}{2}+\frac{1}{2} & j+\frac{q}{2}+\frac{1}{2} \\ m+p & i+q \end{vmatrix} d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}\left(m+n+\frac{1}{2}, i+j+\frac{1}{2}\right) \\ = 2qd_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(m, i) + \begin{vmatrix} n+r+p+1 & j+s+q+1 \\ m+p & i+q \end{vmatrix} d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}\left(n+\frac{1}{2}, j+\frac{1}{2}\right),$$

$$(3-58) \quad 4qd_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(m+n+1, i+j+1) \\ = \begin{vmatrix} n+\frac{p}{2}+\frac{1}{2} & j+\frac{q}{2}+\frac{1}{2} \\ m+r+p+1 & i+s+q+1 \end{vmatrix} d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}\left(m+\frac{1}{2}, i+\frac{1}{2}\right) \\ - \begin{vmatrix} n+r+p+1 & j+s+q+1 \\ m+\frac{p}{2}+\frac{1}{2} & i+\frac{q}{2}+\frac{1}{2} \end{vmatrix} d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}\left(n+\frac{1}{2}, j+\frac{1}{2}\right).$$

*Proof.* By [Definition 2.1](#),  $\varphi_{r+\frac{1}{2},s+\frac{1}{2}} \in \Delta^{\bar{1}}(\mathcal{S}(p, q))$  if and only if the following three expressions are valid:

$$(3-59) \quad 2 \begin{vmatrix} n+p & j+q \\ m+p & i+q \end{vmatrix} \varphi_{r+\frac{1}{2},s+\frac{1}{2}}(L_{m+n,i+j}) \\ = [\varphi_{r+\frac{1}{2},s+\frac{1}{2}}(L_m, i), L_n, j] + [L_m, i + \varphi_{r+\frac{1}{2},s+\frac{1}{2}}(L_n, j)],$$

$$(3-60) \quad 2 \begin{vmatrix} n+\frac{1}{2}+\frac{p}{2} & j+\frac{1}{2}+\frac{q}{2} \\ m+p & i+q \end{vmatrix} \varphi_{r+\frac{1}{2},s+\frac{1}{2}}(G_{m+n+\frac{1}{2},i+j+\frac{1}{2}}) \\ = [\varphi_{r+\frac{1}{2},s+\frac{1}{2}}(L_m, i), G_{n+\frac{1}{2},j+\frac{1}{2}}] + [L_m, i, \varphi_{r+\frac{1}{2},s+\frac{1}{2}}(G_{n+\frac{1}{2},j+\frac{1}{2}})],$$

$$(3-61) \quad 4q\varphi_{r+\frac{1}{2},s+\frac{1}{2}}(L_{m+n+1,i+j+1}) \\ = [\varphi_{r+\frac{1}{2},s+\frac{1}{2}}(G_{m+\frac{1}{2},i+\frac{1}{2}}, G_{n+\frac{1}{2},j+\frac{1}{2}})] - [G_{m+\frac{1}{2},i+\frac{1}{2}}, \varphi_{r+\frac{1}{2},s+\frac{1}{2}}(G_{n+\frac{1}{2},j+\frac{1}{2}})].$$

Due to [\(3-59\)](#), we can see that

$$2 \begin{vmatrix} n+p & j+q \\ m+p & i+q \end{vmatrix} d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(m+n, i+j)G_{m+n+r+\frac{1}{2},i+j+s+\frac{1}{2}} \\ = 2 \begin{vmatrix} n+p & j+q \\ m+p & i+q \end{vmatrix} \varphi_{r+\frac{1}{2},s+\frac{1}{2}}(L_{m+n,i+j}) \\ = [\varphi_{r+\frac{1}{2},s+\frac{1}{2}}(L_m, i), L_n, j] + [L_m, i, \varphi_{r+\frac{1}{2},s+\frac{1}{2}}(L_n, j)] \\ = [d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(m, i)G_{m+\frac{1}{2},i+\frac{1}{2}}, L_n, j] + [L_m, i, d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(n, j)G_{n+r+\frac{1}{2},j+r+\frac{1}{2}}] \\ = \begin{vmatrix} n+p & j+q \\ m+r+\frac{p}{2}+\frac{1}{2} & i+s+\frac{q}{2}+\frac{1}{2} \end{vmatrix} d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(m, i)G_{m+n+r+\frac{1}{2},i+j+s+\frac{1}{2}} \\ + \begin{vmatrix} n+r+\frac{p}{2}+\frac{1}{2} & j+s+\frac{q}{2}+\frac{1}{2} \\ m+p & i+q \end{vmatrix} d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(n, j)G_{m+n+r+\frac{1}{2},i+j+s+\frac{1}{2}}.$$

Thus, [\(3-56\)](#) is obtained. By [\(3-60\)](#), we have

$$2 \begin{vmatrix} n+\frac{p}{2}+\frac{1}{2} & j+\frac{q}{2}+\frac{1}{2} \\ m+p & i+q \end{vmatrix} d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{1}}(m+n+\frac{1}{2}, i+j+\frac{1}{2})L_{m+n+r+1,i+j+s+1} \\ = 2 \begin{vmatrix} n+\frac{1}{2}+\frac{p}{2} & j+\frac{1}{2}+\frac{q}{2} \\ m+p & i+q \end{vmatrix} \varphi_{r+\frac{1}{2},s+\frac{1}{2}}(G_{m+n+\frac{1}{2},i+j+\frac{1}{2}}) \\ = [\varphi_{r+\frac{1}{2},s+\frac{1}{2}}(L_m, i), G_{n+\frac{1}{2},j+\frac{1}{2}}] + [L_m, i, \varphi_{r+\frac{1}{2},s+\frac{1}{2}}(G_{n+\frac{1}{2},j+\frac{1}{2}})] \\ = [d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(m, i)G_{m+\frac{1}{2},i+\frac{1}{2}}, G_{n+\frac{1}{2},j+\frac{1}{2}}] \\ + [L_m, i, d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{1}}(n+\frac{1}{2}, j+\frac{1}{2})L_{n+r+1,j+s+1}] \\ = 2qd_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(m, i)L_{m+n+r+1,i+j+s+1} \\ + \begin{vmatrix} n+r+p+1 & j+s+q+1 \\ m+p & i+q \end{vmatrix} d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{1}}(n+\frac{1}{2}, j+\frac{1}{2})L_{m+n+r+1,i+j+s+1}.$$

Consequently, we come to (3-57). Because of (3-61), we observe

$$\begin{aligned}
 & 4qd_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(m+n+1, i+j+1)G_{m+n+r+\frac{3}{2},i+j+s+\frac{3}{2}} \\
 &= 4q\varphi_{r+\frac{1}{2},s+\frac{1}{2}}(L_{m+n+1,i+j+1}) \\
 &= [\varphi_{r+\frac{1}{2},s+\frac{1}{2}}(G_{m+\frac{1}{2},i+\frac{1}{2}}), G_{n+\frac{1}{2},j+\frac{1}{2}}] - [G_{m+\frac{1}{2},i+\frac{1}{2}}, \varphi_{r+\frac{1}{2},s+\frac{1}{2}}(G_{n+\frac{1}{2},j+\frac{1}{2}})] \\
 &= [d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{1}}(G_{m+\frac{1}{2},i+\frac{1}{2}}), G_{n+\frac{1}{2},j+\frac{1}{2}}] + [G_{m+\frac{1}{2},i+\frac{1}{2}}, d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{1}}(G_{n+\frac{1}{2},j+\frac{1}{2}})] \\
 &= \left| \begin{array}{cc} n+\frac{p}{2}+\frac{1}{2} & j+\frac{q}{2}+\frac{1}{2} \\ m+r+p+1 & i+s+q+1 \end{array} \right| d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{1}}(m+\frac{1}{2}, i+\frac{1}{2})G_{m+n+r+\frac{3}{2},i+j+s+\frac{3}{2}} \\
 &\quad - \left| \begin{array}{cc} n+r+p+1 & j+s+q+1 \\ m+\frac{p}{2}+\frac{1}{2} & i+\frac{q}{2}+\frac{1}{2} \end{array} \right| d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{1}}(n+\frac{1}{2}, j+\frac{1}{2})G_{m+n+r+\frac{3}{2},i+j+s+\frac{3}{2}}.
 \end{aligned}$$

Hence, it follows that (3-58) holds truly.  $\square$

Let  $\varphi \in \Delta^{\bar{1}}(\mathcal{S}(p, q))$  and  $\psi_{m+\frac{1}{2},i+\frac{1}{2}}$  be the left multiplication by  $G_{m+\frac{1}{2},i+\frac{1}{2}}$  in  $\mathcal{S}(p, q)$ ,  $m, i, r, s \in \mathbb{Z}$ . Since  $|\psi_{m+\frac{1}{2},i+\frac{1}{2}}| = 1$  and we have, for all  $x, y \in \{L_{n,j}, G_{n+\frac{1}{2},j+\frac{1}{2}} \mid n, j \in \mathbb{Z}\}$ ,

$$\begin{aligned}
 \psi_{m+\frac{1}{2},i+\frac{1}{2}}([x, y]) &= [G_{m+\frac{1}{2},i+\frac{1}{2}}, [x, y]] \\
 &= [[G_{m+\frac{1}{2},i+\frac{1}{2}}, x], y] + (-1)^{|x|}[x, [G_{m+\frac{1}{2},i+\frac{1}{2}}, y]] \\
 &= [\psi_{m+\frac{1}{2},i+\frac{1}{2}}(x), y] + (-1)^{|x|}[x, \psi_{m+\frac{1}{2},i+\frac{1}{2}}(y)],
 \end{aligned}$$

$\psi_{m+\frac{1}{2},i+\frac{1}{2}}$  is an odd superderivation of  $\mathcal{S}(p, q)$ .

Since the supercommutator

$$[\varphi_{r+\frac{1}{2},s+\frac{1}{2}}, \psi_{m+\frac{1}{2},i+\frac{1}{2}}] = \varphi_{r+\frac{1}{2},s+\frac{1}{2}}\psi_{m+\frac{1}{2},i+\frac{1}{2}} + \psi_{m+\frac{1}{2},i+\frac{1}{2}}\varphi_{r+\frac{1}{2},s+\frac{1}{2}}$$

satisfies

$$\begin{aligned}
 (3-62) \quad & [\varphi_{r+\frac{1}{2},s+\frac{1}{2}}, \psi_{m+\frac{1}{2},i+\frac{1}{2}}](L_{n,j}) \\
 &= \varphi_{r+\frac{1}{2},s+\frac{1}{2}}([G_{m+\frac{1}{2},i+\frac{1}{2}}, L_{n,j}]) + [G_{m+\frac{1}{2},i+\frac{1}{2}}, \varphi_{r+\frac{1}{2},s+\frac{1}{2}}(L_{n,j})] \\
 &= \left| \begin{array}{cc} n+p & j+q \\ m+\frac{p}{2}+\frac{1}{2} & i+\frac{q}{2}+\frac{1}{2} \end{array} \right| d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{1}}(m+n+\frac{1}{2}, i+j+\frac{1}{2})L_{m+n+r+1,i+j+s+1} \\
 &\quad + 2qd_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{1}}(n, j)L_{m+n+r+1,i+j+s+1},
 \end{aligned}$$

$$\begin{aligned}
 (3-63) \quad & [\varphi_{r+\frac{1}{2},s+\frac{1}{2}}, \psi_{m+\frac{1}{2},i+\frac{1}{2}}](G_{n+\frac{1}{2},j+\frac{1}{2}}) \\
 &= \varphi_{r+\frac{1}{2},s+\frac{1}{2}}([G_{m+\frac{1}{2},i+\frac{1}{2}}, G_{n+\frac{1}{2},j+\frac{1}{2}}]) + [G_{m+\frac{1}{2},i+\frac{1}{2}}, \varphi_{r+\frac{1}{2},s+\frac{1}{2}}(G_{n+\frac{1}{2},j+\frac{1}{2}})] \\
 &= 2qd_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(m+n+1, i+j+1)G_{m+n+r+\frac{3}{2},i+j+s+\frac{3}{2}} \\
 &\quad + \left| \begin{array}{cc} n+r+p+1 & j+s+q+1 \\ m+\frac{p}{2}+\frac{1}{2} & i+\frac{q}{2}+\frac{1}{2} \end{array} \right| d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{1}}(n+\frac{1}{2}, j+\frac{1}{2})G_{m+n+r+\frac{3}{2},i+j+s+\frac{3}{2}},
 \end{aligned}$$

for all  $x, y \in \{L_{n,j}, G_{n+\frac{1}{2},j+\frac{1}{2}} \mid n, j \in \mathbb{Z}\}$ , it follows that

$$\begin{aligned} & [\varphi_{r+\frac{1}{2},s+\frac{1}{2}}, \psi_{m+\frac{1}{2},i+\frac{1}{2}}]([x, y]) \\ &= \varphi_{r+\frac{1}{2},s+\frac{1}{2}}(\psi_{m+\frac{1}{2},i+\frac{1}{2}}([x, y])) + \psi_{m+\frac{1}{2},i+\frac{1}{2}}(\varphi_{r+\frac{1}{2},s+\frac{1}{2}}([x, y])) \\ &= \frac{1}{2}([\varphi_{r+\frac{1}{2},s+\frac{1}{2}}, \psi_{m+\frac{1}{2},i+\frac{1}{2}}](x), y) + [x, [\varphi_{r+\frac{1}{2},s+\frac{1}{2}}, \psi_{m+\frac{1}{2},i+\frac{1}{2}}](y)]. \end{aligned}$$

Therefore, we can conclude that  $[\varphi_{r+\frac{1}{2},s+\frac{1}{2}}, \psi_{m+\frac{1}{2},i+\frac{1}{2}}]$  is an even  $\frac{1}{2}$ -superderivation of  $\mathfrak{S}(p, q)$ , which was given in [Corollary 3.19](#).

So, if  $q \neq 0$ , then we arrive at

$$(3-64) \quad [\varphi_{r+\frac{1}{2},s+\frac{1}{2}}, \psi_{m+\frac{1}{2},i+\frac{1}{2}}](L_{n,j}) = \begin{cases} 0, & (m+r+1, i+s+1) \neq (0, 0), \\ cL_{n,j}, & (m+r+1, i+s+1) = (0, 0), \end{cases}$$

$$(3-65) \quad [\varphi_{r+\frac{1}{2},s+\frac{1}{2}}, \psi_{m+\frac{1}{2},i+\frac{1}{2}}](G_{n+\frac{1}{2},j+\frac{1}{2}}) = \begin{cases} 0, & (m+r+1, i+s+1) \neq (0, 0), \\ cL_{n,j}, & (m+r+1, i+s+1) = (0, 0) \end{cases}$$

for some constant  $c \in \mathbb{C}$ .

If  $q = 0$  and  $p \notin \mathbb{Z}$ , then we get

$$(3-66) \quad [\varphi_{r+\frac{1}{2},s+\frac{1}{2}}, \psi_{m+\frac{1}{2},i+\frac{1}{2}}](L_{n,j}) = \begin{cases} 0, & (m+r+1, i+s+1) \neq (0, 0), \\ c_1L_{n,j}, & (m+r+1, i+s+1) = (0, 0), \end{cases}$$

$$(3-67) \quad [\varphi_{r+\frac{1}{2},s+\frac{1}{2}}, \psi_{m+\frac{1}{2},i+\frac{1}{2}}](G_{n+\frac{1}{2},j+\frac{1}{2}}) = \begin{cases} 0, & (m+r+1, i+s+1) \neq (0, 0), \\ c_1L_{n,j}, & (m+r+1, i+s+1) = (0, 0) \end{cases}$$

for some constant  $c_1 \in \mathbb{C}$ .

If  $q = 0$  and  $p \in \mathbb{Z}^*$ , then we have

$$(3-68) \quad [\varphi_{r+\frac{1}{2},s+\frac{1}{2}}, \psi_{m+\frac{1}{2},i+\frac{1}{2}}](L_{n,j}) = \begin{cases} 0, & (m+r+1, i+s+1) \neq (0, 0), (0, p), \\ c_2L_{n,j}, & (m+r+1, i+s+1) = (0, 0), \\ 0, & (m+r+1, i+s+1) = (0, p) \text{ and } (n, j) \neq (-2p, 0), \\ c_3L_{-p,0}, & (m+r+1, i+s+1) = (0, p) \text{ and } (n, j) = (-2p, 0), \end{cases}$$

$$(3-69) \quad [\varphi_{r+\frac{1}{2},s+\frac{1}{2}}, \psi_{m+\frac{1}{2},i+\frac{1}{2}}](G_{n+\frac{1}{2},j+\frac{1}{2}}) = \begin{cases} 0, & (m+r+1, i+s+1) \neq (0, 0), \\ c_2L_{n,j}, & (m+r+1, i+s+1) = (0, 0) \end{cases}$$

for some constants  $c_2, c_3 \in \mathbb{C}$ .

If  $p = q = 0$ , then we obtain

$$(3-70) \quad [\varphi_{r+\frac{1}{2},s+\frac{1}{2}}, \psi_{m+\frac{1}{2},i+\frac{1}{2}}](L_{n,j}) = \begin{cases} 0, & (m+r+1, i+s+1) \neq (0, 0), \\ c_4L_{n,j}, & (m+r+1, i+s+1) = (0, 0) \text{ and } (n, j) \neq (0, 0), \\ c_5L_{n,j}, & (m+r+1, i+s+1) = (0, 0) \text{ and } (n, j) = (0, 0), \end{cases}$$

$$(3-71) \quad [\varphi_{r+\frac{1}{2},s+\frac{1}{2}}, \psi_{m+\frac{1}{2},i+\frac{1}{2}}](G_{n+\frac{1}{2},j+\frac{1}{2}}) = \begin{cases} 0, & (m+r+1, i+s+1) \neq (0, 0), \\ c_4L_{n,j}, & (m+r+1, i+s+1) = (0, 0) \end{cases}$$

for some constants  $c_4, c_5 \in \mathbb{C}$ .

**3.2.1.** *The case  $p, q \neq 0$ .* By (3-62)–(3-65), we have

$$(3-72) \quad \left| \begin{array}{cc} n+p & j+q \\ m+\frac{p}{2}+\frac{1}{2} & i+\frac{q}{2}+\frac{1}{2} \end{array} \right| d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}(m+n+\frac{1}{2}, i+j+\frac{1}{2}) \\ + 2qd_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}(n, j) = 0,$$

$$(3-73) \quad 2qd_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(m+n+1, i+j+1) \\ + \left| \begin{array}{cc} n+r+p+1 & j+s+q+1 \\ m+\frac{p}{2}+\frac{1}{2} & i+\frac{q}{2}+\frac{1}{2} \end{array} \right| d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}(n+\frac{1}{2}, j+\frac{1}{2}) = 0$$

for all  $(m, i) \neq (-r-1, -s-1)$ .

**Lemma 3.21.** *Let  $r, s \in \mathbb{Z}$ , and  $\varphi \in \Delta^{\bar{1}}(\mathcal{S}(p, q))$ .*

(1) *If  $p, q \in 2\mathbb{Z} + 1$ , then*

$$\varphi_{r+\frac{1}{2}, s+\frac{1}{2}} = 0 \quad \text{for } (r, s) \neq \left(\frac{p}{2} - \frac{1}{2}, \frac{q}{2} - \frac{1}{2}\right), \\ \varphi_{\frac{p}{2}, \frac{q}{2}}(L_{m, i}) = 0 \quad \text{for all } (m, i) \in \mathbb{Z} \times \mathbb{Z}, \\ \varphi_{\frac{p}{2}, \frac{q}{2}}(G_{m+\frac{1}{2}, i+\frac{1}{2}}) = 0 \quad \text{for } (m, i) \neq \left(-\frac{3p}{2} - \frac{1}{2}, -\frac{3q}{2} - \frac{1}{2}\right).$$

(2) *Otherwise,  $\varphi_{r+\frac{1}{2}, s+\frac{1}{2}} = 0$ .*

*Proof.* Taking  $m = i = 0$  in (3-56), we obtain

$$(3-74) \quad (q(n-r) - p(j-s))d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}(n+\frac{1}{2}, j+\frac{1}{2}) = 2qd_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(0, 0).$$

Then taking  $n = r$  and  $j = s$  in (3-74), it gives  $d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(0, 0) = 0$ . Hence, (3-74) can be given by

$$(3-75) \quad (p(i-s) - q(m-r))d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}(m+\frac{1}{2}, i+\frac{1}{2}) = 0 \quad \text{for all } (r, s), (m, i) \in \mathbb{Z} \times \mathbb{Z}.$$

Taking  $n = j = 0$  in (3-72), it yields

$$(3-76) \quad \left(p\left(i+\frac{1}{2}\right) - q\left(m+\frac{1}{2}\right)\right)d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}\left(m+\frac{1}{2}, i+\frac{1}{2}\right) = 0$$

for all  $(r, s) \in \mathbb{Z} \times \mathbb{Z}$  and  $(m, i) \neq (-r-1, -s-1)$ . When  $(m, i) = (-r-1, -s-1)$ , (3-75) can be written

$$\left(p\left(s+\frac{1}{2}\right) - q\left(r+\frac{1}{2}\right)\right)d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}\left(-r-\frac{1}{2}, -s-\frac{1}{2}\right) = 0.$$

In addition, subtracting (3-75) from (3-76), we get

$$\left(p\left(s+\frac{1}{2}\right) - q\left(r+\frac{1}{2}\right)\right)d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}\left(m+\frac{1}{2}, i+\frac{1}{2}\right) = 0$$

for all  $(r, s), (m, i) \in \mathbb{Z} \times \mathbb{Z}$ . Clearly, we will analyze whether  $p\left(s+\frac{1}{2}\right) - q\left(r+\frac{1}{2}\right)$  is zero or not.

Case 1:  $p(s + \frac{1}{2}) - q(r + \frac{1}{2}) \neq 0$ . We have

$$d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}(m + \frac{1}{2}, i + \frac{1}{2}) = 0 \quad \text{for all } (m, i) \in \mathbb{Z} \times \mathbb{Z}.$$

So (3-58) becomes

$$d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(m + n + 1, i + j + 1) = 0 \quad \text{for all } (m, i), (n, j) \in \mathbb{Z} \times \mathbb{Z}.$$

Letting  $u = m + n + 1$  and  $v = i + j + 1$ , we obtain

$$d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(u, v) = 0 \quad \text{for all } (u, v) \in \mathbb{Z} \times \mathbb{Z}.$$

Case 2:  $p(s + \frac{1}{2}) - q(r + \frac{1}{2}) = 0$ . There exists some  $k \in \mathbb{C}$  satisfying  $r = kp - \frac{1}{2}$ ,  $s = kq - \frac{1}{2} \in \mathbb{Z}$ . Now taking into (3-75), we have

$$(p(i + \frac{1}{2}) - q(m + \frac{1}{2}))d_{kp, kq}^{\bar{1}}(m + \frac{1}{2}, i + \frac{1}{2}) = 0.$$

Accordingly, it follows that

$$(3-77) \quad d_{kp, kq}^{\bar{1}}(m + \frac{1}{2}, i + \frac{1}{2}) = 0 \quad \text{for } (m, i) \in \mathbb{Z} \times \mathbb{Z} \text{ and } \frac{m + \frac{1}{2}}{i + \frac{1}{2}} \neq \frac{p}{q}.$$

Letting  $u = m + n + 1$  and  $v = i + j + 1$  with  $\frac{m+\frac{1}{2}}{i+\frac{1}{2}} \neq \frac{p}{q}$  and  $\frac{n+\frac{1}{2}}{j+\frac{1}{2}} \neq \frac{p}{q}$ , we can obtain, by (3-58) and (3-77),

$$d_{kp, kq}^{\bar{0}}(u, v) = 0 \quad \text{for all } (u, v) \in \mathbb{Z} \times \mathbb{Z}.$$

In conclusion, we obtain

$$(3-78) \quad d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(u, v) = 0 \quad \text{for all } (u, v), (r, s) \in \mathbb{Z} \times \mathbb{Z}.$$

Next, we will calculate

$$d_{kp, kq}^{\bar{1}}(m + \frac{1}{2}, i + \frac{1}{2}) = 0 \quad \text{with } \frac{m + \frac{1}{2}}{i + \frac{1}{2}} = \frac{p}{q}.$$

We choose  $(m, i) \in \mathbb{Z} \times \mathbb{Z}$  with  $\frac{m+\frac{1}{2}}{i+\frac{1}{2}} = \frac{p}{q}$ , then there exists some  $l \in \mathbb{C}$  satisfying  $m = lp - \frac{1}{2}$ ,  $i = lq - \frac{1}{2} \in \mathbb{Z}$ . When  $p \neq q$ , taking  $n = lp - \frac{1}{2}$ ,  $j = lq - \frac{1}{2}$ ,  $(m, i) \in \mathbb{Z} \times \mathbb{Z}$  with  $\frac{m}{i} = \frac{p}{q}$  and  $m \neq -r - 1$  in (3-73), we have

$$(k + l + 1)(p - q)d_{kp, kq}^{\bar{1}}(lp, lq) = 0.$$

So, we get

$$d_{kp, kq}^{\bar{1}}(lp, lq) = 0 \quad \text{for } k + l + 1 \neq 0 \text{ and } p \neq q.$$

When  $p = q$ , taking  $n = lp - \frac{1}{2}$ ,  $j = lq - \frac{1}{2}$  and  $m \neq i \in \mathbb{Z}$  in (3-73), we have

$$(k + l + 1)d_{kp, kp}^{\bar{1}}(lp, lp) = 0.$$

Therefore, we can conclude

$$d_{kp, kp}^{\bar{1}}(lp, lp) = 0 \quad \text{for } k+l+1 \neq 0.$$

If  $k, l$  satisfy  $k+l+1=0$ , i.e.,  $(r, s) = (kp - \frac{1}{2}, kq - \frac{1}{2})$  and  $(m, i) = (lp - \frac{1}{2}, lq - \frac{1}{2}) = (-kp - p - \frac{1}{2}, -kq - q - \frac{1}{2})$ , then we can deduce  $p, q \in \mathbb{Z}$ . Furthermore, taking  $n = -p, m \neq -\frac{p+1}{2}, j \neq -q$  and  $i \neq -s - 1$  in (3-72), we have

$$d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}(m - p + \frac{1}{2}, i + j + \frac{1}{2}) = 0.$$

Letting  $u = m - p$  and  $v = i + j$ , we get  $d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}(u + \frac{1}{2}, v + \frac{1}{2}) = 0$  for  $u \neq -\frac{3p}{2} - \frac{1}{2}$ . And hence, we can arrive at

$$d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}(u + \frac{1}{2}, v + \frac{1}{2}) = 0$$

for  $(r, s) \neq (\frac{p}{2} - \frac{1}{2}, \frac{q}{2} - \frac{1}{2})$  and  $(u, v) \neq (-\frac{3p}{2} - \frac{1}{2}, -\frac{3q}{2} - \frac{1}{2})$ .  $\square$

According to this conclusion, we can obtain the following lemma.

**Lemma 3.22.** *Let  $p, q \in 2\mathbb{Z} + 1$ . Then the linear map  $\gamma : \mathcal{S}(p, q) \rightarrow \mathcal{S}(p, q)$  is a  $\frac{1}{2}$ -derivation of  $\mathcal{S}(p, q)$  such that*

$$\begin{aligned} \gamma(L_{m,i}) &= 0, \\ \gamma(G_{m+\frac{1}{2}, i+\frac{1}{2}}) &= \begin{cases} 0, & (m, i) \neq (-\frac{3p}{2} - \frac{1}{2}, -\frac{3q}{2} - \frac{1}{2}), \\ L_{-p, -q}, & (m, i) = (-\frac{3p}{2} - \frac{1}{2}, -\frac{3q}{2} - \frac{1}{2}) \end{cases} \end{aligned}$$

for  $(m, i) \in \mathbb{Z} \times \mathbb{Z}$ .

*Proof.* We observe that

$$\gamma = \sum_{r, s \in \mathbb{Z}} \gamma_{r+\frac{1}{2}, s+\frac{1}{2}} = \gamma_{\frac{p}{2}, \frac{q}{2}}.$$

Given Lemma 3.20, we need to check (3-56)–(3-58) for  $(r, s) = (\frac{p}{2} - \frac{1}{2}, \frac{q}{2} - \frac{1}{2})$  and

$$(3-79) \quad d_{\frac{p}{2}, \frac{q}{2}}^{\bar{0}}(m, i) = 0, \quad (m, i) \in \mathbb{Z} \times \mathbb{Z},$$

$$(3-80) \quad d_{\frac{p}{2}, \frac{q}{2}}^{\bar{1}}(m + \frac{1}{2}, i + \frac{1}{2}) = \begin{cases} 0, & (m, i) \neq (-\frac{3p}{2} - \frac{1}{2}, -\frac{3q}{2} - \frac{1}{2}), \\ 1, & (m, i) = (-\frac{3p}{2} - \frac{1}{2}, -\frac{3q}{2} - \frac{1}{2}). \end{cases}$$

Evidently, (3-56) is satisfied. The next step is to check (3-57).

Case 1:  $(n, j), (m+n, i+j) \neq (-\frac{3p}{2} - \frac{1}{2}, -\frac{3q}{2} - \frac{1}{2})$ . Clearly, both sides of (3-57) are zero by (3-80).

Case 2:  $(n, j) = (-\frac{3p}{2} - \frac{1}{2}, -\frac{3q}{2} - \frac{1}{2})$ . Then (3-57) becomes

$$(pi - qm)d_{\frac{p}{2}, \frac{q}{2}}^{\bar{1}}(m - \frac{3p}{2}, i - \frac{3q}{2}) = 0.$$

If  $(m, i) = (0, 0)$ , then it is trivially satisfied, otherwise both sides are zero by (3-80).

Case 3:  $(m + n, i + j) = (-\frac{3p}{2} - \frac{1}{2}, -\frac{3q}{2} - \frac{1}{2})$ , and  $(n, j) \neq (-\frac{3p}{2} - \frac{1}{2}, -\frac{3q}{2} - \frac{1}{2})$ . Then (3-57) becomes

$$\left( (n + \frac{p}{2} + \frac{1}{2})(-j - \frac{q}{2} - \frac{1}{2}) - (-n - \frac{p}{2} - \frac{1}{2})(j + \frac{q}{2} + \frac{1}{2}) \right) d_{\frac{p}{2}, \frac{q}{2}}^{\bar{1}} \left( -\frac{3p}{2}, -\frac{3q}{2} \right) = 0.$$

Both sides of (3-57) are zero by (3-80).

It remains to check (3-58). Due to (3-79) and (3-80), (3-58) becomes

$$\begin{aligned} & \left| \begin{array}{cc} n + \frac{p}{2} + \frac{1}{2} & j + \frac{q}{2} + \frac{1}{2} \\ m + \frac{3p}{2} + \frac{1}{2} & i + \frac{3q}{2} + \frac{1}{2} \end{array} \right| d_{\frac{p}{2}, \frac{q}{2}}^{\bar{1}} \left( m + \frac{1}{2}, i + \frac{1}{2} \right) \\ & - \left| \begin{array}{cc} n + \frac{3p}{2} + \frac{1}{2} & j + \frac{3q}{2} + \frac{1}{2} \\ m + \frac{p}{2} + \frac{1}{2} & i + \frac{q}{2} + \frac{1}{2} \end{array} \right| d_{\frac{p}{2}, \frac{q}{2}}^{\bar{1}} \left( n + \frac{1}{2}, j + \frac{1}{2} \right) = 0. \end{aligned}$$

It evidently holds by (3-80).  $\square$

With Lemma 3.22, we can directly infer the proposition below.

**Proposition 3.23.** *Let  $p, q \in \mathbb{C}^*$ . Then*

$$\Delta^{\bar{1}}(\mathcal{S}(p, q)) = \begin{cases} \langle \gamma \rangle, & p, q \in 2\mathbb{Z} + 1, \\ \{0\}, & \text{otherwise,} \end{cases}$$

where  $\gamma$  is as in Lemma 3.22.

**3.2.2. The case  $p \neq 0, q = 0$ .** In this case, (3-56) becomes

$$(3-81) \quad 2 \left| \begin{array}{cc} n+p & j \\ m+p & i \end{array} \right| d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(m+n, i+j) = \left| \begin{array}{cc} n+p & j \\ m+r+\frac{p}{2}+\frac{1}{2} & i+s+\frac{1}{2} \end{array} \right| d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(m, i) \\ + \left| \begin{array}{cc} n+r+\frac{p}{2}+\frac{1}{2} & j+s+\frac{1}{2} \\ m+p & i \end{array} \right| d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(n, j).$$

On the other hand, by using (3-62)–(3-63) and (3-66)–(3-69), respectively, we have

$$(3-82) \quad \left| \begin{array}{cc} n+p & j \\ m+\frac{p}{2}+\frac{1}{2} & i+\frac{1}{2} \end{array} \right| d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}} \left( m+n+\frac{1}{2}, i+j+\frac{1}{2} \right) = 0$$

for  $(m, i) \neq (-r-1, -s-1)$ ,  $(-p-r-1, -s-1)$ , and

$$\left| \begin{array}{cc} n+r+p+1 & j+s+1 \\ m+\frac{p}{2}+\frac{1}{2} & i+\frac{1}{2} \end{array} \right| d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}} \left( n+\frac{1}{2}, j+\frac{1}{2} \right) = 0$$

for  $(m, i) \neq (-r-1, -s-1)$ .

**Lemma 3.24.** *Let  $r, s \in \mathbb{Z}$  and  $\varphi \in \Delta^{\bar{1}}(\mathcal{S}(p, 0))$ . Then*

$$\varphi_{r+\frac{1}{2}, s+\frac{1}{2}} = 0.$$

*Proof.* Taking  $n \notin \{-p, 0\}$ ,  $j = 0$  and  $m \notin \{-p-r-1, -r-1\}$  in (3-82), we have

$$d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}\left(m+n+\frac{1}{2}, i+\frac{1}{2}\right) = 0.$$

Since any  $k \in \mathbb{Z}$  can be written as  $m+n$  with  $m \notin \{-p-r-1, -r-1\}$  and  $n \notin \{-p, 0\}$  (by choosing  $n \notin \{-p, 0, k+p+r+1, k+r+1\}$ ), we obtain

$$d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}\left(k+\frac{1}{2}, i+\frac{1}{2}\right) = 0 \quad \text{for all } (k, i) \in \mathbb{Z} \times \mathbb{Z}.$$

Choosing  $n = j = 0$  in (3-81), we obtain

$$(3-83) \quad p\left(i-s-\frac{1}{2}\right)d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(m, i) = \left(i\left(r+\frac{p}{2}+\frac{1}{2}\right) - (m+p)\left(s+\frac{1}{2}\right)\right)d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(0, 0).$$

Substituting  $j = -i \neq 0$  and  $m = n = 0$  into (3-81), it leads to

$$4pi d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(0, 0) = \left(p\left(i+s+\frac{1}{2}\right) + i\left(r+\frac{p}{2}+\frac{1}{2}\right)\right)d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(0, i) \\ + \left(p\left(i-s-\frac{1}{2}\right) + i\left(r+\frac{p}{2}+\frac{1}{2}\right)\right)d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(0, -i).$$

Given (3-83), then we get

$$4pip\left(i-s-\frac{1}{2}\right)\left(-i-s-\frac{1}{2}\right)d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(0, 0) \\ = \left(\left(p\left(i+s+\frac{1}{2}\right) + i\left(r+\frac{p}{2}+\frac{1}{2}\right)\right)\left(i\left(r+\frac{p}{2}+\frac{1}{2}\right) - p\left(s+\frac{1}{2}\right)\right)\left(-i-s-\frac{1}{2}\right)\right. \\ \left.+ \left(p\left(i-s-\frac{1}{2}\right) + i\left(r+\frac{p}{2}+\frac{1}{2}\right)\right)\left(-i\left(r+\frac{p}{2}+\frac{1}{2}\right) - p\left(s+\frac{1}{2}\right)\right)\left(i-s-\frac{1}{2}\right)\right)d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(0, 0).$$

Thus, we have

$$\left(r+\frac{5p}{2}+\frac{1}{2}\right)\left(r-\frac{p}{2}+\frac{1}{2}\right)d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(0, 0) = 0.$$

We will discuss whether  $r$  is an element of  $\left\{-\frac{5p}{2}-\frac{1}{2}, \frac{p}{2}-\frac{1}{2}\right\}$ .

Case 1:  $r \neq \left\{-\frac{5p}{2}-\frac{1}{2}, \frac{p}{2}-\frac{1}{2}\right\}$ . It follows that

$$d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(0, 0) = 0 \quad \text{for } r \notin \left\{-\frac{5p}{2}-\frac{1}{2}, \frac{p}{2}-\frac{1}{2}\right\}.$$

From (3-83), we can get

$$d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(m, i) = 0 \quad \text{for } r \notin \left\{-\frac{5p}{2}-\frac{1}{2}, \frac{p}{2}-\frac{1}{2}\right\}.$$

Case 2:  $r = -\frac{5p}{2}-\frac{1}{2}$ . Then (3-81) becomes

$$(3-84) \quad 2(i(n+p) - j(m+p))d_{-\frac{5p}{2}, s+\frac{1}{2}}^{\bar{0}}(m+n, i+j) \\ = \left((n+p)\left(i+s+\frac{1}{2}\right) - j(m-2p)\right)d_{-\frac{5p}{2}, s+\frac{1}{2}}^{\bar{0}}(m, i) \\ + \left(i(n-2p) - (m+p)\left(j+s+\frac{1}{2}\right)\right)d_{-\frac{5p}{2}, s+\frac{1}{2}}^{\bar{0}}(n, j).$$

And (3-83) becomes

$$(3-85) \quad p\left(i - s - \frac{1}{2}\right)d_{-\frac{5p}{2}, s+\frac{1}{2}}^{\bar{0}}(m, i) = -(2pi + (m + p)\left(s + \frac{1}{2}\right))d_{-\frac{5p}{2}, s+\frac{1}{2}}^{\bar{0}}(0, 0).$$

Taking  $m = n = 0$  and  $i, j, i \pm j \neq 0$  in (3-84), we have

$$2(i - j)d_{-\frac{5p}{2}, s+\frac{1}{2}}^{\bar{0}}(0, i + j) = \left(i + 2j + s + \frac{1}{2}\right)d_{-\frac{5p}{2}, s+\frac{1}{2}}^{\bar{0}}(0, i) \\ - \left(2i + j + s + \frac{1}{2}\right)d_{-\frac{5p}{2}, s+\frac{1}{2}}^{\bar{0}}(0, j).$$

Because of (3-85), we can obtain

$$d_{-\frac{5p}{2}, s+\frac{1}{2}}^{\bar{0}}(0, 0) = 0.$$

Then we use (3-85), it follows that

$$d_{-\frac{5p}{2}, s+\frac{1}{2}}^{\bar{0}}(m, i) = 0 \quad \text{for all } (m, i) \in \mathbb{Z} \times \mathbb{Z}.$$

Case 3:  $r = \frac{p}{2} - \frac{1}{2}$ . In this case, (3-81) becomes

$$(3-86) \quad 2(i(n + p) - j(m + p))d_{\frac{p}{2}, s+\frac{1}{2}}^{\bar{0}}(m + n, i + j) \\ = \left((n + p)\left(i + s + \frac{1}{2}\right) - j(m + p)\right)d_{\frac{p}{2}, s+\frac{1}{2}}^{\bar{0}}(m, i) \\ + \left(i(n + p) - (m + p)\left(j + s + \frac{1}{2}\right)\right)d_{\frac{p}{2}, s+\frac{1}{2}}^{\bar{0}}(n, j).$$

And (3-83) can be expressed as

$$(3-87) \quad p\left(i - s - \frac{1}{2}\right)d_{\frac{p}{2}, s+\frac{1}{2}}^{\bar{0}}(m, i) = \left(pi - (m + p)\left(s + \frac{1}{2}\right)\right)d_{\frac{p}{2}, s+\frac{1}{2}}^{\bar{0}}(0, 0).$$

Choosing  $m = 0$  in (3-87) and utilizing  $i - s - \frac{1}{2} \neq 0$ , we can see

$$(3-88) \quad d_{\frac{p}{2}, s+\frac{1}{2}}^{\bar{0}}(0, i) = d_{\frac{p}{2}, s+\frac{1}{2}}^{\bar{0}}(0, 0) \quad \text{for all } i \in \mathbb{Z}.$$

Substituting  $m = j = 0$  in (3-86), we have

$$(3-89) \quad 2i(n + p)d_{\frac{p}{2}, s+\frac{1}{2}}^{\bar{0}}(n, i) = (n + p)\left(i + s + \frac{1}{2}\right)d_{\frac{p}{2}, s+\frac{1}{2}}^{\bar{0}}(0, i) \\ + \left(i(n + p) - p\left(s + \frac{1}{2}\right)\right)d_{\frac{p}{2}, s+\frac{1}{2}}^{\bar{0}}(n, 0).$$

Furthermore, taking  $i = 0$ , it follows that

$$(3-90) \quad pd_{\frac{p}{2}, s+\frac{1}{2}}^{\bar{0}}(n, 0) = (n + p)d_{\frac{p}{2}, s+\frac{1}{2}}^{\bar{0}}(0, 0).$$

Multiplying (3-89) by  $p$  and using (3-88) and (3-90), we get

$$2pi(n+p)d_{\frac{p}{2},s+\frac{1}{2}}^{\bar{0}}(n,i) = p(n+p)(i+s+\frac{1}{2})d_{\frac{p}{2},s+\frac{1}{2}}^{\bar{0}}(0,0) \\ + (n+p)(i(n+p) - p(s+\frac{1}{2}))d_{\frac{p}{2},s+\frac{1}{2}}^{\bar{0}}(0,0).$$

Additionally, we arrive at

$$2pid_{\frac{p}{2},s+\frac{1}{2}}^{\bar{0}}(n,i) = i(2p+n)d_{\frac{p}{2},s+\frac{1}{2}}^{\bar{0}}(0,0) \quad \text{for } n \neq -p.$$

And then, we can deduce that

$$(3-91) \quad 2pd_{\frac{p}{2},s+\frac{1}{2}}^{\bar{0}}(n,i) = (2p+n)d_{\frac{p}{2},s+\frac{1}{2}}^{\bar{0}}(0,0) \quad \text{for } n \neq -p \text{ and } i \neq 0.$$

Taking  $n \notin \{-p, 0\}$  and  $i \neq 0$  and combining (3-87) and (3-91), we have

$$d_{\frac{p}{2},s+\frac{1}{2}}^{\bar{0}}(0,0) = 0.$$

Due to (3-87), we obtain

$$d_{\frac{p}{2},s+\frac{1}{2}}^{\bar{0}}(m,i) = 0$$

for all  $(m, i) \in \mathbb{Z} \times \mathbb{Z}$ . □

By Lemma 3.24, we directly obtain the following proposition.

**Proposition 3.25.** *Let  $p \in \mathbb{C}^*$  and  $q = 0$ . Then*

$$\Delta^{\bar{1}}(\mathcal{S}(p, 0)) = 0.$$

**3.2.3. The case  $p = 0, q \neq 0$ .** By (3-62)–(3-63) and (3-64)–(3-65), respectively, we have

$$(3-92) \quad \left| \begin{array}{cc} n & j+q \\ m+\frac{1}{2} & i+\frac{q}{2}+\frac{1}{2} \end{array} \right| d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{1}}(m+n+\frac{1}{2}, i+j+\frac{1}{2}) + 2qd_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{1}}(n, j) = 0,$$

$$2qd_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(m+n+1, i+j+1) + \left| \begin{array}{cc} n+r+1 & j+s+q+1 \\ m+\frac{1}{2} & i+\frac{q}{2}+\frac{1}{2} \end{array} \right| d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{1}}(n+\frac{1}{2}, j+\frac{1}{2}) = 0$$

for all  $(m, i) \neq (-r-1, -s-1)$ .

**Lemma 3.26.** *Let  $r, s \in \mathbb{Z}$  and  $\varphi \in \Delta^{\bar{1}}(\mathcal{S}(0, q))$ . Then*

$$\varphi_{r+\frac{1}{2},s+\frac{1}{2}} = 0.$$

*Proof.* Applying the proof of Lemma 3.24 concerning  $d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(m+\frac{1}{2}, i+\frac{1}{2})$ , we can conclude that

$$d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(m, i) = 0 \quad \text{for all } (m, i) \in \mathbb{Z} \times \mathbb{Z}.$$

Then (3-92) becomes

$$\left| \begin{array}{cc} n & j+q \\ m+\frac{1}{2} & i+\frac{q}{2}+\frac{1}{2} \end{array} \right| d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{1}}(m+n+\frac{1}{2}, i+j+\frac{1}{2}) = 0.$$

Taking  $n = 0$ ,  $j \neq -q$  and  $i \neq -s - 1$ , we have

$$d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}\left(m + \frac{1}{2}, i + j + \frac{1}{2}\right) = 0.$$

Since any  $k \in \mathbb{Z}$  can be written as  $i + j$  with  $i \neq -s - 1$  and  $j \neq -q$ , by choosing  $j \notin \{-q, k + s + 1\}$ , we obtain

$$d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}\left(m + \frac{1}{2}, k + \frac{1}{2}\right) = 0$$

for all  $(m, k) \in \mathbb{Z} \times \mathbb{Z}$ . □

A direct result is the subsequent proposition.

**Proposition 3.27.** *Let  $p = 0$  and  $q \in \mathbb{C}^*$ . Then*

$$\Delta^{\bar{1}}(\mathcal{S}(p, 0)) = 0.$$

**3.2.4. The case  $p = q = 0$ .** In this case, (3-56) becomes

$$(3-93) \quad 2 \begin{vmatrix} n & j \\ m & i \end{vmatrix} d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(m+n, i+j) = \begin{vmatrix} n & j \\ m+r+\frac{1}{2} & i+s+\frac{1}{2} \end{vmatrix} d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(m, i) \\ + \begin{vmatrix} n+r+\frac{1}{2} & j+s+\frac{1}{2} \\ m & i \end{vmatrix} d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(n, j).$$

On the other hand, by using (3-62)–(3-63) and (3-70)–(3-71), respectively, we have

$$(3-94) \quad \begin{vmatrix} n & j \\ m+\frac{1}{2} & i+\frac{1}{2} \end{vmatrix} d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}\left(m+n+\frac{1}{2}, i+j+\frac{1}{2}\right) = 0, \quad (m, i) \neq (-r-1, -s-1),$$

$$(3-95) \quad \begin{vmatrix} n+r+1 & j+s+1 \\ m+\frac{1}{2} & i+\frac{1}{2} \end{vmatrix} d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}\left(n+\frac{1}{2}, j+\frac{1}{2}\right) = 0, \quad (m, i) \neq (-r-1, -s-1).$$

**Lemma 3.28.** *Let  $r, s \in \mathbb{Z}$  and  $\varphi \in \Delta^{\bar{1}}(\mathcal{S}(0, 0))$ . Then*

$$\varphi_{r+\frac{1}{2}, s+\frac{1}{2}} = 0.$$

*Proof.* If  $j \neq -s - 1$ , then we take  $m \notin \left\{-r - 1, \frac{(n+r+1)(i+\frac{1}{2})}{j+s+1} - \frac{1}{2}\right\}$  in (3-95). It follows that

$$d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}\left(n + \frac{1}{2}, j + \frac{1}{2}\right) = 0 \quad \text{for all } n \in \mathbb{Z} \text{ and } j \neq -s - 1.$$

If  $n \neq -r - 1$ , then we choose  $i \notin \left\{-s - 1, \frac{(j+s+1)(m+\frac{1}{2})}{n+r+1} - \frac{1}{2}\right\}$  in (3-95). We can arrive at

$$d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}\left(n + \frac{1}{2}, j + \frac{1}{2}\right) = 0 \quad \text{for all } j \in \mathbb{Z} \text{ and } n \neq -r - 1.$$

Hence, we can conclude

$$d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}\left(n + \frac{1}{2}, j + \frac{1}{2}\right) = 0 \quad \text{for } (n, j) \neq (-r - 1, -s - 1).$$

Substituting  $m = -n - r - 1$ ,  $j = -i - s - 1$  and  $i \neq -s - 1$  into (3-94), it leads to

$$\begin{vmatrix} -r - \frac{1}{2} & -s - \frac{1}{2} \\ -n - r - \frac{1}{2} & i + \frac{1}{2} \end{vmatrix} d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}(-r - \frac{1}{2}, -s - \frac{1}{2}) = 0.$$

Since  $r \in \mathbb{Z}$  and  $r + \frac{1}{2} \neq 0$ , then we take  $i \notin \{-\frac{n(s+\frac{1}{2})}{r+\frac{1}{2}} - s - 1, -s - 1\}$ . It follows that

$$d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}(-r - \frac{1}{2}, -s - \frac{1}{2}) = 0.$$

Therefore, we arrive at

$$d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{1}}(n + \frac{1}{2}, j + \frac{1}{2}) = 0 \quad \text{for all } (n, j), (r, s) \in \mathbb{Z} \times \mathbb{Z}.$$

Choosing  $n = j = 0$  in (3-93), we have

$$(i(r + \frac{1}{2}) - m(s + \frac{1}{2}))d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(0, 0) = 0.$$

Since  $s + \frac{1}{2} \neq 0$ , then taking  $m \neq \frac{i(r+\frac{1}{2})}{s+\frac{1}{2}}$ , we have

$$d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(0, 0) = 0.$$

Substituting  $m = j = 0$  into (3-93), we obtain

$$(3-96) \quad 2ni d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(n, i) = n(i + s + \frac{1}{2})d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(0, i) + i(n + r + \frac{1}{2})d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(n, 0).$$

Putting  $m = -n$  and  $j = 0$  into (3-93), we get

$$(3-97) \quad 2ni d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(0, i) = n(i + s + \frac{1}{2})d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(-n, i) \\ + (i(n + r + \frac{1}{2}) + n(s + \frac{1}{2}))d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(n, 0).$$

Taking  $i = 0$  in (3-97), we can conclude

$$(3-98) \quad d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(-n, 0) = -d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(n, 0) \quad \text{for all } n \in \mathbb{Z}.$$

Substituting  $j = -i$  and  $m = 0$  into (3-93), it follows that

$$(3-99) \quad 2ni d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(n, 0) = (n(i + s + \frac{1}{2}) + i(r + \frac{1}{2}))d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(0, i) \\ + i(n + r + \frac{1}{2})d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(n, -i).$$

Letting  $n = 0$  in (3-99), we have

$$(3-100) \quad d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(0, -i) = -d_{r+\frac{1}{2}, s+\frac{1}{2}}^{\bar{0}}(0, i) \quad \text{for all } i \in \mathbb{Z}.$$

Choosing  $n = -(2r + 1)$  in (3-96), we have

$$(3-101) \quad 4id_{r+\frac{1}{2},s+\frac{1}{2}}^0(-2r-1, i) \\ = (2i + 2s + 1)d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(0, i) + id_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(-2r-1, 0).$$

Then we let  $n = 2r + 1$  in (3-97) together with (3-98). It follows that

$$(3-102) \quad 4id_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(0, i) = (2i + 2s + 1)d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(-2r-1, i) \\ + (3i + 2s + 1)d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(2r+1, 0), \\ 4id_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(0, i) = (2i + 2s + 1)d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(-2r-1, i) \\ - (3i + 2s + 1)d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(-2r-1, 0).$$

Replacing  $i$  by  $-i$  and taking  $n = -2r - 1$  in (3-99) together with (3-100), it gives

$$(3-103) \quad 4id_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(-2r-1, 0) \\ = -(2s - i + 1)d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(0, -i) + id_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(-2r-1, i), \\ 4id_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(-2r-1, 0) \\ = (2s - i + 1)d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(0, i) + id_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(-2r-1, i).$$

Combining (3-101) and (3-103), we can deduce

$$(3-104) \quad 15id_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(-2r-1, 0) = (10s - 2i + 5)d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(0, i),$$

$$(3-105) \quad 15id_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(-2r-1, i) = (10s + 7i + 5)d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(0, i).$$

If  $i \neq 0$ , taking (3-104) and (3-105) in (3-102), we have

$$(10i - 2s - 1)d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(0, i) = 0.$$

Since  $s \in \mathbb{Z}$ , then we obtain  $\frac{1}{5}(s + \frac{1}{2}) \notin \mathbb{Z}$  and  $i \neq \frac{1}{5}(s + \frac{1}{2})$ , i.e.,  $10i - 2s - 1 \neq 0$ . Hence, we arrive at

$$d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(0, i) = 0 \quad \text{for all } i \in \mathbb{Z}.$$

We take  $i = -2s - 1$  in (3-96). It follows that

$$(3-106) \quad 4nd_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(n, -2s-1) \\ = nd_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(0, -2s-1) + (2n + 2r + 1)d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(n, 0).$$

Replacing  $n$  by  $-n$  and taking  $i = -2s - 1$  in (3-97) together with (3-98), it gives

$$(3-107) \quad 4nd_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(0, -2s - 1) \\ = nd_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(n, -2s - 1) + (2r - n + 1)d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(n, 0).$$

Letting  $i = 2s + 1$  in (3-99), we obtain

$$(3-108) \quad 4nd_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(n, 0) = (2n + 2r + 1)d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(n, -2s - 1) \\ - (3n + 2r + 1)d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(0, -2s - 1).$$

Combining (3-106)–(3-108), we have

$$(10n - 2r - 1)d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(n, 0) = 0.$$

As the similar reason above, we can get

$$d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(n, 0) = 0 \quad \text{for all } n \in \mathbb{Z}.$$

In this end, by (3-96), we obtain

$$d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(n, i) = 0 \quad \text{for } n, i \neq 0.$$

Therefore, we can conclude

$$d_{r+\frac{1}{2},s+\frac{1}{2}}^{\bar{0}}(n, i) = 0$$

for  $(n, j) \in \mathbb{Z} \times \mathbb{Z}$ . □

It leads directly to the following proposition.

**Proposition 3.29.** *Let  $p = q = 0$ . Then*

$$\Delta^{\bar{1}}(\mathcal{S}(0, 0)) = 0.$$

By integrating Propositions 3.23, 3.25, 3.27 and 3.29, we can deduce the following corollary.

**Corollary 3.30.** *Let  $p, q \in \mathbb{C}$ . Then*

$$\Delta^{\bar{1}}(\mathcal{S}(p, q)) = \begin{cases} \langle \gamma \rangle, & p, q \in 2\mathbb{Z} + 1, \\ \{0\}, & \text{otherwise,} \end{cases}$$

where  $\gamma$  is as in Lemma 3.22.

#### 4. Transposed Poisson superalgebra structures on $\mathcal{S}(p, q)$

**Theorem 4.1.** *Let  $p$  and  $q$  be fixed complex numbers.*

- (1) *If  $p \notin \mathbb{Z}$  or  $q \neq 0$ , then all the transposed Poisson superalgebra structures on  $(\mathcal{S}(p, q), [\cdot, \cdot])$  are trivial.*
- (2) *If  $p \in \mathbb{Z}$  and  $q = 0$ , then the nontrivial transposed Poisson superalgebra structure  $(\mathcal{S}(p, 0), \cdot, [\cdot, \cdot])$  on  $(\mathcal{S}(p, 0), [\cdot, \cdot])$  is, up to an isomorphism,*

$$(4-1) \quad L_{-2p,0} \cdot L_{-2p,0} = L_{-p,0}.$$

*Proof.* Let  $(\mathcal{S}(p, q), \cdot, [\cdot, \cdot])$  be a transposed Poisson superalgebra, i.e.,  $(\mathcal{S}(p, q), \cdot)$  is supercommutative and (2-2) holds. Given  $(m, i) \in \mathbb{Z} \times \mathbb{Z}$ , we denote by  $\varphi^{m,i}$  and  $\psi^{m+\frac{1}{2},i+\frac{1}{2}}$  the left multiplication by  $L_{m,i}$  and  $G_{m+\frac{1}{2},i+\frac{1}{2}}$ , respectively, in  $(\mathcal{S}(p, q), \cdot)$ , that is,

$$(4-2) \quad \begin{aligned} L_{m,i} \cdot L_{n,j} &= \varphi^{m,i}(L_{n,j}), \\ L_{m,i} \cdot G_{n+\frac{1}{2},j+\frac{1}{2}} &= \varphi^{m,i}(G_{n+\frac{1}{2},j+\frac{1}{2}}), \\ G_{m+\frac{1}{2},i+\frac{1}{2}} \cdot L_{n,j} &= \psi^{m+\frac{1}{2},i+\frac{1}{2}}(L_{n,j}), \\ (4-3) \quad G_{m+\frac{1}{2},i+\frac{1}{2}} \cdot G_{n+\frac{1}{2},j+\frac{1}{2}} &= \psi^{m+\frac{1}{2},i+\frac{1}{2}}(G_{n+\frac{1}{2},j+\frac{1}{2}}). \end{aligned}$$

In view of supercommutativity of  $(\mathcal{S}(p, q), \cdot)$ , we have

$$(4-4) \quad \begin{aligned} L_{m,i} \cdot L_{n,j} &= \varphi^{n,j}(L_{m,i}), \\ L_{m,i} \cdot G_{n+\frac{1}{2},j+\frac{1}{2}} &= \psi^{n+\frac{1}{2},j+\frac{1}{2}}(L_{m,i}), \\ G_{m+\frac{1}{2},i+\frac{1}{2}} \cdot L_{n,j} &= \varphi^{n,j}(G_{m+\frac{1}{2},i+\frac{1}{2}}), \\ (4-5) \quad G_{m+\frac{1}{2},i+\frac{1}{2}} \cdot G_{n+\frac{1}{2},j+\frac{1}{2}} &= -\psi^{n+\frac{1}{2},j+\frac{1}{2}}(G_{m+\frac{1}{2},i+\frac{1}{2}}). \end{aligned}$$

By using (2-2), we have  $\varphi^{m,i} \in \Delta^{\bar{0}}(\mathcal{S}(p, q))$  and  $\psi^{m+\frac{1}{2},i+\frac{1}{2}} \in \Delta^{\bar{1}}(\mathcal{S}(p, q))$ . Furthermore, based on the Corollaries 3.19 and 3.30, we proceed to the following discussion.

Case 1:  $p, q \neq 0$  and  $p, q \notin 2\mathbb{Z} + 1$ . It is clear that  $\varphi^{m,i} = a^{m,i}id$  for some  $a^{m,i} \in \mathbb{C}$  by Proposition 3.4 and  $\psi^{m+\frac{1}{2},i+\frac{1}{2}} = 0$  by Proposition 3.23. It follows from (4-2) and (4-4) that  $a^{m,i} = 0$  for all  $(m, i) \in \mathbb{Z} \times \mathbb{Z}$ . So,  $\cdot$  is trivial whenever  $p, q \neq 0$  and  $p, q \notin 2\mathbb{Z} + 1$ .

Case 2:  $p, q \neq 0$  and  $p, q \in 2\mathbb{Z} + 1$ . We have  $\varphi^{m,i} = a^{m,i}id$  for some  $a^{m,i} \in \mathbb{C}$  by Proposition 3.4. Equations (4-2) and (4-4) imply  $a^{m,i} = 0$ , whence  $\varphi^{m,i} = 0$ . We arrive at  $L_{m,i} \cdot L_{n,j} = L_{m,i} \cdot G_{n,j} = 0$ . From Proposition 3.23, it is easy to observe that

$$\psi^{m+\frac{1}{2},i+\frac{1}{2}} = b^{m+\frac{1}{2},i+\frac{1}{2}}\gamma,$$

with  $b^{m+\frac{1}{2}, i+\frac{1}{2}} \in \mathbb{C}$ . On the one hand, it follows from (3-80) and (4-3) that

$$\begin{aligned} G_{m+\frac{1}{2}, i+\frac{1}{2}} \cdot G_{n+\frac{1}{2}, j+\frac{1}{2}} &= b^{m+\frac{1}{2}, i+\frac{1}{2}} \gamma(G_{n+\frac{1}{2}, j+\frac{1}{2}}) \\ &= \begin{cases} 0, & (n, j) \neq \left(-\frac{3p}{2} - \frac{1}{2}, -\frac{3q}{2} - \frac{1}{2}\right), \\ b^{m+\frac{1}{2}, i+\frac{1}{2}} L_{-p, -q}, & (n, j) = \left(-\frac{3p}{2} - \frac{1}{2}, -\frac{3q}{2} - \frac{1}{2}\right). \end{cases} \end{aligned}$$

On the other hand, by using (3-80) and (4-5), we arrive at

$$\begin{aligned} G_{m+\frac{1}{2}, i+\frac{1}{2}} \cdot G_{n+\frac{1}{2}, j+\frac{1}{2}} &= -b^{n+\frac{1}{2}, j+\frac{1}{2}} \gamma(G_{m+\frac{1}{2}, i+\frac{1}{2}}) \\ &= \begin{cases} 0, & (m, i) \neq \left(-\frac{3p}{2} - \frac{1}{2}, -\frac{3q}{2} - \frac{1}{2}\right), \\ -b^{n+\frac{1}{2}, j+\frac{1}{2}} L_{-p, -q}, & (m, i) = \left(-\frac{3p}{2} - \frac{1}{2}, -\frac{3q}{2} - \frac{1}{2}\right). \end{cases} \end{aligned}$$

Thus, the product  $G_{m+\frac{1}{2}, i+\frac{1}{2}} \cdot G_{n+\frac{1}{2}, j+\frac{1}{2}}$  is zero unless

$$(m, i) = (n, j) = \left(-\frac{3p}{2} - \frac{1}{2}, -\frac{3q}{2} - \frac{1}{2}\right).$$

But  $G_{-\frac{3p}{2}, -\frac{3q}{2}} \cdot G_{-\frac{3p}{2}, -\frac{3q}{2}} = -G_{-\frac{3p}{2}, -\frac{3q}{2}} \cdot G_{-\frac{3p}{2}, -\frac{3q}{2}}$ , then  $G_{-\frac{3p}{2}, -\frac{3q}{2}} \cdot G_{-\frac{3p}{2}, -\frac{3q}{2}} = 0$ . So,  $\cdot$  is trivial.

Case 3:  $p \notin \mathbb{Z}$  and  $q = 0$ . We obtain that  $\varphi^{m,i} = a^{m,i} id$  for some  $a^{m,i} \in \mathbb{C}$  by Proposition 3.14 and  $\psi^{m+\frac{1}{2}, i+\frac{1}{2}} = 0$  by Proposition 3.25. It follows from (4-2) and (4-4) that  $a^{m,i} = 0$  for all  $(m, i) \in \mathbb{Z} \times \mathbb{Z}$ . So,  $\cdot$  is trivial.

Case 4:  $p \in \mathbb{Z}^*$  and  $q = 0$ . We have  $\psi^{m+\frac{1}{2}, i+\frac{1}{2}} = 0$  by Proposition 3.25. It leads to  $G_{m+\frac{1}{2}, i+\frac{1}{2}} \cdot L_{n,j} = 0$  and  $G_{m+\frac{1}{2}, i+\frac{1}{2}} \cdot G_{n+\frac{1}{2}, j+\frac{1}{2}} = 0$ .

Because of Proposition 3.14, it is clear to show that  $\varphi^{m,i} = f^{m,i} id + g^{m,i} \alpha$  for  $f^{m,i}, g^{m,i} \in \mathbb{C}$ . We can get from (3-36) and (4-2) that

$$\begin{aligned} L_{m,i} \cdot L_{n,j} &= f^{m,i} L_{n,j} + g^{m,i} \alpha(L_{n,j}) \\ &= \begin{cases} f^{m,i} L_{n,j}, & (n, j) \neq (-2p, 0), \\ f^{m,i} L_{-2p,0} + g^{m,i} L_{-p,0}, & (n, j) = (-2p, 0). \end{cases} \end{aligned}$$

On the other hand, by using (3-36) and (4-4), we have

$$\begin{aligned} L_{m,i} \cdot L_{n,j} &= f^{m,i} L_{n,j} + g^{m,i} \alpha(L_{n,j}) \\ &= \begin{cases} f^{n,j} L_{m,i}, & (m, i) \neq (-2p, 0), \\ f^{n,j} L_{-2p,0} + g^{n,j} L_{-p,0}, & (m, i) = (-2p, 0). \end{cases} \end{aligned}$$

Thus, we can discuss it into the following cases.

Subcase 1:  $(m, i), (n, j) \neq (-2p, 0)$ . We get  $f^{m,i} L_{n,j} = f^{n,j} L_{m,i}$ . So taking  $(m, i) \neq (n, j)$  we conclude that  $f^{m,i} = f^{n,j} = 0$ . Thus,  $L_{m,i} \cdot L_{n,j} = 0$ .

Subcase 2:  $(m, i) = (-2p, 0), (n, j) \neq (-2p, 0)$ . We have

$$f^{-2p,0} L_{n,j} = f^{n,j} L_{-2p,0} + g^{n,j} L_{-p,0} = g^{n,j} L_{-p,0},$$

because of  $f^{n,j} = 0$  for  $(n, j) \neq (-2p, 0)$ . So, taking  $(n, j) \neq (-p, 0)$ , we conclude that  $f^{-2p,0} = 0$ , whence  $L_{-2p,0} \cdot L_{n,j} = 0$ .

Subcase 3:  $(m, i) \neq (-2p, 0)$ ,  $(n, j) = (-2p, 0)$ . It implies  $L_{m,i} \cdot L_{-2p,0} = 0$ .

Subcase 4:  $(m, i) = (n, j) = (-2p, 0)$ . It leads to

$$L_{-2p,0} \cdot L_{-2p,0} = f^{-2p,0} L_{-2p,0} + g^{-2p,0} L_{-p,0} = g^{-2p,0} L_{-p,0},$$

because of  $f^{-2p,0} = 0$ .

Thus, we can conclude that

$$L_{m,i} \cdot L_{n,j} = \begin{cases} 0, & (m, i), (n, j) \neq (-2p, 0), \\ g^{-2p,0} L_{-p,0}, & (m, i) = (n, j) = (-2p, 0). \end{cases}$$

Therefore, the product  $(\mathcal{S}(p, 0), \cdot)$  is of the form

$$(4-6) \quad L_{-2p,0} \cdot L_{-2p,0} = c L_{-p,0},$$

where  $c \in \mathbb{C}$ . Assume that  $c \neq 0$ , otherwise the transposed Poisson superalgebra structure is trivial. We observe that  $L_{-p,0} \in Z(\mathcal{S}(p, 0))$ , where  $Z(\mathcal{S}(p, 0))$  is the center of the Lie superalgebra  $(\mathcal{S}(p, 0), [\cdot, \cdot])$ . Indeed,

$$[L_{m,i}, L_{-p,0}] = ((-p + p) \cdot 0 - (m + p) \cdot 0) L_{m+n,i+j} = 0$$

for all  $(m, i) \in \mathbb{Z} \times \mathbb{Z}$ . Hence the linear map  $\phi$  such that  $\phi(L_{m,i}) = L_{m,i}$  for  $(m, i) \neq (-p, 0)$  and  $\phi(L_{-p,0}) = k L_{-p,0}$  is an automorphism of  $(\mathcal{S}(p, 0), [\cdot, \cdot])$  for any  $k \in \mathbb{C}^*$ . If  $p \neq 0$ , then we take  $k = c^{-1}$ . Furthermore, we obtain an isomorphic transposed Poisson superalgebra structure  $*$  on  $(\mathcal{S}(p, 0), [\cdot, \cdot])$  in which the only nonzero product is

$$\begin{aligned} L_{-2p,0} * L_{-2p,0} &= \phi(L_{-2p,0}) * \phi(L_{-2p,0}) = \phi(L_{-2p,0} \cdot L_{-2p,0}) \\ &= \phi(c L_{-2p,0}) = c^{-1} \cdot c L_{-p,0} = L_{-p,0}. \end{aligned}$$

So, up to an isomorphism, we may consider  $c = 1$  in (4-6).

Case 5:  $p = q = 0$ . We have  $\psi^{m+\frac{1}{2}, i+\frac{1}{2}} = 0$  by [Proposition 3.29](#). It implies

$$G_{m+\frac{1}{2}, i+\frac{1}{2}} \cdot L_{n,j} = 0 \quad \text{and} \quad G_{m+\frac{1}{2}, i+\frac{1}{2}} \cdot G_{n+\frac{1}{2}, j+\frac{1}{2}} = 0.$$

It is easy to show that  $\varphi^{m,i} = x^{m,i} id + y^{m,i} \beta$  for  $x^{m,i}, y^{m,i} \in \mathbb{C}$  by [Proposition 3.18](#). We can get from (3-52) and (4-2) that

$$L_{m,i} \cdot L_{n,j} = x^{m,i} L_{n,j} + y^{m,i} \alpha(L_{n,j}) = \begin{cases} x^{m,i} L_{n,j}, & (n, j) \neq (0, 0), \\ (x^{m,i} + y^{m,i}) L_{0,0}, & (n, j) = (0, 0). \end{cases}$$

On the other hand, by using (3-52) and (4-4), we obtain

$$L_{m,i} \cdot L_{n,j} = x^{m,i} L_{n,j} + y^{m,i} \alpha(L_{n,j}) = \begin{cases} x^{n,i} L_{m,i}, & (m, i) \neq (0, 0), \\ (x^{n,j} + y^{n,j}) L_{0,0}, & (m, i) = (0, 0). \end{cases}$$

Thus, we can discuss it into the following cases.

Subcase 1:  $(m, i), (n, j) \neq (0, 0)$ . We get  $x^{m,i}L_{n,j} = x^{n,j}L_{m,i}$ . So taking  $(m, i) \neq (n, j)$  we conclude that  $x^{m,i} = x^{n,j} = 0$ . Thus, we arrive at  $L_{m,i} \cdot L_{n,j} = 0$ .

Subcase 2:  $(m, i) = (0, 0), (n, j) \neq (0, 0)$ . We have  $x^{0,0}L_{n,j} = (x^{n,j} + y^{n,j})L_{0,0} = y^{n,j}L_{0,0}$ . So, we obtain  $x^{0,0} = y^{n,j} = 0$ , whence  $L_{0,0} \cdot L_{n,j} = 0$ .

Subcase 3:  $(m, i) \neq (0, 0), (n, j) = (0, 0)$ . It implies  $L_{m,i} \cdot L_{0,0} = 0$ .

Subcase 4:  $(m, i) = (n, j) = (0, 0)$ . It leads to  $L_{0,0} \cdot L_{0,0} = (x^{0,0} + y^{0,0})L_{0,0} = y^{0,0}L_{0,0}$ , because of  $x^{0,0} = 0$ .

Thus, we have

$$L_{m,i} \cdot L_{n,j} = \begin{cases} 0, & (m, i), (n, j) \neq (0, 0), \\ y^{0,0}L_{0,0}, & (m, i), (n, j) = (0, 0). \end{cases}$$

Therefore, the product  $(\mathcal{S}(0, 0), \cdot)$  is of the form

$$(4-7) \quad L_{0,0} \cdot L_{0,0} = cL_{0,0},$$

where  $c \in \mathbb{C}$ . Assume that  $c \neq 0$ , otherwise the transposed Poisson superalgebra structure is trivial. Observe that  $L_{0,0} \in Z(\mathcal{S}(0, 0))$ , where  $Z(\mathcal{S}(0, 0))$  is the center of the Lie superalgebra  $(\mathcal{S}(0, 0), [\cdot, \cdot])$ . Indeed,

$$[L_{m,i}, L_{0,0}] = 0$$

for all  $(m, i) \in \mathbb{Z} \times \mathbb{Z}$ . Hence the linear map  $\phi$  such that  $\phi(L_{m,i}) = L_{m,i}$  for  $(m, i) \neq (0, 0)$  and  $\phi(L_{0,0}) = kL_{-p,0}$  is an automorphism of  $(\mathcal{S}(0, 0), [\cdot, \cdot])$  for any  $k \in \mathbb{C}^*$ . Then taking  $k = c$ , we obtain an isomorphic transposed Poisson superalgebra structure  $*$  on  $(\mathcal{S}(0, 0), [\cdot, \cdot])$  in which the only nonzero product is

$$\begin{aligned} L_{0,0} * L_{0,0} &= c^{-1}\phi(L_{0,0}) * c^{-1}\phi(L_{0,0}) = c^{-2}\phi(L_{0,0} \cdot L_{0,0}) \\ &= c^{-2}\phi(cL_{0,0}) = c^{-2} \cdot c^2L_{0,0} = L_{0,0}. \end{aligned}$$

So, up to an isomorphism, we may consider  $c = 1$  in (4-7).

Conversely, each of two associative and supercommutative multiplication (4-1) defines a transposed Poisson superalgebra structure on  $\mathcal{S}(p, 0)$ ,  $p \in \mathbb{Z}$ . If  $p \in \mathbb{Z}^*$ , we can observe that  $\mathcal{S}(p, 0) \cdot \mathcal{S}(p, 0) \subseteq \langle L_{-p,0} \rangle \subseteq \mathbb{Z}(\mathcal{S}(p, 0))$ . Hence, the right-hand side of (2-2) is always zero. In fact, the left-hand side of (2-2) is zero as well, because of  $[\mathcal{S}(p, 0), \mathcal{S}(p, 0)] \subseteq \text{Ann}(\mathcal{S}(p, 0))$ , where  $\text{Ann}(\mathcal{S}(p, 0))$  is the annihilator of  $(\mathcal{S}(p, 0), \cdot)$ . Assuming  $[L_{m,i}, L_{n,j}] \in \langle L_{-2p,0} \rangle$ , we obtain from (2-1) that  $m + n = -2p$  and  $i + j = 0$ . But it leads to

$$-j(n + p) - j(m + p) = -j(n + p) - j(-2p - n + p) = 0,$$

so we have  $[L_{m,i}, L_{n,j}] = 0$ . Thus, we have  $[L_{m,i}, L_{n,j}] \subseteq \text{Ann}(\mathcal{S}(p, 0))$  for all  $(m, i), (n, j) \in \mathbb{Z} \times \mathbb{Z}$ , as needed. If  $p = 0$ , since  $L_{0,0} \in \mathbb{Z}(\mathcal{S}(0, 0))$ , then it leads to

$\mathcal{S}(0, 0) \cdot \mathcal{S}(0, 0) \subseteq \mathbb{Z}(\mathcal{S}(0, 0))$ . Hence the right-hand side of (2-2) is zero. Assuming  $[L_{m,i}, L_{n,j}] \in \langle L_{0,0} \rangle$ , we obtain from (2-1) that  $m = -n$  and  $j = -i$ . But then we get

$$ni - (-n)(-i) = ni - ni = 0.$$

So  $[L_{m,i}, L_{n,j}] = 0$ . We get  $[L_{m,i}, L_{n,j}] \subseteq \text{Ann}(\mathcal{S}(0, 0))$  for all  $(m, i), (n, j) \in \mathbb{Z} \times \mathbb{Z}$ . Then the left-hand side of (2-2) is zero as needed.  $\square$

### Acknowledgments

Y. Gao is supported by the Natural Sciences and Engineering Research Council of Canada and the National Natural Science Foundation of China (No. 11931009). J.-C. Sun is thankful for the support of the National Natural Science Foundation of China (Nos. 12071276, 11931009 and 12226402).

### References

- [1] C. Bai, R. Bai, L. Guo, and Y. Wu, “Transposed Poisson algebras, Novikov–Poisson algebras and 3-Lie algebras”, *J. Algebra* **632** (2023), 535–566. [MR](#) [Zbl](#)
- [2] K. Bhaskara and K. Viswanath, *Poisson algebras and Poisson manifolds*, Pitman Research Notes in Math. **174**, Longman Scientific & Technical, Harlow, 1988. [Zbl](#)
- [3] R. Block, “On torsion-free abelian groups and Lie algebras”, *Proc. Amer. Math. Soc.* **9** (1958), 613–620. [MR](#) [Zbl](#)
- [4] V. Chari and A. Pressley, *A guide to quantum groups*, Cambridge Univ. Press, 1994. [MR](#) [Zbl](#)
- [5] R. Fehlbberg Júnior and I. Kaygorodov, “On the Kantor product, II”, *Carpathian Math. Publ.* **14**:2 (2022), 543–563. [MR](#) [Zbl](#)
- [6] B. L. M. Ferreira, I. Kaygorodov, and V. Lopatkin, “ $\frac{1}{2}$ -derivations of Lie algebras and transposed Poisson algebras”, *Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Mat. RACSAM* **115**:3 (2021), art. id. 142. [MR](#) [Zbl](#)
- [7] V. T. Filippov, “On  $\delta$ -derivations of Lie algebras”, *Sibirsk. Mat. Zh.* **39**:6 (1998), 1409–1422. In Russian; translated in *Sib. Math. J.* **39**:6 (1998), 1218–1230. [MR](#) [Zbl](#)
- [8] I. B. Kaygorodov, “On  $\delta$ -derivations of classical Lie superalgebras”, *Sibirsk. Mat. Zh.* **50**:3 (2009), 547–565. In Russian; translated in *Sib. Math. J.* **50**:3 (2009), 434–449. [MR](#) [Zbl](#)
- [9] I. B. Kaygorodov, “ $\delta$ -superderivations of simple finite-dimensional Jordan and Lie superalgebras”, *Algebra Logika* **49**:2 (2010), 195–215. In Russian; translated in *Algebra Logic* **49**:2 (2010), 130–144. [MR](#) [Zbl](#)
- [10] I. Kaygorodov and M. Khrypchenko, “Transposed Poisson structures on Block Lie algebras and superalgebras”, *Linear Algebra Appl.* **656** (2023), 167–197. [MR](#) [Zbl](#)
- [11] I. Laraiedh and S. Silvestrov, “Constructions of BiHom-X algebras and bimodules of some BiHom-dialgebras”, *Algebra Discrete Math.* **34**:2 (2022), 273–316. [MR](#) [Zbl](#)
- [12] L.-C. Li, “Classical  $r$ -matrices and compatible Poisson structures for Lax equations on Poisson algebras”, *Comm. Math. Phys.* **203**:3 (1999), 573–592. [MR](#) [Zbl](#)
- [13] T. Ma and B. Li, “Transposed BiHom-Poisson algebras”, *Comm. Algebra* **51**:2 (2023), 528–551. [MR](#) [Zbl](#)

- [14] J. M. Osborn and K. Zhao, “Infinite-dimensional Lie algebras of generalized Block type”, *Proc. Amer. Math. Soc.* **127**:6 (1999), 1641–1650. [MR](#) [Zbl](#)
- [15] D. Ž. Đoković and K. Zhao, “Derivations, isomorphisms and second cohomology of generalized Block algebras”, *Algebra Colloq.* **3**:3 (1996), 245–272. [MR](#) [Zbl](#)
- [16] C. Xia, “Representations of twisted infinite Lie conformal superalgebras”, *J. Algebra* **596** (2022), 155–176. [MR](#) [Zbl](#)
- [17] C. Xia, “Classification of finite irreducible conformal modules over  $N = 2$  Lie conformal superalgebras of Block type”, *Algebr. Represent. Theory* **26**:5 (2023), 1731–1757. [MR](#) [Zbl](#)
- [18] C. Xia, T. You, and L. Zhou, “Structure of a class of Lie algebras of Block type”, *Comm. Algebra* **40**:8 (2012), 3113–3126. [MR](#) [Zbl](#)
- [19] Q. Xie and J. Sun, “Non-weight modules over  $N = 1$  Lie superalgebras of Block type”, *Forum Math.* **35**:5 (2023), 1279–1300. [MR](#) [Zbl](#)
- [20] Q. Xie, J. Sun, and C. Xia, “Non-weight modules over the Block type algebra  $\mathcal{B}'(p, q)$ ”, *J. Geom. Phys.* **193** (2023), art. id. 104988. [MR](#) [Zbl](#)
- [21] X. Xu, “Generalizations of the Block algebras”, *Manuscripta Math.* **100**:4 (1999), 489–518. [MR](#) [Zbl](#)
- [22] L. Yuan and Q. Hua, “ $\frac{1}{2}$ -(bi)derivations and transposed Poisson algebra structures on Lie algebras”, *Linear Multilinear Algebra* **70**:22 (2022), 7672–7701. [MR](#) [Zbl](#)

Received May 17, 2024. Revised November 25, 2024.

ANQI HUANG  
DEPARTMENT OF MATHEMATICS  
SHANGHAI UNIVERSITY  
SHANGHAI  
CHINA  
[anqi\\_huang@shu.edu.cn](mailto:anqi_huang@shu.edu.cn)

YUN GAO  
DEPARTMENT OF MATHEMATICS AND STATISTICS  
YORK UNIVERSITY  
TORONTO, ON  
CANADA  
[ygao@yorku.ca](mailto:ygao@yorku.ca)

JIANCAI SUN  
DEPARTMENT OF MATHEMATICS  
SHANGHAI UNIVERSITY  
SHANGHAI  
CHINA  
[jcsun@shu.edu.cn](mailto:jcsun@shu.edu.cn)

# PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

[msp.org/pjm](http://msp.org/pjm)

## EDITORS

Don Blasius (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[blasius@math.ucla.edu](mailto:blasius@math.ucla.edu)

Matthias Aschenbrenner  
Fakultät für Mathematik  
Universität Wien  
Vienna, Austria  
[matthias.aschenbrenner@univie.ac.at](mailto:matthias.aschenbrenner@univie.ac.at)

Robert Lipshitz  
Department of Mathematics  
University of Oregon  
Eugene, OR 97403  
[lipshitz@uoregon.edu](mailto:lipshitz@uoregon.edu)

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Kefeng Liu  
School of Sciences  
Chongqing University of Technology  
Chongqing 400054, China  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Atsushi Ichino  
Department of Mathematics  
Kyoto University  
Riverside, CA 90095-1555  
[atsushi.ichino@gmail.com](mailto:atsushi.ichino@gmail.com)

Dimitri Shlyakhtenko  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[shlyakht@ipam.ucla.edu](mailto:shlyakht@ipam.ucla.edu)

Ruixiang Zhang  
Department of Mathematics  
University of California  
Berkeley, CA 94720-3840  
[ruixiang@berkeley.edu](mailto:ruixiang@berkeley.edu)

## PRODUCTION

Silvio Levy, Scientific Editor, [production@msp.org](mailto:production@msp.org)

---

See inside back cover or [msp.org/pjm](http://msp.org/pjm) for submission instructions.

---

The subscription price for 2025 is US \$677/year for the electronic version, and \$917/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).


---

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

---

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2025 Mathematical Sciences Publishers

# PACIFIC JOURNAL OF MATHEMATICS

Volume 335    No. 1    March 2025

---

Central nilpotency of left skew braces and solutions of the Yang–Baxter equation	1
ADOLFO BALLESTER-BOLINCHES, RAMÓN ESTEBAN-ROMERO, MARIA FERRARA, VICENT PÉREZ-CALABUIG and MARCO TROMBETTI	
Hyperbolic $L$ -space knots not concordant to algebraic knots	33
MACIEJ BORODZIK and MASAKAZU TERAGAITO	
Weighted total variation minimization problem with mixed Dirichlet–Neumann boundary conditions	53
SAMER DWEIK	
Positive knots and ribbon concordance	81
JOE BONINGER	
Existence for some degenerate Hessian-type equations arising in conformal geometry	97
YAN HE, QIANG TU and NI XIANG	
Transposed Poisson superalgebra structures on twisted $N = 1$ Block–Lie superalgebra	119
ANQI HUANG, YUN GAO and JIANCAI SUN	
The CR Yamabe constant and inequivalent CR structures	163
CHANYOUNG SUNG and YUYA TAKEUCHI	
Weighted low-lying zeros of $L$ -functions attached to Siegel modular forms	183
SHIFAN ZHAO	