

*Pacific
Journal of
Mathematics*

AN EXTENSION OF GOW'S THEOREM

ROBERT M. GURALNICK AND PHAM HUU TIEP

AN EXTENSION OF GOW'S THEOREM

ROBERT M. GURALNICK AND PHAM HUU TIEP

In memory of our good friend and esteemed colleague Gary Seitz

We extend Gow's theorem to finite groups G whose generalized Fitting subgroup is $Z(G)S$ for a quasisimple Lie-type group S of simply connected type in characteristic p , and whose center $Z(G)$ has p' -order.

A result of Rod Gow [2000, Theorem 2] asserts that the product $a^G b^G$ of any two regular semisimple classes in a finite simple group of Lie type G contains every nontrivial semisimple element $x \in G$. This result has been used in many applications. It has also been extended to any quasisimple Lie-type group of simply connected type: the product $a^G b^G$ of any two regular semisimple classes in G contains every noncentral semisimple element $x \in G$; see [Guralnick and Tiep 2015, Lemma 5.1].

We will further extend Gow's theorem. Let p be a prime and let \underline{G} be a simple, simply connected algebraic group defined over $\overline{\mathbb{F}}_p$. Let $F : \underline{G} \rightarrow \underline{G}$ be a Steinberg endomorphism, so that

$$S := \underline{G}^F$$

is quasisimple. (In particular, we do not view $\mathrm{PSL}_2(9)$ as $\mathrm{Sp}_4(2)'$, $\mathrm{SU}_3(3)$ as $G_2(2)'$, or $\mathrm{SL}_2(8)$ as ${}^2G_2(3)'$.)

We will consider finite groups G with

$$(1) \quad F^*(G) = Z(G)S \quad \text{and} \quad p \nmid |Z(G)|,$$

(so $C_G(S) = C_G(F^*(G)) = Z(G)Z(S)$ is a p' -group), and aim to show that the product $a^G b^G$ of two particular conjugacy classes in G will cover all elements $g \in G$ of a certain kind. Before going on we state a special case of a consequence of our main result which is less technical. Versions of this result have already been used in [Acciarri et al. 2023; Guralnick et al. 2025].

Guralnick was partially supported by the NSF (grant DMS-1901595) and a Simons Foundation Fellowship 609771. Tiep was partially supported by the NSF (grant DMS-2200850) and the Joshua Barlaz Chair in Mathematics. The authors are grateful to the referee for a careful reading and helpful comments on the paper.

MSC2020: 20C15, 20C33, 20G40.

Keywords: finite groups of Lie type, products of conjugacy classes.

Corollary 1. *Let G be as above. Assume that $a \in G$ has order prime to p , $|\mathbf{C}_S(a)|$ has order prime to p and that $s \in S \setminus \mathbf{Z}(S)$ is semisimple. Then $s^t = [a, u]$ for some $t, u \in S$.*

In fact, one need not assume that g has order prime to p . See our main result [Theorem 6](#).

Let a and b be elements of G , and set $G_1 := \langle S, a, b \rangle$. Then $S \triangleleft G_1$, and hence $S \triangleleft F^*(G_1)$. It follows that $F(G_1)$, as well as any quasisimple subnormal subgroup $T \neq S$ of G_1 , centralizes S . Hence by [\(1\)](#), $F^*(G_1) = \mathbf{Z}(G_1)S$. Furthermore,

$$\mathbf{Z}(G_1) \leq \mathbf{C}_G(S) = \mathbf{Z}(G)\mathbf{Z}(S)$$

is also a p' -subgroup. Hence, for our purposes, we may assume

$$(2) \quad G = \langle S, a, b \rangle.$$

Let St denote the Steinberg character of S , and for $x \in G$, write x_p for the p -part of x . By [\[Feit 1993, Corollary D\]](#), St extends to a rational-valued character St_G of G (called the *basic p -Steinberg character* of G). By [\[Feit 1995, Theorem C\]](#), there is a Sylow p -subgroup P and a p -subgroup D of G , of order

$$|D| = p^d = |G/S|_p,$$

such that $P = Q \rtimes D$ for a Sylow p -subgroup Q of S and the following statement holds. For any element $x \in G$, $\text{St}_G(x) \neq 0$ if and only if $x_p \in D$ (up to conjugation), in which case

$$\text{St}_G(x) = \pm |\mathbf{C}_S(x)|_p.$$

In view of these results, the proper generalization to G of regular semisimple classes in S will be that $a, b \in G$ satisfy

$$(3) \quad a_p \in D \text{ up to conjugation in } G \quad \text{and} \quad p \nmid |\mathbf{C}_S(a)|,$$

and

$$(4) \quad b_p \in D \text{ up to conjugation in } G \quad \text{and} \quad p \nmid |\mathbf{C}_S(b)|.$$

Certainly, $g \in G$ can belong to $a^G b^G$ only when it does so in the solvable group G/S , so we will assume

$$(5) \quad gS \in (aS)^{G/S} (bS)^{G/S} \quad \text{and} \quad g_p \in D \text{ up to conjugation in } G.$$

For instance, if G/S is abelian, then the first condition in [\(5\)](#) is equivalent to $g \in abS$.

Proposition 2. (i) *If $p > 2$ and $Q \in \text{Syl}_p(S)$, then $\mathbf{C}_G(Q) = \mathbf{Z}(G)\mathbf{Z}(S)\mathbf{Z}(Q)$.*

(ii) *If $g \in G \setminus \mathbf{Z}(G)\mathbf{Z}(S)$ and $g_p \in D$, then p divides $[S : \mathbf{C}_S(g)]$.*

Proof. Since $Z := \mathbf{Z}(G)\mathbf{Z}(S)$ is a p' -group centralizing Q and normal in G , we may work in $\bar{G} := G/Z$ and identify Q with $\bar{Q} := QZ/Z$. Note that $Z \cap S = \mathbf{Z}(S)$. Moreover, as S is perfect, we have $C_G(S/\mathbf{Z}(S)) = C_G(S) = Z$. It follows that

$$\bar{S} := S/\mathbf{Z}(S) \triangleleft \bar{G} \leq \text{Aut}(\bar{S}),$$

i.e., \bar{G} is almost simple.

Consider any element $x \in \bar{C} := C_{\bar{G}}(\bar{Q})$. Then $H := \langle \bar{S}, x \rangle \leq \bar{G}$ is also almost simple, whence $\mathbf{O}_{p'}(H) = 1$, and $R := \langle \bar{Q}, x_p \rangle$ is a Sylow p -subgroup of H centralized by x . It follows that $R = \mathbf{O}_p(N_H(R))$. By [Gorenstein et al. 1998, Corollary 3.1.4], $\mathbf{O}_p(N_H(R)) = F^*(N_H(R))$, whence x belongs to $C_{N_H(R)}(R) \leq R$ and so $x = x_p$ is a p -element. Thus \bar{C} is a p -group.

Similarly, $\bar{Q} = \mathbf{O}_p(N_{\bar{S}}(\bar{Q})) = F^*(N_{\bar{S}}(\bar{Q}))$, and so

$$\bar{C} \cap \bar{S} = C_{\bar{S}}(\bar{Q}) = C_{N_{\bar{S}}(\bar{Q})}(\bar{Q}) \leq \bar{Q}.$$

It follows that $\bar{C} \cap \bar{S} = \mathbf{Z}(\bar{Q})$.

(i) Now we assume $p > 2$ and show that $\bar{C} \leq \bar{S}$, which implies that $C_G(Q) = \mathbf{Z}(Q)Z$. Assume the contrary: $\bar{C} \not\leq \bar{S}$. Since \bar{C} is a p -group, we can find a p -element $x \in \bar{C} \setminus \bar{S}$; in particular, $[x, \bar{Q}] = 1$. Now $R := \langle x, \bar{Q} \rangle$ is Sylow p -subgroup of $H := \langle \bar{S}, x \rangle$ and $x \in \mathbf{Z}(R)$. As $\bar{S} \triangleleft H \leq \bar{G} \leq \text{Aut}(\bar{S})$, we still have $\mathbf{O}_{p'}(H) = 1$. Now, if $p > 2$ then $\mathbf{Z}(R) \leq F^*(H) = \bar{S}$ by [Glauberman et al. 2020, Corollary 1.2], and hence $x \in \bar{S}$, contrary to the choice of x .

(ii) Assume the contrary that $p \nmid [S : C_S(g)]$. Conjugating g suitably, we may assume that $g \in C_G(Q)$ with $Q \in \text{Syl}_p(S)$ as before.

Suppose first that $p > 2$. Then $g \in \mathbf{Z}(G)\mathbf{Z}(S)\mathbf{Z}(Q)$ by (i), and so $g_p \in S$. But g_p is conjugate to an element in D by assumption and $D \cap S = 1$, so $g_p = 1$. It follows that $g \in \mathbf{Z}(G)\mathbf{Z}(S)$, a contradiction.

Thus we have $p = 2$. Then

$$\text{St}_G(1) = |Q| = |C_S(g)|_p = \pm \text{St}_G(g).$$

On the other hand, St is trivial at $\mathbf{Z}(S)$, so the generalized center of St_G contains $Z = \mathbf{Z}(G)\mathbf{Z}(S)$ and hence equals G as G/Z is almost simple with socle \bar{S} . As the generalized center of St_G contains g , we conclude that $g \in Z$, again a contradiction. \square

Fix any element $g \in G$ satisfying (5). Then $SC_G(g) \leq G$, so

$$(6) \quad Z \ni [G : SC_G(g)] = \frac{|G| \cdot |C_S(g)|}{|S| \cdot |C_G(g)|} = \frac{[G : C_G(g)]}{[S : C_S(g)]}.$$

Write

$$(7) \quad \frac{[G : C_G(g)]_p}{[S : C_S(g)]_p} = p^e.$$

Lemma 3. *Let X be a finite group, which is abelian-by-cyclic, that is, X has a normal abelian subgroup $A \triangleleft X$ such that X/A is cyclic. Suppose $x, y, z \in X$ are such that*

$$X = \langle x, y \rangle \quad \text{and} \quad z \equiv xy \pmod{[X, X]}.$$

Then

$$\sum_{\alpha \in \text{Irr}(X)} \frac{\alpha(x)\alpha(y)\overline{\alpha(z)}}{\alpha(1)} = |X/[X, X]|.$$

Proof. The condition $z \equiv xy \pmod{[X, X]}$ implies that

$$\sum_{\alpha \in \text{Irr}(X), \alpha(1)=1} \frac{\alpha(x)\alpha(y)\overline{\alpha(z)}}{\alpha(1)} = |X/[X, X]|.$$

Hence it suffices to show that the contribution of any nonlinear $\alpha \in \text{Irr}(X)$ to the sum in the statement is 0. Consider any irreducible constituent λ of $\alpha|_A$. Suppose λ is not X -invariant. As $X = \langle x, y \rangle$, we may assume that λ is not x -invariant, in which case $\alpha(x) = 0$ by Clifford's theorem and the contribution is 0 as claimed.

Suppose now that λ is X -invariant. Then for any $a \in A$ and $t \in X$, as $\lambda(1) = 1$ we have

$$\lambda(tat^{-1}a^{-1}) = \lambda(tat^{-1})/\lambda(a) = 1,$$

whence $[t, a] \in \text{Ker}(\lambda)$ and $\text{Ker}(\lambda) \triangleleft X$. It follows that $A/\text{Ker}(\lambda) \leq \mathbf{Z}(X/\text{Ker}(\lambda))$. But X/A is cyclic, so $X/\text{Ker}(\lambda)$ is abelian. Now λ is the unique irreducible constituent of α_A , so $\text{Ker}(\lambda) \leq \text{Ker}(\alpha)$, and hence α , viewed as an irreducible character of $X/\text{Ker}(\lambda)$, must be linear, contrary to the assumption $\alpha(1) > 1$. \square

Proposition 4. *Under the assumptions (1)–(5), assume in addition that G/S is abelian-by-cyclic. Then*

$$\Sigma_1 := \sum_{\chi \in \text{Irr}(G|\text{St})} \frac{\chi(a)\chi(b)\overline{\chi(g)}}{\chi(1)} \cdot |g^G|$$

is a rational integer whose p -part is at most p^{d+e} .

Proof. As mentioned above, St extends to St_G . Hence, by Gallagher's theorem [Isaacs 2006, (6.17)], any $\chi \in \text{Irr}(G|\text{St})$ is of the form

$$\chi = \text{St}_G \alpha$$

with $\alpha \in \text{Irr}(G/S)$. Using (2), (5) and Lemma 3, we see that

$$\sum_{\alpha \in \text{Irr}(G/S)} \frac{\alpha(a)\alpha(b)\overline{\alpha(g)}}{\alpha(1)}$$

is a rational integer whose p -part is at most p^d .

On the other hand, by (3) and (4) we see that

$$\begin{aligned} \frac{\text{St}_G(a)\text{St}_G(b)\overline{\text{St}_G(g)}}{\text{St}_G(1)} \cdot |g^G| &= \pm \frac{|C_S(g)|_p \cdot |G|_p \cdot |G|_{p'}}{|S|_p \cdot |C_G(g)|_p \cdot |C_G(g)|_{p'}} \\ &= \pm \frac{[G : C_G(g)]_p}{[S : C_S(g)]_p} \cdot [G : C_G(g)]_{p'} \end{aligned}$$

is p^e times a p' -integer. Hence the statement follows. □

Recall that St is the only p -defect zero character of S . By the main result of [Humphreys 1971], all the remaining characters of S belong to p -blocks of maximal defect. The next result deals with these characters.

Proposition 5. *Under the assumptions (1)–(5), assume in addition that G/S has a cyclic Sylow p -subgroup and a normal p -complement, and that $g \notin \mathbf{Z}(G)\mathbf{Z}(S)$. Then*

$$\Sigma_2 := \sum_{\chi \in \text{Irr}(G) \setminus \text{Irr}(G|\text{St})} \frac{\chi(a)\chi(b)\overline{\chi(g)}}{\chi(1)} \cdot |g^G|$$

is p^{d+e+1} times an algebraic integer.

Proof. By the hypothesis we can write $G/S = (H/S) \rtimes D$ for some normal subgroup $H \geq S$ of G . Note that any $\chi \in \text{Irr}(G) \setminus \text{Irr}(G|\text{St})$ lies above some $\theta \in \text{Irr}(H)$ which does not lie above St . Suppose θ is not G -invariant. As $G = \langle H, a, b \rangle$, we may assume that θ is not a -invariant, in which case $\chi(a) = 0$ by Clifford's theorem, and the contribution of χ to Σ_2 is 0.

Hence we need to count the total contribution to Σ_2 of the characters $\chi \in \text{Irr}(G|\theta)$, where $\theta \notin \text{Irr}(H|\text{St})$ is G -invariant. Since G/H is cyclic, any such θ extends to a character χ_1 of G , and we may write

$$\text{Irr}(G|\theta) = \{\chi_1\mu \mid \mu \in \text{Irr}(G/H)\}.$$

By the assumption $\theta \notin \text{Irr}(H|\text{St})$, every irreducible constituent of $\theta|_S$ belongs to an S -block B_S of maximal p -defect.

Conjugating g suitably, we may assume that $g_p \in D$. Note that $g_{p'} \in H$, so $g = g_p g_{p'}$ belongs to

$$K := \langle H, g_p \rangle \triangleleft G.$$

Set

$$\chi_2 := (\chi_1)|_K \in \text{Irr}(K|\theta).$$

Now the p -block B of H that contains θ covers B_S , and $p \nmid |H/S|$, so B has maximal defect; see, e.g., [Navarro 1998, Theorem 9.26]. But $K/H \hookrightarrow D$ is a p -group, so by [Navarro 1998, Corollary 9.6] there is a unique p -block B_2 of K that covers B . In particular,

$$\text{Irr}(K|\theta) \subseteq \text{Irr}(B_2).$$

Moreover, B is K -invariant as θ is K -invariant, whence B_2 is of maximal defect by [Navarro 1998, Theorem 9.17]. It follows that B_2 contains a character χ_0 of height zero, and so of p' -degree.

As χ_0 and χ_2 belong to the same block, we know that the two algebraic integers

$$\omega_{\chi_i}(g) = \frac{\chi_i(g)}{\chi_i(1)} \cdot |g^K|$$

for $i \in \{0, 2\}$ are congruent modulo p . By Proposition 2, $|g^S|$ is divisible by p , so $|g^K|$ is divisible by p as well; see the computation in (6). But $p \nmid \chi_0(1)$, so $p \mid \omega_{\chi_0}(g)$. It follows that

$$(8) \quad p \text{ divides } \omega_{\chi_2}(g) = \frac{\chi_2(g)}{\chi_2(1)} \cdot |g^K| = \frac{\chi_1(g)}{\chi_1(1)} \cdot |g^K|.$$

Next, (6) applied to $S \triangleleft H$ with $p \nmid |H/S|$ shows that

$$|g^S|_p = |g^H|_p.$$

On the other hand, g_p centralizes g , and $K = \langle H, g_p \rangle$, so $HC_K(g) = K$, showing that $g^K = g^H$. Hence $|g^S|_p = |g^K|_p$, and (7) becomes

$$p^e = \frac{|g^G|_p}{|g^K|_p}.$$

Together with (8), we now obtain

$$p^{e+1} \text{ divides } \omega_{\chi_1}(g) = \frac{\chi_1(g)}{\chi_1(1)} \cdot |g^G|.$$

Now, (5) implies that $g \equiv ab \pmod{G/H}$, and so

$$\sum_{\mu \in \text{Irr}(G/H)} \frac{\mu(a)\mu(b)\overline{\mu(g)}}{\mu(1)} = |G/H| = p^d.$$

It follows that

$$\sum_{\chi \in \text{Irr}(G|\theta)} \frac{\chi(a)\chi(b)\overline{\chi(g)}}{\chi(1)} \cdot |g^G| = \omega_{\chi_1}(g) \sum_{\mu \in \text{Irr}(G/H)} \frac{\mu(a)\mu(b)\overline{\mu(g)}}{\mu(1)} = p^d \omega_{\chi_1}(g)$$

is p^{d+e+1} times an algebraic integer. □

Theorem 6. *Under the assumptions (1)–(5), assume in addition that all the following conditions hold:*

- (a) G/S is abelian-by-cyclic.
- (b) G/S has cyclic Sylow p -subgroups and a normal p -complement.
- (c) $g \notin \mathbf{Z}(G)\mathbf{Z}(S)$.

Then $g \in a^G b^G$.

Proof. By Propositions 4 and 5,

$$|g^G| \sum_{\chi \in \text{Irr}(G)} \frac{\chi(a)\chi(b)\overline{\chi(g)}}{\chi(1)} = \Sigma_1 + \Sigma_2$$

is $p^s(u + p^t v)$, where $u \in \mathbb{Z} \setminus p\mathbb{Z}$, v is an algebraic integer, $0 \leq s \leq d + e$, and $t \geq 1$. Now if $u + p^t v = 0$, then $v = -u/p^t$ is rational and an algebraic integer, so $v \in \mathbb{Z}$ and $u \in p\mathbb{Z}$, a contradiction. Thus

$$\sum_{\chi \in \text{Irr}(G)} \frac{\chi(a)\chi(b)\overline{\chi(g)}}{\chi(1)} \neq 0,$$

and so $g \in a^G b^G$ by Frobenius' character formula. □

Proof of Corollary 1. To prove this, we may replace G by $\langle S, a, b \rangle$ with $b := a^{-1}$. Then (3)–(4) hold with $g := s$. Now G/S is cyclic, so $s \in a^G b^G$ by Theorem 6. But $a^G = a^S$ and $b^G = b^S$ since $G = \langle S, a \rangle = \langle S, b \rangle$, so the statement follows. □

In what follows, q is always a power of the prime p . We will use the structure of $\text{Aut}(\bar{S})$ as described in [Gorenstein et al. 1998, Theorem 2.5.12], in particular the notation $\text{Inndiag}(\bar{S})$ and $\text{Outdiag}(\bar{S})$.

Theorem 7. *Under the assumptions (1)–(5), assume in addition that all the following conditions hold for g , $\bar{S} = S/\mathbf{Z}(S)$, and $\bar{G} = G/\mathbf{Z}(G)\mathbf{Z}(S)$:*

- (a) $g \notin \mathbf{Z}(G)\mathbf{Z}(S)$.
- (b) *If $\bar{S} = \text{PSL}_n(q)$ with $n \geq 3$, or $\bar{S} = P\Omega_{2n}^+(q)$ with $n \geq 4$, or $S = E_6(q)$, then the quotient $\bar{G}/(\bar{G} \cap \text{Inndiag}(\bar{S}))$ is cyclic.*

Then $g \in a^G b^G$.

Proof. Recall that $G/S \cong \bar{G}/\bar{S}$ is a subgroup of $O := \text{Out}(\bar{S})$. By Theorem 6, we need to show that $A = G/S$ satisfies both of the conditions (a) and (b) listed therein. Note that both (a) and (b) in Theorem 6 follow from the condition

- (9) A admits a normal abelian p' -subgroup B with A/B being cyclic.

In turn, (9) is a consequence of the condition

- (10) $O := \text{Out}(\bar{S})$ admits a normal abelian p' -subgroup J with O/J being cyclic.

(Indeed, taking $B := A \cap J$ we have $A/B \hookrightarrow O/J$.)

Set $J := \text{Outdiag}(\bar{S}) := \text{Inndiag}(\bar{S})/\bar{S}$. Now, if \bar{S} is a *twisted* group, i.e., the parameter d for $\bar{S} \cong {}^d\Sigma(q)$ in [Gorenstein et al. 1998, Theorem 2.5.12] is greater than one, then (10) holds (for this choice of J). It remains to consider the untwisted groups, that is, the ones with $d = 1$.

If $\bar{S} = \text{PSL}_2(q)$ then (10) holds. Suppose that $\bar{S} = \text{PSL}_n(q)$ with $n \geq 3$, or $\bar{S} = P\Omega_{2n}^+(q)$ with $n \geq 4$, or $S = E_6(q)$. Then taking $B := (\bar{G} \cap \text{Inndiag}(\bar{S}))/\bar{S}$, we see that A/B is cyclic by assumption (b) in Theorem 7, hence (9) holds.

In the remaining cases, \bar{S} is of type B_n , C_n , G_2 , F_4 , E_7 , or E_8 , hence O/J is cyclic, and so (10) holds. \square

Next we deduce another consequence of Theorem 6. For the definition of the reduced Clifford group $\Gamma^+(\mathbb{F}_q^n) = \text{CSpin}_n^\epsilon(q)$, see, for example, [Tiep and Zalesski 2005, §6]; in particular, it contains $\text{Spin}_n^\epsilon(q)$ as a normal subgroup with factor C_{q-1} .

Theorem 8. *Let q be a prime power, and let (G, S) be any of the following pairs of groups:*

- (a) $G = \text{GL}_n(q)$ with $n \geq 2$, $(n, q) \neq (2, 2), (2, 3)$, and $S = \text{SL}_n(q)$.
- (b) $G = \text{GU}_n(q)$ with $n \geq 2$, $(n, q) \neq (2, 2), (2, 3), (3, 2)$, and $S = \text{SU}_n(q)$.
- (c) $G = \text{CSp}_{2n}(q)$ with $n \geq 2$, $(n, q) \neq (2, 2)$, and $S = \text{Sp}_{2n}(q)$.
- (d) $G = \text{CSpin}_n^\epsilon(q)$ with $n \geq 5$, $2 \nmid q$, and $\epsilon = \pm$, and $S = \text{Spin}_n^\epsilon(q)$.
- (e) $G = \text{GO}_n^\epsilon(q)$ or $\text{SO}_n^\epsilon(q)$ with $n \geq 5$, $2 \nmid q$, and $\epsilon = \pm$, and $S = \Omega_n^\epsilon(q)$.

Suppose that $a, b \in G$ are such that $p \nmid |C_S(a)|$ and $p \nmid |C_S(b)|$. If $g \in G$ is any noncentral p' -element such that $g \in abS$, then $g \in a^G b^G$.

Proof. For all of the above pairs but (e), we have that S is a quasisimple Lie-type group of simply connected type, $S \triangleleft G$, $F^*(G) = \mathbf{Z}(G)S$, and $\mathbf{Z}(S) \leq \mathbf{Z}(G)$. Furthermore, G/S is abelian of p' -order, and (3), (4), and (5) are all fulfilled. Hence the statement follows from Theorem 6. In the case of (e), the same proof of Theorem 6 applies. \square

Note that Theorem 6 also applies to $\text{GO}_{2n}^\epsilon(q)$ with $2 \mid q$ and $n \geq 3$. But we do not include them in Theorem 8 since the subgroup D is now of order 2 and so conditions (3)–(5) are more complicated than those formulated in Theorem 8.

References

- [Acciarri et al. 2023] C. Acciarri, R. M. Guralnick, and P. Shumyatsky, “Criteria for solubility and nilpotency of finite groups with automorphisms”, *Bull. Lond. Math. Soc.* **55**:3 (2023), 1340–1346. [MR](#) [Zbl](#)
- [Feit 1993] W. Feit, “Extending Steinberg characters”, pp. 1–9 in *Linear algebraic groups and their representations* (Los Angeles, CA, 1992), edited by R. S. Elman et al., Contemp. Math. **153**, Amer. Math. Soc., Providence, RI, 1993. [MR](#) [Zbl](#)
- [Feit 1995] W. Feit, “Steinberg characters”, *Bull. London Math. Soc.* **27**:1 (1995), 34–38. [MR](#) [Zbl](#)
- [Glauberman et al. 2020] G. Glauberman, R. Guralnick, J. Lynd, and G. Navarro, “Centers of Sylow subgroups and automorphisms”, *Israel J. Math.* **240**:1 (2020), 253–266. [MR](#) [Zbl](#)

- [Gorenstein et al. 1998] D. Gorenstein, R. Lyons, and R. Solomon, *The classification of the finite simple groups, III*, Math. Surveys and Monographs **40.3**, Amer. Math. Soc., Providence, RI, 1998. [MR](#)
- [Gow 2000] R. Gow, “Commutators in finite simple groups of Lie type”, *Bull. London Math. Soc.* **32**:3 (2000), 311–315. [MR](#) [Zbl](#)
- [Guralnick and Tiep 2015] R. M. Guralnick and P. H. Tiep, “Lifting in Frattini covers and a characterization of finite solvable groups”, *J. Reine Angew. Math.* **708** (2015), 49–72. [MR](#) [Zbl](#)
- [Guralnick et al. 2025] R. M. Guralnick, H. P. Tong-Viet, and G. Tracey, “Weakly subnormal subgroups and variations of the Baer-Suzuki theorem”, *J. Lond. Math. Soc. (2)* **111**:1 (2025), art. id. e70057, 29 pp. [MR](#)
- [Humphreys 1971] J. E. Humphreys, “Defect groups for finite groups of Lie type”, *Math. Z.* **119** (1971), 149–152. [MR](#) [Zbl](#)
- [Isaacs 2006] I. M. Isaacs, *Character theory of finite groups*, AMS Chelsea, Providence, RI, 2006. [MR](#) [Zbl](#)
- [Navarro 1998] G. Navarro, *Characters and blocks of finite groups*, London Mathematical Society Lecture Note Series **250**, Cambridge University Press, 1998. [MR](#) [Zbl](#)
- [Tiep and Zalesski 2005] P. H. Tiep and A. E. Zalesski, “Real conjugacy classes in algebraic groups and finite groups of Lie type”, *J. Group Theory* **8**:3 (2005), 291–315. [MR](#) [Zbl](#)

Received April 8, 2024. Revised October 20, 2024.

ROBERT M. GURALNICK
DEPARTMENT OF MATHEMATICS
UNIVERSITY OF SOUTHERN CALIFORNIA
LOS ANGELES, CA
UNITED STATES
guralnic@usc.edu

PHAM HUU TIEP
DEPARTMENT OF MATHEMATICS
RUTGERS UNIVERSITY
PISCATAWAY, NJ
UNITED STATES
tiep@math.rutgers.edu

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Fakultät für Mathematik
Universität Wien
Vienna, Austria
matthias.aschenbrenner@univie.ac.at

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Atsushi Ichino
Department of Mathematics
Kyoto University
Riverside, CA 92521-0135
atsushi.ichino@gmail.com

Robert Lipshitz
Department of Mathematics
University of Oregon
Eugene, OR 97403
lipshitz@uoregon.edu

Kefeng Liu
School of Sciences
Chongqing University of Technology
Chongqing 400054, China
liu@math.ucla.edu

Sucharit Sarkar
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
sucharit@math.ucla.edu

Dimitri Shlyakhtenko
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
shlyakht@ipam.ucla.edu

Ruixiang Zhang
Department of Mathematics
University of California
Berkeley, CA 94720-3840
ruixiang@berkeley.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2025 is US \$677/year for the electronic version, and \$917/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2025 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 336 No. 1-2 May 2025

In memoriam Gary Seitz

Gary Seitz (1943–2023): In memoriam	1
MARTIN W. LIEBECK, GERHARD RÖHRLE and DONNA TESTERMAN	
Intrinsic components in involution centralizers of fusion systems	7
MICHAEL ASCHBACHER	
On good A_1 subgroups, Springer maps, and overgroups of distinguished unipotent elements in reductive groups	29
MICHAEL BATE, SÖREN BÖHM, BENJAMIN MARTIN and GERHARD RÖHRLE	
The q -Schur category and polynomial tilting modules for quantum GL_n	63
JONATHAN BRUNDAN	
The binary actions of simple groups of Lie type of characteristic 2	113
NICK GILL, PIERRE GUILLOT and MARTIN W. LIEBECK	
Finite simple groups have many classes of p -elements	137
MICHAEL GIUDICI, LUKE MORGAN and CHERYL E. PRAEGER	
Monogamous subvarieties of the nilpotent cone	161
SIMON M. GOODWIN, RACHEL PENGELLY, DAVID I. STEWART and ADAM R. THOMAS	
An extension of Gow's theorem	181
ROBERT M. GURALNICK and PHAM HUU TIEP	
On dimensions of RoCK blocks of cyclotomic quiver Hecke superalgebras	191
ALEXANDER KLESHCHEV	
Representation growth of Fuchsian groups and modular forms	217
MICHAEL J. LARSEN, JAY TAYLOR and PHAM HUU TIEP	
D_4 -type subgroups of $F_4(q)$	249
R. LAWThER	
Constructible representations and Catalan numbers	339
GEORGE LUSZTIG and ERIC SOMMERS	
A reduction theorem for simple groups with $e(G) = 3$	351
RICHARD LYONS and RONALD SOLOMON	
Decomposition numbers in the principal block and Sylow normalisers	367
GUNTER MALLE and NOELIA RIZO	
Levi decompositions of linear algebraic groups and nonabelian cohomology	379
GEORGE J. MCNINCH	
On the intersection of principal blocks	399
GABRIEL NAVARRO, A. A. SCHAEFFER FRY and PHAM HUU TIEP	
Hesselink strata in small characteristic and Lusztig–Xue pieces	415
ALEXANDER PREMÉT	
Multiplicity-free representations of the principal A_1 -subgroup in a simple algebraic group	433
ALUNA RIZZOLI and DONNA TESTERMAN	