

*Pacific
Journal of
Mathematics*

**DECOMPOSITION NUMBERS IN THE PRINCIPAL BLOCK
AND SYLOW NORMALISERS**

GUNTER MALLE AND NOELIA RIZO

DECOMPOSITION NUMBERS IN THE PRINCIPAL BLOCK AND SYLOW NORMALISERS

GUNTER MALLE AND NOELIA RIZO

To the memory of Gary Seitz

If G is a finite group and p is a prime number, we investigate the relationship between the p -modular decomposition numbers of characters of height zero in the principal p -block of G and the p -local structure of G . In particular we prove that, under certain conditions on the nonabelian composition factors of G , $d_{\chi 1_G} \neq 0$ for all irreducible characters χ of degree prime to p in the principal p -block of G if, and only if, the normaliser of a Sylow p -subgroup of G has a normal p -complement.

1. Introduction

Let G be a finite group, p a prime number, $\text{Irr}(G)$ the set of irreducible ordinary characters of G and $\text{IBr}(G)$ the set of irreducible p -Brauer characters of G . Then the restriction χ° of any $\chi \in \text{Irr}(G)$ to the set of p -regular elements of G can be written as

$$\chi^\circ = \sum_{\varphi \in \text{IBr}(G)} d_{\chi\varphi} \varphi,$$

where the $d_{\chi\varphi}$ are uniquely determined nonnegative integers. These are called the p -modular *decomposition numbers of G* , and a great deal of literature is devoted to understanding them.

Navarro and Tiep [2020; 2022] initiated the investigation on relations between p -decomposition numbers and properties of Sylow p -normalisers, considering two different settings. In [Navarro and Tiep 2020] they conjectured that if $p > 3$ then $d_{\chi 1_G} \neq 0$ for all $\chi \in \text{Irr}_{p'}(G)$, that is, for all irreducible characters of G of degree prime to p , if and only if G has self-normalising Sylow p -subgroups, and that this happens if and only if $d_{\chi 1_G} = 1$ for all $\chi \in \text{Irr}_{p'}(G)$. Note that irreducible characters

Malle gratefully acknowledges financial support by the DFG—Project-ID 286237555. Rizo is supported by Ministerio de Ciencia e Innovación (Grant PID2019-103854GB-I00 and Grant PID2022-137612NBI00 funded by MCIN/AEI/10.13039/501100011033 and “ERDF A way of making Europe”) and a CDEIGENT grant CIDEIG/2022/29 funded by Generalitat Valenciana.

MSC2020: 20C15, 20C20, 20C33, 20D20.

Keywords: decomposition numbers, principal blocks, Sylow normalisers.

of degree prime to p lie in p -blocks of maximal defect, so this situation can only happen when the principal block is the only Brauer p -block of maximal defect of G . It is then natural to wonder what can be said when this is not the case and one restricts attention to the principal p -block $B_0(G)$. In this sense, in [Navarro and Tiep 2022] they conjectured that $d_{\chi_{1G}} \neq 0$ for all $\chi \in \text{Irr}(B_0(G))$ if and only if G has a normal p -complement. In this paper we consider the intersection of the two conditions in [Navarro and Tiep 2020] and [2022], namely what happens if $d_{\chi_{1G}} = 1$ just for all $\chi \in \text{Irr}_{p'}(B_0(G))$. Our main result is:

Theorem A. *Let G be a finite group and $p > 3$ be a prime. Assume that all non-abelian simple composition factors of G of order divisible by p satisfy **Property (*)** below. The following are equivalent:*

- (i) *For every $\chi \in \text{Irr}_{p'}(B_0(G))$ we have $d_{\chi_{1G}} \neq 0$.*
- (ii) *For every $\chi \in \text{Irr}_{p'}(B_0(G))$ we have $d_{\chi_{1G}} = 1$.*
- (iii) *For $P \in \text{Syl}_p(G)$ we have $N_G(P) = P \times K$ for some $K \leq G$.*

Moreover, if G is p -solvable, this equivalence holds for every prime p .

Notice that, as pointed out in [Navarro and Tiep 2020], all irreducible characters of odd degree of the alternating group \mathfrak{A}_5 contain the trivial character in their 2-modular reduction, and similarly, all irreducible characters of the Ree group ${}^2G_2(27)$ of nonzero 3-defect contain the trivial character in their 3-modular reduction, while the respective Sylow p -normalisers have no normal p -complement, so the equivalence in **Theorem A** fails for nonsolvable groups with $p \leq 3$.

Theorem A involves the following property that a finite nonabelian simple group might, or might not, satisfy:

Property (*). Let S be nonabelian simple and $p > 3$ a prime dividing $|S|$. Then for all almost simple groups H with socle S and $|H : S|$ a p -power, there exists $\chi \in \text{Irr}_{p'}(B_0(H))$ such that $d_{\chi_{1H}} = 0$.

We prove **Property (*)** does hold for many simple groups: for sporadic groups, alternating groups, and simple groups of Lie type in characteristic different from p . We also show it for some groups of Lie type in characteristic p . (The general case of groups of Lie type in their defining characteristic was also left open in [Navarro and Tiep 2020; 2022].) As the knowledge on p -decomposition numbers for groups of Lie type in their own characteristic is too weak at present we refrain from making a general conjecture and just leave it as a question as to whether **Property (*)** holds for all nonabelian finite simple groups.

Structure of the paper. In **Section 2** we prove **Property (*)** for S a sporadic simple group, an alternating group, a simple group of Lie type in characteristic different from p as well as for some groups of Lie type in characteristic p . In **Section 3** we

show that [Theorem A](#) holds for p -solvable groups and in [Theorem 3.4](#) we reduce the general case to the validity of [Property \(*\)](#) on composition factors, thus completing the proof of [Theorem A](#).

2. Almost simple groups

In this section, we discuss instances of [Property \(*\)](#) from the introduction.

Alternating and sporadic groups. We start out with simple groups not of Lie type.

Proposition 2.1. *Property (*) holds for S a sporadic simple group or the Tits group.*

Proof. Let S be as in the assumption and $H \geq S$ as in [Property \(*\)](#). Since $p > 2$ this means $H = S$. By [[Navarro and Tiep 2022](#), Proposition 3.2] there exists $\chi \in \text{Irr}(B_0(H))$ such that $d_{\chi 1_H} = 0$. If S has abelian Sylow p -subgroups, by one direction of Brauer’s height zero conjecture [[Kessar and Malle 2013](#)], we then have $\chi \in \text{Irr}_{p'}(B_0(H))$ as required. Now assume Sylow p -subgroups are nonabelian. By [[Navarro and Tiep 2020](#), Lemma 3.2], there is $\chi \in \text{Irr}_{p'}(H) = \text{Irr}_{p'}(B_0(H))$ with $d_{\chi 1_H} = 0$, and again we are done since by inspection in [[GAP 2020](#)], S has just one p -block of maximal defect. □

Proposition 2.2. *Property (*) holds for S an alternating group.*

Proof. Let $S = \mathfrak{A}_n$ with $n \geq 5$. As $p > 2$ again we have $H = S$. Set $G := \mathfrak{S}_n$. Recall that the irreducible characters of \mathfrak{S}_n are naturally labelled by partitions of n . If p divides n , then let $\chi = \chi^{(n-2, 1^2)} \in \text{Irr}(B_0(G))$, as in [[Navarro and Tiep 2022](#), Proposition 3.1]. Then χ° is the sum of two irreducible Brauer characters of degrees $n - 2$ and $\frac{1}{2}(n - 2)(n - 3)$ and hence it is of degree prime to p and does not contain any irreducible Brauer character of degree 1. Now, any $\theta \in \text{Irr}(S)$ under χ lies in $\text{Irr}_{p'}(B_0(S))$ and $d_{\theta 1_S} = 0$, as wanted.

So we may assume that p does not divide n . Let $n = a_k p^k + \dots + a_1 p + a_0$ be the p -adic expansion of n . Since p does not divide n , we have $a_0 > 0$. Suppose first that $a_0 > 1$. Consider $\chi = \chi^{(a_0, 1^{n-a_0})}$. By Peel’s theorem (see [[James 1978](#), Theorem 24.1]), $\chi^\circ \in \text{IBr}(G)$. Hence, it is enough to show that $\chi(1) \neq 1$, $\chi(1)$ is p' , and $\chi \in \text{Irr}(B_0(G))$. By the hook length formula, χ has degree

$$\chi(1) = \frac{n!}{n \cdot (a_0 - 1)!(n - a_0)!} = \binom{n - 1}{n - a_0}.$$

Then $\chi(1) \neq 1$ since $1 < a_0 < n$. Moreover

$$n - 1 = a_k p^k + \dots + a_1 p + (a_0 - 1)$$

and

$$n - a_0 = a_k p^k + \dots + a_1 p,$$

so by Lucas' theorem we have

$$\chi(1) = \binom{n-1}{n-a_0} \equiv \prod_{i=1}^k \binom{a_i}{a_i} \binom{a_0-1}{0} \equiv 1 \pmod{p}.$$

Thus, $\chi(1)$ is not divisible by p . Finally since the p -core of $\lambda = (a_0, 1^{n-a_0})$ is (a_0) , we have that χ lies in the principal p -block by the Nakayama conjecture, as desired. Now take $\theta \in \text{Irr}(S)$ under χ , so $\theta \in \text{Irr}_{p'}(B_0(S))$ and $d_{\theta 1_S} = 0$.

Finally, suppose that $a_0 = 1$, so p divides $n - 1$. In this case consider $\chi = \chi^{(n-3, 2, 1)}$, so χ lies in $B_0(G)$. This is the character in the proof of [Navarro and Tiep 2020, Lemma 3.1(ii)], so we are done in this case as well. \square

Groups of Lie type in nondefining characteristic. For groups of Lie type in cross characteristic we consider the following setup. Let \mathbf{G} be a simple linear algebraic group of adjoint type over an algebraically closed field of characteristic r and $F : \mathbf{G} \rightarrow \mathbf{G}$ a Steinberg map, with group of fixed points $G := \mathbf{G}^F$. It is well known that any simple group of Lie type can be obtained as $S = [G, G]$ for \mathbf{G}, F chosen suitably. Moreover, if \mathbf{G}_{sc} denotes a simply connected covering of \mathbf{G} , with corresponding Steinberg map also denoted F , then $S \cong \mathbf{G}_{\text{sc}}^F / \mathbf{Z}(\mathbf{G}_{\text{sc}}^F)$, if S is not the Tits simple group, which was already discussed in Proposition 2.1. Let (\mathbf{G}^*, F) be dual to (\mathbf{G}, F) and $G^* := \mathbf{G}^{*F}$.

We let $\mathbf{B} \leq \mathbf{G}$ denote an F -stable Borel subgroup of \mathbf{G} , and set $B := \mathbf{B}^F$.

We recall that outer automorphisms of prime order $p \geq 5$ of simple groups of Lie type are either field automorphisms, or diagonal automorphisms for groups of types $\text{PSL}_n(\epsilon q)$, with $\epsilon \in \{\pm 1\}$.

Proposition 2.3. *Property (*) holds for S as above if $|B|$ is prime to p .*

Proof. By assumption p divides $|S|$, hence also $|G| = |G^*|$. Let $1 \neq s \in G^*$ be a (semisimple) p -element in the centre of a Sylow p -subgroup of G^* , and let $\chi \in \text{Irr}(G)$ be the semisimple character in the Lusztig series $\mathcal{E}(G, s)$, unique since \mathbf{G} has connected centre, see [Geck and Malle 2020, Definition 2.6.9]. By the degree formula for Jordan decomposition [Geck and Malle 2020, Corollary 2.6.6], $\chi(1)$ is then prime to p , and also $\chi(1) > 1$ as p does not divide $|\mathbf{Z}(G^*)|$ and so $C_{G^*}(s) < G^*$. Furthermore, by [Hiss 1990b, Corollary 3.4], the semisimple character χ lies in the same p -block of G as the semisimple character in $\mathcal{E}(G, 1)$, i.e., the trivial character, so in the principal p -block. Since p does not divide $|B|$, the permutation module 1_B^G is projective, and thus contains the projective cover of the trivial module. But all constituents of 1_B^G are unipotent, so lie in $\mathcal{E}(G, 1)$; see [Geck and Malle 2020, Example 3.2.6]. Hence, by Brauer reciprocity, 1_G° does not occur as a constituent of χ° . Taking for θ any character of S below χ we see that $\theta \in \text{Irr}_{p'}(B_0(S))$ and $d_{\theta 1_S} = 0$.

Next observe that the order of any outer diagonal automorphism of S divides the order of B . Thus, H is an extension of S by a p -power order field automorphism γ . Note that any such automorphism of S extends to G and then also induces a dual automorphism on G^* , which we denote γ^* ; see, e.g., [Taylor 2018, §5.6]. Now choose s more precisely to be also γ^* -invariant (which is possible as γ^* is a p -element, so necessarily has nontrivial fixed points on the centre of a γ^* -stable Sylow p -subgroup of G^*). Then χ is also γ -invariant by [Taylor 2018, Proposition 7.2]. Since the index $|G : S|$ is a divisor of $|B|$, hence prime to p , there exists a γ -invariant $\theta \in \text{Irr}_{p'}(B_0(S))$ below χ , with $d_{\theta|_S} = 0$. Let $\tilde{\theta}$ be an extension of θ to $H = S\langle\gamma\rangle$ in $B_0(H)$. Then $\tilde{\theta}$ is as desired. \square

Proposition 2.4. *Property (*) holds for S as above if F is a Frobenius map with respect to an \mathbb{F}_q -structure and p divides $q - 1$.*

Proof. Let W be the Weyl group of G , that is, the F -fixed points of the Weyl group of G . As p divides $q - 1$, the p -decomposition matrix of the group algebra of W embeds into the p -modular decomposition matrix of G ; see [Dipper 1990, Corollary 4.10]. Let $\epsilon \in \text{Irr}_{p'}(W)$ be the (linear) sign character. Then $\epsilon^\circ \neq 1_W^\circ$ since $p > 2$, whence $d_{\epsilon|_W} = 0$. Now ϵ corresponds to the Steinberg character St of G . Then $d_{\text{St}|_G} = d_{\epsilon|_W} = 0$, St lies in the principal p -block of G (e.g., by [Enguehard 2000, Theorem A]) and its degree is a power of the defining characteristic, so prime to p . Now note that any p -automorphism of S is realised inside the extension of G (which induces all diagonal automorphisms) by a generator γ of the cyclic group of p -power order field automorphisms, so we may assume $H \leq \tilde{G} := G\langle\gamma\rangle$. By [Geck and Malle 2020, Theorem 4.5.11], St is invariant under γ . Let $\tilde{\text{St}}$ be an extension of St to \tilde{G} in $B_0(\tilde{G})$, so $d_{\tilde{\text{St}}|\tilde{G}} = 0$. Then $\tilde{\text{St}}|_H$ is irreducible, since St restricts irreducibly to S , hence lies in $B_0(H)$ and so is as required. \square

Theorem 2.5. *Property (*) holds for S of Lie type when p is not the defining characteristic.*

Proof. By Proposition 2.3 we may assume that p divides the order of a Borel subgroup B of G . If F is a Frobenius map with respect to an \mathbb{F}_q -structure and p divides $q - 1$, we are done by Proposition 2.4. If G is a Suzuki or Ree group and p divides $|B|$, then $p | (q^2 - 1)$ where F^2 defines an \mathbb{F}_{q^2} -structure, and the exact same arguments as in the proof of Proposition 2.4 apply.

We are reduced to the case that F is a Frobenius map with respect to an \mathbb{F}_q -structure and p divides $|B|$ but not $q - 1$. Since p is not the defining prime, this implies that G is a twisted group of Lie type ${}^2A_{n-1}$, 2D_n , or 2E_6 and p divides $q + 1$, respectively of type 3D_4 and p divides $q^2 + q + 1$. Let $d = 2, 3$ in the respective cases. Then the centraliser of a Sylow d -torus of G is a maximal torus, so has a unique d -cuspidal unipotent character. Thus, by [Enguehard 2000, Theorem A] there is a unique unipotent block of G of maximal defect, the principal block, which

hence contains the Steinberg character St of G . By [Hiss 1990a, Theorem B] we have $d_{\text{St}_1G} = 0$ in our case unless $G = \text{PGU}_3(q)$. Except for that latter case, we can now argue as in the proof of Proposition 2.4 to conclude. For $G = \text{PGU}_3(q)$ let χ be the cuspidal unipotent character of degree $q(q-1)$, prime to p . By [Geck 1990, Theorem 4.3(a)] it lies in $B_0(G)$ and satisfies $d_{\chi_{1G}} = 0$. Again, χ restricts irreducibly to S and is invariant under all automorphisms, so we can argue as before. \square

Groups of Lie type in defining characteristic. We do not see how to approach Property (*) for groups of Lie type in their defining characteristic in general. All characters of positive defect lie in the principal block and decomposition numbers tend to be large and little is known.

Proposition 2.6 (Navarro and Tiep). *Property (*) holds if $S = \text{PSp}_{2n}(p^f)$, $n \geq 1$, with either $p > 3$ or $p = 3$ and n even.*

Proof. In this case the principal block of S is the only p -block of positive defect. Let H be almost simple with H/S a p -group. By [Navarro 1998, Corollary 9.6] there is just one p -block of H covering $B_0(S)$, necessarily the principal block $B_0(H)$. Now the irreducible character $\chi \in \text{Irr}_{p'}(H)$ with $d_{\chi_{1H}} = 0$ constructed in [Navarro and Tiep 2020, Proposition 3.11] lies above a character of S of positive defect, hence in the principal p -block of H and we are done. \square

Observe that this does not extend to $p = 3$ and n odd: the group $H = \text{PSp}_2(3^3)$ has no irreducible character $\chi \in \text{Irr}_{3'}(H)$ with $d_{\chi_{1H}} = 0$.

[Navarro and Tiep 2022] also contains results for special linear and unitary groups but these are not applicable here as the considered characters are not of p' -degree. Nevertheless, we can follow their general approach.

For $G = \text{SL}_n(q)$ we let τ_j , $j = 1, \dots, q-2$ denote the nonunipotent Weil characters of degree $(q^n - 1)/(q - 1)$, ordered so that τ_j is trivial on the centre $Z(\text{SL}_n(q))$ of order $z := \gcd(n, q - 1)$ if and only if $z \mid j$.

Proposition 2.7. *Let $S = \text{PSL}_n(q)$ with $q = p^f$, $p \neq 2$ and $n \geq 3$.*

- (a) *If either $\gcd(p - 1, (q - 1)/\gcd(n, q - 1)) > 1$ or $2^f < (q - 1)/z - 1$ then there is $\chi \in \text{Irr}_{p'}(B_0(S))$ such that $d_{\chi_{1S}} = 0$.*
- (b) *Write $f = p^a f'$ with $\gcd(p, f') = 1$ and set $q' := p^{f'}$. If*

$$a = 0 \quad \text{or} \quad 2^f < (q' - 1)/\gcd(q' - 1, n) - 1$$

then Property () holds for $S = \text{PSL}_n(q)$.*

Proof. Let $G := \text{SL}_n(q)$ and set $z = \gcd(n, q - 1)$. We are interested in the characters τ_j of G that are trivial on $Z(G)$, that is, for which $j = zj'$ for some integer $1 \leq j' \leq (q - 1 - z)/z$. By [Zaleski and Suprunenko 1990, Theorem 1.11]

the decomposition number $d_{\tau_j 1_G}$ equals the number of solutions $x_s \in \{0, 1\}$ of the congruence

$$n(p-1) \sum_{s=0}^{f-1} x_s p^s \equiv j \pmod{(q-1)}.$$

Dividing by z , we need to count solutions to

$$\frac{n}{z(p-1)} \sum_{s=0}^{f-1} x_s p^s \equiv j' \pmod{(q-1)/z}.$$

If there is a prime ℓ dividing $p-1$ and $(q-1)/z$, then reducing modulo ℓ we see there is no solution for $j' = 1$. Also, the left hand side can take at most 2^f distinct values. Since there are $(q-1)/z - 1$ admissible values for j' , there is j' with no solutions whenever $2^f < (q-1)/z - 1$. Thus, under either of our assumptions we find j' with $\tau_j = \tau_{zj'} \in \text{Irr}_{p'}(G)$ with $d_{\tau_j 1_G} = 0$. Since G has a single p -block of positive defect, τ_j lies in the principal block. Furthermore, by construction $Z(\text{SL}_n(q))$ lies in the kernel of τ_j and hence τ_j deflates to a character of $S = \text{PSL}_n(q)$. Thus we get (a).

(b) Since $p > 2$ does not divide $q-1$, the p -power order automorphisms of S are field automorphisms, of order dividing p^a where $f = p^a f'$ is as in the statement. If $a = 0$, **Property (*)** follows from (a) since necessarily $H = S$. For $a > 0$ let γ be a field automorphism of G (and hence of S) of order p^a . There are exactly $q' - 2$ Weil characters of G invariant under γ , which hence extend to $G\langle\gamma\rangle$. Of these, $(q' - 1)/\text{gcd}(q' - 1, n) - 1$ are trivial on $Z(G)$, so define characters in $\text{Irr}_{p'}(S)$ invariant under γ . By the argument above, if this number is bigger than 2^f then there exists such a character χ with $d_{\chi 1_S} = 0$. Hence any character of H in $B_0(H)$ lying above it verifies **Property (*)**. \square

Note that the case $n = 2$ is addressed in **Proposition 2.6**. Observe that the condition in **Proposition 2.7(a)** is satisfied if $\text{gcd}(n, q-1) = 1$, for example. It also holds when $p > n+1$ (since $p-1$ always divides $q-1$), or if $q > n(2^f + 1)$. Thus, **Proposition 2.7** extends and complements [Navarro and Tiep 2022, Proposition 3.3(ii)].

Corollary 2.8. *Let $S = \text{PSL}_n(q)$ with $q = p^f$, $p \neq 2$ and $3 \leq n \leq 9$. Then there is $\chi \in \text{Irr}_{p'}(B_0(S))$ with $d_{\chi 1_S} = 0$ unless possibly S is one of*

$$\text{PSL}_4(5), \text{PSL}_6(3), \text{PSL}_6(7), \text{PSL}_8(3), \text{PSL}_8(9), \text{PSL}_8(5), \text{PSL}_8(25).$$

Proof. For the groups $\text{PSL}_n(q)$, $n \leq 9$, considered here, either the conditions in **Proposition 2.7(a)** are satisfied, or if not, a direct checking with the Zalesski–Suprunenko formula shows the claim, except for the groups listed in the conclusion and for $\text{PSL}_4(3)$. The decomposition matrix of $\text{PSL}_4(3)$ is available in [GAP 2020] from which the claim can be verified for that group. \square

For $G = \mathrm{SU}_n(q)$ we let τ_j , $j = 1, \dots, q$, denote the nonunipotent Weil characters, constructed by Seitz [1975], of degree $(q^n - (-1)^n)/(q + 1)$, again ordered so that τ_j is trivial on the centre $Z(\mathrm{SU}_n(q))$ of order $z := \gcd(n, q + 1)$ if and only if $z | j$.

Proposition 2.9. *Let $S = \mathrm{PSU}_n(q)$ with $q = p^f$, $p \neq 2$ and $n \geq 3$.*

- (a) *If $2^f < (q + 1)/z - 1$ then there is $\chi \in \mathrm{Irr}_{p'}(B_0(S))$ such that $d_{\chi 1_S} = 0$.*
 (b) *Write $f = p^a f'$ with $\gcd(p, f') = 1$ and set $q' := p^{f'}$. If*

$$a = 0 \quad \text{or} \quad 2^f < (q' + 1)/\gcd(q' + 1, n) - 1$$

then Property () holds for $S = \mathrm{PSU}_n(q)$.*

Proof. The argument is very similar to the one for the special linear groups. Let $G = \mathrm{SU}_n(q)$ and set $z = \gcd(n, q + 1) = |Z(G)|$. Again, we consider characters τ_j trivial on $Z(G)$, that is, for which $j = zj'$ for some integer $1 \leq j' \leq (q + 1 - z)/z$. By [Zaleski 1990, Main Theorem] the decomposition number $d_{\tau_j 1_G}$ equals the number of solutions $x_s \in \{0, 1\}$ of the congruence

$$n \left((p - 1) \sum_{s=0}^{f-1} x_s p^s - 1 \right) \equiv j \pmod{(q + 1)}.$$

(In fact, [Zaleski 1990] has an additional summand of $\frac{1}{2}(q + 1)$ on the right-hand side, but this disappears here due to a different numbering of the τ_j , see [Navarro and Tiep 2022, p. 612]; in any case, this difference will not matter for our argument here.) Since the left-hand side can take at most 2^f distinct values, while there are $(q + 1)/z - 1$ admissible values for j' the assertion in (a) follows. For (b) we can argue exactly as in the proof of Proposition 2.7. \square

As in [Navarro and Tiep 2020; 2022] we have no general results for orthogonal or exceptional type groups in their defining characteristic.

3. The reduction

In this section we prove Theorem A. We will need the following results, which we collect here for the reader's convenience.

Lemma 3.1 [Murai 1994, Lemma 4.3]. *Let $N \triangleleft G$ and let $\theta \in \mathrm{Irr}_{p'}(B_0(N))$. Suppose that θ extends to PN , where $P \in \mathrm{Syl}_p(G)$. Then there exists $\chi \in \mathrm{Irr}_{p'}(B_0(G))$ satisfying $[\theta^G, \chi] \neq 0$.*

The following argument is inside the proof of [Navarro and Tiep 2020, Theorem 2.6].

Lemma 3.2. *Let G be a finite group and suppose that $d_{\chi 1_G} \neq 0$ for every $\chi \in \mathrm{Irr}_{p'}(B_0(G))$. Let $M \triangleleft G$ and let $P \in \mathrm{Syl}_p(G)$. Then $d_{\psi 1_{MP}} \neq 0$ for every $\psi \in \mathrm{Irr}_{p'}(B_0(MP))$.*

Proof. Since MP/M is a p -group, we have that $\psi_M = \tau \in \text{Irr}_{p'}(B_0(M))$. By [Lemma 3.1](#) there exists $\chi \in \text{Irr}_{p'}(B_0(G))$ lying over τ . By hypothesis, $d_{\chi 1_G} \neq 0$, and then χ_M° contains 1_M , so $d_{\tau 1_M} \neq 0$. Since MP/M is a p -group, we have that $(MP)^\circ = M^\circ$ and then $d_{\psi 1_{MP}} \neq 0$, as wanted. \square

We next prove [Theorem A](#) for p -solvable groups.

Theorem 3.3. *Let G be a p -solvable group. Then the following are equivalent:*

- (i) *For every $\chi \in \text{Irr}_{p'}(B_0(G))$ we have $d_{\chi 1_G} \neq 0$.*
- (ii) *For every $\chi \in \text{Irr}_{p'}(B_0(G))$ we have $d_{\chi 1_G} = 1$.*
- (iii) *For $P \in \text{Syl}_p(G)$ we have $N_G(P) = P \times K$ for some $K \leq G$.*

Proof. We first prove (iii) implies (ii). By [\[Navarro et al. 2014, Theorem 3.2\]](#) we have $K \subseteq \mathbf{O}_{p'}(G) =: X$. Now, let $\bar{G} = G/X$ and $\bar{P} = PX/X$, then (iii) implies that $N_{\bar{G}}(\bar{P}) \cong \bar{P}$. Since $\text{Irr}_{p'}(B_0(G)) = \text{Irr}_{p'}(B_0(G/X))$ we know by [\[Navarro and Tiep 2020, Theorem B\]](#) that (i) and (ii) hold.

Since (ii) implies (i) trivially, we just need to show that (i) implies (iii). We proceed by induction on $|G|$. Let $N = \mathbf{O}_{p'}(G)$ and use the bar notation. Since $\text{Irr}_{p'}(B_0(G)) = \text{Irr}_{p'}(B_0(\bar{G}))$, if $N > 1$, we have by induction that $N_{\bar{G}}(\bar{P}) = \bar{P} \times \bar{K}$. By [\[Navarro et al. 2014, Theorem 3.2\]](#) we have that $\bar{K} \subseteq \mathbf{O}_{p'}(\bar{G}) = 1$ so $\bar{K} = 1$ and hence $N_{\bar{G}}(\bar{P}) = \bar{P}$. This implies that $N_G(P) = P \times C_N(P)$ and we are done. So we may assume that $N = 1$. But in this case the principal p -block is the only p -block of G , and we are done by [\[Navarro and Tiep 2020, Theorem B\]](#). \square

We finally prove [Theorem A](#).

Theorem 3.4. *Let G be a finite group. Assume that $p > 3$ and all nonabelian composition factors of G of order divisible by p satisfy [Property \(*\)](#). Then the following are equivalent:*

- (i) *For every $\chi \in \text{Irr}_{p'}(B_0(G))$ we have $d_{\chi 1_G} \neq 0$.*
- (ii) *For every $\chi \in \text{Irr}_{p'}(B_0(G))$ we have $d_{\chi 1_G} = 1$.*
- (iii) *For $P \in \text{Syl}_p(G)$ we have $N_G(P) = P \times K$ for some $K \leq G$.*

Proof. Since [\[Navarro et al. 2014, Theorem 3.2\]](#) holds for odd primes, we can argue as in the first part of the proof of [Theorem 3.3](#) to show that (iii) \implies (ii) \implies (i) (notice that, as happens in [\[Navarro and Tiep 2020\]](#), (iii) \implies (ii) \implies (i) is always true if $p > 3$, with no extra conditions on the composition factors of G). So we just need to prove that (i) implies (iii). We work by induction on $|G|$. Arguing as in the second paragraph of the proof of [Theorem 3.3](#) we may then assume that $\mathbf{O}_{p'}(G) = 1$.

Let $M \triangleleft G$ be the largest p -solvable normal subgroup of G . We claim that $M = 1$. Let $\bar{G} = G/M$ and use the bar notation. Suppose $M > 1$. Since $\text{Irr}_{p'}(B_0(\bar{G}))$ is

contained in $\text{Irr}_{p'}(B_0(G))$ we have by induction that $N_{\bar{G}}(\bar{P}) = \bar{P} \times \bar{K}$. By [Navarro and Tiep 2020, Theorem 3.2] this implies that $\bar{K} \subseteq \mathbf{O}_{p'}(\bar{G})$. Since M is the largest normal p -solvable subgroup of G , $\mathbf{O}_{p'}(\bar{G})$ is trivial and hence $N_{\bar{G}}(\bar{P}) = \bar{P}$. By [Guralnick et al. 2004, Theorem 1.1] this forces \bar{G} to be solvable. Hence G is p -solvable and we are done by Theorem 3.3.

Let N be a minimal normal subgroup of G ; thus $N = S_1 \times \cdots \times S_t$ where $S_i \cong S$ is a nonabelian simple group of order divisible by p and the S_i are transitively permuted by G . Let $X/N = \mathbf{O}_{p'}(G/N)$ and let $H = \bigcap N_G(S_i)$. We claim that $G = (H \cap X)P$. Write $Y = (H \cap X)P$. We first show that $\mathbf{O}_{p'}(Y) = 1$. Indeed, since $\mathbf{O}_{p'}(G)$, we have that $\mathbf{O}_{p'}(Y) \cap (H \cap X) \subseteq \mathbf{O}_{p'}(H \cap X) = 1$. Now, $\mathbf{O}_{p'}(Y) \cong (H \cap X)\mathbf{O}_{p'}(Y)/(H \cap X)$ is a p -group, so $\mathbf{O}_{p'}(Y) = 1$ as wanted. By Lemma 3.2 applied to $H \cap X$ in place of M , we have that Y satisfies (i). Since the nonabelian composition factors of Y are composition factors of G , if $Y < G$, by induction this gives $N_Y(P) = P \times K$ for some K . But then $K \subseteq \mathbf{O}_{p'}(Y)$ by [Navarro et al. 2014, Theorem 3.2], so $K = 1$. This means that $N_Y(P) = P$ and then by [Guralnick et al. 2004, Theorem 1.1] the group Y is solvable. But then N is solvable, a contradiction. Hence $Y = G$, as wanted.

Since $G = (H \cap X)P$ and H acts trivially on $\{S_1, \dots, S_t\}$, P must act transitively on the set $\{S_1, \dots, S_t\}$. Write $S = S_1$ and, for $i = 2, \dots, t$, write $S_i = S^{x_i}$ with $x_i \in P$. We proceed now as in the proof of [Navarro and Tiep 2020, Theorem 2.6]. Let $R = N_P(S)$. If $SR = G$, then $S = N$ and $C_G(S)$ is a normal p -subgroup of G . Since there are no nontrivial p -solvable normal subgroups of G , we conclude that $C_G(S) = 1$ and hence G is almost simple with socle S and $|G : S|$ is a power of p . Since S satisfies Property (*) by assumption, we have a contradiction. Hence we may assume that $SR < G$.

Let $Q = P \cap N$ and let $R_1 = R \cap S = Q \cap S = P \cap S \in \text{Syl}_p(S)$. Let $\gamma \in \text{Irr}_{p'}(B_0(SR))$ and notice that $\gamma_S = \psi \in \text{Irr}_{p'}(B_0(S))$ since SR/S is a p -group. For $i = 2, \dots, t$, let $\psi_i = \psi^{x_i} \in \text{Irr}_{p'}(B_0(S_i))$ and let $\eta = \psi \times \psi_2 \times \cdots \times \psi_t \in \text{Irr}_{p'}(B_0(N))$, which is P -invariant by [Navarro et al. 2007, Lemma 4.1(ii)]. Then η extends to PN . By Lemma 3.1 and hypothesis we have $d_{\eta_{1_N}} \neq 0$. By [Navarro and Tiep 2020, Lemma 2.3] we have that $d_{\psi_{1_S}} \neq 0$ and then, since SR/S is a p -group, we conclude that $d_{\gamma_{1_{SR}}} \neq 0$. Since $SR < G$ and S is a composition factor of G , this implies $N_{SR}(R) = R \times K$ for some K . Then $K \subseteq \mathbf{O}_{p'}(SR)$. Arguing as before, we have that $\mathbf{O}_{p'}(SR) = 1$, so $K = 1$ and then $N_{SR}(R) = R$. Now by [Guralnick et al. 2004, Theorem 1.1], SR is solvable, and hence S is solvable, which is our final contradiction. \square

Acknowledgements

Rizo would like to thank Gabriel Navarro for bringing this topic to her attention.

References

- [Dipper 1990] R. Dipper, “On quotients of Hom-functors and representations of finite general linear groups, I”, *J. Algebra* **130**:1 (1990), 235–259. [MR](#) [Zbl](#)
- [Enguehard 2000] M. Enguehard, “Sur les l -blocs unipotents des groupes réductifs finis quand l est mauvais”, *J. Algebra* **230**:2 (2000), 334–377. [MR](#) [Zbl](#)
- [GAP 2020] The GAP Group, “GAP: groups, algorithms, and programming”, 2020, available at <http://www.gap-system.org>. Version 4.11.0.
- [Geck 1990] M. Geck, “Irreducible Brauer characters of the 3-dimensional special unitary groups in nondefining characteristic”, *Comm. Algebra* **18**:2 (1990), 563–584. [MR](#) [Zbl](#)
- [Geck and Malle 2020] M. Geck and G. Malle, *The character theory of finite groups of Lie type: a guided tour*, Cambridge Stud. Adv. Math. **187**, Cambridge Univ. Press, 2020. [MR](#) [Zbl](#)
- [Guralnick et al. 2004] R. M. Guralnick, G. Malle, and G. Navarro, “Self-normalizing Sylow subgroups”, *Proc. Amer. Math. Soc.* **132**:4 (2004), 973–979. [MR](#) [Zbl](#)
- [Hiss 1990a] G. Hiss, “The number of trivial composition factors of the Steinberg module”, *Arch. Math. (Basel)* **54**:3 (1990), 247–251. [MR](#) [Zbl](#)
- [Hiss 1990b] G. Hiss, “Regular and semisimple blocks of finite reductive groups”, *J. Lond. Math. Soc. (2)* **41**:1 (1990), 63–68. [MR](#) [Zbl](#)
- [James 1978] G. D. James, *The representation theory of the symmetric groups*, Lecture Notes in Math. **682**, Springer, 1978. [MR](#) [Zbl](#)
- [Kessar and Malle 2013] R. Kessar and G. Malle, “Quasi-isolated blocks and Brauer’s height zero conjecture”, *Ann. of Math. (2)* **178**:1 (2013), 321–384. [MR](#) [Zbl](#)
- [Murai 1994] M. Murai, “Block induction, normal subgroups and characters of height zero”, *Osaka J. Math.* **31**:1 (1994), 9–25. [MR](#) [Zbl](#)
- [Navarro 1998] G. Navarro, *Characters and blocks of finite groups*, Lond. Math. Soc. Lect. Note Ser. **250**, Cambridge Univ. Press, 1998. [MR](#) [Zbl](#)
- [Navarro and Tiep 2020] G. Navarro and P. H. Tiep, “Decomposition numbers and local properties”, *J. Algebra* **558** (2020), 620–639. [MR](#) [Zbl](#)
- [Navarro and Tiep 2022] G. Navarro and P. H. Tiep, “Decomposition numbers and global properties”, *J. Algebra* **607** (2022), 607–617. [MR](#) [Zbl](#)
- [Navarro et al. 2007] G. Navarro, P. H. Tiep, and A. Turull, “ p -rational characters and self-normalizing Sylow p -subgroups”, *Represent. Theory* **11** (2007), 84–94. [MR](#) [Zbl](#)
- [Navarro et al. 2014] G. Navarro, P. H. Tiep, and C. Vallejo, “McKay natural correspondences on characters”, *Algebra Number Theory* **8**:8 (2014), 1839–1856. [MR](#) [Zbl](#)
- [Seitz 1975] G. M. Seitz, “Some representations of classical groups”, *J. Lond. Math. Soc. (2)* **10** (1975), 115–120. [MR](#) [Zbl](#)
- [Taylor 2018] J. Taylor, “Action of automorphisms on irreducible characters of symplectic groups”, *J. Algebra* **505** (2018), 211–246. [MR](#) [Zbl](#)
- [Zalesski 1990] A. E. Zalesskiĭ, “A fragment of the decomposition matrix of the special unitary group over a finite field”, *Izv. Akad. Nauk SSSR Ser. Mat.* **54**:1 (1990), 26–41. In Russian; translated in *Math. USSR-Izv.* **36** (1991), 23–39. [MR](#) [Zbl](#)
- [Zalesski and Suprunenko 1990] A. E. Zalesskii and I. D. Suprunenko, “Permutation representations and a fragment of the decomposition matrix of symplectic and special linear groups over a finite field”, *Sibirsk. Mat. Zh.* **31**:5 (1990), 46–60. In Russian; translated in *Siberian Math. J.* **31**:5 (1990), 744–755. [MR](#) [Zbl](#)

Received May 21, 2024. Revised July 19, 2024.

GUNTER MALLE
FACHBEREICH MATHEMATIK
RPTU KAISERSLAUTERN
KAISERSLAUTERN
GERMANY

malle@mathematik.uni-kl.de

NOELIA RIZO
DEPARTAMENT DE MATEMÀTIQUES
UNIVERSITAT DE VALÈNCIA
BURJASSOT
SPAIN

noelia.rizo@uv.es

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Fakultät für Mathematik
Universität Wien
Vienna, Austria
matthias.aschenbrenner@univie.ac.at

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Atsushi Ichino
Department of Mathematics
Kyoto University
Riverside, CA 92521-0135
atsushi.ichino@gmail.com

Robert Lipshitz
Department of Mathematics
University of Oregon
Eugene, OR 97403
lipshitz@uoregon.edu

Kefeng Liu
School of Sciences
Chongqing University of Technology
Chongqing 400054, China
liu@math.ucla.edu

Sucharit Sarkar
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
sucharit@math.ucla.edu

Dimitri Shlyakhtenko
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
shlyakht@ipam.ucla.edu

Ruixiang Zhang
Department of Mathematics
University of California
Berkeley, CA 94720-3840
ruixiang@berkeley.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2025 is US \$677/year for the electronic version, and \$917/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2025 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 336 No. 1-2 May 2025

In memoriam Gary Seitz

Gary Seitz (1943–2023): In memoriam	1
MARTIN W. LIEBECK, GERHARD RÖHRLE and DONNA TESTERMAN	
Intrinsic components in involution centralizers of fusion systems	7
MICHAEL ASCHBACHER	
On good A_1 subgroups, Springer maps, and overgroups of distinguished unipotent elements in reductive groups	29
MICHAEL BATE, SÖREN BÖHM, BENJAMIN MARTIN and GERHARD RÖHRLE	
The q -Schur category and polynomial tilting modules for quantum GL_n	63
JONATHAN BRUNDAN	
The binary actions of simple groups of Lie type of characteristic 2	113
NICK GILL, PIERRE GUILLOT and MARTIN W. LIEBECK	
Finite simple groups have many classes of p -elements	137
MICHAEL GIUDICI, LUKE MORGAN and CHERYL E. PRAEGER	
Monogamous subvarieties of the nilpotent cone	161
SIMON M. GOODWIN, RACHEL PENGELLY, DAVID I. STEWART and ADAM R. THOMAS	
An extension of Gow's theorem	181
ROBERT M. GURALNICK and PHAM HUU TIEP	
On dimensions of RoCK blocks of cyclotomic quiver Hecke superalgebras	191
ALEXANDER KLESHCHEV	
Representation growth of Fuchsian groups and modular forms	217
MICHAEL J. LARSEN, JAY TAYLOR and PHAM HUU TIEP	
D_4 -type subgroups of $F_4(q)$	249
R. LAWThER	
Constructible representations and Catalan numbers	339
GEORGE LUSZTIG and ERIC SOMMERS	
A reduction theorem for simple groups with $e(G) = 3$	351
RICHARD LYONS and RONALD SOLOMON	
Decomposition numbers in the principal block and Sylow normalisers	367
GUNTER MALLE and NOELIA RIZO	
Levi decompositions of linear algebraic groups and nonabelian cohomology	379
GEORGE J. MCNINCH	
On the intersection of principal blocks	399
GABRIEL NAVARRO, A. A. SCHAEFFER FRY and PHAM HUU TIEP	
Hesselink strata in small characteristic and Lusztig–Xue pieces	415
ALEXANDER PREMÉT	
Multiplicity-free representations of the principal A_1 -subgroup in a simple algebraic group	433
ALUNA RIZZOLI and DONNA TESTERMAN	