

*Pacific
Journal of
Mathematics*

**REPRODUCING KERNEL FUNCTIONS
AND BOUNDARY LIMITS IN MODEL SPACES**

YONGJIANG DUAN, YUFEI LI AND JAVAD MASHREGHI

REPRODUCING KERNEL FUNCTIONS AND BOUNDARY LIMITS IN MODEL SPACES

YONGJIANG DUAN, YUFEI LI AND JAVAD MASHREGHI

We investigate the boundary behavior of functions in model spaces, emphasizing their approach towards boundary points through arbitrary but fixed regions. We generalize classical results to encompass *any* approach region, thereby resolving an open question about reproducing kernel functions. The study connects the geometric nature of approach regions to the analytic properties of functions in the model space as they approach the unit circle. Key conditions are established to fully characterize boundedness, weak convergence, and the existence of limits within the approach region for functions in these spaces. Furthermore, precise criteria for norm convergence of reproducing kernel functions are provided, offering deeper insights into how boundary geometry influences analytic behavior. These results extend classical boundary value problems to an ultimately broader framework, highlighting the interplay between geometry and analysis.

1. Introduction

Let \mathbb{D} denote the open unit disk of the complex plane \mathbb{C} , and H^2 denote the Hardy space on \mathbb{D} . The space H^2 can also be identified with a closed subspace of $L^2(\mathbb{T})$ via nontangential limits. A bounded analytic function u on the unit disk is said to be inner if $|u(\xi)| = 1$ a.e. on \mathbb{T} . It is well known [2] that every inner function u admits the factorization

$$(1-1) \quad u(z) = cB(z)s(z)\Delta(z)$$

with

$$B(z) = \prod_{n \geq 1} \frac{\bar{a}_n}{|a_n|} \frac{a_n - z}{1 - \bar{a}_n z},$$

$$s(z) = \exp \left\{ - \int_0^{2\pi} \frac{e^{i\theta} + z}{e^{i\theta} - z} d\sigma(\theta) \right\},$$

$$\Delta(z) = \exp \left\{ - \sum_{n \geq 1} \rho_n \frac{e^{i\zeta_n} + z}{e^{i\zeta_n} - z} \right\},$$

MSC2020: 30J05, 46E22.

Keywords: model space, boundary behavior, inner function.

where c is a unimodular constant, $\{a_n\}$ is a Blaschke sequence which means it satisfies $\sum (1 - |a_n|) < \infty$, (we assume $a_n/|a_n| \equiv 1$ if $a_n = 0$), σ is a finite positive continuous singular measure with respect to Lebesgue measure, and $\zeta_n \in [0, 2\pi)$, $\rho_n > 0$, $\sum \rho_n < \infty$.

For each nonconstant inner function u , define the model space $K_u = H^2 \ominus uH^2$, which is a reproducing kernel Hilbert space whose kernel has the form

$$k_\lambda^u(z) = \frac{1 - \overline{u(\lambda)}u(z)}{1 - \bar{\lambda}z}, \quad \lambda, z \in \mathbb{D}.$$

In particular, when $u = B$, let $B_1 = 1$, $B_n(z) = \prod_{1 \leq i \leq n-1} (a_i - z)/(1 - \bar{a}_i z)$, $n = 2, 3, \dots$, and

$$(1-2) \quad e_n(z) = \frac{\sqrt{1 - |a_n|^2}}{1 - \bar{a}_n z} B_n(z), \quad n \geq 1.$$

Then $\{e_n\}$ is an orthonormal basis for K_B . Such a basis is the so-called Takenaka–Malmquist–Walsh basis (see [9, Proposition 5.25]).

For $\xi \in \mathbb{T}$ and $\alpha > 1$, the Stolz domain at ξ is defined by

$$\Gamma_\alpha(\xi) = \{z \in \mathbb{D} : |z - \xi| < \alpha(1 - |z|)\}.$$

A function $f : \mathbb{D} \rightarrow \mathbb{C}$ is said to have a nontangential limit at ξ if $\lim_{z \rightarrow \xi} f(z)$ exists whenever z converges to ξ in every fixed Stolz domain $\Gamma_\alpha(\xi)$. The function f is said to have angular derivative in the sense of Carathéodory (ADC) at the point $\xi \in \mathbb{T}$, denoted by $f'(\xi)$, if the difference quotient $(f(z) - \eta)/(z - \xi)$ has a nontangential limit for some $\eta \in \mathbb{T}$. This is the celebrated Julia–Carathéodory theorem as follows [5; 11].

Theorem A. *For $b : \mathbb{D} \rightarrow \mathbb{D}$ analytic and $\xi \in \mathbb{T}$, the following are equivalent:*

- (1) $\liminf_{z \rightarrow \xi} (1 - |b(z)|)/(1 - |z|) < \infty$, where the limit is taken as z tends to ξ unrestrictedly in \mathbb{D} .
- (2) b has an ADC at ξ .
- (3) Both b and b' have (finite) nontangential limits at ξ , with $|\lim_{r \rightarrow 1} b(r\xi)| = 1$.

There are many extensions for this result to higher dimensions for holomorphic mappings of several complex variables; see, e.g., [1; 4; 13; 14; 16]. In particular, if b is an inner function, due to Ahern and Clark, the Julia–Carathéodory theorem can be more accurately described as follows (see [3] or see [9, Theorem 7.24]).

Theorem B. *Let u be an inner function factorized according to (1-1). Then the following are equivalent:*

- (i) u has an ADC at ξ .
- (ii) $\liminf_{z \rightarrow \xi} \|k_z^u\|^2 = \liminf_{z \rightarrow \xi} (1 - |u(z)|)/(1 - |z|) < \infty$.
- (iii) For every fixed Stolz domain $\Gamma_\alpha(\xi)$ at ξ , $\sup\{\|k_z^u\| : z \in \Gamma_\alpha(\xi)\} < \infty$.

- (iv) Each $f \in K_u$ has a nontangential limit at $\xi \in \mathbb{T}$.
- (v)

$$(1-3) \quad \sum_{n \geq 1} \frac{1 - |a_n|}{|\xi - a_n|^2} + \int_0^{2\pi} \frac{d\sigma(\theta)}{|\xi - e^{i\theta}|^2} + \sum_{n \geq 1} \frac{\rho_n}{|\xi - e^{i\zeta_n}|^2} < \infty.$$

Moreover, when these conditions hold, $k_\xi^u = (1 - \overline{u(\xi)}u)/(1 - \bar{\xi}z) \in K_u$ and $u'(\xi) = \bar{\xi}u(\xi)|u'(\xi)|$, where $u(\xi)$ and $u'(z)$ are the nontangential limits of $u(z)$ and $u'(z)$ at ξ , respectively, and $k_z^u \rightarrow k_\xi^u$ in the norm of K_u as z tends to ξ nontangentially from \mathbb{D} .

To extend **Theorem B** to a broader context of convergence, let Ω_ξ be an *approach region* with vertex at $\xi \in \mathbb{T}$. By the term ‘‘approach region’’, we mean an open connected subset of \mathbb{D} whose closure intersects \mathbb{T} only at ξ . For example, an approach region might be the interior of a triangle whose vertex is at ξ (a Stolz approach region), or the interior of an internally tangent circle with contact point ξ (an oricyclic approach region), or perhaps one might want to consider other tangential approach regions. In fact, Hartmann and Ross [10, Theorem 4.1] gave an alternate proof of (iii) \iff (iv) of **Theorem B**, which can be used to prove the following result: for any approach region Ω_ξ , every $f \in K_u$ has a limit as $z \rightarrow \xi$ within $z \in \Omega_\xi$ if and only if $\sup\{\|k_z^u\| : z \in \Omega_\xi\} < \infty$. Therefore, it is natural to explore the following question.

Question 1.1. Is there a condition, similar to Frostman’s condition (1-3), that one could exploit to determine whether or not $\sup\{\|k_z^u\| : z \in \Omega_\xi\} < \infty$?

This question, posed in [9, pp. 168–169], serves as the foundation for our investigation in this work. The structure of the paper is as follows. In **Section 2**, we outline the origin of our approach and present the main results, **Theorems 2.1** and **2.2**. The first theorem extends **Theorem B** to the context of weak convergence, but crucially, it applies to arbitrary approach regions — addressing the central objective of this work. The second theorem provides a complete characterization of norm convergence, thereby fully resolving **Question 1.1**. **Sections 3** and **4** include several technical lemmas that are noteworthy in their own right and also serve as essential components for the respective proofs of the main results. Finally, in **Section 5**, we explore deeper into the convergence results within two specific approach regions, further illustrating the robustness and applicability of the main theorems.

2. Our approach and the main results

For $c > 0$ and $\gamma \geq 1$, define

$$R(c, \gamma) := \{z \in \mathbb{D} \setminus \{0\} : 1 - |z| \geq c|\arg z|^\gamma\} \quad (\text{see Figure 1}),$$

$$\Gamma_{c, \gamma} := \{z \in \mathbb{D} \setminus \{0\} : 1 - |z| > c|1 - z|^\gamma\},$$




Figure 1. $R(1, 1)$ (left), $R(1, 2)$ (middle) and $R(1, 3)$ (right).

where $\lfloor \arg z \rfloor$ means the length of the shorter one of the two arcs on \mathbb{T} joining $z/|z|$ and 1. If $z \in \mathbb{D}$ with $\operatorname{Re} z > 0$ and $|z| > \frac{1}{2}$, then $\lfloor \arg z \rfloor \in [0, \pi/2]$ and hence $\sin \arg z \sim \lfloor \arg z \rfloor$, so that

$$|1 - z|^\gamma \sim (1 - |z|)^\gamma + \lfloor \arg z \rfloor^\gamma,$$

which implies that there exist two positive numbers $c_1(\gamma)$ and $c_2(\gamma)$, such that

$$\Gamma_{c_1(\gamma), \gamma} \subset R(c, \gamma) \subset \Gamma_{c_2(\gamma), \gamma}.$$

A function f defined on \mathbb{D} is said to have a T_γ -limit at 1 if f has a limit as $z \rightarrow 1$ within $R(c, \gamma)$ for every $c > 0$. In particular, f has a T_1 -limit at 1 if and only if f has a nontangential limit at 1. For $\gamma > 1$, Protas [15, Theorem 3, Example 2] gave a sufficient condition that each $f \in K_u$ has a T_γ -limit at 1 and showed that this condition is not necessary for some model space K_B with a Blaschke product B . Leung [12, Theorem 4] proved the following result concerned with existence of T_γ -limits of functions in K_u .

Theorem C. *Let u be an inner function factorized according to (1-1) and let $\gamma \geq 1$. Then each $f \in K_u$ has a T_γ -limit at 1 if and only if (1-3) holds for $\xi = 1$ and*

$$\limsup_{t \rightarrow 0^+} F_{t, \gamma} < \infty,$$

where

$$(2-1) \quad F_{t, \gamma} = \left\{ \sum_{t/2 < \lfloor \arg a_n \rfloor < 2t} \frac{1 - |a_n|}{t^{2\gamma} + (t - \lfloor \arg a_n \rfloor)^2} + \left(\int_{t/2}^{2t} + \int_{2\pi - 2t}^{2\pi - t/2} \right) \frac{d\sigma(\lfloor \theta \rfloor)}{t^{2\gamma} + (t - \lfloor \theta \rfloor)^2} + \sum_{t/2 < \lfloor \xi_n \rfloor < 2t} \frac{\rho_n}{t^{2\gamma} + (t - \lfloor \xi_n \rfloor)^2} \right\}.$$

In this paper, we follow Leung’s footsteps and generalize this result to *any approach region* and answer [Question 1.1](#).

For a finite Blaschke product B , the dimension of the model space K_B is finite and any $f \in K_B$ is analytic on $c\mathbb{D}$ for some $|c| > 1$. So, in this paper, we always

assume that B is an infinite Blaschke product. Throughout this paper, for $\omega \in \overline{\mathbb{D}}$, $\arg \omega$ is always restricted to the interval $[0, 2\pi)$ ($\arg w = 0$ if $w = 0$). For $z \in \mathbb{D}$, write

$$E_z = \begin{cases} \emptyset & \text{if } \arg z = 0, \\ ((\arg z)/2, 2 \arg z) & \text{if } \arg z \in (0, \pi), \\ (2(\arg z - \pi), \pi + (\arg z)/2) & \text{if } \arg z \in [\pi, 2\pi), \end{cases}$$

and

$$F_z = \left\{ \sum_{\arg a_n \in E_z} \frac{1 - |a_n|}{(1 - |z|)^2 + (\arg z - \arg a_n)^2} + \int_{E_z} \frac{d\sigma(\theta)}{(1 - |z|)^2 + (\arg z - \theta)^2} + \sum_{\zeta_n \in E_z} \frac{\rho_n}{(1 - |z|)^2 + (\arg z - \zeta_n)^2} \right\}.$$

Intuitively, when $\arg z \in [\pi, 2\pi)$, $\{e^{i\theta} : \theta \in E_z\} = \overline{\{e^{i\theta} : \theta \in E_{\bar{z}}\}}$, i.e., they are symmetric about the x -axis.

We can now state the main results of this work.

Theorem 2.1. *Let u be an inner function factorized according to (1-1). Suppose Ω_1 is any approach region with vertex at 1. Then the following are equivalent.*

(i) *We have*

$$\sup_{z \in \Omega_1} \|k_z^u\| < \infty.$$

(ii) *k_z^u converges weakly as $z \rightarrow 1$ within Ω_1 .*

(iii) *For each $f \in K_u$, f has a limit as $z \rightarrow 1$ within Ω_1 .*

(iv) *For each $f \in K_u$, f is bounded on Ω_1 .*

(v) *(1-3) holds for $\xi = 1$ and*

$$(2-2) \quad \limsup_{\substack{z \in \Omega_1 \\ z \rightarrow 1}} F_z < \infty.$$

Note that as z tends to 1 nontangentially from \mathbb{D} , k_z^u converges weakly if and only if k_z^u converges in the norm of K_u . So it is natural to consider for any approach region Ω_1 , when k_z^u converges in the norm of K_u as $z \rightarrow 1$ within Ω_1 . Fortunately, we get a complete characterization.

Theorem 2.2. *Let u be an inner function factorized according to (1-1). Suppose Ω_1 is an approach region with vertex at 1. Then $k_z^u \rightarrow k_1^u$ in the norm of K_u as $z \rightarrow 1$ within Ω_1 if and only if (1-3) holds for $\xi = 1$ and*

$$(2-3) \quad \lim_{\substack{z \in \Omega_1 \\ z \rightarrow 1}} F_z = 0.$$

3. Weak convergence

In this section, we establish some technical lemmas and then use them to give the proof of [Theorem 2.1](#). First we show that the convergence of all functions in K_u is a stronger condition than the convergence of any divisor of u .

Lemma 3.1. *Let u be an inner function, and let Ω_1 be an approach region with vertex at 1. If each $f \in K_u$ has a limit as $z \rightarrow 1$ within Ω_1 , then any divisor v of u has a limit of modulus 1 as $z \rightarrow 1$ within Ω_1 .*

Proof. Since v is a divisor of u , $K_v \subseteq K_u$ and for any reproducing kernel k_z^v in K_v , we have

$$\|k_z^v\|^2 = k_z^v(z) = \frac{1 - |v(z)|^2}{1 - |z|^2}, \quad z \in \mathbb{D}.$$

The hypothesis together with the uniform boundedness principle yields

$$\sup_{z \in \Omega_1} \|k_z^v\| \leq \sup_{z \in \Omega_1} \|k_z^u\| < \infty,$$

which implies $|v(z)| \rightarrow 1$ as $z \rightarrow 1$ within Ω_1 .

Again by hypothesis, each $g \in K_v$ has a limit as $z \rightarrow 1$ within Ω_1 . In particular, choose $z_0 \in \Omega_1$ such that $v(z_0) \neq 0$; then $k_{z_0}^v$ has a limit as $z \rightarrow 1$ within Ω_1 and hence v has a limit of modulus 1 as $z \rightarrow 1$ within Ω_1 . □

Remark 3.2. From the proof of [Lemma 3.1](#), it is easy to see that if we replace the approach region Ω_1 by $\partial\Omega_1 \setminus \{1\}$, $\bar{\Omega}_1 \setminus \{1\}$ or any arc lying in \mathbb{D} and terminating at 1, [Lemma 3.1](#) still holds.

Ahern and Clark [\[2\]](#) proved that K_u is unitarily equivalent to the direct sum of three L^2 spaces. In particular,

$$(3-1) \quad \|k_z^s\|^2 = 2 \int_0^{2\pi} \frac{|s_\theta(z)|^2}{|1 - e^{-i\theta}z|^2} d\sigma(\theta), \quad \|k_z^\Delta\|^2 = 2 \sum_{n=1}^\infty \rho_n \frac{\int_{n-1}^n |\Delta_\lambda(z)|^2 d\lambda}{|1 - e^{-i\xi_n}z|^2},$$

with

$$s_\theta(z) = \exp \left\{ - \int_0^\theta \frac{e^{it} + z}{e^{it} - z} d\sigma(t) \right\}, \quad \theta \in [0, 2\pi),$$

and

$$\Delta_\lambda(z) = \exp \left\{ - \sum_{j=1}^n \rho_j \frac{e^{i\xi_j} + z}{e^{i\xi_j} - z} - (\lambda - n) \rho_{n+1} \frac{e^{i\xi_{n+1}} + z}{e^{i\xi_{n+1}} - z} \right\},$$

where $\lambda \in (0, \infty)$ and n is the integral part of λ (see [\[2, pp. 194–195\]](#)). Moreover, recall that $\{e_n\}$ is an orthonormal basis for K_B (see [\(1-2\)](#)) and hence by Parseval’s

equality, we have

$$(3-2) \quad \|k_z^B\|^2 = \sum_{n=1}^{\infty} |\langle k_z^B, e_n \rangle|^2 = \sum_{n=1}^{\infty} |e_n(z)|^2 = \sum_{n=1}^{\infty} \frac{1 - |a_n|^2}{|1 - \bar{a}_n z|^2} |B_n(z)|^2.$$

Moreover, [2, Lemma 4.1] showed that for any $f \in K_u$, f admits an orthogonal decomposition

$$f = P_B f + B P_s \bar{B} f + B s P_\Delta \bar{B} s f,$$

where P_B , P_s and P_Δ are the projections onto K_B , K_s and K_Δ . In particular, if $f = k_z^u$, then

$$\begin{aligned} k_z^u(w) &= \langle P_B k_z^u + B P_s \bar{B} k_z^u + B s P_\Delta \bar{B} s k_z^u, k_w^u \rangle \\ &= \langle P_B k_z^u, k_w^u \rangle + \langle B P_s \bar{B} k_z^u, k_w^u \rangle + \langle B s P_\Delta \bar{B} s k_z^u, k_w^u \rangle \\ &= \langle P_B k_z^u, k_w^B \rangle + B(w) \langle P_s \bar{B} k_z^u, k_w^s \rangle + B(w) s(w) \langle P_\Delta \bar{B} s k_z^u, k_w^\Delta \rangle \\ &= \langle k_z^u, k_w^B \rangle + B(w) \langle k_z^u, B k_w^s \rangle + B(w) s(w) \langle k_z^u, B s k_w^\Delta \rangle \\ &= k_z^B(w) + B(w) \bar{B}(z) k_z^s(w) + B(w) s(w) \overline{B(z) s(z)} k_z^\Delta(w). \end{aligned}$$

Therefore, we get the following lemma.

Lemma 3.3. *Let u be an inner function factorized according to (1-1). Then*

$$\begin{aligned} \|k_z^u\|^2 &= \|k_z^B\|^2 + |B(z)|^2 \|k_z^s\|^2 + |B(z) s(z)|^2 \|k_z^\Delta\|^2 \\ &= \sum_{n=1}^{\infty} \frac{1 - |a_n|^2}{|1 - \bar{a}_n z|^2} |B_n(z)|^2 + 2 |B(z)|^2 \int_0^{2\pi} \frac{|s_\theta(z)|^2}{|1 - e^{-i\theta} z|^2} d\sigma(\theta) \\ &\quad + 2 |B(z) s(z)|^2 \sum_{n=1}^{\infty} \rho_n \frac{\int_{n-1}^n |\Delta_\lambda(z)|^2 d\lambda}{|1 - e^{-i\zeta_n} z|^2}. \end{aligned}$$

Lemma 3.4. *Let $a \in \bar{\mathbb{D}}$ and $z \in \mathbb{D}$ with $|z| > \frac{1}{2}$. Then the following hold:*

(i) *We have*

$$|1 - \bar{a}z|^2 \leq 2((1 - |z|)^2 + (1 - |a|)^2 + (\arg z - \arg a)^2).$$

(ii) *If $\arg a \notin E_z$, then*

$$|1 - \bar{a}z|^2 \geq \frac{|1 - a|^2}{8}.$$

(iii) *If $\arg a \in E_z$ and $|a| > \frac{1}{2}$, then*

$$(1 - |z|)^2 + (1 - |a|)^2 + (\arg z - \arg a)^2 \leq \pi^2 |1 - \bar{a}z|^2.$$

Proof. (i): This follows a direct calculation:

$$\begin{aligned}
 (3-3) \quad |1 - \bar{a}z|^2 &= (1 - |az|)^2 + 4|az| \sin^2 \frac{\arg(\bar{a}z)}{2} \\
 &= (1 - |az|)^2 + 4|az| \sin^2 \frac{\arg z - \arg a}{2} \\
 &\leq 2((1 - |z|)^2 + (1 - |a|)^2) + (\arg z - \arg a)^2 \\
 &\leq 2((1 - |z|)^2 + (1 - |a|)^2 + (\arg z - \arg a)^2).
 \end{aligned}$$

(ii): We divide this part into three cases.

Case 1: $\arg z = 0$. In this case, combining (3-3) and $|z| > \frac{1}{2}$, we have

$$|1 - \bar{a}z|^2 > (1 - |a|)^2 + 2|a| \sin^2 \frac{\arg a}{2} \geq \frac{|1 - a|^2}{2}.$$

Case 2: $\arg z \in (0, \pi)$ and $\arg a \notin E_z$. In this case either $0 \leq \arg a \leq (\arg z)/2$ or $\arg a \geq 2\arg z$. In the first situation, we have $0 \leq \arg a \leq \arg z - \arg a \leq \pi$. In the second situation, we have

$$\begin{cases} (\arg a)/2 \leq \arg a - \arg z \leq \pi & \text{if } 2\arg z \leq \arg a \leq \pi + \arg z, \\ 2\pi - \arg a \leq 2\pi - \arg a + \arg z \leq \pi & \text{if } \pi + \arg z \leq \arg a < 2\pi. \end{cases}$$

If $0 \leq \arg a \leq \arg z - \arg a \leq \pi$, then

$$\sin^2 \frac{\arg z - \arg a}{2} \geq \sin^2 \frac{\arg a}{2} \geq \sin^2 \frac{\arg a}{4}.$$

If $2\arg z \leq \arg a \leq \pi + \arg z$, then

$$\sin^2 \frac{\arg z - \arg a}{2} \geq \sin^2 \frac{\arg a}{4}.$$

If $\pi + \arg z \leq \arg a < 2\pi$, then

$$\begin{aligned}
 \sin^2 \frac{\arg z - \arg a}{2} &= \sin^2 \frac{2\pi - \arg a + \arg z}{2} \geq \sin^2 \frac{\arg a}{2} \\
 &= 4 \sin^2 \frac{\arg a}{4} \cos^2 \frac{\arg a}{4} \geq 2 \cos^2 \frac{\arg a}{4} \\
 &\geq \cos^2 \frac{\arg a}{4}.
 \end{aligned}$$

So in both situations, combining (3-3) and $|z| > \frac{1}{2}$, we get

$$|1 - \bar{a}z|^2 \geq (1 - |a|)^2 + 2|a| \min \left\{ \sin^2 \frac{\arg a}{4}, \cos^2 \frac{\arg a}{4} \right\}.$$

Again by $\sin^2 \frac{\arg a}{2} = 4 \sin^2 \frac{\arg a}{4} \cos^2 \frac{\arg a}{4}$, we have

$$\min \left\{ \sin^2 \frac{\arg a}{4}, \cos^2 \frac{\arg a}{4} \right\} \geq \frac{1}{4} \sin^2 \frac{\arg a}{2}.$$

This leads to

$$|1 - \bar{a}z|^2 \geq (1 - |a|)^2 + \frac{|a|}{2} \sin^2 \frac{\arg a}{2} \geq \frac{|1 - a|^2}{8}.$$

Case 3: $\arg z \in [\pi, 2\pi)$ and $\arg a \notin E_z$. By the symmetry, i.e., $\{e^{i\theta} : \theta \in E_z\} = \{e^{i\theta} : \theta \in E_{\bar{z}}\}$, the desired result immediately follows from Case 2.

(iii): The condition $\arg a \in E_z$ implies $\arg z \in (0, 2\pi)$ and then

$$(3-4) \quad \begin{cases} |\arg z - \arg a| \leq \arg z \leq \pi & \text{if } \arg z \in (0, \pi), \\ |\arg z - \arg a| \leq 2\pi - \arg z \leq \pi & \text{if } \arg z \in [\pi, 2\pi). \end{cases}$$

Combining (3-3) and $|z| > \frac{1}{2}, |a| > \frac{1}{2}$, we have

$$|1 - \bar{a}z|^2 \geq (1 - |az|)^2 + \sin^2 \frac{\arg z - \arg a}{2}.$$

Moreover, it follows from $(1 - xy) \geq (2 - x - y)/2$ for $0 \leq x, y \leq 1$ that

$$|1 - \bar{a}z|^2 \geq \frac{(2 - |a| - |z|)^2}{4} + \sin^2 \frac{\arg z - \arg a}{2}.$$

This together with $(2 - x - y)^2 = ((1 - x) + (1 - y))^2 \geq (1 - x)^2 + (1 - y)^2$ for $0 \leq x, y \leq 1, \sin x \geq 2x/\pi$ for $x \in [0, \pi/2]$ and (3-4) gives

$$\begin{aligned} |1 - \bar{a}z|^2 &\geq \frac{(1 - |a| + 1 - |z|)^2}{4} + \frac{(\arg z - \arg a)^2}{\pi^2} \\ &\geq \frac{(1 - |a|)^2 + (1 - |z|)^2 + (\arg z - \arg a)^2}{\pi^2}, \end{aligned}$$

as desired. □

Lemma 3.5. *Let B be a Blaschke product with zeros $\{a_n\}$ and Ω_1 be an approach region with vertex at 1. For $z \in \mathbb{D}$, write*

$$\begin{aligned} G_z &= \sum_{\substack{\arg a_n \in E_z \\ |a_n| > \frac{1}{2}}} \frac{1 - |a_n|}{(1 - |z|)^2 + (1 - |a_n|)^2 + (\arg z - \arg a_n)^2}, \\ H_z &= \sum_{\arg a_n \in E_z} \frac{1 - |a_n|}{(1 - |z|)^2 + (\arg z - \arg a_n)^2}. \end{aligned}$$

If $\sum_{n \geq 1} (1 - |a_n|)/|1 - a_n|^2 < \infty$, then there exists a constant C , with $0 < C < 1$, such that

$$C \limsup_{\substack{z \in \Omega_1 \\ z \rightarrow 1}} H_z \leq \limsup_{\substack{z \in \Omega_1 \\ z \rightarrow 1}} G_z \leq \limsup_{\substack{z \in \Omega_1 \\ z \rightarrow 1}} H_z.$$

Proof. If $\arg z = 0$, then $E_z = \emptyset$ and the desired result is obvious. So we can assume that $\arg z \in (0, 2\pi)$. It is clear that $G_z \leq H_z$ and hence it suffices to show that

$$C \limsup_{\substack{z \in \Omega_1 \\ z \rightarrow 1}} H_z \leq \limsup_{\substack{z \in \Omega_1 \\ z \rightarrow 1}} G_z < \infty$$

for some $C \in (0, 1)$.

Now assume $M = \limsup_{z \in \Omega_1, z \rightarrow 1} G_z < \infty$, for each $\epsilon > 0$, there exist $r \in (0, 1)$ and $t_0 \in (0, \frac{1}{2})$ such that

$$\frac{1 - |a_n|}{(1 - |z|)^2 + (1 - |a_n|)^2 + (\arg z - \arg a_n)^2} \leq M + \epsilon,$$

when $z \in \Omega_1$, $|z| \in (r, 1)$, $\arg z \in (0, t_0) \cup (2\pi - t_0, 2\pi)$, $\arg a_n \in E_z$ and $|a_n| > \frac{1}{2}$.

If $\sum_{n \geq 1} (1 - |a_n|)/|1 - a_n|^2 < \infty$, then we conclude from [6, p. 336] that

$$\sum_{\arg a_n \in (0, \pi)} \frac{1 - |a_n|}{(\arg a_n)^2} + \sum_{\arg a_n \in [\pi, 2\pi)} \frac{1 - |a_n|}{(2\pi - \arg a_n)^2} < \infty.$$

So there exists $t \in (0, t_0)$ such that

$$1 - |a_n| \leq \frac{1}{M + \epsilon + 2},$$

and hence $|a_n| > \frac{1}{2}$ for $\arg a_n \in (0, t) \cup (2\pi - t, 2\pi)$. This implies

$$\begin{aligned} (1 - |a_n|)^2 &\leq \frac{1 - |a_n|}{(M + \epsilon + 2)} \\ &\leq \frac{M + \epsilon}{M + \epsilon + 2} ((1 - |z|)^2 + (1 - |a_n|)^2 + (\arg z - \arg a_n)^2), \end{aligned}$$

when $z \in \Omega_1$, $|z| \in (r, 1)$, $\arg z \in (0, t/2) \cup (2\pi - t/2, 2\pi)$ and $\arg a_n \in E_z$. This can be rewritten as

$$(1 - |a_n|)^2 \leq \frac{M + \epsilon}{2} ((1 - |z|)^2 + (\arg z - \arg a_n)^2),$$

when $z \in \Omega_1$, $|z| \in (r, 1)$, $\arg z \in (0, t/2) \cup (2\pi - t/2, 2\pi)$ and $\arg a_n \in E_z$. Therefore,

$$\frac{1 - |a_n|}{(1 - |z|)^2 + (1 - |a_n|)^2 + (\arg z - \arg a_n)^2} \geq \frac{2}{M + \epsilon + 2} \frac{1 - |a_n|}{(1 - |z|)^2 + (\arg z - \arg a_n)^2},$$

when $z \in \Omega_1$, $|z| \in (r, 1)$, $\arg z \in (0, t/2) \cup (2\pi - t/2, 2\pi)$ and $\arg a_n \in E_z$. Choose $C = 2/(M + \epsilon + 2)$ and we complete the proof. \square

Now we are at the position to prove [Theorem 2.1](#).

The proof of Theorem 2.1. (ii) \iff (iii) and (iii) \implies (iv) are obvious and (iv) \implies (i) follows from the uniform boundedness principle.

(i) \implies (iii): This follows from the proof of [10, Theorem 4.1].

(i) \iff (v): This is the delicate part. It follows from $\sup_{z \in \Omega_1} \|k_z^u\| < \infty$ that $\liminf_{z \rightarrow 1} \|k_z^u\| < \infty$, which is equivalent to (1-3) for $\xi = 1$ by Theorem B. Therefore, since in (v) we already assume (1-3) for $\xi = 1$, to finish the proof, it suffices to show that if (1-3) holds for $\xi = 1$, then $\sup_{z \in \Omega_1} \|k_z^u\| < \infty$ if and only if (2-2) holds.

If (1-3) holds for $\xi = 1$ and $|z| > \frac{1}{2}$, then by Lemma 3.4(ii), we have

$$(3-5) \quad \sum_{\arg a_n \notin E_z} \frac{1 - |a_n|^2}{|1 - \bar{a}_n z|^2} |B_n(z)|^2 + \int_{\theta \notin E_z} \frac{|s_\theta(z)|^2}{|1 - e^{-i\theta} z|^2} d\sigma(\theta) + \sum_{\zeta_n \notin E_z} \rho_n \frac{\int_{n-1}^n |\Delta_\lambda(z)|^2 d\lambda}{|1 - e^{-i\zeta_n} z|^2} \\ \leq 16 \sum_{n=1}^\infty \frac{1 - |a_n|}{|1 - a_n|^2} + 8 \int_0^{2\pi} \frac{d\sigma(\theta)}{|1 - e^{i\theta}|^2} + 8 \sum_{n=1}^\infty \frac{\rho_n}{|1 - e^{-i\zeta_n}|^2} < \infty.$$

Also it is clear that

$$(3-6) \quad \sum_{\substack{\arg a_n \in E_z \\ |a_n| \leq \frac{1}{2}}} \frac{1 - |a_n|^2}{|1 - \bar{a}_n z|^2} \leq 4 \sum_{n=1}^\infty (1 - |a_n|^2) < \infty.$$

It follows from Lemma 3.3 that

$$\|k_z^u\|^2 = \|k_z^B\|^2 + |B(z)|^2 \|k_z^s\|^2 + |B(z)s(z)|^2 \|k_z^\Delta\|^2,$$

which together with $\sup_{z \in \Omega_1} \|k_z^u\| < \infty$ and Lemma 3.1 implies $|u(z)| \rightarrow 1$ as $z \rightarrow 1$ within Ω_1 . Hence we can find $r \in (\frac{1}{2}, 1)$ such that $|u(z)| > \frac{1}{2}$ whenever $z \in \Omega_1$ and $|z| > r$. Recall that when $|u(z)| > \frac{1}{2}$, each of its factors has to be bounded from below by $\frac{1}{2}$. So $\sup_{z \in \Omega_1} \|k_z^u\| < \infty$ is equivalent to

$$(3-7) \quad \sup_{\substack{z \in \Omega_1 \\ |z| > r}} (\|k_z^B\|^2 + \|k_z^s\|^2 + \|k_z^\Delta\|^2) < \infty.$$

Combining (3-5), (3-6) and Lemma 3.3 together, (3-7) in turn is equivalent to

$$\sum_{\substack{\arg a_n \in E_z \\ |a_n| > \frac{1}{2}}} \frac{1 - |a_n|^2}{|1 - \bar{a}_n z|^2} + \int_{\theta \in E_z} \frac{d\sigma(\theta)}{|1 - e^{-i\theta} z|^2} + \sum_{\zeta_n \in E_z} \frac{\rho_n}{|1 - e^{-i\zeta_n} z|^2} < \infty,$$

when $z \in \Omega_\xi$ and $|z| > r$. This is equivalent to (2-2) by the combination of Lemmas 3.4 and 3.5. This completes the proof. \square

Remark 3.6. From the proof of Theorem 2.1, it is easy to see that if we replace the approach region Ω_1 by $\partial\Omega_1 \setminus \{1\}$, $\bar{\Omega}_1 \setminus \{1\}$ or any arc lying in \mathbb{D} and terminating at 1, Theorem 2.1 still holds.

4. Norm convergence

We establish another set of technical lemmas which are used to prove [Theorem 2.2](#).

By [Theorem B](#) and [[7](#), Corollary 1.7.14, p. 29], we have the following lemma.

Lemma 4.1. *Let u be an inner function factorized according to (1-1). If (1-3) holds for $\xi = 1$, then*

$$\begin{aligned} \|k_1^u\|^2 &= \|k_1^B\|^2 + \|k_1^\xi\|^2 + \|k_1^\Delta\|^2 \\ &= \sum_{n=1}^\infty \frac{1 - |a_n|^2}{|1 - a_n|^2} + 2 \int_0^{2\pi} \frac{d\sigma(\theta)}{|1 - e^{i\theta}|^2} + 2 \sum_{n=1}^\infty \frac{\rho_n}{|1 - e^{i\xi_n}|^2}. \end{aligned}$$

Lemma 4.2. *Let B be a Blaschke product with zeros $\{a_n\}$ such that*

$$\sum_{n=1}^\infty \frac{1 - |a_n|}{|1 - a_n|^2} < \infty.$$

Then

$$\|k_1^B\|^2 = \lim_{\substack{z \in \mathbb{D} \\ z \rightarrow 1}} \sum_{\arg a_n \notin E_z} \frac{1 - |a_n|^2}{|1 - \bar{a}_n z|^2} |B_n(z)|^2.$$

Proof. By hypothesis, it is clear that $\{n \in \mathbb{N} : \arg a_n = 0\}$ is a finite set. Hence B admits a decomposition $B = f_0 f_1$, where f_0 and f_1 are two Blaschke products with zeros $\{a_n : \arg a_n = 0\}$ and $\{a_n : \arg a_n \in (0, 2\pi)\}$, respectively.

Note that $k_z^B = k_z^{f_0} + \overline{f_0(z)} f_0 k_z^{f_1}$ and hence

$$\|k_z^B\|^2 = \|k_z^{f_0}\|^2 + |f_0(z)|^2 \|k_z^{f_1}\|^2.$$

Observe that f_0 is a finite Blaschke product and for each a_n with $\arg a_n = 0$, we have $\arg a_n \notin E_z$ for any $z \in \mathbb{D}$, so by (3-3), we conclude that

$$\|k_1^{f_0}\|^2 = \lim_{\substack{z \in \mathbb{D} \\ z \rightarrow 1}} \|k_z^{f_0}\|^2 = \lim_{\substack{z \in \mathbb{D} \\ z \rightarrow 1}} \sum_{\arg a_n \notin E_z} \frac{1 - |a_n|^2}{|1 - \bar{a}_n z|^2} |B_n(z)|^2.$$

Moreover, it is clear that $\lim_{z \in \mathbb{D}, z \rightarrow 1} |f_0(z)| = 1$. So to end the proof, it suffices to consider the case that the zeros $\{a_n\}$ of B satisfy $\arg a_n \in (0, 2\pi)$.

Now again by hypothesis, there exists a natural number N such that

$$\sum_{n=N+1}^\infty \frac{1 - |a_n|^2}{|1 - a_n|^2} < \epsilon.$$

There also exist two positive numbers $r \in (\frac{1}{2}, 1)$ and $t_0 \in (0, \pi/2)$ such that

$$\left| \sum_{n=1}^N \frac{1 - |a_n|^2}{|1 - \bar{a}_n z|^2} |B_n(z)|^2 - \sum_{n=1}^N \frac{1 - |a_n|^2}{|1 - a_n|^2} \right| < \epsilon,$$

when $|z| \in (r, 1)$ and $\arg z \in [0, t_0) \cup (2\pi - t_0, 2\pi)$.

For $|z| \in (r, 1)$, Lemma 3.4(ii) shows that

$$\sum_{\substack{n=N+1 \\ \arg a_n \notin E_z}}^{\infty} \frac{1 - |a_n|^2}{|1 - \bar{a}_n z|^2} |B_n(z)|^2 \leq 8 \sum_{n=N+1}^{\infty} \frac{1 - |a_n|^2}{|1 - a_n|^2} < 8\epsilon.$$

Put

$$t = \min\{t_0, \min\{(\arg a_n)/2 : 1 \leq n \leq N, \arg a_n \in (0, \pi)\}, \min\{(2\pi - \arg a_n)/2 : 1 \leq n \leq N, \arg a_n \in [\pi, 2\pi)\}\}.$$

Here we claim that, for $1 \leq n \leq N$, if $\arg z \in [0, t) \cup (2\pi - t, 2\pi)$, then $\arg a_n \notin E_z$. To do this, three cases arise. If $\arg z = 0$, then $E_z = \emptyset$, so that $\arg a_n \notin E_z$ for every natural number n . But if $\arg z \in (0, t)$, then for $\arg a_n \in E_z$, we have $0 < (\arg a_n)/2 < \arg z < t$, which implies $n > N$. Finally, if $\arg z \in (2\pi - t, 2\pi)$, whenever $\arg a_n \in E_z$, we have $0 < (2\pi - \arg a_n)/2 < (2\pi - \arg z) < t$, which implies $n > N$. Thus the claim is proved.

Let $|z| \in (r, 1)$ and $\arg z \in [0, t) \cup (2\pi - t, 2\pi)$. Then the above discussion implies

$$\begin{aligned} & \left| \sum_{\arg a_n \notin E_z} \frac{1 - |a_n|^2}{|1 - \bar{a}_n z|^2} |B_n(z)|^2 - \sum_{n=1}^{\infty} \frac{1 - |a_n|^2}{|1 - a_n|^2} \right| \\ & \leq \left| \sum_{\arg a_n \notin E_z} \frac{1 - |a_n|^2}{|1 - \bar{a}_n z|^2} |B_n(z)|^2 - \sum_{n=1}^N \frac{1 - |a_n|^2}{|1 - \bar{a}_n z|^2} |B_n(z)|^2 \right| \\ & \quad + \left| \sum_{n=1}^N \frac{1 - |a_n|^2}{|1 - \bar{a}_n z|^2} |B_n(z)|^2 - \sum_{n=1}^N \frac{1 - |a_n|^2}{|1 - a_n|^2} \right| + \left| \sum_{n=1}^N \frac{1 - |a_n|^2}{|1 - a_n|^2} - \sum_{n=1}^{\infty} \frac{1 - |a_n|^2}{|1 - a_n|^2} \right| \\ & < 10\epsilon, \end{aligned}$$

which together with Lemma 4.1 gives the desired result. □

Lemma 4.3. *Let Ω_1 be an approach region with vertex at 1 and let $u = s\Delta$ be a singular inner function factorized according to (1-1). If $\sup_{z \in \Omega_1} \|k_z^s\| < \infty$, then*

$$(4-1) \quad \|k_1^s\|^2 = 2 \lim_{\substack{z \in \Omega_1 \\ z \rightarrow 1}} \int_{\theta \notin E_z} \frac{|s_\theta(z)|^2}{|1 - e^{-i\theta}z|^2} d\sigma(\theta)$$

and

$$(4-2) \quad \|k_1^\Delta\|^2 = 2 \lim_{\substack{z \in \Omega_1 \\ z \rightarrow 1}} \sum_{\zeta_n \notin E_z} \rho_n \frac{\int_{n-1}^n |\Delta_\lambda(z)|^2 d\lambda}{|1 - e^{-i\zeta_n}z|^2}.$$

Proof. To begin with, we first prove (4-1). It is clear that

$$\sup_{z \in \Omega_1} \|k_z^s\| \leq \sup_{z \in \Omega_1} \|k_z^u\| < \infty.$$

So [Lemma 3.1](#) together with [Theorem 2.1](#) implies that s has a limit of modulus 1 as $z \rightarrow 1$ within Ω_1 and (1-3) holds for $\xi = 1$. Consequently,

$$(4-3) \quad M = \int_0^{2\pi} \frac{d\sigma(\theta)}{|1 - e^{i\theta}|^2} < \infty.$$

Recall that σ is a finite positive continuous singular measure with respect to Lebesgue measure, so by (4-3), we conclude that for each $\epsilon > 0$, there exists $t_0 \in (0, \pi/2)$ such that

$$(4-4) \quad \int_{\theta \in B_{t_0}} \frac{d\sigma(\theta)}{|1 - e^{i\theta}|^2} < \epsilon,$$

where $B_{t_0} = [0, t_0) \cup (2\pi - t_0, 2\pi)$. There also exist two positive numbers $r \in (\frac{1}{2}, 1)$ and $t \in (0, t_0/2)$ such that $1 - |s(z)|^2 < \epsilon$ and

$$(4-5) \quad \left| \int_{\theta \notin B_{t_0}} \frac{d\sigma(\theta)}{|1 - e^{-i\theta}z|^2} - \int_{\theta \notin B_{t_0}} \frac{d\sigma(\theta)}{|1 - e^{i\theta}|^2} \right| < \epsilon,$$

when $z \in \Omega_1$, $|z| \in (r, 1)$ and $\arg z \in [0, t) \cup (2\pi - t, 2\pi)$.

Now let $z \in \Omega_1$, $|z| \in (r, 1)$ and $\arg z \in [0, t) \cup (2\pi - t, 2\pi)$. Then $E_z \subseteq B_{t_0}$ and hence [Lemma 3.4\(2\)](#) yields

$$(4-6) \quad \int_{\theta \in (B_{t_0} \setminus E_z)} \frac{|s_\theta(z)|^2}{|1 - e^{-i\theta}z|^2} d\sigma(\theta) \leq 8 \int_{\theta \in B_{t_0}} \frac{d\sigma(\theta)}{|1 - e^{i\theta}|^2} < 8\epsilon$$

and

$$(4-7) \quad \int_{\theta \notin B_{t_0}} \frac{1 - |s_\theta(z)|^2}{|1 - e^{-i\theta}z|^2} d\sigma(\theta) \leq 8(1 - |s(z)|^2) \int_{\theta \notin E_z} \frac{d\sigma(\theta)}{|1 - e^{i\theta}|^2} \leq 8M\epsilon.$$

Putting (4-4)–(4-7) together, it follows that

$$\begin{aligned} & \left| \int_{\theta \notin E_z} \frac{|s_\theta(z)|^2}{|1 - e^{-i\theta}z|^2} d\sigma(\theta) - \int_0^{2\pi} \frac{d\sigma(\theta)}{|1 - e^{i\theta}|^2} \right| \\ & \leq \left| \int_{\theta \notin E_z} \frac{|s_\theta(z)|^2}{|1 - e^{-i\theta}z|^2} d\sigma(\theta) - \int_{\theta \notin B_{t_0}} \frac{|s_\theta(z)|^2}{|1 - e^{-i\theta}z|^2} d\sigma(\theta) \right| \\ & \quad + \left| \int_{\theta \notin B_{t_0}} \frac{|s_\theta(z)|^2}{|1 - e^{-i\theta}z|^2} d\sigma(\theta) - \int_{\theta \notin B_{t_0}} \frac{d\sigma(\theta)}{|1 - e^{-i\theta}z|^2} \right| \\ & \quad + \left| \int_{\theta \notin B_{t_0}} \frac{d\sigma(\theta)}{|1 - e^{-i\theta}z|^2} - \int_{\theta \notin B_{t_0}} \frac{d\sigma(\theta)}{|1 - e^{i\theta}|^2} \right| \\ & \quad + \left| \int_{\theta \notin B_{t_0}} \frac{d\sigma(\theta)}{|1 - e^{i\theta}|^2} - \int_0^{2\pi} \frac{d\sigma(\theta)}{|1 - e^{i\theta}|^2} \right| \\ & \leq (10 + 8M)\epsilon, \end{aligned}$$

which together with [Lemma 4.1](#) shows that (4-1) holds.

The proof of (4-2) is similar to the proof of (4-1) and hence it will be omitted. This completes the proof. \square

The proof of Theorem 2.2. To begin with, we show that $k_z^u \rightarrow k_1^u$ in the norm of K_u as $z \rightarrow 1$ within Ω_1 if and only if $\|k_1^u\| = \lim_{z \in \Omega_1, z \rightarrow 1} \|k_z^u\|$. Indeed, if $\|k_1^u\| = \lim_{z \in \Omega_1, z \rightarrow 1} \|k_z^u\|$, then Theorem B implies that $u(1)$ exists and Lemma 3.1 implies that

$$u_{\Omega_1}(1) := \lim_{\substack{z \in \Omega_1 \\ z \rightarrow 1}} u(z), \quad |u_{\Omega_1}(1)| = 1.$$

A theorem of Lindelöf [8, Theorem 2.3, p. 19] says that if $h \in H^\infty$ and $h(z) \rightarrow A$ as $z \rightarrow 1$ along a continuous path $G \subset \mathbb{D}$ terminating at 1, then $h(z) \rightarrow A$ as $z \rightarrow 1$ in any Stolz domain $\Gamma_\alpha(1)$. This implies $u_{\Omega_1}(1) = u(1)$. So for every $w \in \mathbb{D}$,

$$(4-8) \quad k_1^u(w) = \frac{1 - \overline{u(1)}u(w)}{1 - w} = \lim_{\substack{z \in \Omega_1 \\ z \rightarrow 1}} \frac{1 - \overline{u(z)}u(w)}{1 - \bar{z}w} = \lim_{\substack{z \in \Omega_1 \\ z \rightarrow 1}} k_z^u(w).$$

Moreover, by Theorem 2.1, it follows that k_z^u weakly as $z \rightarrow 1$ within Ω_1 and we denote this limit by k_{1, Ω_1}^u . Combining this and (4-8), we have $k_{1, \Omega_1}^u = k_1^u$.

Therefore,

$$\lim_{\substack{z \in \Omega_1 \\ z \rightarrow 1}} \|k_z^u - k_1^u\|^2 = \lim_{\substack{z \in \Omega_1 \\ z \rightarrow 1}} (\|k_z^u\|^2 - 2 \operatorname{Re}\langle k_z^u, k_1^u \rangle + \|k_1^u\|^2) = 0.$$

Consequently, to end the proof, it suffices to show that if $\sup_{z \in \Omega_1} \|k_z^u\| < \infty$ and (1-3) holds for $\xi = 1$, then $\|k_1^u\| = \lim_{z \in \Omega_1, z \rightarrow 1} \|k_z^u\|$ if and only if (2-3) holds.

It is clear that $\{n \in \mathbb{N} : |a_n| \leq \frac{1}{2}\}$ is a finite set. Hence we assume that $|a_j| \leq \frac{1}{2}$ for $1 \leq j \leq N_1$ and $|a_j| > \frac{1}{2}$ for all $j > N_1$, where N_1 is some positive integer. Now write $B(z) = f_0(z) f_1(z)$, where f_0 is a finite Blaschke product with zeros $\{a_n : 1 \leq n \leq N_1\}$ and f_1 is a Blaschke product with zeros $\{a_n : n > N_1\}$. Note that $k_z^B = k_z^{f_0} + \overline{f_0(z)} f_0 k_z^{f_1}$ and hence

$$\|k_z^u\|^2 = \|k_z^{f_0}\|^2 + |f_0(z)|^2 \|k_z^{f_1}\|^2 + |B(z)|^2 \|k_z^s\|^2 + |B(z)s(z)|^2 \|k_z^\Delta\|^2.$$

Now Lemma 4.1 implies

$$\|k_1^u\|^2 = \|k_1^{f_0}\|^2 + \|k_1^{f_1}\|^2 + \|k_1^s\|^2 + \|k_1^\Delta\|^2.$$

Since f_0 is a finite Blaschke product, it is clear that

$$\|k_1^{f_0}\| = \lim_{\substack{z \in \mathbb{D} \\ z \rightarrow 1}} \|k_z^{f_0}\|.$$

Lemma 3.1 shows that both B and s have limits of modulus 1 as $z \rightarrow 1$ within Ω_1 . This together with Lemmas 3.4, 3.5, 4.2 and 4.3 shows that $\|k_1^u\| = \lim_{z \in \Omega_1, z \rightarrow 1} \|k_z^u\|$ if and only if (2-3) holds. This completes the proof. \square

5. Special approach regions

In this section, we consider two special approach regions:

(i) For $c > 0$, an oricyclic approach region is defined by

$$E(c, 1) = \{z \in \mathbb{D} : |1 - z|^2 < c(1 - |z|^2)\}.$$

A simple computation shows that $E(c, 1)$ is an open disk internally tangent to the circle at 1 with center $1/(1 + c)$ and radius $c/(1 + c)$.

(ii) For $c > 0$ and $\gamma \geq 1$, let

$$(5-1) \quad R'(c, \gamma) = \{z \in \mathbb{D} \setminus \{0\} : 1 - |z| > c \lfloor \arg z \rfloor^\gamma\},$$

where $\lfloor \arg z \rfloor$ means the length of the shorter one of the two arcs on \mathbb{T} joining $z/|z|$ and 1.

In what follows, we first show that f has a limit as $z \rightarrow 1$ within $R'(c, \gamma)$ if and only if f has a limit as $z \rightarrow 1$ within $R(c, \gamma)$. In fact, we can prove this equivalence in any approach region.

Lemma 5.1. *Let u be an inner function and Ω_1 be an approach region with vertex at 1. Then*

$$\sup_{z \in \Omega_1} \|k_z^u\| < \infty \iff \sup_{\omega \in \bar{\Omega}_1 \setminus \{1\}} \|k_\omega^u\| < \infty.$$

Proof. It is clear that

$$\sup_{\omega \in \bar{\Omega}_1 \setminus \{1\}} \|k_\omega^u\| < \infty \implies \sup_{z \in \Omega_1} \|k_z^u\| < \infty$$

and since every $f \in K_u$ is continuous at every point $\eta \in \partial\Omega_1 \setminus \{1\}$,

$$\sup_{\eta \in \partial\Omega_1 \setminus \{1\}} |f(\eta)| \leq \sup_{z \in \Omega_1} |f(z)|,$$

so that

$$\sup_{\eta \in \Omega_1} \|k_\eta^u\| < \infty \implies \sup_{z \in \bar{\Omega}_1 \setminus \{1\}} \|k_z^u\| < \infty$$

by [Theorem 2.1](#) and [Remark 3.6](#). □

In particular, when $\Omega_1 = R'(c, \gamma)$, we recover [Theorem C \[12, Theorem 4\]](#).

Corollary 5.2. *Let u be an inner function factorized according to (1-1). Then*

$$\sup_{z \in R(c, \gamma)} \|k_z^u\| < \infty$$

if and only if (1-3) holds for $\xi = 1$ and

$$\limsup_{t \rightarrow 0^+} F_{t, \gamma} < \infty,$$

where $F_{t, \gamma}$ is seen in (2-1).

Proof. By Remark 3.6 for $\Omega_1 = R'(c, \gamma)$, to finish the proof, it suffices to show that

$$\limsup_{\substack{z \in R(c, \gamma) \\ z \rightarrow 1}} F_z < \infty \iff \limsup_{t \rightarrow 0^+} F_{t, \gamma} < \infty.$$

\Leftarrow : For $z \in \mathbb{D}$ with $[\arg z] \in [0, \pi/4)$, write $t = \lfloor \arg z \rfloor$. If $\arg z \in [0, \pi/4)$, then $\theta \in E_z$ implies

$$(5-2) \quad \lfloor \theta \rfloor = \theta \in (t/2, 2t) \quad \text{and} \quad (\arg z - \theta)^2 = (t - \lfloor \theta \rfloor)^2.$$

If $\arg z \in (7\pi/4, 2\pi)$, then $\theta \in E_z$ implies

$$(5-3) \quad \begin{aligned} \lfloor \theta \rfloor &= 2\pi - \theta \in (2\pi - (\pi + (\arg z)/2), 2\pi - 2(\arg z - \pi)) = (t/2, 2t), \\ (\arg z - \theta)^2 &= ((2\pi - t) - (2\pi - \theta))^2 = (t - \lfloor \theta \rfloor)^2. \end{aligned}$$

For $z \in R(c, \gamma)$ with $[\arg z] \in [0, \pi/4)$, we have $1 - |z| \geq c \lfloor \arg z \rfloor^\gamma$ and then by (5-2) and (5-3), we have

$$F_z \leq \frac{1}{\min\{c^2, 1\}} F_{t, \gamma},$$

which implies

$$\limsup_{\substack{z \in R(c, \gamma) \\ z \rightarrow 1}} F_z \leq \frac{1}{\min\{c^2, 1\}} \limsup_{t \rightarrow 0^+} F_{t, \gamma} < \infty.$$

\Rightarrow : It is easy to see that

$$\limsup_{\substack{z \in \partial R(c, \gamma) \\ z \rightarrow 1}} F_z \leq \limsup_{\substack{z \in R(c, \gamma) \\ z \rightarrow 1}} F_z < \infty.$$

For $t \in [0, t_0]$ for some $t_0 \in (0, \pi/4)$, there exist $z_1, z_2 \in \partial R(c, \gamma)$ such that $\arg z_1 = t$ and $\arg z_2 = 2\pi - t$. Thus $\lfloor \theta \rfloor \in (t/2, 2t)$ implies that either $\theta \in (t/2, 2t)$ if $\theta \in [0, \pi)$ or $2\pi - \theta \in (t/2, 2t)$ if $\theta \in [\pi, 2\pi)$. In the first situation,

$$(5-4) \quad \theta \in E_{z_1} \quad \text{and} \quad (t - \lfloor \theta \rfloor)^2 = (\arg z_1 - \theta)^2.$$

In the second situation,

$$(5-5) \quad \begin{aligned} \theta \in \left(2\pi - 2t, 2\pi - \frac{t}{2} \right) &= \left(2\pi - 2(2\pi - \arg z_2), 2\pi - \frac{2\pi - \arg z_2}{2} \right) \\ &= \left(2(\arg z_2 - \pi), \pi + \frac{\arg z_2}{2} \right) \\ &= E_{z_2}, \\ (t - \lfloor \theta \rfloor)^2 &= (2\pi - \arg z_2 - (2\pi - \theta))^2 = (\arg z_2 - \theta)^2. \end{aligned}$$

For $z \in \partial R'(c, \gamma) \setminus \{1\}$, we have $1 - |z| = c \lfloor \arg z \rfloor^\gamma$ and then by (5-4) and (5-5), we have

$$F_{t,\gamma} \leq \frac{1}{\min\{1/c^2, 1\}} \max\{F_{z_1}, F_{z_2}\},$$

and if $t \rightarrow 0^+$, then both z_1 and z_2 tend to 1. Therefore,

$$\limsup_{t \rightarrow 0^+} F_{t,\gamma} \leq \frac{1}{\min\{1/c^2, 1\}} \limsup_{\substack{z \in \partial R(c,\gamma) \\ z \rightarrow 1}} F_z < \infty. \quad \square$$

Combining Corollary 5.2 and Lemma 5.1, we immediately get the following result.

Corollary 5.3. *Let u be an inner function factorized according to (1-1). Then $\sup_{z \in R'(c,\gamma)} \|k_z^u\| < \infty$ if and only if $\sup_{z \in R(c,\gamma)} \|k_z^u\| < \infty$. In particular,*

$$\sup_{z \in E(c,1)} \|k_z^u\| < \infty$$

if and only if (1-3) holds for $\xi = 1$ and

$$\limsup_{t \rightarrow 0^+} F_{t,2} < \infty,$$

where $F_{t,2}$ is seen in (2-1).

Note that (2-2) is not easy to calculate. So in what follows, for the approach region $R'(c, \gamma)$, we give a simple example showing that (2-2) can be greatly simplified.

Example 5.4. Let B be a Blaschke product with zeros $\{a_n = r_n e^{i/2^n} : n \in \mathbb{N}\}$ and $R'(c, \gamma)$ be defined by (5-1). In a moment, we will show that k_z^B converges weakly as $z \rightarrow 1$ within $R'(c, \gamma)$ if and only if

$$\sum_{n=1}^{\infty} 4^n (1 - r_n) < \infty \quad \text{and} \quad \limsup_{n \rightarrow \infty} 4^{n\gamma} (1 - r_n) < \infty.$$

It has been shown by Cargo [6, p. 336] that $\sum_{n=1}^{\infty} (1 - r_n) / |1 - r_n e^{i/2^n}|^2 < \infty$ is equivalent to $\sum_{n=1}^{\infty} 4^n (1 - r_n) < \infty$. So by Corollary 5.2, it suffices to show that

$$\limsup_{\arg z \rightarrow 0} \sum_{\frac{1}{2^n} \in E_z} \frac{1 - r_n}{(\arg z)^{2\gamma} + \left(\arg z - \frac{1}{2^n}\right)^2} < \infty \iff \limsup_{n \rightarrow \infty} 4^{n\gamma} (1 - r_n) < \infty.$$

\implies : This follows by choosing the subsequence $\{\arg z = 1/2^n\}$.

\impliedby : For each $\arg z \in (0, \pi/4)$, $1/2^n \in E_z$ if and only if $-\log_2(\arg z) - 1 < n < -\log_2(\arg z) + 1$, which means that there are at most two natural numbers n_1, n_2

such that $1/2^{n_1}, 1/2^{n_2} \in E_z$, and hence $\arg z > 1/2^{n_i+1}$ for $i = 1, 2$. Therefore,

$$\begin{aligned} \limsup_{\arg z \rightarrow 0} \sum_{\frac{1}{2^n} \in E_z} \frac{1 - r_n}{(\arg z)^{2\gamma} + \left(\arg z - \frac{1}{2^n}\right)^2} \\ \leq \limsup_{n_1, n_2 \rightarrow \infty} \{4^{(n_1+1)\gamma} (1 - r_{n_1}) + 4^{(n_2+1)\gamma} (1 - r_{n_2})\} < \infty, \end{aligned}$$

as desired.

Acknowledgements

The authors thank the anonymous referees for careful reading of the manuscript and for constructive suggestions which improved the exposition of the paper and certainly helped to make it more readable. The research is partially supported by National Natural Science Foundation of China (No.12171075; No.12001089) and Science and Technology Research Project of Education Department of Jilin Province (No. JJKH20241406KJ), and as well by the Canada Research Chairs program.

References

- [1] J. Agler, J. E. McCarthy, and N. J. Young, “A Carathéodory theorem for the bidisk via Hilbert space methods”, *Math. Ann.* **352**:3 (2012), 581–624. [MR](#) [Zbl](#)
- [2] P. R. Ahern and D. N. Clark, “On functions orthogonal to invariant subspaces”, *Acta Math.* **124** (1970), 191–204. [MR](#) [Zbl](#)
- [3] P. R. Ahern and D. N. Clark, “Radial limits and invariant subspaces”, *Amer. J. Math.* **92** (1970), 332–342. [MR](#) [Zbl](#)
- [4] F. Bracci and D. Shoikhet, “Boundary behavior of infinitesimal generators in the unit ball”, *Trans. Amer. Math. Soc.* **366**:2 (2014), 1119–1140. [MR](#) [Zbl](#)
- [5] C. Carathéodory, “Über die Winkelderivierten von beschränkten analytischen Funktionen”, *Sitzungsberichte Akad. Berlin* **1929** (1929), 39–52. [Zbl](#)
- [6] G. T. Cargo, “Angular and tangential limits of Blaschke products and their successive derivatives”, *Canadian J. Math.* **14** (1962), 334–348. [MR](#) [Zbl](#)
- [7] J. A. Cima, A. L. Matheson, and W. T. Ross, *The Cauchy transform*, Mathematical Surveys and Monographs **125**, Amer. Math. Soc., Providence, RI, 2006. [MR](#) [Zbl](#)
- [8] E. F. Collingwood and A. J. Lohwater, *The theory of cluster sets*, Cambridge Tracts in Mathematics and Mathematical Physics **56**, Cambridge Univ. Press, 1966. [MR](#) [Zbl](#)
- [9] S. R. Garcia, J. Mashreghi, and W. T. Ross, *Introduction to model spaces and their operators*, Cambridge Studies in Advanced Mathematics **148**, Cambridge Univ. Press, 2016. [MR](#) [Zbl](#)
- [10] A. Hartmann and W. T. Ross, “Truncated Toeplitz operators and boundary values in nearly invariant subspaces”, *Complex Anal. Oper. Theory* **7**:1 (2013), 261–273. [MR](#) [Zbl](#)
- [11] G. Julia, “Extension nouvelle d’un lemme de Schwarz”, *Acta Math.* **42**:1 (1920), 349–355. [MR](#) [Zbl](#)
- [12] K. K. Leung, “On the boundary behaviour of functions orthogonal to invariant subspaces”, *Trans. Amer. Math. Soc.* **244** (1978), 67–84. [MR](#) [Zbl](#)

- [13] J. E. McCarthy and J. E. Pascoe, “The Julia–Carathéodory theorem on the bidisk revisited”, *Acta Sci. Math. (Szeged)* **83**:1-2 (2017), 165–175. [MR](#) [Zbl](#)
- [14] J. E. Pascoe, M. Sargent, and R. Tully-Doyle, “A controlled tangential Julia–Carathéodory theory via averaged Julia quotients”, *Anal. PDE* **14**:6 (2021), 1773–1795. [MR](#) [Zbl](#)
- [15] D. Protas, “Tangential limits of functions orthogonal to invariant subspaces”, *Trans. Amer. Math. Soc.* **166** (1972), 163–172. [MR](#) [Zbl](#)
- [16] G. Ren and X. Wang, “Julia theory for slice regular functions”, *Trans. Amer. Math. Soc.* **369**:2 (2017), 861–885. [MR](#) [Zbl](#)

Received January 22, 2025. Revised May 9, 2025.

YONGJIANG DUAN
DEPARTMENT OF MATHEMATICS
JINAN UNIVERSITY
GUANGZHOU
CHINA
yjduan@jnu.edu.cn

YUFEI LI
SCHOOL OF MATHEMATICS AND STATISTICS
NORTHEAST NORMAL UNIVERSITY
CHANGCHUN
CHINA
liyf495@nenu.edu.cn

JAVAD MASHREGHI
DÉPARTEMENT DE MATHÉMATIQUES ET DE STATISTIQUE
UNIVERSITÉ DE LAVAL
QUÉBEC
CANADA
javad.mashreghi@mat.ulaval.ca

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Fakultät für Mathematik
Universität Wien
Vienna, Austria
matthias.aschenbrenner@univie.ac.at

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Atsushi Ichino
Department of Mathematics
Kyoto University
Riverside, CA 92521-0135
atsushi.ichino@gmail.com

Robert Lipshitz
Department of Mathematics
University of Oregon
Eugene, OR 97403
lipshitz@uoregon.edu

Kefeng Liu
School of Sciences
Chongqing University of Technology
Chongqing 400054, China
liu@math.ucla.edu

Sucharit Sarkar
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
sucharit@math.ucla.edu

Dimitri Shlyakhtenko
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
shlyakht@ipam.ucla.edu

Ruixiang Zhang
Department of Mathematics
University of California
Berkeley, CA 94720-3840
ruixiang@berkeley.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2025 is US \$677/year for the electronic version, and \$917/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2025 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 337 No. 2 August 2025

| | |
|------------------------------------------------------------------------------------------|-----|
| A characterization of quasihomogeneous bivariate polynomials | 201 |
| DAVID BRADLEY-WILLIAMS, PABLO CUBIDES KOVACSICS and IMMANUEL HALUPCZOK | |
| Quartic del Pezzo surfaces without quadratic points | 215 |
| BRENDAN CREUTZ and BIANCA VIRAY | |
| Reproducing kernel functions and boundary limits in model spaces | 225 |
| YONGJIANG DUAN, YUFEI LI and JAVAD MASHREGHI | |
| A note on infinite partitions of free products of Boolean algebras | 245 |
| MARIO JARDÓN SANTOS | |
| On Matlis reflexive modules | 257 |
| HENNING KRAUSE | |
| The ramification tree and almost Dedekind domains of prescribed SP-rank | 273 |
| BALINT RAGO and DARIO SPIRITO | |
| The Gysin braid for S^3-actions on manifolds | 297 |
| JOSÉ IGNACIO ROYO PRIETO and MARTINTXO SARALEGI-ARANGUREN | |
| A conjecture on multiplicities for strongly tempered spherical varieties | 339 |
| CHEN WAN and LEI ZHANG | |
| Curvatures of real connections on Hermitian manifolds | 365 |
| JUN WANG and XIAOKUI YANG | |