

*Pacific
Journal of
Mathematics*

**MAPPING CLASSES FIXING AN ISOTROPIC HOMOLOGY
CLASS OF MINIMAL GENUS 0 IN RATIONAL 4-MANIFOLDS**

SERAPHINA EUN BI LEE

MAPPING CLASSES FIXING AN ISOTROPIC HOMOLOGY CLASS OF MINIMAL GENUS 0 IN RATIONAL 4-MANIFOLDS

SERAPHINA EUN BI LEE

For any $N \geq 1$, let M_N denote the rational 4-manifold $\mathbb{C}P^2 \# N \overline{\mathbb{C}P^2}$. We study the stabilizer $\text{Stab}(w)$ of a primitive, isotropic class $w \in H_2(M_N; \mathbb{Z})$ of minimal genus 0 under the natural action of the topological mapping class group $\text{Mod}(M_N)$ on $H_2(M_N; \mathbb{Z})$. Although most elements of $\text{Stab}(w)$ cannot be represented by homeomorphisms that preserve any Lefschetz fibration $M_N \rightarrow \Sigma$, we show that every element of $\text{Stab}(w)$ can be represented by a diffeomorphism that *almost preserves* a holomorphic, genus-0 Lefschetz fibration $\text{proj} : M_N \rightarrow \mathbb{C}P^1$ whose generic fibers represent the homology class w . We also answer the Nielsen realization problem for a certain maximal torsion-free, abelian subgroup Λ_w of $\text{Mod}(M_N)$ by finding a lift of Λ_w to $\text{Diff}^+(M_N) \leq \text{Homeo}^+(M_N)$ under the quotient map $q : \text{Homeo}^+(M_N) \rightarrow \text{Mod}(M_N)$. This lift of Λ_w can be made to almost preserve $\text{proj} : M_N \rightarrow \mathbb{C}P^1$. All results of this paper also hold for every primitive, isotropic class $w \in H_2(M_N; \mathbb{Z})$ if $N \leq 8$ because any such class has minimal genus 0.

1. Introduction

The (topological) *mapping class group* $\text{Mod}(M)$ of a closed, oriented manifold M is the group

$$\text{Mod}(M) := \pi_0(\text{Homeo}^+(M))$$

of isotopy classes of orientation-preserving homeomorphisms of M . There is a natural action of $\text{Mod}(M)$ on $H_2(M; \mathbb{Z})$ preserving the intersection form Q_M and we consider the stabilizer $\text{Stab}(w) \leq \text{Mod}(M)$ of any class $w \in H_2(M; \mathbb{Z})$.

Suppose M is a smooth, simply connected 4-manifold. If $w \in H_2(M; \mathbb{Z})$ is a nonzero homology class with self-intersection 0 then w is called *isotropic*. One way in which isotropic classes arise are as the homology class of the generic fibers of a Lefschetz fibration $p : M \rightarrow \Sigma$ where Σ is a closed, oriented surface. We say that a diffeomorphism φ of M *preserves* p if there exists some diffeomorphism ψ of Σ such that $p \circ \varphi = \psi \circ p$.

MSC2020: 57K40, 57S25.

Keywords: rational surface, mapping class group, Nielsen realization problem, Lefschetz fibration.

In some settings, elements $g \in \text{Stab}(w) \leq \text{Mod}(M)$ are known to admit representative maps φ that preserve some Lefschetz fibration $p : M \rightarrow \Sigma$ whose generic fibers represent the homology class w . For example, Gizatullin ([12]) showed that any parabolic automorphism of a compact Kähler surface M must preserve some elliptic fibration $M \rightarrow \Sigma$ (also see [4, Proposition 1.4] or [7, Theorem 4.3, Appendix]). In the smooth setting, Farb–Looijenga ([9, Theorem 1.11]) showed that on a K3 manifold, any $g \in \text{Stab}(w)$ can be represented by a diffeomorphism preserving the fibers of some holomorphic elliptic fibration $M \rightarrow \mathbb{C}\mathbb{P}^1$; this result is an application of their study of the moduli space of genus one fibered K3 surfaces with only nodal singular fibers. As another application, Farb–Looijenga ([9, Corollary 1.12]) study the Nielsen realization problem for a certain rank-20 free abelian subgroup of $\text{Stab}(w)$ by diffeomorphisms preserving the fibers of a given genus one fibration of a K3 manifold.

In this paper we study representative maps of the stabilizers of isotropic classes of *rational manifolds* M and their relationships to genus-0 Lefschetz fibrations $M \rightarrow \Sigma$. More specifically, we study manifolds of the form

$$M_N := \mathbb{C}\mathbb{P}^2 \# N \overline{\mathbb{C}\mathbb{P}^2} \quad \text{for } N \geq 1,$$

which are the underlying smooth 4-manifolds of the blowup of $\mathbb{C}\mathbb{P}^2$ at N points. The total space M of a nontrivial genus-0 Lefschetz fibration $M \rightarrow \Sigma$ is a rational manifold M_N for some N if M is simply connected. If $N \leq 8$, all primitive, isotropic classes $w \in H_2(M_N; \mathbb{Z})$ are represented by generic fibers of a genus-0, holomorphic Lefschetz fibration $p : M_N \rightarrow \mathbb{C}\mathbb{P}^1$. We sometimes refer to such a Lefschetz fibration as a *conic bundle structure on M_N* . Note that these Lefschetz fibrations are not relatively minimal unless $N = 1$. See [Section 2.3](#).

Representing $\text{Stab}(w)$ by diffeomorphisms. Let $N \geq 1$ and let $w \in H_2(M_N; \mathbb{Z})$ be any primitive, isotropic class of minimal genus 0. Although any such class w is represented by a generic fiber of a genus-0 Lefschetz fibration $p : M_N \rightarrow \mathbb{C}\mathbb{P}^1$, the following proposition shows that there does not exist any $\varphi \in \text{Homeo}^+(M_N)$ with $[\varphi] \in \text{Stab}(w)$ that preserves such a fibration p if $[\varphi]$ has infinite order in $\text{Mod}(M_N)$.

Proposition 1.1. *Let $N \geq 1$ and let $w \in H_2(M_N; \mathbb{Z})$ be a primitive, isotropic class of minimal genus 0. Let $\varphi \in \text{Homeo}^+(M_N)$ represent an infinite-order mapping class $[\varphi] \in \text{Stab}(w) \leq \text{Mod}(M_N)$. There does not exist any Lefschetz fibration $p : M_N \rightarrow \Sigma$ where Σ is a closed, oriented surface and where the generic fiber represents w such that φ preserves p .*

For a proof, see [Section 2.3](#). In this paper we ask instead that any diffeomorphism representing any infinite-order mapping class $f \in \text{Stab}(w) \leq \text{Mod}(M_N)$ *almost preserves* some Lefschetz fibration $p : M_N \rightarrow \Sigma$.

Definition 1.2 (almost preserving a Lefschetz fibration). A group of diffeomorphisms $G \leq \text{Diff}^+(M)$ *almost preserves* a Lefschetz fibration $p : M \rightarrow \Sigma$ if the elements of G act on the fibers of p outside of disjoint neighborhoods of the singular fibers of p . More precisely, there exist

- (a) disjoint, open neighborhoods $V_1, \dots, V_m \subseteq \Sigma$ of the images of the singular points $z_1, \dots, z_m \in \Sigma$, and
- (b) a homomorphism $i : G \rightarrow \text{Diff}^+(\Sigma - \bigcup_{k=1}^m V_k)$

such that for all $\varphi \in G$, the following commutes:

$$\begin{array}{ccc}
 M - \bigcup_{k=1}^m p^{-1}(V_k) & \xrightarrow{\varphi} & M - \bigcup_{k=1}^m p^{-1}(V_k) \\
 \downarrow p & & \downarrow p \\
 \Sigma - \bigcup_{k=1}^m V_k & \xrightarrow{i(\varphi)} & \Sigma - \bigcup_{k=1}^m V_k
 \end{array}$$

A diffeomorphism $\varphi \in \text{Diff}^+(M)$ *almost preserves* a Lefschetz fibration $p : M \rightarrow \Sigma$ if the group $\langle \varphi \rangle \leq \text{Diff}^+(M)$ almost preserves $p : M \rightarrow \Sigma$.

On the other hand, any element of $\text{Stab}(w) \leq \text{Mod}(M_N)$ with $N \geq 2$ must preserve the following subgroup of $H_2(M_N; \mathbb{Z})$:

$$w^\perp := \{w_0 \in H_2(M_N; \mathbb{Z}) : \mathcal{Q}_{M_N}(w, w_0) = 0\} \cong \mathbb{Z}^N.$$

Thus $\text{Stab}(w)$ acts on the lattice $(w^\perp/\mathbb{Z}\{w\}, \bar{\mathcal{Q}}_{M_N})$ where $\bar{\mathcal{Q}}_{M_N}$ is the unimodular, symmetric, bilinear form on $w^\perp/\mathbb{Z}\{w\}$ induced by \mathcal{Q}_{M_N} . Since $(H_2(M_N; \mathbb{Z}), \mathcal{Q}_{M_N})$ has signature $(1, N)$, $(w^\perp/\mathbb{Z}\{w\}, \bar{\mathcal{Q}}_{M_N})$ must be negative definite of rank $N - 1$.

Definition 1.3. Let Λ_w be the kernel of the map $\text{Stab}(w) \rightarrow \text{Aut}(w^\perp/\mathbb{Z}\{w\}, \bar{\mathcal{Q}}_{M_N})$.

There is an identification of Λ_w with the subgroup of even elements of the lattice $(w^\perp/\mathbb{Z}\{w\}, \bar{\mathcal{Q}}_{M_N})$, and $\text{Stab}(w)$ fits into a split short exact sequence

$$(1) \quad 0 \rightarrow \underbrace{\Lambda_w}_{\cong \mathbb{Z}^{N-1} \leq w^\perp/\mathbb{Z}\{w\}} \rightarrow \text{Stab}(w) \rightarrow \text{Aut}(w^\perp/\mathbb{Z}\{w\}, \bar{\mathcal{Q}}_{M_N}) \rightarrow 0.$$

Two properties of Λ_w are that it is a maximal torsion-free, abelian subgroup of $\text{Mod}(M_N)$ and that it has finite index in $\text{Stab}(w)$. See Lemmas 2.5 and 2.6.

With the preliminaries above in hand, we state our main result concerning the Nielsen realization problem for Λ_w .

Theorem 1.4 (realizing Λ_w by diffeomorphisms). *Let $N \geq 2$ and let $w \in H_2(M_N; \mathbb{Z})$ be a primitive, isotropic class of minimal genus 0. There exists a homomorphism*

$\rho_w : \Lambda_w \rightarrow \text{Diff}^+(M_N)$ such that the following diagram commutes:

$$\begin{array}{ccc}
 & & \text{Diff}^+(M_N) \\
 & \nearrow \rho_w & \downarrow q \\
 \Lambda_w & \hookrightarrow & \text{Mod}(M_N)
 \end{array}$$

The image $\rho_w(\Lambda_w)$ almost preserves a holomorphic genus-0 Lefschetz fibration $p : M_N \rightarrow \mathbb{C}\mathbb{P}^1$ whose generic fiber represents the homology class w .

If $N \leq 8$, [Theorem 1.4](#) holds for any primitive, isotropic class $w \in H_2(M_N; \mathbb{Z})$ because any such class has minimal genus 0. See [Corollary 3.10](#).

Compare [Theorem 1.4](#) to the case of the K3 manifold M for which the subgroup $\Lambda_w \leq \text{Mod}(M)$ is isomorphic to \mathbb{Z}^{20} , where $w \in H_2(M; \mathbb{Z})$ is a fiber class of a genus-1 fibration of M with only nodal fibers. As mentioned above, Farb–Looijenga ([\[9, Corollary 1.12\]](#)) showed that Λ_w lifts to the group of diffeomorphisms preserving the fibers of the given genus-1 fibration. In contrast to the case of the K3 manifold, [Theorem 1.4](#) shows that Nielsen realization for Λ_w holds in our setting despite the fact that no element of Λ_w can preserve any genus-0 Lefschetz fibration of M_N ([Proposition 1.1](#)).

The next theorem uses the short exact sequence [\(1\)](#) and the diffeomorphisms constructed in the proof of [Theorem 1.4](#) to find a diffeomorphism representative of any element of $\text{Stab}(w)$ that almost preserves a genus-0 Lefschetz fibration.

Theorem 1.5 (mapping classes fixing an isotropic class). *Let $N \geq 1$ and let $w \in H_2(M_N; \mathbb{Z})$ be a primitive, isotropic class of minimal genus 0. For any $h \in \text{Stab}(w)$, there exists $\varphi \in \text{Diff}^+(M_N)$ almost preserving a holomorphic genus-0 Lefschetz fibration $p : M_N \rightarrow \mathbb{C}\mathbb{P}^1$ whose generic fiber represents the homology class w such that $[\varphi] = h \in \text{Mod}(M_N)$.*

Similarly as with [Theorem 1.4](#), [Theorem 1.5](#) holds for any primitive, isotropic class $w \in H_2(M_N; \mathbb{Z})$ if $2 \leq N \leq 8$. See [Corollary 4.2](#).

A large part of the work of this paper is to ensure that the diffeomorphisms constructed in [Section 3](#) commute as diffeomorphisms of M_N . We point out that the calculations of [Section 3](#) are essential to the proof of [Theorem 1.4](#) regarding the Nielsen realization problem for Λ_w although [Theorem 1.5](#) alone may be proven more succinctly. For the sake of concreteness, we give explicit constructions of all diffeomorphisms used in this paper.

One way to interpret the results of this paper is via the natural action of (an index-2 subgroup of) $\text{Mod}(M_N)$ on \mathbb{H}^N and the classification of hyperbolic isometries into three types: elliptic, parabolic, and hyperbolic. Infinite-order elements of the stabilizer $\text{Stab}(w)$ for an isotropic class $w \in H_2(M_N; \mathbb{Z})$ are precisely the elements

of $\text{Mod}(M_N)$ acting by parabolic isometries on \mathbb{H}^N (Lemma 2.2). Therefore the following is an immediate corollary of Theorem 1.5.

Corollary 1.6. *Let $2 \leq N \leq 8$. If $g \in \text{Mod}(M_N)$ acts by a parabolic isometry on \mathbb{H}^N then there exists $\varphi \in \text{Diff}^+(M_N)$ that almost preserves a holomorphic genus-0 Lefschetz fibration $p : M_N \rightarrow \mathbb{C}P^1$ such that $[\varphi] = g$.*

Related work. The relationship between mapping classes of 4-manifolds fixing an isotropic class and Lefschetz fibrations with the prescribed generic fiber has been studied in some settings. As mentioned above, see Gizatullin [12] and Cantat [4] for the case of compact, Kähler surfaces and elliptic fibrations and Farb–Looijenga [9] for the case of K3 manifolds; [9] was an inspiration for this current paper.

Automorphisms preserving a genus-0 Lefschetz fibration (or a conic bundle structure) also play an important role in the study of finite groups of automorphisms of M_N . An example of such a complex automorphism is the *de Jonquières* involution, which is a main tool for this paper. Some examples of work in this direction include the classification of order-2 birational automorphisms of $\mathbb{C}P^2$ up to conjugacy (Bertini [2], Bayle–Beauville [1]) and finite subgroups of birational automorphisms of $\mathbb{C}P^2$ in general (Dolgachev–Iskovskikh [8], Blanc [3]) in the complex category and a study of finite groups of symplectomorphisms of rational surfaces (Chen–Li–Wu [5]) in the symplectic category.

Outline. In Section 2, we recall relevant facts about the mapping class group $\text{Mod}(M_N)$ of rational manifolds and deduce basic facts about isotropic classes $w \in H_2(M_N; \mathbb{Z})$, including the proof of Proposition 1.1. In Section 3, we prove Theorem 1.4 by explicitly constructing the necessary diffeomorphisms. Using these diffeomorphisms from Section 3, we prove Theorem 1.5 in Section 4.

2. Isotropic homology classes and their stabilizers in $\text{Mod}(M_N)$

We collect useful properties of the mapping class groups of 4-manifolds, isotropic classes in $H_2(M_N; \mathbb{Z})$, and certain Lefschetz fibrations.

2.1. The mapping class group of M_N . For any 4-manifold M , let Q_M denote the intersection form on $H_2(M; \mathbb{Z})$. The form Q_M is an integral, unimodular, nondegenerate, symmetric bilinear form on $H_2(M; \mathbb{Z})/\text{Torsion}$; we denote the lattice $(H_2(M; \mathbb{Z})/\text{Torsion}, Q_M)$ by H_M . The automorphism group of the lattice H_M is denoted $O(H_M)$.

The mapping class group $\text{Mod}(M) := \pi_0(\text{Homeo}^+(M))$ of a closed, oriented, simply connected 4-manifold M is computable due to the following theorems of Freedman [10], Perron [18], Quinn [19], Cochran–Habegger [6], and Gabai–Gay–Hartman–Krushkal–Powell [11]. (For a more detailed history of this theorem, see [11, Section 1.3].)

Theorem 2.1. *Let M^4 be a closed, oriented, and simply connected manifold. The map*

$$\Phi : \text{Mod}(M) \rightarrow \text{O}(H_M)$$

given by $\Phi : [\varphi] \mapsto \varphi_*$ is an isomorphism of groups.

By the Mayer–Vietoris sequence, $H_2(M_N; \mathbb{Z}) = H_2(\mathbb{C}\mathbb{P}^2; \mathbb{Z}) \oplus H_2(\overline{\mathbb{C}\mathbb{P}^2}; \mathbb{Z})^{\oplus N}$, and we have the usual \mathbb{Z} -basis $\{H, E_1, \dots, E_N\}$. The intersection form Q_{M_N} is given by the diagonal, $(N + 1) \times (N + 1)$ matrix

$$\text{diag}(1, -1, \dots, -1)$$

with respect to the \mathbb{Z} -basis $\{H, E_1, \dots, E_N\}$. On the other hand, there is a natural \mathbb{Z} -basis

$$(2) \quad \{s, v, e_1, \dots, e_{N-1}\}$$

of $H_2((\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1) \# (N - 1)\overline{\mathbb{C}\mathbb{P}^2}; \mathbb{Z})$ via the Mayer–Vietoris sequence; here, s and v correspond to the first and second factors of $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$ respectively. There is a diffeomorphism $(\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1) \# (N - 1)\overline{\mathbb{C}\mathbb{P}^2} \cong M_N$ for all $N \geq 2$ giving an identification

$$v = H - E_1, \quad s = H - E_2, \quad e_1 = H - E_1 - E_2, \quad e_k = E_{k+1} \text{ for all } 2 \leq k \leq N - 1.$$

We will mostly work with the \mathbb{Z} -basis $\{s, v, e_1, \dots, e_{N-1}\}$ of $H_2(M_N; \mathbb{Z})$.

Therefore by [Theorem 2.1](#),

$$\text{Mod}(M_N) \cong \text{O}(1, N)(\mathbb{Z}) := \text{O}(H_{M_N})$$

We will identify $\text{O}(H_{M_N})$ and $\text{Mod}(M_N)$ throughout this paper.

On the other hand, consider $E^{1,N} := (\mathbb{R}^{N+1}, Q_N)$, where Q_N is the diagonal bilinear symmetric form of signature $(1, N)$:

$$Q_N((x_0, x_1, \dots, x_N), (y_0, y_1, \dots, y_N)) = x_0 y_0 - x_1 y_1 - \dots - x_N y_N.$$

There is a natural identification of the \mathbb{R} -span of the \mathbb{Z} -basis $\{H, E_1, \dots, E_N\}$ of $H_2(M_N; \mathbb{Z})$ with \mathbb{R}^{N+1} , under which the \mathbb{R} -bilinear extension of Q_{M_N} coincides with Q_N . The hyperboloid model for \mathbb{H}^N sits in $E^{1,N}$ by

$$\mathbb{H}^N = \{w = (w_0, w_1, \dots, w_N) \in \mathbb{R}^{N+1} : Q_N(w, w) = 1, w_0 > 0\}.$$

where the Riemannian metric is defined by the restriction of $-Q_N$ to \mathbb{H}^N (see [\[20, Chapter 2\]](#)). Because $\text{O}(1, N)(\mathbb{Z})$ acts on \mathbb{R}^{N+1} and preserves Q_N , it contains an index-2 subgroup $\text{O}^+(1, N)(\mathbb{Z})$ acting by isometries on \mathbb{H}^N .

The boundary sphere of \mathbb{H}^N corresponds to

$$\partial\mathbb{H}^N = \{w = (w_0, w_1, \dots, w_N) \in \mathbb{R}^{N+1} : Q_N(w, w) = 0, w_0 > 0\} / \sim$$

where $aw \sim w$ for all $a \in \mathbb{R}_{>0}$. Parabolic isometries of \mathbb{H}^N are those that fix a unique point of $\partial\mathbb{H}^n$ and no point in \mathbb{H}^n . By [20, Problem 2.5.24(g)], parabolic isometries not only preserve some line in \mathbb{R}^{N+1} but fix it pointwise. Moreover, parabolic isometries in $O^+(1, N)(\mathbb{Z})$ must fix a nonzero, isotropic vector with integral entries, i.e., some nonzero $w \in H_2(M_N; \mathbb{Z})$ with $Q_{M_N}(w, w) = 0$, and have infinite order ([20, Exercise 2.5.20]). The converse is true as well:

Lemma 2.2. *Let $N \geq 2$. An element $f \in O^+(1, N)(\mathbb{Z})$ acts by a parabolic isometry if and only if f has infinite order and there exists some primitive, isotropic class $w \in H_2(M_N; \mathbb{Z})$ such that $f \in \text{Stab}(w) \leq O(H_{M_N})$.*

Proof. One direction holds by the discussion preceding the statement of the lemma, so it suffices to prove that if $f \in \text{Stab}(w)$ has infinite order then f acts on \mathbb{H}^N by a parabolic isometry.

Note that f fixes the point of $\partial\mathbb{H}^N$ corresponding to $w \in H_2(M_N; \mathbb{Z})$. Because stabilizers of points in \mathbb{H}^N in $O(H_{M_N})$ have finite order, we only need to show that this is the unique point of $\partial\mathbb{H}^N$ fixed by f . To do this, let $w_0 \in \mathbb{E}^{1,N}$ be an isotropic vector such that $f(w_0) = \lambda w_0$ for some $\lambda \in \mathbb{R}$. If $w_0 \in \mathbb{R}\{w\}^\perp$ then w_0 must be a scalar multiple of w because the restriction of Q_{M_N} to $\mathbb{R}\{w\}^\perp/\mathbb{R}\{w\}$ is negative definite. If $Q_{M_N}(w, w_0) =: a \neq 0$ then $\lambda = 1$ because

$$a = Q_{M_N}(w, w_0) = Q_{M_N}(f(w), f(w_0)) = Q_{M_N}(w, \lambda w_0) = \lambda a.$$

Then $f(aw + w_0) = aw + w_0$ and

$$Q_{M_N}(aw + w_0, aw + w_0) = 2aQ_{M_N}(w, w_0) = 2a^2 > 0.$$

A scalar multiple of $aw + w_0$ lies in \mathbb{H}^N , meaning f acts on \mathbb{H}^N by an elliptic isometry, and all such isometries of \mathbb{H}^N in $O(H_{M_N})$ have finite order. Therefore, w_0 must be a scalar multiple of w and hence f fixes a unique point in $\partial\mathbb{H}^N$. \square

2.2. Primitive, isotropic classes $w \in H_2(M_N; \mathbb{Z})$ and $\text{Stab}(w) \leq \text{Mod}(M_N)$. Consider lattices (L, Q) , where $L \cong \mathbb{Z}^r$ as an abelian group for some $r \in \mathbb{N}$ and Q is an integral, unimodular, nondegenerate, symmetric, bilinear form on L . For each primitive, isotropic vector $w \in L$, there exists $u \in L$ such that $Q(w, u) = 1$ by unimodularity of Q . There is an orthogonal decomposition

$$L = \mathbb{Z}\{u, w\} \oplus \mathbb{Z}\{u, w\}^\perp$$

to which Q restricts to a unimodular form on each factor. The restriction of Q to $\mathbb{Z}\{u, w\}$ has signature $(1, 1)$. Note that $\mathbb{Z}\{u, w\}^\perp$ is a lift of $w^\perp/\mathbb{Z}\{w\}$ under the natural quotient $w^\perp \rightarrow w^\perp/\mathbb{Z}\{w\}$. This means that $(\mathbb{Z}\{u, w\}^\perp, Q|_{\mathbb{Z}\{u, w\}^\perp})$ is isometric (i.e., isomorphic as a lattice) to $(w^\perp/\mathbb{Z}\{w\}, \bar{Q})$ via this quotient, where \bar{Q} is the induced bilinear form on $w^\perp/\mathbb{Z}\{w\}$. We fix the above notation throughout this subsection.

Lemma 2.3. *Let $w \in L$ be a primitive, isotropic vector. If $h_1, h_2 \in \text{Stab}(w) \leq \text{O}(L, Q)$ and $h_1|_{w^\perp} = h_2|_{w^\perp}$ then $h_1 = h_2$. In particular, for any $h_1, h_2 \in \text{Stab}(v) \leq \text{O}(H_{M_N})$, where v is the homology class as given in (2) and $N \geq 2$, if $h_1(e_k) = h_2(e_k)$ for all $1 \leq k \leq N - 1$ then $h_1 = h_2$.*

Proof. Observe that $h_1^{-1} \circ h_2$ acts as the identity on $\mathbb{Z}\{u, w\}^\perp \leq w^\perp$. We claim that $h_1^{-1} \circ h_2(u) = u$. To see this, write $h_1^{-1} \circ h_2(u) = au + bw + v_0$ for some $v_0 \in \mathbb{Z}\{u, w\}^\perp$ and some $a, b \in \mathbb{Z}$. For any $v_1 \in \mathbb{Z}\{u, w\}^\perp$,

$$0 = Q(u, v_1) = Q((h_1^{-1} \circ h_2)(u), (h_1^{-1} \circ h_2)(v_1)) = Q(au + bw + v_0, v_1) = Q(v_0, v_1).$$

Therefore, $v_0 = 0$ by unimodularity of $Q|_{\mathbb{Z}\{u, w\}^\perp}$. Moreover, $(h_1^{-1} \circ h_2)(w) = w$, so

$$\begin{aligned} 1 &= Q((h_1^{-1} \circ h_2)(u), (h_1^{-1} \circ h_2)(w)) = Q(au + bw, w) = a, \\ 0 &= Q((h_1^{-1} \circ h_2)(u), (h_1^{-1} \circ h_2)(u)) = Q(u + bw, u + bw) = 2b. \end{aligned}$$

Therefore, $(h_1^{-1} \circ h_2)(u) = u$ and $h_1^{-1} \circ h_2$ restricts to the identity on $\mathbb{Z}\{u, w\}$. In the case of $v \in H_2(M_N; \mathbb{Z})$ for any $N \geq 2$, apply the above argument with $w = v$, $u = s$ and $\mathbb{Z}\{u, w\}^\perp = \mathbb{Z}\{e_1, \dots, e_{N-1}\}$. □

Let Λ_w denote the kernel of the natural map $h_w : \text{Stab}(w) \rightarrow \text{O}(w^\perp/\mathbb{Z}\{w\}, \bar{Q})$ (cf. Definition 1.3). In order to describe Λ_w , we introduce an important type of element of $\text{O}(H_{M_N})$ used throughout this paper.

Definition 2.4. Let $N \geq 2$ and $u \in H_2(M_N; \mathbb{Z})$ satisfy $Q_{M_N}(u, u) = \pm 1$ or ± 2 . The reflection Ref_u about u is an element of $\text{O}(H_{M_N})$ defined by

$$\text{Ref}_u(x) = x - \frac{2Q_{M_N}(x, u)}{Q_{M_N}(u, u)}u.$$

We now use reflections and Eichler transformations to give generators for Λ_w .

Lemma 2.5. *Let (L, Q) be any lattice and $w \in L$ be a primitive, isotropic vector. Let $A \leq w^\perp/\mathbb{Z}\{w\}$ denote the subgroup of even elements with respect to \bar{Q} . Then there is an isomorphism of groups*

$$E(w, \cdot) : A \rightarrow \Lambda_w.$$

In the case that $(L, Q) = H_{M_N}$ for any $N \geq 2$ and $w = v$, the group Λ_w is generated by

$$f_k := \text{Ref}_{e_k} \circ \text{Ref}_{e_{k+1}} \circ \text{Ref}_{v - e_k - e_{k+1}} \circ \text{Ref}_{e_k - e_{k+1}}$$

for $1 \leq k \leq N - 2$ and $g := \text{Ref}_{e_1} \circ \text{Ref}_{v - e_1}$.

Proof. For any $f \in \Lambda_w$, there exists $c(f) \in w^\perp/\mathbb{Z}\{w\}$ such that for any $e \in w^\perp$,

$$f(e) = e - \bar{Q}(c(f), e)w$$

by the definition of Λ_w and the unimodularity of \bar{Q} . This defines a homomorphism $c : \Lambda_w \rightarrow w^\perp/\mathbb{Z}\{w\}$ which is injective by [Lemma 2.3](#).

For any $f \in \Lambda_w$, there exists $a \in \mathbb{Z}$ and $e \in \mathbb{Z}\{u, w\}^\perp$ such that

$$f(u) = u + aw + e$$

because $Q(f(u), w) = 1$. (Here, $u \in L$ is as chosen at the beginning of this subsection.) Moreover,

$$Q(u, u) = Q(f(u), f(u)) = Q(u, u) + 2a + Q(e, e)$$

and so $Q(e, e)$ is even. Because $c(f), e$ are contained in $\mathbb{Z}\{u, w\}^\perp$ where $c(f) \in w^\perp/\mathbb{Z}\{w\}$ is identified with its lift in $\mathbb{Z}\{u, w\}^\perp$,

$$\begin{aligned} 0 &= Q(f(u), f(c(f))) = Q(f(u), c(f) - Q(c(f), c(f))w) \\ &= Q(e, c(f)) - Q(c(f), c(f)), \end{aligned}$$

$$0 = Q(f(u), f(e)) = Q(f(u), e - Q(e, c(f))w) = Q(e, e) - Q(e, c(f)).$$

By the second string of equalities, $Q(e, c(f))$ is even, and by the first, $Q(c(f), c(f)) = \bar{Q}(c(f), c(f))$ is even. Hence $c(\Lambda_w) \leq A$.

Consider the homomorphism $E(w, \cdot) : A \rightarrow \Lambda_w$ defined by

$$E(w, e) : x \mapsto x + Q(w, x)e - Q(e, x)w - \frac{1}{2}Q(e, e)Q(w, x)w$$

for each $[e] \in A \leq w^\perp/\mathbb{Z}\{w\}$ with $e \in w^\perp$, where $E(w, e)$ is an *Eichler transformation*. A computation shows that $E(w, \cdot)$ does not depend on the choice of lift $e \in w^\perp$, and hence descends to a well-defined homomorphism on $A \leq w^\perp/\mathbb{Z}\{w\}$. Another computation shows that $c \circ E(w, \cdot) = \text{Id}|_A$. Finally, if $(L, Q) = (H_2(M_N; \mathbb{Z}), Q_{M_N})$ and $w = v$, compute that $f_k = E(w, e_k + e_{k+1})$ for each $1 \leq k \leq N - 2$ and $g = E(w, 2e_1)$, which together generate Λ_w as $e_k + e_{k+1}$ with $1 \leq k \leq N - 2$ and $2e_1$ generate A . \square

We combine the results of this subsection and record an important algebraic property of $\text{Stab}(w)$. Below, $\text{O}(r)(\mathbb{Z})$ denotes the automorphism group of the diagonal lattice $(\mathbb{Z}^r, \text{diag}(1, \dots, 1))$, or equivalently, the automorphism group of the diagonal lattice $(\mathbb{Z}^r, \text{diag}(-1, \dots, -1))$.

Lemma 2.6. *For any primitive, isotropic vector $w \in L$, there is a split short exact sequence*

$$0 \rightarrow \Lambda_w \rightarrow \text{Stab}(w) \xrightarrow{h_w} \text{O}(w^\perp/\mathbb{Z}\{w\}, \bar{Q}) \rightarrow 0.$$

In the case that $(L, Q) = \text{H}_{M_N}$ for any $N \geq 2$ and $w = f(v)$ for any $f \in \text{O}(\text{H}_{M_N})$, the split short exact sequence above is isomorphic to

$$0 \rightarrow \mathbb{Z}^{N-1} \rightarrow \text{Stab}(w) \xrightarrow{h_w} \text{O}(N-1)(\mathbb{Z}) \rightarrow 0.$$

There is an equality of subgroups $\Lambda_w = f \circ \Lambda_v \circ f^{-1}$. Moreover, $\Lambda_w \cong \mathbb{Z}^{N-1}$ is a finite-index maximal torsion-free subgroup of $\text{Stab}(w)$ and a maximal torsion-free, abelian subgroup of $\text{O}(\text{H}_{M_N})$.

Proof. There is a section ℓ of h_w defined by

$$\ell : f \mapsto \text{Id} \oplus f \in \text{O}(\mathbb{Z}\{u, w\} \oplus \mathbb{Z}\{u, w\}^\perp, Q) = \text{O}(L, Q)$$

which shows that h_w is surjective and the sequence is split.

In the case of $(L, Q) = \text{H}_{M_N}$ with $N \geq 2$ and $w = f(v)$ for any $f \in \text{O}(\text{H}_{M_N})$, we can let $u = f(s)$, in which case

$$(w^\perp/\mathbb{Z}\{w\}, \bar{Q}) \cong (\mathbb{Z}\{f(e_1), \dots, f(e_{N-1})\}, Q_{M_N})$$

and so $\text{O}(w^\perp/\mathbb{Z}\{w\}, \bar{Q}) \cong \text{O}(N-1)(\mathbb{Z})$ is finite. The subgroup $A \leq w^\perp/\mathbb{Z}\{w\}$ of even elements with respect to \bar{Q} has index 2 in $w^\perp/\mathbb{Z}\{w\}$ which has rank $N-1$, and so $\Lambda_w \cong A \cong \mathbb{Z}^{N-1}$. Because the sequence is split, the subgroup $\langle \Lambda_w, h \rangle$ of $\text{Stab}(w)$ generated by Λ_w and h must have torsion for any $h \in \text{Stab}(w)$ with $h \notin \Lambda_w$ and so Λ_w is a maximal torsion-free subgroup of $\text{Stab}(w)$.

To see that $\Lambda_w = f \circ \Lambda_v \circ f^{-1}$, compute for any $h \in \Lambda_v$ and $e \in v^\perp$ that

$$(f \circ h \circ f^{-1})(f(e)) = f(h(e)) = f(e - \bar{Q}(c(h), e)v) = f(e) - \bar{Q}(c(h), e)w$$

for some $c(h) \in v^\perp/\mathbb{Z}\{v\}$ as in the proof of [Lemma 2.5](#) and where \bar{Q} is the bilinear form on $v^\perp/\mathbb{Z}\{v\}$ induced by Q . Because $w^\perp = f(v^\perp)$, we see that $f \circ h \circ f^{-1}$ induces the identity map on $w^\perp/\mathbb{Z}\{w\}$, showing that $f \circ \Lambda_v \circ f^{-1} \subseteq \Lambda_w$. By symmetry, it follows that $f \circ \Lambda_v \circ f^{-1} = \Lambda_w$. Each of the generators of Λ_v given in [Lemma 2.5](#) is contained in $\text{O}^+(1, N)(\mathbb{Z})$. Because $\Lambda_w = f \circ \Lambda_v \circ f^{-1}$ and $\text{O}^+(1, N)(\mathbb{Z})$ is a normal subgroup of $\text{O}(\text{H}_{M_N})$, we conclude that Λ_w is contained in $\text{O}^+(1, N)(\mathbb{Z})$.

It remains to show that Λ_w is a maximal torsion-free, abelian subgroup of $\text{O}(\text{H}_{M_N})$. To this end, consider any $h \in \Lambda_w$ with $h \neq \text{Id}$. Because $\Lambda_w \leq \text{O}^+(1, N)(\mathbb{Z})$ and Λ_w is torsion-free, [Lemma 2.2](#) shows that h is parabolic and $w \in H_2(M_N; \mathbb{Z})$ is the unique isotropic element of $H_2(M_N; \mathbb{Z})$ fixed by h , up to scaling. Suppose $k \in \text{O}(\text{H}_{M_N})$ commutes with some $h \in \Lambda_w$ and that $\langle k, \Lambda_w \rangle$ is torsion-free. Note that then $\langle -k, \Lambda_w \rangle$ is also torsion-free because $-\text{Id} \in \text{O}(\text{H}_{M_N})$ is in the center of $\text{O}(\text{H}_{M_N})$ and has order 2. Moreover, $k(w) = \pm w$ because h fixes $k(w)$, so $k \in \text{Stab}(w)$ or $-k \in \text{Stab}(w)$. If $-k \in \text{Stab}(w)$ then $\langle -k, \Lambda_w \rangle = \Lambda_w$ because $\langle -k, \Lambda_w \rangle$ is torsion-free and Λ_w is a maximal torsion-free subgroup of $\text{Stab}(w)$. However, $-k \notin \Lambda_w$ because $-k \circ k^{-1} = -\text{Id}$ is torsion and $\langle k, \Lambda_w \rangle$ is torsion-free. Therefore, $k \in \text{Stab}(w)$ and $k \in \Lambda_w$ since $\langle k, \Lambda_w \rangle$ is a torsion-free subgroup of $\text{Stab}(w)$. □

To use [Lemma 2.6](#), we apply a theorem of Li–Li [[15](#), Theorem 4.2] which says that for any $N \geq 2$ and any primitive, isotropic class $w \in H_2(M_N; \mathbb{Z})$ of minimal genus 0, there exists $\varphi \in \text{Diff}^+(M_N)$ such that $[\varphi](v) = w$. Moreover, following elementary lemma strengthens this theorem in the case $2 \leq N \leq 8$. Recall the fixed \mathbb{Z} -basis $\{s, v, e_1, \dots, e_{N-1}\}$ of $H_2(M_N; \mathbb{Z})$ given in [\(2\)](#).

Lemma 2.7. *If $2 \leq N \leq 8$ and $w \in H_2(M_N; \mathbb{Z})$ is an isotropic class, then*

- (a) *there exists $f \in \text{O}(H_{M_N})$ such that $f(v) = w$ if w is primitive, and*
- (b) *w has minimal genus 0.*

Proof. To prove [\(a\)](#), suppose w is primitive and $u \in H_{M_N}$ is chosen as in the beginning of this subsection. The restriction of Q_{M_N} to $\mathbb{Z}\{w, u\}$ is unimodular and indefinite so $\mathbb{Z}\{w, u\}^\perp$ is negative definite of rank $N - 1 < 8$. There exists a unique unimodular and negative definite lattice of rank r if $r \leq 7$; see [[16](#), p. 1], for example. Therefore, $(\mathbb{Z}\{w, u\}^\perp, Q_{M_N}|_{\mathbb{Z}\{w, u\}^\perp})$ is isometric to $(\mathbb{Z}\{e_1, \dots, e_{N-1}\}, Q_{M_N}|_{\mathbb{Z}\{e_1, \dots, e_{N-1}\}})$; let $w_0 \in \mathbb{Z}\{w, u\}^\perp$ satisfy $Q_{M_N}(w_0, w_0) = -1$.

With $a := Q_{M_N}(u, u)$, we have $Q_{M_N}(w, u - aw_0) = 1$ and

$$Q_{M_N}(u - aw_0, u - aw_0) = a - a^2 \equiv 0 \pmod{2}.$$

So $\mathbb{Z}\{w, u - aw_0\}$ is unimodular, even, and indefinite. Again, $\mathbb{Z}\{w, u - aw_0\}^\perp$ is negative definite of rank $N - 1 < 8$, and so $(\mathbb{Z}\{w, u - aw_0\}^\perp, Q_{M_N}|_{\mathbb{Z}\{w, u - aw_0\}^\perp})$ is isometric to $(\mathbb{Z}\{e_1, \dots, e_{N-1}\}, Q_{M_N}|_{\mathbb{Z}\{e_1, \dots, e_{N-1}\}})$. There exists $f \in \text{O}(H_{M_N})$ that preserves the orthogonal direct sums below

$$f : \mathbb{Z}\{v, s\} \oplus \mathbb{Z}\{e_1, \dots, e_{N-1}\} \rightarrow \mathbb{Z}\{w, u - aw_0\} \oplus \mathbb{Z}\{w, u - aw_0\}^\perp$$

such that $f(v) = w$. This proves [\(a\)](#).

To prove [\(b\)](#), we may assume that $w \neq 0$. Suppose $w_1 \in H_2(M_N; \mathbb{Z})$ is a primitive isotropic class such that $aw_1 = w$ for some $a \in \mathbb{Z}$. By [\(a\)](#), there exists some $f \in \text{Mod}(M_N)$ such that $f(v) = w_1$. Because $N \leq 9$, there exists a diffeomorphism $\varphi \in \text{Diff}^+(M_N)$ such that $[\varphi] = f$ by [[21](#), Theorem 2], and so the minimal genus of w and the minimal genus of av are equal, and the minimal genus of $av = a(H - E_1)$ is 0 (cf. [[15](#), Theorem 4.2]). \square

2.3. Lefschetz fibrations, conic bundles, and de Jonquières involutions. Let $N = 2m + 1 \geq 3$ be odd and fix some distinct complex numbers $a_1, \dots, a_{2m} \in \mathbb{C}$. Consider the birational map $\mathcal{J}_0 : \mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1 \dashrightarrow \mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$ given by

$$([X_1 : X_2], [Y_1 : Y_2]) \mapsto \left([X_1 : X_2], \left[Y_2 \prod_{i=m+1}^{2m} (X_1 - a_i X_2) : Y_1 \prod_{i=1}^m (X_1 - a_i X_2) \right] \right).$$




Figure 1. Each line represents a copy of $\mathbb{C}\mathbb{P}^1$ and is labeled with its homology class in M_N . The rightmost fiber, for $1 \leq i \leq N - 1$, is a singular fiber. Each singular fiber is a union of two (-1) -spheres intersecting transversely once.

Then \mathcal{J}_0 lifts to an automorphism \mathcal{J} of order 2 called a *de Jonquieres involution* of $X := \text{Bl}_P(\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1)$ where

$$P := \{([a_i : 1], [1 : 0]) : 1 \leq i \leq m\} \cup \{([a_i : 1], [0 : 1]) : m + 1 \leq i \leq 2m\}$$

is a set in $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$ with $2m$ points. Note that X is diffeomorphic to M_N . Under this identification, $e_k \in H_2(M_N; \mathbb{Z})$ is the class of the exceptional fiber above $([a_k : 1], [1 : 0])$ for each $1 \leq k \leq m$ and the class of the exceptional fiber above $([a_k : 1], [0 : 1])$ for each $m + 1 \leq k \leq 2m$.

The projection map $\text{pr}_0 : \mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1 \rightarrow \mathbb{C}\mathbb{P}^1$ onto the first coordinate extends to a map $\text{pr} : X \rightarrow \mathbb{C}\mathbb{P}^1$ defining a holomorphic genus-0 Lefschetz fibration (in other words, a conic bundle). By construction, $\text{pr} \circ \mathcal{J} = \text{pr}$.

If $z \neq z_k := [a_k : 1] \in \mathbb{C}\mathbb{P}^1$ for any k , the fiber of pr over a point $z \in \mathbb{C}\mathbb{P}^1$ is $\{z\} \times \mathbb{C}\mathbb{P}^1$ which is in the homology class $v \in H_2(M_N; \mathbb{Z})$. Because \mathcal{J} acts on each such $\text{pr}^{-1}(z)$ in an orientation-preserving way, $[\mathcal{J}] \in \text{Stab}(v) \leq \text{Mod}(M_N)$. Moreover for all $1 \leq k \leq 2m$ and all $([a_k : 1], [Y_1 : Y_2]) \notin P$,

$$\mathcal{J}_0 : ([a_k : 1], [Y_1 : Y_2]) \mapsto \begin{cases} ([a_k : 1], [1 : 0]) & \text{if } 1 \leq k \leq m, \\ ([a_k : 1], [0 : 1]) & \text{if } m + 1 \leq k \leq 2m. \end{cases}$$

Therefore, $[\mathcal{J}]$ must send the homology class $v - e_k$ of the strict transform of $\text{pr}^{-1}([a_k : 1])$ in X to the exceptional divisor e_k . See [Figure 1](#) for an illustration of the action of \mathcal{J} on the fibers of pr .

The maps pr and \mathcal{J} will be used in the explicit constructions in [Sections 3 and 4](#). The goal of the rest of this section is to show that it suffices to only consider the Lefschetz fibration $\text{pr} : M_N \rightarrow \mathbb{C}\mathbb{P}^1$ for our setting and to prove [Proposition 1.1](#).

Proposition 2.8. *Let $p : M_N \rightarrow \Sigma$ be a Lefschetz fibration where Σ is a closed, oriented surface and the generic fiber F satisfies $[F] \neq 0 \in H_2(M_N; \mathbb{Z})$. If $[F]$ has minimal genus 0 then $\Sigma = \mathbb{C}\mathbb{P}^1$ and $F = \mathbb{C}\mathbb{P}^1$.*

Proof. Because M_N is closed, a generic fiber F is a compact submanifold of M and has finitely many connected components, i.e., $\pi_0(F)$ is finite. By the exact sequence of the fibration [14, Proposition 8.1.9], there is a bijection $\pi_1(\Sigma) \rightarrow \pi_0(F)$ because $\pi_1(M_N) = 0$. Therefore, $\pi_1(\Sigma)$ is finite because $\pi_0(F)$ is finite. Because Σ is a closed, oriented surface, this implies that $\Sigma = \mathbb{C}\mathbb{P}^1$. Furthermore, F is connected since $\pi_0(F) = \pi_1(\mathbb{C}\mathbb{P}^1) = 0$.

Because $[F]$ is nontrivial in $H_2(M_N; \mathbb{R})$ since $H_2(M_N; \mathbb{Z})$ has no torsion, M_N can be given a symplectic structure such that F is a symplectic submanifold (Gompf [14, Theorem 10.2.18], [13, Theorem 1.2]) and so F must achieve the minimal genus in its homology class by the solution to the symplectic Thom conjecture (Oszváth–Szabó [17, Theorem 1.1]). \square

Proof of Proposition 1.1. Suppose there exists such a Lefschetz fibration $p : M_N \rightarrow \Sigma$ whose generic fiber represents w and a homeomorphism $h : \Sigma \rightarrow \Sigma$ with $p \circ \varphi = h \circ p$. Because w is nonzero and has minimal genus 0, Proposition 2.8 says that $\Sigma = \mathbb{C}\mathbb{P}^1$ and the generic fiber of p has genus 0. After blowing down the (-1) -spheres contained in the fibers of p , we see that p must be a $\mathbb{C}\mathbb{P}^1$ -bundle over Σ by [14, Proposition 8.1.7]. Because all $\mathbb{C}\mathbb{P}^1$ -bundles over $\mathbb{C}\mathbb{P}^1$ are holomorphic, M_N gets a complex structure as a rational surface and p is holomorphic.

We prove by induction on N that if some homeomorphism $\varphi \in \text{Homeo}^+(M_N)$ preserves a genus-0 Lefschetz fibration $p : M_N \rightarrow \mathbb{C}\mathbb{P}^1$ then $[\varphi] \in \text{Mod}(M_N)$ has finite order. If $N = 1$ then it is easy to see $\text{Mod}(M_N) = \text{O}(H_{M_N})$ is finite. Now assume for some $N_0 > 1$ that the claim holds for any $1 \leq N < N_0$.

Let $N = N_0$ and suppose $\varphi \in \text{Homeo}^+(M_N)$ preserves a genus-0 Lefschetz fibration $p : M_N \rightarrow \mathbb{C}\mathbb{P}^1$. Then φ must permute the singular fibers because none of the singular fibers are homeomorphic to a generic fiber $\mathbb{C}\mathbb{P}^1$. There are finitely many singular fibers, so some power φ^k must preserve each singular fiber. Each singular fiber F of p is a union of finitely many spheres of negative self-intersection intersecting transversely at finitely many points q_1, \dots, q_m . Because φ^k restricts to a homeomorphism of each singular fiber, φ^k must permute the points q_1, \dots, q_m . Moreover, φ^k also restricts to a homeomorphism on $F - \{q_1, \dots, q_m\}$, a disjoint union of finitely many spheres with punctures. Therefore, a further power $\varphi^{k\ell}$ must preserve each component of $F - \{q_1, \dots, q_m\}$ and its orientation.

Let $S \subseteq F$ be an embedded (-1) -sphere in M_N . Because the homeomorphism $\varphi^{k\ell}$ fixes each point q_1, \dots, q_m and preserves $S - (S \cap \{q_1, \dots, q_m\}) \subseteq F - \{q_1, \dots, q_m\}$, it must preserve $S \subseteq M_N$. Let $b : M_N \rightarrow M$ be the map that blows down S to a point $q \in M$. Being a rational surface, M is diffeomorphic to M_{N-1} or $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$.




Figure 2. The sets U_k , B_k , and V_k in the case $n = 7$. The set B_4 , shaded in blue, surrounds z_4 and z_5 . The annulus U_1 , shaded in orange, is the collar neighborhood of B_1 surrounding z_1 and z_2 . The disk V_6 , shaded in red, contains z_6 and contained in $(B_5 - U_5) \cap (B_6 - U_6)$.

Because $\varphi^{k\ell}$ defines a homeomorphism on $M_N - S$, it induces a homeomorphism of $M - q$ that extends to a homeomorphism ψ of M and preserves the Lefschetz fibration $p' : M \rightarrow \mathbb{C}\mathbb{P}^1$ such that $p = p' \circ b$. If M is diffeomorphic to M_{N-1} then $[\psi]$ has finite order in $\text{Mod}(M)$ by the inductive hypothesis. Otherwise, M is diffeomorphic to $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$ and so $\text{Mod}(M)$ is finite. Therefore, $[\psi]$ also has finite order in $\text{Mod}(M)$.

Finally, note that $b_* : H_2(M_N; \mathbb{Z}) \rightarrow H_2(M; \mathbb{Z})$ induces the quotient map

$$H_2(M_N; \mathbb{Z}) \cong \mathbb{Z}\{[S]\}^\perp \oplus \mathbb{Z}\{[S]\} \rightarrow \mathbb{Z}\{[S]\}^\perp \cong H_2(M; \mathbb{Z}).$$

Because $\psi \circ b = b \circ \varphi^{k\ell}$ and $\varphi_*^{k\ell}([S]) = [S]$, the restriction of $\varphi_*^{k\ell}$ to $\mathbb{Z}\{[S]\}^\perp$ must have the same order as ψ_* . Finally, this shows that $[\varphi^{k\ell}]$, and therefore $[\varphi]$, has finite order in $\text{Mod}(M_N)$. □

3. Theorem 1.4: lifting Λ_w to $\text{Diff}^+(M_N)$

We turn to the proof of Theorem 1.4, after fixing some notation regarding certain subsets of $\mathbb{C}\mathbb{P}^1$ illustrated in Figure 2. Let $N = n + 1$ and $m = \lceil \frac{n}{2} \rceil$; thus $N = 2m + 1$ if n is even and $N = 2m$ if n is odd. Fix distinct complex numbers $a_1, \dots, a_{2m} \in \mathbb{C}$ and let $z_k := [a_k : 1]$ for all $k = 1, \dots, n$. Then:

- (a) For each $1 \leq k \leq n - 1$, let $B_k \cong \mathbb{D}^2$ denote a closed disk in $\mathbb{C}\mathbb{P}^1 - \{[1 : 0]\}$ containing z_k and z_{k+1} and no other points z_j for $j \neq k, k + 1$ so that $B_k \cap B_{k'} = \emptyset$ if $|k - k'| > 1$.
- (b) For each $1 \leq k \leq n - 1$, let $U_k \cong [0, 1] \times \mathbb{S}^1 \subseteq B_k$ denote a collar neighborhood of B_k that does not contain z_k and z_{k+1} where $\{0\} \times \mathbb{S}^1$ corresponds to ∂B_k .
- (c) For each $1 \leq k \leq n$, let $V_k \cong \mathbb{D}^2$ denote a closed disk containing z_k in $B_1 - U_1 - B_2$ if $k = 1$, in $(B_k - U_k) \cap (B_{k-1} - U_{k-1})$ if $2 \leq k \leq n - 1$, or in $B_{n-1} - U_{n-1} - B_{n-2}$ if $k = n$.

As in Section 2.3, let

$$P = \{([a_i : 1], [1 : 0]) : 1 \leq i \leq m\} \cup \{([a_i : 1], [0 : 1]) : m + 1 \leq i \leq 2m\}$$

and consider the de Jonquières involution \mathcal{J} on $\mathrm{Bl}_P(\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1)$. Identify M_{n+1} with

- (a) $\mathrm{Bl}_P(\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1)$ if n is even, and
- (b) $\mathrm{Bl}_{P-\{([a_{2m}:1], [0:1])\}}(\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1)$ if n is odd.

In both cases, consider $\mathrm{pr} : M_{n+1} \rightarrow \mathbb{C}\mathbb{P}^1$ defined in [Section 2.3](#). There is a natural inclusion

$$\mathrm{pr}^{-1}(B_k) \hookrightarrow \mathrm{Bl}_P(\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1)$$

that is preserved by \mathcal{J} on $\mathrm{Bl}_P(\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1)$ for all $1 \leq k \leq n-1$. We use this inclusion to define $\mathcal{J}|_{\mathrm{pr}^{-1}(B_k)}$ on each $\mathrm{pr}^{-1}(B_k) \subseteq M_{n+1}$ regardless of the parity of n . Note that $\mathcal{J}|_{\mathrm{pr}^{-1}(B_k)} = \mathcal{J}|_{\mathrm{pr}^{-1}(B_{k+1})}$ when restricted to $\mathrm{pr}^{-1}(B_k) \cap \mathrm{pr}^{-1}(B_{k+1})$ for all $1 \leq k \leq n-2$.

There are four main steps to the proof of [Theorem 1.4](#).

- (1) Constructing commuting diffeomorphisms $\gamma_1, \dots, \gamma_{n-1} \in \mathrm{Diff}^+(M_N)$ that preserve the genus-0 holomorphic Lefschetz fibration $\mathrm{pr} : M_N \rightarrow \mathbb{C}\mathbb{P}^1$ such that $\mathrm{supp}(\gamma_k) \subseteq \mathrm{pr}^{-1}(B_k)$ and γ_k agrees with \mathcal{J} on $\mathrm{pr}^{-1}(B_k - U_k)$ for each $1 \leq k \leq n-1$. These maps should be thought of as *local de Jonquières maps*.
- (2) Constructing commuting diffeomorphisms $r_1, \dots, r_n \in \mathrm{Diff}^+(M_N)$ satisfying $\mathrm{supp}(r_k) \subseteq \mathrm{pr}^{-1}(V_k)$ and $[r_k] = \mathrm{Ref}_{e_k}$ for each $1 \leq k \leq n$.
- (3) Defining a homomorphism $\rho_v : \Lambda_v \rightarrow \mathrm{Diff}^+(M_N)$ using the diffeomorphisms above so that ρ_v is a section of $q : \mathrm{Diff}^+(M_N) \rightarrow \mathrm{Mod}(M_N)$ restricted to Λ_v and $\rho_v(\Lambda_v)$ almost preserves $\mathrm{pr} : M_N \rightarrow \mathbb{C}\mathbb{P}^1$.
- (4) Defining a homomorphism $\rho_w : \Lambda_w \rightarrow \mathrm{Diff}^+(M_N)$ for any other primitive, isotropic class w of minimal genus 0 by pre- and post-composing ρ_v by conjugation in $\mathrm{Mod}(M_N)$ and $\mathrm{Diff}^+(M_N)$.

Step 1: Constructing local de Jonquières maps $\gamma_1, \dots, \gamma_{n-1} \in \mathrm{Diff}^+(M_N)$. First, recall by construction that for each $1 \leq k \leq n-1$, the disk B_k is a closed subset of $\mathbb{C}\mathbb{P}^1 - \{[1:0]\}$. Throughout this section, we identify $\mathbb{C}\mathbb{P}^1 - \{[1:0]\}$ with \mathbb{C} by the diffeomorphism $[a:1] \mapsto a$. Then the disk B_k and the annulus U_k are subsets of \mathbb{C} and the point $z_k = [a_k:1] \in \mathbb{C}\mathbb{P}^1 - \{[0:1]\}$ corresponds to $a_k \in \mathbb{C}$ under this identification.

For each $1 \leq k \leq n-1$, define $\lambda_k : U_k \rightarrow \mathbb{C}^\times$ by

$$\lambda_k(x) := \sqrt{\frac{\prod_{i=1}^m (x - a_i)}{\prod_{i=m+1}^{2m} (x - a_i)}}$$

with any smooth choice of square root. A computation of fundamental groups shows that such a choice exists. For completeness, we include a proof below.

Lemma 3.1. *For each $1 \leq k \leq n-1$, there exists a smooth map $\lambda_k : U_k \rightarrow \mathbb{C}^\times$ so that*

$$\lambda_k(x)^2 = \frac{\prod_{i=1}^m (x - a_i)}{\prod_{i=m+1}^{2m} (x - a_i)}.$$

Proof. Consider a function $\mu_k : B_k - \{a_k, a_{k+1}\} \rightarrow \mathbb{C}^\times$ defined by

$$\mu_k(x) := \left(\prod_{i=1}^m (x - a_i) \right) \left(\prod_{i=m+1}^{2m} (x - a_i)^{-1} \right).$$

Then $\mu_k(x)$ is well-defined and nonzero for any $x \in B_k - \{a_k, a_{k+1}\}$ because $x - a_i \neq 0$ for any $1 \leq i \leq n$. It suffices to show that there exists a lift $\lambda_k : U_k \rightarrow \mathbb{C}^\times$ of the restriction $\mu_k|_{U_k} : U_k \rightarrow \mathbb{C}^\times$ under the double cover $\mathbb{C}^\times \rightarrow \mathbb{C}^\times$ given by $a \mapsto a^2$. In other words, we will show that $(\mu_k)_*(\pi_1(U_k))$ is contained in $2\mathbb{Z} \leq \mathbb{Z} \cong \pi_1(\mathbb{C}^\times)$.

Let $\delta_k, \delta_{k+1} \in \pi_1(B_k - \{a_k, a_{k+1}\})$ be generators so that $\delta_k \delta_{k+1}$ is a generator of $\pi_1(U_k) \leq \pi_1(B_k - \{a_k, a_{k+1}\})$ and so that δ_k (resp. δ_{k+1}) is freely homotopic in $B_k - \{a_k, a_{k+1}\}$ to a loop $S^1 \rightarrow B_k - \{a_k, a_{k+1}\}$ given by $\theta \mapsto a_k + \varepsilon e^{2\pi\theta\sqrt{-1}} \in V_k$ (resp. $\theta \mapsto a_{k+1} + \varepsilon e^{2\pi\theta\sqrt{-1}} \in V_{k+1}$) for some $0 < \varepsilon \ll 1$.

The restrictions of μ_k to $V_k - \{a_k\}$ and $V_{k+1} - \{a_{k+1}\}$ take the forms

$$\mu_k|_{V_k - \{a_k\}}(x) = \eta_k(x)(x - a_k)^{\pm 1}, \quad \mu_k|_{V_{k+1} - \{a_{k+1}\}}(x) = \eta_{k+1}(x)(x - a_{k+1})^{\pm 1},$$

where $\eta_i : V_i \rightarrow \mathbb{C}^\times$ are nonvanishing functions for each $1 \leq i \leq n$. Each of $\mu_k(\delta_k)$ and $\mu_k(\delta_{k+1})$ is freely homotopic to the loop $S^1 \rightarrow \mathbb{C}^\times$ given by $\theta \mapsto e^{2\pi\theta\sqrt{-1}}$ or by $\theta \mapsto e^{-2\pi\theta\sqrt{-1}}$, depending on the exponent of $(x - a_k)$ and $(x - a_{k+1})$ in $\mu_k(x)$. Therefore, $(\mu_k)_*(\delta_k \delta_{k+1})$ is an element of $2\mathbb{Z} \leq \pi_1(\mathbb{C}^\times)$. \square

For such a choice of λ_k , consider the map $M_{\lambda_k} : U_k \rightarrow \mathrm{PGL}_2(\mathbb{C})$ given by

$$M_{\lambda_k}(x) = \begin{pmatrix} 1 & 1 \\ -\lambda_k(x) & \lambda_k(x) \end{pmatrix} \in \mathrm{PGL}_2(\mathbb{C}).$$

We also record the inverse of $M_{\lambda_k}(x)$ for later use:

$$M_{\lambda_k}(x)^{-1} = \begin{pmatrix} 1 & -1/\lambda_k(x) \\ 1 & 1/\lambda_k(x) \end{pmatrix} \in \mathrm{PGL}_2(\mathbb{C}).$$

Viewing $M_{\lambda_k}(x)$ and $M_{\lambda_k}(x)^{-1}$ as automorphisms of $\mathbb{C}\mathbb{P}^1$, define a diffeomorphism u_{λ_k} of $\mathrm{pr}^{-1}(U_k) = U_k \times \mathbb{C}\mathbb{P}^1$ by

$$u_{\lambda_k}([x : 1], [Y_1 : Y_2]) = ([x : 1], M_{\lambda_k}(x) \cdot [Y_1 : Y_2]).$$

Let $T : [0, 1] \rightarrow [0, 1]$ be a smooth, nondecreasing function such that $T|_{[0, \varepsilon]} \equiv 0$ and $T|_{[1-\varepsilon, 1]} \equiv 1$ for some $0 < \varepsilon \ll 1$. Identifying $\mathrm{pr}^{-1}(U_k) = U_k \times \mathbb{C}\mathbb{P}^1$ with $[0, 1] \times \partial B_k \times \mathbb{C}\mathbb{P}^1$ (cf. Figure 2), we define a diffeomorphism j_k of $\mathrm{pr}^{-1}(U_k)$ by




Figure 3. Illustrating the action of j_k on $\text{pr}^{-1}(U_k) \cong [0, 1] \times \partial B_k \times \mathbb{C}\mathbb{P}^1$. The horizontal line represents $B_k \subseteq \mathbb{C}\mathbb{P}^1$ and the orange portion represents the annulus $U_k \subseteq B_k \subseteq \mathbb{C}\mathbb{P}^1$ whose width is parametrized by $t \in [0, 1]$. The two blue points represent $\partial B_k \cong \mathbb{S}^1$. The diffeomorphism j_k acts by rotation-by- $\pi T(t)$ on the sphere lying above a point $(t, x) \in [0, 1] \times \partial B_k \cong U_k$.

$$j_k(t, \theta, [Y_1 : Y_2]) = (t, \theta, [e^{\sqrt{-1}\pi T(t)} Y_1 : Y_2]).$$

Roughly, j_k is a map on $[0, 1] \times \mathbb{S}^1 \times \mathbb{C}\mathbb{P}^1$ induced by an isotopy of $\mathbb{S}^1 \times \mathbb{C}\mathbb{P}^1$ from the $\text{Id} \times \text{Id}$ to $\text{Id} \times R(\pi)$, where $R(\pi)$ is a rotation-by- π map on $\mathbb{C}\mathbb{P}^1$. See Figure 3.

In the next lemma, we show that the de Jonquières map is conjugate to $\text{Id} \times R(\pi)$ on each $\{t\} \times \partial B_k \times \mathbb{C}\mathbb{P}^1$, which will be used to modify $\mathcal{J}|_{\text{pr}^{-1}(U_k)}$ to be the identity near the boundary $\text{pr}^{-1}(\partial B_k)$.

Lemma 3.2. *Let $1 \leq k \leq n - 1$. On $\text{pr}^{-1}(U_k) = [0, 1] \times \partial B_k \times \mathbb{C}\mathbb{P}^1$,*

$$u_{\lambda_k} \circ j_k \circ u_{\lambda_k}^{-1} = \begin{cases} \mathcal{J} & \text{on } \text{pr}^{-1}([1 - \varepsilon, 1] \times \partial B_k), \\ \text{Id} & \text{on } \text{pr}^{-1}([0, \varepsilon] \times \partial B_k). \end{cases}$$

Proof. On $\text{pr}^{-1}([0, \varepsilon] \times \partial B_k)$, note that $j_k \equiv \text{Id}$. On $\text{pr}^{-1}([1 - \varepsilon, 1] \times \partial B_k)$,

$$j_k([x : 1], [Y_1 : Y_2]) = ([x : 1], [-Y_1 : Y_2]).$$

For all $[x : 1] \in [1 - \varepsilon, 1] \times \partial B_k \subseteq U_k$

$$M_{\lambda_k}(x) \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} M_{\lambda_k}(x)^{-1} = \begin{pmatrix} 0 & 1 \\ \lambda_k(x)^2 & 0 \end{pmatrix} \in \text{PGL}_2(\mathbb{C}),$$

and so

$$u_{\lambda_k} \circ j_k \circ u_{\lambda_k}^{-1}([x : 1], [Y_1 : Y_2]) = ([x : 1], [Y_2 : \lambda_k(x)^2 Y_1]) = \mathcal{J}([x : 1], [Y_1 : Y_2]). \quad \square$$

The diffeomorphisms γ_k below should be thought of as *local* de Jonquières maps, acting only on a single pair of singular fibers of pr .

Definition 3.3. For $1 \leq k \leq n - 1$, let γ_k be the diffeomorphism of M_N given by

$$\gamma_k = \begin{cases} \mathcal{J} & \text{on } \text{pr}^{-1}(B_k - U_k), \\ u_{\lambda_k} \circ j_k \circ u_{\lambda_k}^{-1} & \text{on } \text{pr}^{-1}(U_k), \\ \text{Id} & \text{on } \text{pr}^{-1}(\mathbb{C}\mathbb{P}^1 - B_k). \end{cases}$$

Proposition 3.4. *The diffeomorphisms γ_k satisfy the following properties:*

- (a) *The diffeomorphism γ_k preserves pr for all $1 \leq k \leq n-1$. In fact, $\text{pr} \circ \gamma_k = \text{pr}$.*
- (b) *The diffeomorphisms γ_i and γ_j commute for all $1 \leq i, j \leq n-1$.*
- (c) *As mapping classes, $[\gamma_k] = \text{Ref}_{v-e_k-e_{k+1}} \circ \text{Ref}_{e_k-e_{k+1}}$ for all $1 \leq k \leq n-1$.*

Proof. For each k , $\text{pr} \circ u_{\lambda_k} = \text{pr}$ and $\text{pr} \circ j_k = \text{pr}$ by construction of u_{λ_k} and j_k when restricted to $\text{pr}^{-1}(U_k)$. Therefore,

$$(\text{pr} \circ \gamma_k)|_{\text{pr}^{-1}(U_k)} = (\text{pr} \circ (u_{\lambda_k} \circ j_k \circ u_{\lambda_k}^{-1}))|_{\text{pr}^{-1}(U_k)} = \text{pr}|_{\text{pr}^{-1}(U_k)}$$

and γ_k preserves the fibers of pr on $\text{pr}^{-1}(U_k)$ for all k . The same is clearly true on $\text{pr}^{-1}(\mathbb{C}\mathbb{P}^1 - B_k)$ and true on $\text{pr}^{-1}(B_k - U_k)$ by construction of \mathcal{J} . This proves (a).

If $|i - j| > 1$ then $\text{supp}(\gamma_i) \cap \text{supp}(\gamma_j) = \emptyset$ so γ_i and γ_j commute. To show that γ_i and γ_{i+1} commute for $1 \leq i \leq n-2$, we will consider the action of these diffeomorphisms on $\text{pr}^{-1}(B_i \cap B_{i+1})$, which contains $\text{supp}(\gamma_i) \cap \text{supp}(\gamma_{i+1})$. We split $B_i \cap B_{i+1}$ as a union of $\mathcal{C}_i := (B_i \cap B_{i+1}) \cap (U_i \cup U_{i+1})$ and $\mathcal{V}_i := (B_i \cap B_{i+1}) - (U_i \cup U_{i+1})$ (see Figure 4), so

$$\text{pr}^{-1}(B_i \cap B_{i+1}) = \text{pr}^{-1}(\mathcal{C}_i) \cup \text{pr}^{-1}(\mathcal{V}_i).$$

By construction, $\gamma_i|_{\text{pr}^{-1}(\mathcal{V}_i)} = \mathcal{J}|_{\text{pr}^{-1}(\mathcal{V}_i)} = \gamma_{i+1}|_{\text{pr}^{-1}(\mathcal{V}_i)}$, and so γ_i and γ_{i+1} commute on $\text{pr}^{-1}(\mathcal{V}_i)$.

For any $[x : 1] \in \mathcal{C}_i$, both γ_i and γ_{i+1} act on $\text{pr}^{-1}([x : 1])$ by (a). If $[x : 1] \in U_i \cap U_{i+1}$ then for some $t, T \in [0, 1]$ depending on x , we have

$$\gamma_i([x : 1], [Y_1 : Y_2]) = \left([x : 1], \left(M_{\lambda_i}(x) \begin{pmatrix} e^{\sqrt{-1}\pi t} & 0 \\ 0 & 1 \end{pmatrix} M_{\lambda_i}(x)^{-1} \right) \cdot [Y_1 : Y_2] \right),$$

$$\gamma_{i+1}([x : 1], [Y_1 : Y_2]) = \left([x : 1], \left(M_{\lambda_{i+1}}(x) \begin{pmatrix} e^{\sqrt{-1}\pi T} & 0 \\ 0 & 1 \end{pmatrix} M_{\lambda_{i+1}}(x)^{-1} \right) \cdot [Y_1 : Y_2] \right).$$

Moreover, $\lambda_i(x) = \lambda_{i+1}(x)$ or $-\lambda_{i+1}(x)$. In the first case, $M_{\lambda_i}(x) = M_{\lambda_{i+1}}(x)$, so γ_i and γ_{i+1} commute on $\text{pr}^{-1}([x : 1])$. In the second case, we compute for each $[x : 1] \in U_i$ that

$$M_{\lambda_i}(x) = M_{-\lambda_i}(x) \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad M_{\lambda_i}(x)^{-1} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} M_{-\lambda_i}(x)^{-1}$$




Figure 4. For each $1 \leq i \leq n-2$, the sets \mathcal{C}_i (left) and \mathcal{V}_i (right) are contained in $B_i \cap B_{i+1}$.

and so

$$M_{-\lambda_i}(x) \begin{pmatrix} e^{\sqrt{-1}\pi T} & 0 \\ 0 & 1 \end{pmatrix} M_{-\lambda_i}(x)^{-1} = M_{\lambda_i}(x) \begin{pmatrix} 1 & 0 \\ 0 & e^{\sqrt{-1}\pi T} \end{pmatrix} M_{\lambda_i}(x)^{-1}.$$

It is clear that

$$M_{\lambda_i}(x) \begin{pmatrix} e^{\sqrt{-1}\pi t} & 0 \\ 0 & 1 \end{pmatrix} M_{\lambda_i}(x)^{-1} \quad \text{and} \quad M_{\lambda_i}(x) \begin{pmatrix} 1 & 0 \\ 0 & e^{\sqrt{-1}\pi T} \end{pmatrix} M_{\lambda_i}(x)^{-1}$$

commute in $\mathrm{PGL}_2(\mathbb{C})$, which shows that γ_i and γ_{i+1} commute on $\mathrm{pr}^{-1}([x : 1])$ in this case.

If $[x : 1] \in U_i$ and $[x : 1] \notin U_{i+1}$ then for some $t \in [0, 1]$ depending on x and for all $[Y_1 : Y_2] \in \mathbb{C}\mathbb{P}^1$,

$$\gamma_i([x : 1], [Y_1 : Y_2]) = \left([x : 1], \left(M_{\lambda_i}(x) \begin{pmatrix} e^{\sqrt{-1}\pi t} & 0 \\ 0 & 1 \end{pmatrix} M_{\lambda_i}(x)^{-1} \right) \cdot [Y_1 : Y_2] \right)$$

$$\gamma_{i+1}([x : 1], [Y_1 : Y_2]) = \mathcal{J}([x : 1], [Y_1 : Y_2])$$

$$= \left([x : 1], \left(M_{\lambda_{i+1}}(x) \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} M_{\lambda_{i+1}}(x)^{-1} \right) \cdot [Y_1 : Y_2] \right)$$

where the second equality follows from (the proof of) [Lemma 3.2](#). Therefore, we can show that γ_i and γ_{i+1} commute on $\mathrm{pr}^{-1}([x : 1])$ similarly as in the previous case. By analogous computations, γ_i and γ_{i+1} commute on $\mathrm{pr}^{-1}([x : 1])$ if $[x : 1] \in U_{i+1}$ and $[x : 1] \notin U_i$. This proves (b).

Finally, note that for all $j \neq k, k+1$, the map γ_k restricts to the identity on e_j and on $\mathrm{pr}^{-1}(z)$ for any $z \notin B_k$ so $(\gamma_k)_*(e_j) = e_j$ and $(\gamma_k)_*(v) = v$. Moreover, γ_k agrees with \mathcal{J} on $\mathrm{pr}^{-1}(B_k)$, meaning that $(\gamma_k)_*(e_j) = v - e_j$ for $j = k$ and $j = k+1$. This then determines $[\gamma_k] \in \mathrm{Mod}(M_{n+1})$ by [Lemma 2.3](#). A computation shows that the same holds for $\mathrm{Ref}_{v-e_k-e_{k+1}} \circ \mathrm{Ref}_{e_k-e_{k+1}}$. \square

Step 2: Constructing $r_1, \dots, r_n \in \mathrm{Diff}^+(M_N)$. For each $1 \leq k \leq n$, the exceptional divisor e_k has a tubular neighborhood ν_k in $\mathrm{pr}^{-1}(V_k)$ that is diffeomorphic to $\overline{\mathbb{C}\mathbb{P}^2} - \{[0 : 0 : 1]\}$. Let $i_k : \overline{\mathbb{C}\mathbb{P}^2} - \{[0 : 0 : 1]\} \rightarrow \nu_k$ be this diffeomorphism and let τ_0 be a diffeomorphism of $\overline{\mathbb{C}\mathbb{P}^2} - \{[0 : 0 : 1]\}$ given by complex conjugation, $\tau_0 : [X : Y : Z] \mapsto [\bar{X} : \bar{Y} : \bar{Z}]$.

Consider a smooth path $\eta : (0, 1) \rightarrow \mathrm{SO}(4)$ such that

$$\eta(t) = \begin{cases} \mathrm{diag}(1, -1, 1, -1) & \text{if } t \in (1 - \varepsilon, 1) \\ \mathrm{Id} & \text{if } t \in (0, \varepsilon) \end{cases}$$

for some $0 < \varepsilon \ll 1$. Let B denote the punctured ball in $\overline{\mathbb{C}\mathbb{P}^2} - \{[0 : 0 : 1]\}$ given by

$$B := \{[a + b\sqrt{-1} : c + d\sqrt{-1} : 1] \in \overline{\mathbb{C}\mathbb{P}^2} : 0 < \|(a + b\sqrt{-1}, c + d\sqrt{-1})\| < 1\},$$

identifying it with $(0, 1) \times \mathbb{S}^3 \subseteq \mathbb{R}^4$, and define $\tau \in \text{Diff}^+(\overline{\mathbb{C}\mathbb{P}^2} - \{[0 : 0 : 1]\})$ by

$$\tau = \begin{cases} \tau_0 & \text{on } \overline{\mathbb{C}\mathbb{P}^2} - B, \\ (t, x) \mapsto (t, \eta(t)x) & \text{on } B \cong (0, 1) \times \mathbb{S}^3. \end{cases}$$

Then τ is compactly supported in $\overline{\mathbb{C}\mathbb{P}^2} - \{[0 : 0 : 1]\}$.

Definition 3.5. For all $1 \leq k \leq n$, let $r_k \in \text{Diff}^+(M_N)$ be

$$r_k := \begin{cases} i_k \circ \tau \circ i_k^{-1} & \text{on } \nu_k, \\ \text{Id} & \text{on } M_N - \nu_k. \end{cases}$$

Remark 3.6. By construction, the diffeomorphism r_k restricts to an orientation-reversing diffeomorphism of e_k and preserves the homology classes e_i for all $i \neq k$ and v . This forces $[r_k] = \text{Ref}_{e_k} \in \text{Mod}(M_N)$ by [Lemma 2.3](#). Moreover, $\text{supp}(r_k) \subseteq \nu_k \subseteq \text{pr}^{-1}(V_k)$ and r_k preserves $\nu_k \subseteq \text{pr}^{-1}(V_k)$.

Step 3: Constructing $\rho_v : \Lambda_v \rightarrow \text{Diff}^+(M_N)$. The generators f_1, \dots, f_{n-1} of Λ_v (cf. [Lemma 2.5](#)) will be mapped under ρ_v to the following diffeomorphisms.

Lemma 3.7. For each $1 \leq k \leq n - 1$, let

$$\varphi_k := r_k \circ r_{k+1} \circ \gamma_k.$$

Then $\varphi_i \circ \varphi_j = \varphi_j \circ \varphi_i$ for any $1 \leq i, j \leq n - 1$.

Proof. For any $1 \leq i, j \leq n - 1$, the diffeomorphisms φ_i and φ_j commute if $|i - j| > 1$ because they have disjoint support. For any $1 \leq i \leq n - 2$,

$$\varphi_i|_{\text{pr}^{-1}(V_{i+1})} = (r_{i+1} \circ \mathcal{J})|_{\text{pr}^{-1}(V_{i+1})} = \varphi_{i+1}|_{\text{pr}^{-1}(V_{i+1})}$$

so φ_i and φ_{i+1} commute on $\text{pr}^{-1}(V_{i+1})$. Moreover on $S_i := \text{pr}^{-1}(B_i \cap B_{i+1}) - \text{pr}^{-1}(V_{i+1})$

$$\varphi_i|_{S_i} = \gamma_i, \quad \varphi_{i+1}|_{S_i} = \gamma_{i+1}$$

and so φ_i and φ_{i+1} commute on $\text{pr}^{-1}(B_i \cap B_{i+1})$ by [Proposition 3.4\(b\)](#). Finally, φ_i and φ_{i+1} commute on $M_{n+1} - \text{pr}^{-1}(B_i \cap B_{i+1})$ because $\text{supp}(\varphi_i) \cap \text{supp}(\varphi_{i+1})$ is contained in $\text{pr}^{-1}(B_i \cap B_{i+1})$. \square

It remains to construct the image of the last generator g of Λ_v under ρ_v .

Lemma 3.8. The map

$$\psi = \begin{cases} \varphi_1 \circ \varphi_1 & \text{on } \text{pr}^{-1}(V_1) \\ \text{Id} & \text{on } M_{n+1} - \text{pr}^{-1}(V_1). \end{cases}$$

is a well-defined diffeomorphism, which commutes with φ_k for all $1 \leq k \leq n - 1$ and satisfies $[\psi] = g = \text{Ref}_{e_1} \circ \text{Ref}_{v-e_1}$ in $\text{Mod}(M_{n+1})$,

Proof. By definition, r_k has support contained in the interior of $\text{pr}^{-1}(V_k)$ for all $1 \leq k \leq n$. So $r_1|_C = \text{Id}|_C$ on some collar neighborhood C of $\text{pr}^{-1}(V_1)$, and

$$\psi|_C = (\varphi_1 \circ \varphi_1)|_C = (r_1 \circ \mathcal{J} \circ r_1 \circ \mathcal{J})|_C = (\mathcal{J} \circ \mathcal{J})|_C = \text{Id}|_C.$$

Moreover, \mathcal{J} , r_1 , and r_2 all preserve $\text{pr}^{-1}(V_1)$, so the map ψ is indeed a diffeomorphism.

The diffeomorphisms ψ and φ_k have disjoint supports for all $k > 1$. Considering the subsets $\text{pr}^{-1}(V_1)$ and $M_{n+1} - \text{pr}^{-1}(V_1)$ separately shows that φ_1 and ψ commute as well.

Compute for all $2 \leq k \leq n$ that $[\psi](e_k) = e_k$ because $\text{supp}(\psi) \subseteq \text{pr}^{-1}(V_1)$. Moreover, ψ agrees with φ_1^2 on $\text{pr}^{-1}(V_1)$, meaning that

$$[\psi](e_1) = [\varphi_1^2](e_1) = e_1 + 2v.$$

Computing that

$$\text{Ref}_{e_1} \circ \text{Ref}_{v-e_1}(e_k) = [\psi](e_k)$$

for all $1 \leq k \leq n$ and applying [Lemma 2.3](#) shows that $[\psi] = g$. \square

Proposition 3.9. *There is a homomorphism $\rho_v : \Lambda_v \rightarrow \text{Diff}^+(M_N)$ defined by*

$$\rho_v(f_k) := \varphi_k \quad \text{for all } 1 \leq k \leq n-1, \quad \rho_v(g) := \psi,$$

where g and f_k for $1 \leq k \leq n-1$ are the generators of Λ_v as given in [Lemma 2.5](#). Moreover,

- (a) ρ_v is a section of the map $q : \text{Diff}^+(M_N) \rightarrow \text{Mod}(M_N)$ restricted to $\Lambda_v \leq \text{Mod}(M_N)$, and
- (b) for all $\varphi \in \rho_v(\Lambda_v)$,

$$(\text{pr} \circ \varphi)|_{M_N - \bigcup_{i=1}^n \text{pr}^{-1}(V_i)} = \text{pr}|_{M_N - \bigcup_{i=1}^n \text{pr}^{-1}(V_i)}.$$

Hence $\rho_v(\Lambda_v)$ almost preserves the Lefschetz fibration $\text{pr} : M_N \rightarrow \mathbb{C}\mathbb{P}^1$.

Proof. By [Lemma 2.5](#), $\Lambda_v \cong \mathbb{Z}^n$ is generated by f_1, \dots, f_{n-1}, g . By [Lemmas 3.7](#) and [3.8](#), the image of ρ_v is abelian and therefore ρ_v is a well-defined homomorphism.

Compute using [Proposition 3.4\(c\)](#) and [Remark 3.6](#) that

$$[\rho_v(f_k)] = [r_k] \circ [r_{k+1}] \circ [\gamma_k] = \text{Ref}_{e_k} \circ \text{Ref}_{e_{k+1}} \circ \text{Ref}_{v-e_k-e_{k+1}} \circ \text{Ref}_{e_k-e_{k+1}} = f_k \in \Lambda_v.$$

[Lemma 3.8](#) shows that $[\rho_v(g)] = g$. Therefore, ρ_v is a section of the quotient map $q : \text{Diff}^+(M_N) \rightarrow \text{Mod}(M_N)$ restricted to $\Lambda_v \leq \text{Mod}(M_N)$.

Finally, $\text{supp}(r_k) \subseteq \text{pr}^{-1}(V_k)$ for all $1 \leq k \leq n$ (cf. [Remark 3.6](#)). By [Proposition 3.4\(a\)](#), $\text{pr} \circ \gamma_k = \text{pr}$ for all $1 \leq k \leq n-1$, so

$$\varphi_k|_{M_N - \bigcup_{i=1}^n \text{pr}^{-1}(V_i)} = \gamma_k|_{M_N - \bigcup_{i=1}^n \text{pr}^{-1}(V_i)}.$$

By [Lemma 3.8](#),

$$\psi|_{M_N - \bigcup_{i=1}^n \text{pr}^{-1}(V_i)} = \text{Id}|_{M_N - \bigcup_{i=1}^n \text{pr}^{-1}(V_i)}.$$

Hence $\text{pr} \circ \varphi = \text{pr}$ restricted to $M_N - \bigcup_{i=1}^n \text{pr}^{-1}(V_i)$ for all $\varphi \in \rho_v(\Lambda_v)$ and so $\rho_v(\Lambda_v)$ almost preserves pr . \square

Step 4: Extension to any primitive, isotropic class w of minimal genus 0. With the constructions above in hand, we conclude the proof of [Theorem 1.4](#).

Proof of [Theorem 1.4](#). Because w is a primitive, isotropic class of minimal genus 0, there exists some $\alpha \in \text{Diff}^+(M_N)$ such that $[\alpha](w) = v$ by a theorem of Li–Li ([\[15, Theorem 4.2\]](#)). Using the definitions of h_v and h_w (cf. [Lemma 2.6](#)), compute that

$$\Lambda_v = [\alpha] \circ \Lambda_w \circ [\alpha^{-1}]$$

and define $\rho_w : \Lambda_w \rightarrow \text{Diff}^+(M_N)$ by

$$\rho_w(f) = \alpha^{-1} \circ \rho_v([\alpha] \circ f \circ [\alpha^{-1}]) \circ \alpha$$

where $\rho_v : \Lambda_v \rightarrow \text{Diff}^+(M_N)$ is the homomorphism constructed in [Proposition 3.9](#). Compute that for all $x \in M_N - \bigcup_{k=1}^n (\text{pr} \circ \alpha)^{-1}(V_k)$,

$$(\text{pr} \circ \alpha) \circ \rho_w(f)(x) = \text{pr} \circ \rho_v([\alpha] \circ f \circ [\alpha^{-1}]) \circ \alpha(x) = (\text{pr} \circ \alpha)(x)$$

because $\text{pr} \circ \rho_v([\alpha] \circ f \circ [\alpha^{-1}]) = \text{pr}$ on $M_N - \bigcup_{k=1}^n \text{pr}^{-1}(V_k)$ by [Proposition 3.9\(b\)](#). Hence $\rho_w(\Lambda_w)$ almost preserves $\text{pr} \circ \alpha$, which is holomorphic for some complex structure on M_N . Finally, compute by [Proposition 3.9\(a\)](#) that for any $f \in \Lambda_w$,

$$\begin{aligned} q \circ \rho_w(f) &= q(\alpha^{-1} \circ \rho_v([\alpha] \circ f \circ [\alpha^{-1}]) \circ \alpha) \\ &= [\alpha^{-1}] \circ (q \circ \rho_v)([\alpha] \circ f \circ [\alpha^{-1}]) \circ [\alpha] = f. \end{aligned} \quad \square$$

If $2 \leq N \leq 8$, [Theorem 1.4](#) holds for any primitive, isotropic class in $H_2(M_N; \mathbb{Z})$.

Corollary 3.10. *Let $2 \leq N \leq 8$ and let $w \in H_2(M_N; \mathbb{Z})$ be any primitive, isotropic class. There exists a homomorphism $\rho_w : \Lambda_w \rightarrow \text{Diff}^+(M_N)$ such that the following diagram commutes:*

$$\begin{array}{ccc} & & \text{Diff}^+(M_N) \\ & \nearrow \rho_w & \downarrow q \\ \Lambda_w & \hookrightarrow & \text{Mod}(M_N) \end{array}$$

The image $\rho_w(\Lambda_w)$ almost preserves a holomorphic genus-0 Lefschetz fibration $p : M_N \rightarrow \mathbb{C}\mathbb{P}^1$ whose generic fiber represents the homology class w .

Proof. If $2 \leq N \leq 8$ and $w \in H_2(M_N; \mathbb{Z})$ is a primitive, isotropic class then [Lemma 2.7\(b\)](#) says that the minimal genus of w is 0. Now apply [Theorem 1.4](#). \square

4. Theorem 1.5: individual elements of the stabilizer $\text{Stab}(w)$ of w

We next prove [Theorem 1.5](#) using the diffeomorphisms constructed in [Section 3](#). The following lemma considers the subgroup $S_n \leq \text{Stab}(v) \cap \text{Stab}(s) \leq \text{Mod}(M_N)$ given by permuting the classes e_1, \dots, e_n . In other words, we consider the subgroup $\langle \text{Ref}_{e_k - e_{k+1}} : 1 \leq k \leq n-1 \rangle$ of $\text{Mod}(M_N)$.

Lemma 4.1. *For each $1 \leq k \leq n-1$, there exist $s_k \in \text{Diff}^+(M_N)$ and $\tau_k \in \text{Diff}^+(\mathbb{C}\mathbb{P}^1)$ such that*

- (a) $\text{pr} \circ s_k = \tau_k \circ \text{pr}$, and
- (b) $[s_k] = \text{Ref}_{e_k - e_{k+1}} \in \text{Mod}(M_N)$.

Proof. Let $A := \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \in \text{PGL}_2(\mathbb{C})$, so A has order 2 and $A([1 : 1]) = [-1 : 1] \in \mathbb{C}\mathbb{P}^1$. There exists a neighborhood D diffeomorphic to a disk \mathbb{D}^2 of the path $\{[t : 1] : t \in [-1, 1]\} \subseteq \mathbb{C}\mathbb{P}^1$ that is preserved by A . Let $\iota_k : D \hookrightarrow \mathbb{C}\mathbb{P}^1$ be a smooth embedding with image contained in $B_k - U_k$ and

$$\iota_k([-1 : 1]) = [a_k : 1], \quad \iota_k([1 : 1]) = [a_{k+1} : 1],$$

so that ι_k is holomorphic if restricted to small neighborhoods of $[1 : 1]$ and $[-1 : 1]$ in D . Now let $\tau_k \in \text{Diff}^+(\mathbb{C}\mathbb{P}^1)$ be a diffeomorphism such that

$$\tau_k = \begin{cases} \iota_k \circ A \circ \iota_k^{-1} & \text{on } \iota_k(D) \subseteq B_k - U_k, \\ \text{Id} & \text{on } \mathbb{C}\mathbb{P}^1 - B_k. \end{cases}$$

Consider the diffeomorphism $s : (X, Y) \mapsto (\tau_k(X), Y)$ of $\mathbb{C}\mathbb{P}^1 \times \mathbb{C}\mathbb{P}^1$ which extends to a diffeomorphism s_k of M_N because s is holomorphic on a neighborhood of $\text{pr}^{-1}([a_i : 1])$ for all $1 \leq i \leq n$. By construction, $\text{pr} \circ s_k = \tau_k \circ \text{pr}$. Moreover, if $i \neq k$ or $k+1$ then s_k acts as the identity on e_i but $s_k(e_k) = e_{k+1}$ and $s_k(e_{k+1}) = e_k$. Hence $[s_k] = \text{Ref}_{e_k - e_{k+1}}$ by [Lemma 2.3](#) because $[s_k] \in \text{Stab}(v)$. \square

We may assume that for all $1 \leq k \leq n-1$, the choice of $V_k, V_{k+1} \subseteq \mathbb{C}\mathbb{P}^1$ satisfies $V_k, V_{k+1} \subseteq \iota_k(D)$ and $\tau_k(V_k) = V_{k+1}$, where $\iota_k : D \hookrightarrow \mathbb{C}\mathbb{P}^1$ is the embedding defined in the proof of [Lemma 4.1](#). This also implies that $\tau_k(V_{k+1}) = V_k$ because $\tau_k|_{\iota_k(D)}$ has order 2.

Proof of Theorem 1.5. The theorem holds for $N = 1$ because then $\text{Stab}(w) = 1$. Now assume that $N \geq 2$ and that $w = v$. Since $h \in \text{Stab}(v)$, we may write $h = f \circ \sigma$ where $f \in \Lambda_v$ and $\sigma \in \text{Aut}(\mathbb{Z}\{e_1, \dots, e_n\}, \mathcal{Q}_{M_N}) \cong \text{O}(n)(\mathbb{Z})$ by [Lemma 2.6](#). Furthermore, the action of $\text{O}(n)(\mathbb{Z})$ on $\mathbb{Z}\{e_1, \dots, e_n\}$ preserves the set $\{e_1, \dots, e_n, -e_1, \dots, -e_n\}$ of classes of norm -1 . The action of $\text{O}(n)(\mathbb{Z})$ on the set of n unordered pairs $\{e_k, -e_k\}$ with $1 \leq k \leq n$ defines a homomorphism $\text{O}(n)(\mathbb{Z}) \rightarrow S_n$ with kernel $\langle \text{Ref}_{e_k} : 1 \leq k \leq n \rangle \cong (\mathbb{Z}/2\mathbb{Z})^n$. Moreover, this homomorphism admits a section with image $\langle \text{Ref}_{e_k - e_{k+1}} : 1 \leq k \leq n-1 \rangle \leq \text{O}(n)(\mathbb{Z})$. In other words, any

element $\sigma \in \text{Aut}(\mathbb{Z}\{e_1, \dots, e_n\}, \mathcal{Q}_{M_N})$ can be written as a product $[r] \circ [s] \in \text{Aut}(\mathbb{Z}\{e_1, \dots, e_n\}, \mathcal{Q}_{M_N})$ where

$$r \in \langle r_k : 1 \leq k \leq n \rangle \leq \text{Diff}^+(M_N) \quad \text{and} \quad s \in \langle s_k : 1 \leq k \leq n - 1 \rangle \leq \text{Diff}^+(M_N),$$

by [Remark 3.6](#) and [Lemma 4.1\(b\)](#). Let

$$\varphi := \rho_v(f) \circ r \circ s \in \text{Diff}^+(M_N)$$

where $\rho_v : \Lambda_v \rightarrow \text{Diff}^+(M_N)$ is the homomorphism from [Proposition 3.9](#). By construction, $[\varphi] = h$.

Note that

$$\text{pr} \circ \rho_v(f) \circ r \Big|_{M_N - \bigcup_{i=1}^n \text{pr}^{-1}(V_i)} = \text{pr} \Big|_{M_N - \bigcup_{i=1}^n \text{pr}^{-1}(V_i)}$$

by [Proposition 3.9\(b\)](#) and by [Remark 3.6](#). By [Lemma 4.1\(a\)](#), there exists $\tau \in \text{Diff}^+(\mathbb{C}\mathbb{P}^1)$ such that $\text{pr} \circ s = \tau \circ \text{pr}$ and τ preserves $\bigcup_{i=1}^n \text{pr}^{-1}(V_i)$. Hence

$$\text{pr} \circ \varphi \Big|_{M_N - \bigcup_{i=1}^n \text{pr}^{-1}(V_i)} = \text{pr} \circ (\rho_v(f) \circ r \circ s) \Big|_{M_N - \bigcup_{i=1}^n \text{pr}^{-1}(V_i)} = \tau \circ \text{pr} \Big|_{M_N - \bigcup_{i=1}^n \text{pr}^{-1}(V_i)},$$

which shows that φ almost preserves pr .

Take any other primitive, isotropic class $w \in H_2(M_N; \mathbb{Z})$ with minimal genus 0; we proceed similarly as in the proof of [Theorem 1.4](#). Apply Li–Li [[15](#), [Theorem 4.2](#)] to obtain $\alpha \in \text{Diff}^+(M_N)$ such that $[\alpha](w) = v$ and $[\alpha] \circ h \circ [\alpha^{-1}] \in \text{Stab}(v)$. There exists a diffeomorphism $\varphi \in \text{Diff}^+(M_N)$ almost preserving $\text{pr} : M_N \rightarrow \mathbb{C}\mathbb{P}^1$ with $[\varphi] = [\alpha] \circ h \circ [\alpha^{-1}]$. Then $\alpha^{-1} \circ \varphi \circ \alpha$ almost preserves $\text{pr} \circ \alpha$, and $[\alpha^{-1} \circ \varphi \circ \alpha] = h$. □

Corollary 4.2. *Let $2 \leq N \leq 8$ and let $w \in H_2(M_N)$ be any primitive, isotropic class. For any $h \in \text{Stab}(w)$, there exists $\varphi \in \text{Diff}^+(M_N)$ almost preserving some holomorphic genus-0 Lefschetz fibration $p : M_N \rightarrow \mathbb{C}\mathbb{P}^1$ whose generic fiber represents the homology class w such that $[\varphi] = h \in \text{Mod}(M_N)$.*

Proof. If $2 \leq N \leq 8$ and $w \in H_2(M_N; \mathbb{Z})$ is a primitive, isotropic class then [Lemma 2.7\(b\)](#) says that the minimal genus of w is 0. Now apply [Theorem 1.5](#). □

Acknowledgments

I thank Benson Farb for suggesting this problem, for his continuous encouragement, advice, and guidance throughout this project, and for many helpful comments on an earlier draft of this paper. I would also like to thank Farb and Eduard Looijenga for sharing their preliminary results on K3 manifolds (which appeared in [[9](#)]) with me. I would also like to thank Carlos A. Serván for many useful conversations about rational 4-manifolds and Lefschetz fibrations. I thank Dan Margalit for comments on an earlier draft that improved the exposition of this paper. Finally,

I thank anonymous referees for their careful reading, corrections, and extensive suggestions.

References

- [1] L. Bayle and A. Beauville, “Birational involutions of \mathbb{P}^2 ”, *Asian J. Math* **4** (2000), 11–18.
- [2] E. Bertini, “Ricerche sulle trasformazioni univoche involutorie nel piano”, *Ann. Mat. Pura Appl.* **8** (1877), 244–286.
- [3] J. Blanc, “Elements and cyclic subgroups of finite order of the Cremona group”, *Comment. Math. Helv.* **86**:2 (2011), 469–497. [MR](#)
- [4] S. Cantat, “Dynamique des automorphismes des surfaces $K3$ ”, *Acta Math.* **187** (2001), 1–57.
- [5] W. Chen, T.-J. Li, and W. Wu, “Symplectic rational G -surfaces and equivariant symplectic cones”, *J. Differential Geom.* **119**:2 (2021), 221, 260.
- [6] T. D. Cochran and N. Habegger, “On the homotopy theory of simply connected four manifolds”, *Topology* **29**:4 (1990), 419–440.
- [7] J. Diller and C. Favre, “Dynamics of bimeromorphic maps of surfaces”, *Am. J. Math.* **123** (2001), 1135–1169.
- [8] I. V. Dolgachev and V. A. Iskovskikh, “Finite subgroups of the plane Cremona group”, pp. 443–548 in *Algebra, arithmetic, and geometry: in honor of Yu. I. Manin*, vol. 1, edited by Y. Tschinkel and Y. Zarhin, Birkhäuser, 2009.
- [9] B. Farb and E. Looijenga, “Moduli spaces and period mappings of genus one fibered $K3$ surfaces”, preprint, 2023. [arXiv 2303.08214](#)
- [10] M. H. Freedman, “The topology of four-dimensional manifolds”, *J. Differ. Geom.* **17** (1982), 357–453.
- [11] D. Gabai, D. T. Gay, D. Hartman, V. Krushkal, and M. Powell, “Pseudo-isotopies of simply connected 4-manifolds”, preprint, 2023. [arXiv 2311.11196](#)
- [12] M. H. Gizatullin, “Rational G -surfaces”, *Math. USSR Izv.* **16** (1981), 103–134.
- [13] R. E. Gompf, “Locally holomorphic maps yield symplectic structures”, *Comm. Anal. Geom.* **13** (2005), 511–525.
- [14] R. E. Gompf and A. I. Stipsicz, *4-manifolds and Kirby calculus*, American Mathematical Society, 1999.
- [15] B.-H. Li and T.-J. Li, “Symplectic genus, minimal genus and diffeomorphisms”, *Asian J. Math.* **6** (2002), 123–144.
- [16] L. J. Mordell, “The definite quadratic forms in eight variables with determinant unity”, *J. Math. Pures Appl.* **17** (1938), 41–46.
- [17] P. Ozsváth and Z. Szabó, “The symplectic Thom conjecture”, *Ann. Math.* **151** (2000), 93–124.
- [18] B. Perron, “Pseudo-isotopies et isotopies en dimension quatre dans la catégorie topologique”, *Topology* **25**:4 (1986), 381–397. [MR](#)
- [19] F. Quinn, “Isotopy of 4-manifolds”, *J. Differ. Geom.* **24** (1986), 343–372.
- [20] W. P. Thurston, *Three-dimensional geometry and topology*, vol. 1, Princeton University Press, 1997.
- [21] C. Wall, “Diffeomorphisms of 4-manifolds”, *J. London Math. Soc.* **s1-39**:1 (1964), 131–140.

Received July 6, 2023. Revised September 10, 2025.

SERAPHINA EUN BI LEE
DEPARTMENT OF MATHEMATICS
HARVARD UNIVERSITY
UNITED STATES
slee@math.harvard.edu

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Fakultät für Mathematik
Universität Wien
Vienna, Austria
matthias.aschenbrenner@univie.ac.at

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Atsushi Ichino
Department of Mathematics
Kyoto University
Riverside, CA 92521-0135
atsushi.ichino@gmail.com

Robert Lipshitz
Department of Mathematics
University of Oregon
Eugene, OR 97403
lipshitz@uoregon.edu

Kefeng Liu
School of Sciences
Chongqing University of Technology
Chongqing 400054, China
liu@math.ucla.edu

Sucharit Sarkar
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
sucharit@math.ucla.edu

Dimitri Shlyakhtenko
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
shlyakht@ipam.ucla.edu

Ruixiang Zhang
Department of Mathematics
University of California
Berkeley, CA 94720-3840
ruixiang@berkeley.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2025 is US \$677/year for the electronic version, and \$917/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2025 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 339 No. 2 December 2025

| | |
|---|-----|
| Starting the study of outer length billiards | 201 |
| LUCA BARACCO, OLGA BERNARDI and CORENTIN FIEROBE | |
| Mori dream spaces and \mathbb{Q} -homology quadrics | 223 |
| PAOLO CASCINI, FABRIZIO CATANESE, YIFAN CHEN and JONGHAE KEUM | |
| Five-dimensional minimal quadratic and bilinear forms over function fields of conics | 243 |
| ADAM CHAPMAN and AHMED LAGHRIBI | |
| The Manakov equation of mixed type and its matrix generalization | 265 |
| QING DING, CHAOHAO YE and SHIPING ZHONG | |
| Mapping classes fixing an isotropic homology class of minimal genus 0 in rational 4-manifolds | 283 |
| SERAPHINA EUN BI LEE | |
| Lower bounds for fractional Orlicz-type eigenvalues | 309 |
| ARIEL SALORT | |
| Graph thinness: a lower bound and complexity | 333 |
| YAROSLAV SHITOV | |
| Classifying preaisles of derived categories of complete intersections | 345 |
| RYO TAKAHASHI | |