

*Pacific
Journal of
Mathematics*

CLOSURES OF 3-BRAIDS AND DETECTION

FRASER BINNS

Volume 340 No. 1

January 2026

CLOSURES OF 3-BRAIDS AND DETECTION

FRASER BINNS

We give some new link detection results for link Floer homology, Khovanov homology and annular Khovanov homology. The links we detect arise via different closure operations on 3-braids. Examples of our results include that link Floer homology detects the Mazur link, that annular Khovanov homology detects the Mazur pattern, and that Khovanov homology detects L6a2 and L9n15. The Mazur pattern detection result depends on a new bound on the rank of the annular Khovanov homology of certain links.

Braids are of wide mathematical interest; see the survey article [14]. In this paper we will consider the four types of links obtained from braids, as shown in Figure 1.

Let α be a braid. The first two types of link we obtain from α have been widely studied. We have the *braid-closure* of α , $b(\alpha)$, which for the purposes of this paper is the link in the thickened annulus obtained by attaching n parallel strands as in Figure 1(a). Secondly, we have the *augmented braid-closure* of α , $\hat{b}(\alpha)$, which is the link obtained by adding the annular axis to $b(\alpha)$, as shown in Figure 1(b).

For the remaining two types of link we move beyond the usual setting of braid-closures. The *clasp-closure* of α , $c(\alpha)$, can be thought of as the annular link formed by $b(\alpha)$ and replacing two parallel strands in a ball with a *clasp*, as shown in Figure 1(c). Note that the clasp is between the two rightmost strands of α , though this is simply a matter of convention and plays no significant role. The *augmented clasp-closure* of α , $\hat{c}(\alpha)$, is defined analogously to the augmented braid-closure of α ; see Figure 1(d).




Figure 1. The four types of links we study: (a) a braid-closure; (b) an augmented braid-closure; (c) a clasp-closure; (d) an augmented clasp-closure. The red dots indicate the axes.

The author was supported by the Simons Grant *New structures in low-dimensional topology*.
 MSC2020: 57K18.

Keywords: link Floer homology, Khovanov homology, annular Khovanov homology.

As motivation for studying clasp-closures, we mention a result of Martin [32, Proposition 1] stating that the links with the simplest link Floer homology — in an appropriate sense — are augmented braid-closures. A result of the author and Dey [11, Theorem 5.1] says that augmented clasp-closures are examples of links with second simplest link Floer homology, in the same sense. Thus one might reasonably expect that understanding the behavior of categorified link invariants of braid- and clasp-closures of braids might be easier than understanding other types of closure.

(Augmented) braid-closures of 1- and 2-braids are readily classified up to isotopy. Braid-closures of 3-braids were classified completely by Murasugi [33]. In particular, he showed that there are three braid-closures of 3-braids representing the unknot, namely $\sigma_1\sigma_2$, $\sigma_1^{-1}\sigma_2^{-1}$ and $\sigma_1\sigma_2^{-1}$. Here we use the standard Artin generators for the braid group. The augmented braid-closures of these three links are $T(2, 6)$, $T(2, -6)$ and L6a2 respectively. More generally, Birman and Menasco showed that for $|n| \neq 1$ there are two 3-braid representatives of the torus links $T(2, n)$, namely $\sigma_1^{\pm 1}\sigma_2^n$ [15]. Note that $\hat{b}(\sigma_1\sigma_2^3)$ is L9n15 while $\hat{b}(\sigma_1^{-1}\sigma_2^3)$ is L9n16.

One cannot take the clasp-closure of a 1-braid. Clasp-closures of 2-braids are the twisted Whitehead patterns. The case of (augmented) clasp-closures of 3-braids is more complicated. Baldwin and Sivek classified 3-braids with clasp-closures representing the unknot, up to isotopy of the clasp-closure; see the proof of [4, Theorem 6.1]. Up to mirroring and reversal these braids are as follows:

- (1) σ_1^{-1} . The augmentation of this link is L7a6, i.e., the mirror of the Mazur link.
- (2) $\sigma_1^3\sigma_2^{-1}\sigma_1^2\sigma_2$.
- (3) $\sigma_1^n\sigma_2^{-1}\sigma_1\sigma_2$. For $n = 1$ the augmentation of this link is L7a5.

Our goal in this paper is to exploit the Baldwin–Sivek, Murasugi and Birman–Menasco classification results to obtain detection results for various categorified link invariants. There are three different invariants we will study; link Floer homology and two versions of Khovanov homology.

Link Floer homology is an invariant of oriented links defined by Ozsváth and Szabó using symplectic topology [36]. For two-component links it takes value in the category of triply graded vector spaces. Our first results are the following:

Theorem 1.1. *Link Floer homology detects L6a2.*

Theorem 1.3. *Link Floer homology with rational coefficients detects L9n15.*

L6a2 is the augmented braid-closure of $\sigma_1\sigma_2^{-1}$. The author and Martin showed in [13] that link Floer homology detects the augmented braid-closures of the other two braids that represent the unknot; i.e., it was shown that link Floer homology detects $T(2, \pm 6)$ endowed with any orientation. The author and Dey showed in [10] that link Floer homology detects all of the augmented braid-closures of 2-braids.

The augmented braid-closure of the 1-braid is also detected by link Floer homology since it is simply a Hopf link.

The proof strategies for Theorem 1.1 and Theorem 1.3 are that used by the author and Martin in [13]. That is, we use the fact that the link Floer homology of a link L contains various pieces of topological information about L . In particular we appeal to Martin's result that link Floer homology detects braid axes [32, Proposition 1].

We now address 3-braids with unknotted clasp-closures. For the first type we have detection.

Theorem 1.5. *Link Floer homology detects the Mazur link.*

The author and Dey showed that link Floer homology detects the augmented clasp-closures of all but two 2-braids and that the remaining two augmented clasp-closures are the unique links of their link Floer homology type [11, Theorems 6.1 and 6.2].

For the final type of 3-braid with unknotted clasp-closure we do not get detection. Nevertheless we can give the following classification result:

Theorem 1.6. *Let L be a link. $\widehat{\text{HFL}}(L) \cong \widehat{\text{HFL}}(\mathring{c}(\sigma_2^{-1}\sigma_1\sigma_2))$ if and only if L is of the form $\mathring{c}(\sigma_1^n\sigma_2^{-1}\sigma_1\sigma_2)$ for some $n \in \mathbb{Z}$.*

The proof strategies for these two theorems are similar to that used in the proof of Theorem 1.1. The chief difference is that we appeal to the classification of links with link Floer homology of next to minimal rank in certain gradings [11, Theorem 5.1], as opposed to Martin's braid axis detection result, which was a classification of links with link Floer homology of minimal rank in certain gradings [32, Proposition 1].

We now turn to *Khovanov homology*. This is a combinatorial link invariant due to Khovanov that takes values in the category of bigraded vector spaces [26]. We have the following two results:

Theorem 2.1. *Khovanov homology with integer coefficients detects L6a2.*

Theorem 2.2. *Khovanov homology with integer coefficients detects L9n15.*

For context recall that Khovanov homology detects the Hopf link [7]. It also detects the augmented link associated to all 2-braid representatives of the unknot, namely, $T(2, \pm 4)$. This was first proven by using instanton Floer homology [44], but see also [13] for a proof that is more in line with that of Theorem 2.1. Martin showed that Khovanov homology detects $T(2, 6)$, one of the braid-closures of a 3-braid representing the unknot.

The main tool we use to prove these results is Dowlin's spectral sequence [17] from Khovanov homology to knot Floer homology — a version of link Floer homology introduced independently in [35] by Ozsváth and Szabó and in [39] by J. Rasmussen. This allows us to reduce the question of detection for Khovanov homology to problems in link Floer homology.

Finally we study *annular Khovanov homology*, a version of Khovanov homology for links in the thickened annulus due to Asaeda, Przytycki and Sikora [1]. We have the following family of results:

Theorem 3.11. *Annular Khovanov homology with integer coefficients detects $b(\sigma_1\sigma_2^n)$ for $-2 \leq n \leq 5$.*

For context, recall that annular Khovanov homology detects the braid-closure of the identity braids [2], and all braid-closures of 2-braids by a combination of work of Grigsby and Ni [19] and Grigsby, Licata and Wehrli [19]. The author and Martin also showed the $n = 1$ case of Theorem 3.11 in [13]. For the proof of our result we use the Birman–Menasco classification of 3-braids with fixed closures [15].

We can also prove the following:

Theorem 3.13. *Annular Khovanov homology with integer coefficients detects the Mazur pattern.*

Note that annular Khovanov homology detects the clasp-closures of all 2-braids, amongst annular knots [11, Theorem 8.1]. For the proofs of the two preceding theorems we use a version of the following rank bound:

Theorem 3.1. *Let β be an n -braid with $n \geq 2$. Then:*

- (1) $\text{rank}(\text{AKh}(b(\beta); \mathbb{C})) \geq 2n$.¹
- (2) $\text{rank}(\text{AKh}(c(\beta); \mathbb{C})) \geq 4n$.

This result is inspired by the proof of a structurally similar rank bound in knot Floer homology due to Baldwin, Vela and Vick [5]. The proof relies on the left orderability of the braid group. See Lemma 3.2 for the more technical version of the result that we apply to prove Theorem 3.13 and Theorem 3.11. A number of other consequences of Theorem 3.1 are noted in Section 3C. The clasp-closure statement version of Theorem 3.1 is perhaps more interesting because there is currently no analogous result in the link Floer homology context.

Remark. Link Floer homology, Khovanov homology and annular Khovanov homology are invariant under overall orientation reversal. All of the detection and classification results in this paper are thus up to overall orientation reversal, if any relevant link and its reverse are distinct.

We end the introduction with two questions:

Question 1. Is there a complete classification of clasp-closures of 3-braids in the style of the Birman–Menasco classification of braid-closures of 3-braids?

Such a classification might allow one to obtain more classification results for links with categorified link invariants taking certain values.

¹This bound appears in an unpublished note of the author written while he was a graduate student.

Question 2. Does annular Khovanov homology detect all clasp-closures of 3-braids representing the unknot? Does Khovanov homology detect all of their augmentations? Does link Floer homology detect the links $\mathring{b}(\sigma_1\sigma_2^n)$?

Outline. In Section 1 we prove our results for link Floer homology. In Section 2 we prove our Khovanov homology detection results. In Section 3 we prove our annular Khovanov homology detection results as well as our two rank bounds.

1. Link Floer homology

In this section we collect our detection results for link Floer homology. In Section 1B we show that link Floer homology detects L6a2 and L9n15. In Section 1C we give a partial classification of links with the link Floer homology types of augmentations of clasp-closures of index 3-braids that represent the unknot.

1A. A review. Link Floer homology is an invariant introduced by Ozsváth and Szabó [36]. It assigns to each oriented n -component link a finitely generated vector space equipped with $n + 1$ gradings. The first n of these gradings are called *Alexander gradings*, and the last is called the *Maslov grading*. The Alexander gradings takes value in $\frac{1}{2}\mathbb{Z}$, while the Maslov grading takes value in \mathbb{Z} . For each component K of a link L , there is a spectral sequence from $\widehat{\text{HFL}}(L)$ to

$$\widehat{\text{HFL}}(L \setminus K) \otimes V\left[\frac{\ell\text{k}(K, L \setminus L_i)}{2}\right]$$

corresponding to allowing pseudoholomorphic disks to “cross basepoints” in Heegaard diagrams [36, Proposition 7.1]. Throughout this paper we consider link Floer homology with $\mathbb{Z}/2$ coefficients unless explicitly stated otherwise.

Link Floer homology detects the Thurston norm, under mild hypotheses, by a result of Ozsváth and Szabó [37]. It also detects braid closures, by a result of Martin [32, Proposition 1].

1B. Braid-closures.

Theorem 1.1. *Link Floer homology detects L6a2.*

For the reader’s convenience we recall that the link Floer homology of L6a2 is as follows:

(1)

$A_1 \rightarrow$	$-\frac{3}{2}$	$-\frac{1}{2}$	$\frac{1}{2}$	$\frac{3}{2}$
$A_2 \downarrow$				
$\frac{3}{2}$		\mathbb{F}_{-1}	\mathbb{F}_0	
$\frac{1}{2}$	\mathbb{F}_{-3}	\mathbb{F}_{-2}^3	\mathbb{F}_{-1}^3	\mathbb{F}_0
$-\frac{1}{2}$	\mathbb{F}_{-4}	\mathbb{F}_{-3}^3	\mathbb{F}_{-2}^3	\mathbb{F}_{-1}
$-\frac{3}{2}$		\mathbb{F}_{-4}	\mathbb{F}_{-3}	

This can be deduced from, say, the fact that L6a2 is alternating, the multivariable Alexander polynomial of L6a2, the signature of L6a2, and an application of [36, Theorem 1.3].

Proof of Theorem 1.1. Our strategy is to argue that if a link has the link Floer homology type as L6a2 then it is the augmentation of a braid-closure of a 3-braid by applying Martin’s braid axes detection result [32, Proposition 1]. We then appeal to Murasugi’s classification of 3-braids whose braid-closures are unknotted and note that link Floer homology distinguishes the corresponding links.

Suppose L is a link with the link Floer homology of L6a2. Then L cannot be split, since its link Floer homology is not of the correct form. More specifically, if L were split then it could be written as $K_1\#(U \sqcup K_2)$ for K_i knots and U the unknot, where the connect sum is between U and K_1 . Consequently, the Künneth formula for link Floer homology [36, Theorem 1.4] would imply that each Alexander bigrading would have rank at least two, which we can observe is not the case from Table (1). Since the rank of $\widehat{\text{HFL}}(L)$ in the maximal nontrivial A_1 grading is 2, it follows from [32, Proposition 1] that the first component of L , L_1 , is a braid axis. Observe that the Conway polynomial of two-component links — and hence knot Floer homology and link Floer homology — detects the linking number of two-component links [21]. It follows that L is the augmentation of the braid-closure of a 3-braid, L_2 .

For the remainder of this section, let V be a rank-two vector space supported in Alexander grading 0 and Maslov gradings 0 and -1 , and let $[a]$ indicate a shift in Alexander grading by a . There are spectral sequences from $\widehat{\text{HFL}}(L)$ to $\widehat{\text{HFL}}(L_i) \otimes V[\ell k(L)/2]$ for each i corresponding to allowing pseudoholomorphic disks to “cross basepoints” in Heegaard diagrams [36, Proposition 7.1]. Hence $\widehat{\text{HFL}}(L_i)$ can be supported only in Alexander grading zero, so that each L_i is the unknot. Thus, L is the augmentation of a braid-closure of a 3-braid representing the unknot. By Murasugi’s classification of 3-braids up to conjugacy there are exactly three 3-braid representatives of the unknot; namely $(\sigma_1\sigma_2)^{\pm 1}$, and $\sigma_1\sigma_2^{-1}$ [33]. Taking augmentations of the braid-closures of either of the first two braids yields $T(2, \pm 6)$, which have distinct link Floer homology from L . The result follows. \square

Remark 1.2. Of course, a two-component unoriented link can, a priori, be endowed with four distinct orientations. However, $\overline{\text{L6a2}}$ is isotopic to the link obtained from L6a2 by reversing the orientation of either component. Likewise, the reverse of L6a2 is isotopic to L6a2. Thus, Theorem 1.1 holds for oriented links.

We proceed to our next detection result.

Theorem 1.3. *Link Floer homology with rational coefficients detects L9n15.*

We will not compute the link Floer homology of $\mathring{b}(\sigma_1^3\sigma_2)$. Instead, we will rely on

formal properties of link Floer homology. The reason we take rational coefficients is that we will use the Khovanov homology of L9n15 to obtain information about link Floer homology via Dowlin's spectral sequence [17], which is defined over the rational numbers.

Proof. We first study the link Floer homology of $\widehat{\text{HFL}}(\mathring{b}(\sigma_1^3\sigma_2); \mathbb{Q})$. Observe that, perhaps after relabeling components, $\widehat{\text{HFL}}(\mathring{b}(\sigma_1^3\sigma_2); \mathbb{Q})$ has maximal A_2 grading $\frac{3}{2}$, since we may take the second component to be the braid axis for the braid-closure $\mathring{b}(\sigma_1^2\sigma_2)$. Now, $\mathring{b}(\sigma_1^2\sigma_2)$ bounds a 3-punctured torus, so the maximal A_1 -grading is at most $\frac{5}{2}$. In fact, the maximal A_1 grading must be at least $\frac{5}{2}$ since $\widehat{\text{HFL}}(L; \mathbb{Q})$ admits a spectral sequence to

$$\widehat{\text{HFL}}(T(2, -3); \mathbb{Q}) \otimes V[-\frac{3}{2}],$$

so that $\widehat{\text{HFL}}(L; \mathbb{Q})$ must have generators of A_1 grading $\pm\frac{5}{2}$. From Knot atlas [28] we have that $\text{rank}(\text{Kh}(\text{L9n15}; \mathbb{Z}/2)) = 12$ — see also Table (5) — so that $6 = \text{rank}(\text{Khr}(\text{L9n15}; \mathbb{Z}/2)) \geq \text{rank}(\text{Khr}(\text{L9n15}; \mathbb{Q}))$ by the universal coefficient theorem and [40, Corollary 3.2.C]. It follows that $\text{rank}(\widehat{\text{HFL}}(\text{L9n15}; \mathbb{Q})) \leq 12$ by an application of the rank bound from Dowlin's spectral sequence [17] together with some properties of pointed Khovanov homology [6, Lemma 2.11].

Suppose L is a link with the same link Floer homology with rational coefficients as $\mathring{b}(\sigma_1^3\sigma_2)$. Since $\widehat{\text{HFL}}(L; \mathbb{Q})$ determines the Conway polynomial of L and the Conway polynomial of L determines the linking number of two-component links [21], it follows that L has linking number -3 . In particular L is nonsplit. Now, the link Floer polytope of L agrees with that of L9n15. Since the link Floer polytope detects the Thurston polytope [37] for nonsplit links, it follows that the Thurston polytopes of L and L9n15 agree. Consequently L_2 bounds a surface in the exterior of L of Euler characteristic -2 , just as does the corresponding component in L9n15. Since such a surface necessarily has at least four boundary components, it follows that it is, in fact, a 4-punctured disk, so that L_2 is an unknot. Since the rank in the maximum nontrivial Alexander grading is 2, it follows that L_2 is a braid axis by [32, Proposition 1].

We now study the first component of L . From the link Floer polytope of L we can see that L_1 bounds a surface in the exterior of L of Euler characteristic -4 . Since the linking number of L is three, it follows that L_1 has Seifert-genus at most one. If L_1 has a Seifert surface of genus zero, then it is an unknot. Consequently L is an augmented braid closure of a 3-braid representing the unknot. As previously noted, these were classified by Murasugi [33] and are each detected by link Floer homology by [13] and Theorem 1.1. Thus L_1 bounds a genus one surface. In particular, $\widehat{\text{HFL}}(L_1; \mathbb{Q})$ is nontrivial in Alexander gradings ± 1 and trivial in Alexander gradings i with $|i| > 1$. Since $\widehat{\text{HFL}}(L_1; \mathbb{Q})$ has the same rank in Alexander gradings

± 1 and the total rank of $\widehat{\text{HFL}}(L; \mathbb{Q})$ must be odd, it follows that $\widehat{\text{HFL}}(L_1; \mathbb{Q})$ must be nontrivial in Alexander grading zero.

We now prove that L_1 is fibered. By [36, Theorem 1.4], there is a spectral sequence from $\widehat{\text{HFL}}(L; \mathbb{Q})$ to $\widehat{\text{HFL}}(L_1) \otimes V[-\frac{3}{2}]$. This latter space must be of rank at least two for $k = -\frac{1}{2}, -\frac{3}{2}, -\frac{5}{2}$, and so the same is true of $\widehat{\text{HFL}}(L; A_1 = k; \mathbb{Q})$. In fact, by symmetry properties of link Floer homology, $\widehat{\text{HFL}}(L, A_1 = k; \mathbb{Q})$ must also be of rank at least two for $k = \frac{1}{2}, \frac{3}{2}, \frac{5}{2}$. It follows that $\text{rank}(\widehat{\text{HFL}}(L; \mathbb{Q})) = 12$ and indeed that $\widehat{\text{HFL}}(L; \mathbb{Q})$ is of rank two in the maximal nontrivial A_1 grading. Martin's braid axis detection result implies that L_1 is a braid axis for L_2 and so, in particular, fibered [32, Proposition 1]. Indeed, since the maximal nontrivial A_1 grading is $1 + \frac{3}{2}$, L_1 must be a genus one fibered knot. It follows that L_1 is a trefoil or a figure eight knot. To see that L_1 is a left handed trefoil, observe that $\widehat{\text{HFL}}(\text{L9n15}, A_2 = -\frac{5}{2}; \mathbb{Q})$ must be supported in Maslov gradings 0 and -1 since it has a left-handed trefoil component and the linking number is -3 . This in turn implies that L_1 must have a left handed trefoil component, since in the right handed trefoil case and Figure eight case $\widehat{\text{HFL}}(L, A_1 = -\frac{5}{2}; \mathbb{Q})$ would have to be supported in Maslov gradings -2 and -3 or -1 and -2 respectively.

Now by Birman and Menasco's classification theorem for 3-braids [15], there are exactly two 3-braids with braid-closures representing $T(2, -3)$, namely $\sigma_1^{-1}\sigma_2^{-3}$ and $\sigma_1\sigma_2^{-3}$, which is L9n16. These two links are distinguished by their Alexander polynomials, so the result follows. \square

Remark 1.4. Once again there are — a priori — four possible orientations with which L9n15 can be endowed. One pair of these have linking number -3 while the other has linking number 3. It can be checked that each pair with the same linking number are, in fact, isotopic as links. That is, Theorem 1.3 holds as a statement for oriented links.

1C. Clasp-closures. In this section we study the links with the link Floer homology type of augmentations of clasp-closures of 3-braids representing the unknot. We prove two of the theorems advertised in the introduction:

Theorem 1.5. *Link Floer homology detects the Mazur link.*

Theorem 1.6. *Let L be a link. $\widehat{\text{HFL}}(L) \cong \widehat{\text{HFL}}(\hat{c}(\sigma_2^{-1}\sigma_1\sigma_2))$ if and only if L is of the form $\hat{c}(\sigma_1^n\sigma_2^{-1}\sigma_1\sigma_2)$ for some $n \in \mathbb{Z}$.*

We begin by discussing some structural properties of the link Floer homology of links that are augmentations of clasp-closures of 3-braids representing the unknot. Let L be such a link, with the first component of L , L_1 , being the clasp-closure of the 3-braid and the second component, L_2 , its axis. The maximal A_2 grading in which $\widehat{\text{HFL}}(L, A_2)$ has nontrivial support is $\frac{3}{2}$. This follows from [37, Theorem 1.1].




Figure 2. A sutured Heegaard diagram for the sutured manifold obtained by decomposing an augmented clasp-closure along a longitudinal surface for its axis. The outer boundary component of the surface corresponds to a longitude of L_2 , while the inner boundary components correspond to meridians of L_1 .

Lemma 1.7. *Suppose L is the augmentation of a clasp-closure, L_1 , of a braid representing a knot. Then the component of $\widehat{\text{HFL}}(L)$ with maximal nontrivial A_2 grading is given by $\mathbb{F}_{-1}[-1] \oplus \mathbb{F}_0^2[0] \oplus \mathbb{F}_1[1]$ up to overall shifts in the Maslov and A_1 gradings.*

A version of this result without the Maslov grading is given in [11, Lemma 5.9]. The proof of this lemma requires techniques from sutured Floer homology. The reader is directed to Juhász' papers [23; 24; 25] for necessary background.

Proof. Suppose L is as in the statement of the lemma. A sutured Heegaard diagram for the sutured manifold (Y, γ) obtained by decomposing the exterior of L along an appropriate maximal Euler characteristic longitudinal surface for L_2 is shown in Figure 2.

A priori, $\text{SFH}(Y, \gamma)$ only comes with a relative Maslov grading in each spin^c structure. However, in the case at hand these Maslov gradings can be upgraded to a relative Maslov grading that applies across all spin^c structures. To see this, observe that capping off the sutures corresponding to meridians of L_1 results in another sutured manifold, $(\widehat{Y}, \widehat{\gamma})$ in which all four of the generators of $\text{CF}(\widehat{Y}, \widehat{\gamma})$ are supported in a single spin^c -structure. The claim follows. It remains to check that the maps \widehat{p} from [25, Proposition 5.4] respect this relative Maslov grading which applies across all spin^c structures. However, this follows by repeating Juhász' proof of [25, Proposition 5.4]. Specifically, there is a Heegaard diagram for (Y, γ) that can be obtained by doubling a Heegaard diagram for the exterior of L along a certain subsurface [24, Proposition 5.2], and pseudoholomorphic disks from the doubled Heegaard diagram correspond to disks in the Heegaard diagram for L [24, Proposition 7.6]. This correspondence still holds if we fill in boundary components, yielding the desired result. \square

We can now prove that link Floer homology detects augmented clasp-closures of 3-braids. Our proof depends on the much more general classification of links with

link Floer homology of next to minimal rank in the maximal nontrivial Alexander grading of a given component due to the author and Dey [11, Theorem 5.1].

Lemma 1.8. *Suppose that a link L has the link Floer homology type of an augmented clasp-closure of a 3-braid representing the unknot. Then L is an augmented clasp-closure of a 3-braid.*

Proof. Suppose L is as in the statement of the lemma. We first claim that L has linking number ± 1 . Note that augmented clasp-closures of 3-braids have linking number ± 1 . Now recall that the Conway polynomial — and hence link Floer homology — detects the linking number of two-component links [21]. The claim follows.

After relabeling the components of L if necessary, we may assume that the component of $\widehat{\text{HFL}}(L)$ with maximal nontrivial Alexander grading of rank four is L_2 and that the maximal nontrivial A_2 grading is $\frac{3}{2}$ and that $\widehat{\text{HFL}}(L, A_2 = \frac{3}{2})$ is given by $\mathbb{F}_{-1}[-1] \oplus \mathbb{F}_0^2[0] \oplus \mathbb{F}_1[1]$, up to shifts in the A_1 and Maslov gradings by Lemma 1.7. We now bound the genus of the component L_2 . Recall that there is a spectral sequence from $\widehat{\text{HFL}}(L)$ to $\widehat{\text{HFL}}(L_2) \otimes V[\pm\frac{1}{2}]$. It follows that the maximum nontrivial Alexander grading in which L_1 can have nontrivial support is at most one.

By [11, Theorem 5.1] we have four cases to treat:

1. L_2 is a genus one fibered knot and L_1 is a clasp-braid with axis L_2 .
2. L_2 is a genus one nearly fibered knot and L_1 is a braid-closure with axis L_2 .
3. L_2 is a fibered knot and L_1 can be isotoped to a simple closed curve in a minimal genus Seifert surface for L_2 .
4. L_1 is a clasp-closure with L_2 its unknotted axis.

For definitions of “nearly fibered” see [4]. For a definition of what it is to be braided with respect to a nearly fibered knot see [11, Section 3]. We rule out the first three of the four possibilities.

Case 1: The maximal Euler characteristic of a longitudinal surface for L_2 would be -3 , so that the maximal A_2 grading in which $\widehat{\text{HFL}}(L)$ would be nontrivial support would be $\frac{5}{2}$ by [37, Theorem 1.1], a contradiction.

Case 2: Recall that there is a spectral sequence from $\widehat{\text{HFL}}(L)$ to $\widehat{\text{HFK}}(L_2) \otimes V[\pm\frac{1}{2}]$. Since in the maximal nontrivial A_2 grading $\widehat{\text{HFL}}(L)$ is of rank four, as is the rank of the maximal nontrivial Alexander grading of $\widehat{\text{HFK}}(L_2) \otimes V[\pm\frac{1}{2}]$, it follows that this spectral sequence collapses immediately. In particular, it follows that the component of $\widehat{\text{HFK}}(L_2) \otimes V$ in maximal nontrivial Alexander grading is given up to an overall shift in Maslov grading by $\mathbb{F}_{-1} \oplus \mathbb{F}_0^2 \oplus \mathbb{F}_1$. Now, Baldwin and Sivek classified all genus one nearly fibered knots [4]. All such knots have the property




Figure 3. The link $\mathring{c}(\sigma_1^n \sigma_2^{-1} \sigma_1 \sigma_2)$. We consider the unoriented resolution of the crossing highlighted in orange.

that their knot Floer homology in Alexander grading one is supported in exactly one Maslov grading — see [4, Table 1] — a contradiction.

Case 3: This is immediately excluded by the fact that the linking number of L is nonzero. \square

Lemma 1.9. *Suppose L is a link with the link Floer homology of either the Mazur link or $\mathring{c}(\sigma_2^{-1} \sigma_1 \sigma_2)$. Then L has unknotted components each of which is an unknotted clasp closure with respect to the other.*

Proof. Suppose L is as in the statement of the lemma. First observe that for both the Mazur link and $\mathring{c}(\sigma_2^{-1} \sigma_1 \sigma_2)$, each component is a clasp-closure of a 3-braid with respect to the other. The claim now follows from the same argument as given in the previous lemma, but now applied to both components of L . \square

By the preceding lemmas, to complete the proofs of Theorems 1.5 and 1.6 it suffices to show that link Floer homologies distinguishes between appropriate augmentations of clasp-closures of 3-braids representing the unknot.

We first address the links corresponding to the infinite family of braids $\sigma^n \sigma_2^{-1} \sigma_1 \sigma_2$. It can be checked that the unoriented resolution of this link at the crossing shown in Figure 3 is the split sum of a Hopf link and an unknot. J. Wang has shown that if L_b is a band sum of the split union of two links $L_1 \sqcup L_2$ then the link Floer homology of L_b does not change after adding twists to the band [42, Remark 1.18]. Thus the links $\mathring{c}(\sigma^n \sigma_2^{-1} \sigma_1 \sigma_2)$ all have the same link Floer homology.

We can now conclude the proofs of two of the results promised in the introduction.

Proof of Theorems 1.5 and 1.6. Suppose that L is a link as in one the two theorem statements. By Lemma 1.8 and Lemma 1.9, both components of L are clasp-closures of index 3-braids with respect to the other component. In particular, the maximum A_i gradings in which $\widehat{\text{HFL}}(L)$ are nontrivial are $\frac{3}{2}$. On the other hand, $\mathring{c}(\sigma_1^3 \sigma_2^{-1} \sigma_1^2 \sigma_2)$ does not have this property; one of the components — say L_1 — is the clasp-closure of an index 5 braid. Consequently, the maximum A_1 grading in which $\widehat{\text{HFL}}(\mathring{c}(\sigma_1^3 \sigma_2^{-1} \sigma_1^2 \sigma_2))$ is nontrivial is $\frac{5}{2}$. It follows that L cannot be $\mathring{c}(\sigma_1^3 \sigma_2^{-1} \sigma_1^2 \sigma_2)$. By [42, Remark 1.18], $\widehat{\text{HFL}}(\mathring{c}(\sigma_1^n \sigma_2^{-1} \sigma_1 \sigma_2)) \cong \widehat{\text{HFL}}(\mathring{c}(\sigma_2^{-1} \sigma_1 \sigma_2))$

for all n . It thus suffices to show that no two of the Mazur link, $\hat{c}(\sigma_2^{-1}\sigma_1\sigma_2)$, and their mirrors have the same link Floer homology.

Given that link Floer homology detects the linking number of two-component links, the transformation property of link Floer homology under changing the orientation of a link component implies that for a two-component link, L , $\widehat{\text{HFL}}(L)$ determines the Alexander polynomial of L endowed with an arbitrary orientation. For the two links at hand, these are given as follows:

1. $\Delta(\hat{c}(\sigma_1^{-1})) = 2 - 5t + 5t^2 - 2t^3$.
2. $\Delta(\hat{c}'(\sigma_1^{-1})) = 1 - 3t + 3t^2 - 3t^3 + 3t^4 - t^5$.
3. $\Delta(\hat{c}(\sigma_2^{-1}\sigma_1\sigma_2)) = 2 - 7t + 7t^2 - 2t^3$,
4. $\Delta(\hat{c}'(\sigma_2^{-1}\sigma_1\sigma_2)) = 1 - 3t + 5t^2 - 5t^3 + 3t^4 - t^5$.

Here the primes indicate that the orientation of the axis has been reversed. The author used [27] for these computations. Note that the Alexander polynomial of a link is unchanged under mirroring. Thus link Floer homology distinguishes the Mazur link and its mirror, from $\hat{c}(\sigma_2^{-1}\sigma_1\sigma_2)$ and its mirror. Since the Mazur link and its mirror have linking numbers ± 1 , and likewise for $\hat{c}(\sigma_2^{-1}\sigma_1\sigma_2)$, the Alexander polynomial distinguishes each pair of links. Consequently link Floer homology distinguishes between the four links, concluding the proof. \square

Remark 1.10. The Mazur link and its reverse are isotopic, so for this link we have oriented link detection on the nose.

2. Khovanov homology

The goal of this section is to prove the following two results:

Theorem 2.1. *Khovanov homology with integer coefficients detects L6a2.*

Theorem 2.2. *Khovanov homology with integer coefficients detects L9n15.*

From Martin's result that Khovanov homology detects $T(2, 6)$ oriented as a 2-braid closure [32] we know that Khovanov homology detects $T(2, \pm 6)$ with both orientations. Thus Khovanov homology detects all augmentations of braid-closures of 3-braids representing the unknot.

2A. A review. *Khovanov homology* is an invariant of oriented links introduced in [26]. It assigns to each link a \mathbb{Z} -module equipped with two gradings: the *quantum grading* and the *homological grading*. The Khovanov homology of an n -component link is supported in quantum gradings whose parity agrees with that of n . Taking coefficients in a field, the Khovanov homology of L admits a link splitting spectral sequence to the Khovanov homology of the disjoint union of the underlying components of L . This is due to Batson and Seed [9]. Taking coefficients

in \mathbb{Q} , Khovanov homology admits a spectral sequence to an invariant called *Lee homology*, a result from [30]. Lee homology is a finitely generated \mathbb{Q} -vector space equipped with a homology grading compatible with that of Khovanov homology under the Lee spectral sequence. For a two component link L , the Lee homology of L is of rank four, with rank two in homological grading zero and rank four in homological grading the linking number of L .

There is a version of Khovanov homology due to Baldwin, Levine and Sarkar [6], called *pointed Khovanov homology*. Taking coefficients in \mathbb{Q} , pointed Khovanov homology admits a spectral sequence to knot Floer homology, a result of Dowlin [17]. Pointed Khovanov homology is a generalization of an invariant called *reduced Khovanov homology*. Taking coefficients in $\mathbb{Z}/2$, the rank of reduced Khovanov homology is exactly half that of Khovanov homology, a result of Shumakovitch [40, Corollary 3.2.C].

2B. Detection results. For convenience we record the Khovanov homology of L6a2 (see [28]):

$\begin{matrix} h \rightarrow \\ q \downarrow \end{matrix}$	-6	-5	-4	-3	-2	-1	0
-2							\mathbb{Z}
-4						\mathbb{Z}	\mathbb{Z}
-6					$\mathbb{Z} \oplus \mathbb{Z}/2$		
-8				$\mathbb{Z} \oplus \mathbb{Z}/2$	\mathbb{Z}		
-10			$\mathbb{Z} \oplus \mathbb{Z}/2$	\mathbb{Z}			
-12		$\mathbb{Z}/2$	\mathbb{Z}				
-14	\mathbb{Z}	\mathbb{Z}					
-16	\mathbb{Z}						

Lemma 2.3. *Suppose L is a link with the Khovanov homology of the L6a2. Then L is a two-component link with each component an unknot. Moreover $\ell k(L) = -3$.*

Proof. Suppose L is as in the statement of the theorem. Observe that L has an even number of components since the quantum grading is supported in even gradings. The Batson–Seed link splitting spectral sequence [9], together with an application of the universal coefficient theorem, implies that

$$(2) \quad \text{rank}(\text{Kh}(L; \mathbb{Z}/2)) = 20 \geq \prod \text{rank}(\text{Kh}(L_i; \mathbb{Z}/2)).$$

Here, the product is taken over components L_i of L . Since $\text{rank}(\text{Kh}(L_i; \mathbb{Z}/2))$ is of the form $2 + 4k_i$ for some $k_i \geq 0$, we must have that L has at most four components. If L has exactly four components, then each component is an unknot by [29]. To check that this is impossible, we use the refined version of the Batson–Seed link splitting spectral sequence. Equip $\text{Kh}(L; \mathbb{Q})$ with the $l := h - q$ grading. Then [9,

Corollary 4.4] implies that for some constant t

$$(3) \quad \text{rank}^l(\text{Kh}(L; \mathbb{Z}/2)) \geq \text{rank}^{l+t}(W^{\otimes 4}).$$

Here, and for the remainder of this proof, W is the rank-two vector space supported in l gradings 1 and -1 . In particular, there is some l grading in which $\text{rank}(\text{Kh}^l(L; \mathbb{Z}/2)) \geq 6$. This is false by inspection. Thus, L has exactly two components. Observe that equation (2) implies that at least one component of L is an unknot, since the unknot is the unique knot K with $\text{rank}(\text{Kh}(K; \mathbb{Z}/2)) = 2$ [29]. The remaining component K_2 of L has $\text{rank}(\text{Kh}(K_2; \mathbb{Z}/2)) \leq 6$, so that it is either an unknot or a trefoil by [29] and [8].

To see that $\ell k(L) = -3$, recall that $\text{Kh}(L; \mathbb{Q})$ — which can be obtained from $\text{Kh}(L; \mathbb{Z})$ by an application of the universal coefficient theorem — admits a spectral sequence to Lee homology [30]. Lee homology carries a homological grading and the spectral sequence respects this grading. Moreover, the Lee homology of a two-component link L is supported in homological gradings 0 and $2\ell k(L)$, and that each such grading contributes a \mathbb{Q}^2 summand. By inspection of $\text{Kh}(L; \mathbb{Q})$ we see that L must have linking number -3 .

To check that the remaining component of L is also unknotted, we use the refined version of the Batson–Seed link splitting spectral sequence again. Equip $\text{Kh}(L; \mathbb{Q})$ with the $l := h - q$ grading; then [9, Corollary 4.4] implies that

$$(4) \quad \text{rank}^l(\text{Kh}(L; \mathbb{Q})) \geq \text{rank}^{l-3}(\text{Kh}(K_2; \mathbb{Q}) \otimes W),$$

where W is the rank-two vector space supported in l gradings 1 and -1 . Since $\text{Kh}(T(2, 3); \mathbb{Q})$ has a generator in l grading -6 , $\text{Kh}(K_2; \mathbb{Q}) \otimes W$ has a generator of l grading -7 , violating the rank bound. Likewise since $\text{Kh}(T(2, -3); \mathbb{Q})$ has two generators in l grading 3, $\text{Kh}(T(2, -3); \mathbb{Q}) \otimes W$ has two generators in l grading 2 violating the rank bound. \square

The remainder of the proof of Theorem 2.1 amounts to showing that there is a component of L that is a braid axis for the other. To do so, we use Dowlin’s spectral sequence from an appropriate version of Khovanov homology to $\widehat{\text{HFK}}(L; \mathbb{Q})$ to reduce this question to a question about link Floer homology [17].

Proof of Theorem 2.1. Suppose L is as in the statement of the Theorem. By the previous lemma, L has two components. Since L has δ -thin Khovanov homology, $\widehat{\text{HFL}}(L; \mathbb{Q})$ is δ -thin. A result of the author and Dey [12, Proposition 6.1] implies in turn that $\widehat{\text{HFL}}(L; \mathbb{Q})$ decomposes as a direct sum of vector spaces of the form

$$W_a[b, c] := \mathbb{Q}_{a-1}\left[b - \frac{1}{2}, c - \frac{1}{2}\right] \oplus \mathbb{Q}_a\left[b + \frac{1}{2}, c - \frac{1}{2}\right] \oplus \mathbb{Q}_a\left[b - \frac{1}{2}, c + \frac{1}{2}\right] \oplus \mathbb{Q}_{a+1}\left[b + \frac{1}{2}, c + \frac{1}{2}\right].$$

Here $\mathbb{Q}_a[b, c]$ is a \mathbb{Q} summand in (A_1, A_2) grading (b, c) of Maslov grading a . There are at most five of these summands since

$$20 = \text{rank}(\text{Kh}(L; \mathbb{Z}/2)) \geq \text{rank}(\widehat{\text{HFK}}(L; \mathbb{Q})),$$

as follows from Dowlin's spectral sequence [17] together with the same steps applied in the corresponding stage of the proof of Theorem 1.3.

Observe that if the span of an A_i grading is $[\frac{-1}{2}, \frac{1}{2}]$ then L_i is a meridian of the other component, so that L is a Hopf link, since each component is unknotted by Lemma 2.3. Neither Hopf link has the correct Khovanov homology, so the span of each Alexander gradings must be strictly larger than $[\frac{-1}{2}, \frac{1}{2}]$.

If $\widehat{\text{HFL}}(L; \mathbb{Q})$ contains an odd number of $W_a[b, c]$ summands, the symmetry of link Floer homology implies that $\widehat{\text{HFL}}(L; \mathbb{Q})$ contains a $W_b[0, 0]$ summand, for some b . Thus we are in one of the following cases:

1. $\widehat{\text{HFL}}(L; \mathbb{Q})$ has a $W_a[m, n]^{\oplus 2} \oplus W_{a-2m-2n}[-m, -n]^{\oplus 2}$ summand, where $m, n \in \frac{1}{2}\mathbb{Z}$, $m, n \geq \frac{1}{2}$ and $a = b + n + m$, if there is a $W_b[0, 0]$ summand.
2. $\widehat{\text{HFL}}(L; \mathbb{Q})$ has a

$$W_a[m, n] \oplus W_{a-2m}[-m, n] \oplus W_{a-2m-2n}[-m, -n] \oplus W_{a-2n}[m, -n]$$

summand, where $m, n \in \frac{1}{2}\mathbb{Z}$, $m, n \geq \frac{1}{2}$ and $a = b + n + m$, if there is a $W_b[0, 0]$ summand.

3. $\widehat{\text{HFL}}(L)$ is of rank two, in a maximal nontrivial A_i grading,

Suppose that we are in one of the first two cases. L is nonsplit because both components are unknotted and the Khovanov homology of the two-component unlink is of rank four. Thus, we can apply [11, Theorem 5.1]. Since L_1 is unknotted, we deduce that L_2 is a clasp-closure with respect to L_1 . However, Lemma 1.7 then implies that the maximal nontrivial A_1 grading is given, up to affine isomorphism, by $\mathbb{Q}[-1] \oplus \mathbb{Q}^2[0] \oplus \mathbb{Q}[1]$. This is a direct contradiction in case 1. In case 2 $\widehat{\text{HFL}}(L; \mathbb{Q})$ would contain a $W_a[n, \frac{1}{2}] \oplus W_{a-1}[n, -\frac{1}{2}] \oplus W_{a-1-2n}[-n, -\frac{1}{2}] \oplus W_{a-2n}[-n, \frac{1}{2}]$ summand, a contradiction since L has odd linking number so that $\widehat{\text{HFL}}(L; \mathbb{Q})$ must be supported in $\mathbb{Z} + \frac{1}{2}$ valued Alexander gradings.

Thus $\widehat{\text{HFL}}(L; \mathbb{Q})$ is of rank two in one of the maximal nontrivial Alexander gradings. By [32, Proposition 1] one component — say L_1 — is a braid axis for the other — say L_2 . Since the linking number of the two links is -3 , it follows that L_2 is the braid-closure of a 3-braid. Since L_2 represents the unknot, the desired result follows from Murasugi's classification of 3-braids with unknotted braid-closures up to conjugacy [33] and the fact that $T(2, \pm 6)$ has distinct Khovanov homology from L6a2. \square

Remark 2.4. Using the same argument given in Remark 1.2, it can be shown that

Khovanov homology detects L6a2 regardless of the orientation.

We now proceed to our next detection result, for L9n15, whose Khovanov homology we recall [28]:

(5)

$\begin{matrix} h \rightarrow \\ q \downarrow \end{matrix}$	-6	-5	-4	-3	-2	-1	0
-6							\mathbb{Z}
-8					$\mathbb{Z}/2$		\mathbb{Z}
-10					\mathbb{Z}		
-12			\mathbb{Z}				
-14			\mathbb{Z}	\mathbb{Z}			
-16	\mathbb{Z}	\mathbb{Z}					
-18	\mathbb{Z}	\mathbb{Z}					

Proof of Theorem 2.2. Suppose L is a link with $\text{Kh}(L; \mathbb{Z}) \cong \text{Kh}(\text{L9n15}; \mathbb{Z})$. We first determine the components of L . There is an even number of them, since $\text{Kh}(L; \mathbb{Z})$ is supported in even quantum gradings. An application of the universal coefficient theorem gives $\text{Kh}(L; \mathbb{Z}/2) \cong \text{Kh}(\text{L9n15}; \mathbb{Z}/2)$, so $\text{Kh}(L; \mathbb{Z}/2)$ has rank 12. Consider the Batson–Seed link splitting spectral sequence [9]. Every knot has Khovanov homology with $\mathbb{Z}/2$ coefficients of rank $2 + 4m$ for some m , so that L has at most two components. One of them, say L_1 , has $2 \text{rank}(\text{Khr}(L_1; \mathbb{Q}) \leq \text{rank}(\text{Kh}(L_1; \mathbb{Z}/2)) = 2$, thanks to [40, Corollary 3.2.C]; therefore L_1 is unknotted by [29]. The remaining component of L , L_2 , has $\text{rank}(\text{Kh}(L_2; \mathbb{Z}/2)) \leq 6$ and so, in turn, $\text{rank}(\text{Khr}(L_2; \mathbb{Z}/2)) \leq 3$ by [40, Corollary 3.2.C]. It follows from [8] and [29] that L_2 is an unknot or a trefoil.

An application of the universal coefficient theorem shows that $\text{Kh}(L; \mathbb{Q}) \cong \text{Kh}(\text{L9n15}; \mathbb{Q})$. Consider the spectral sequence from Khovanov homology to Lee homology. Since this spectral sequence respects the homological grading and the Lee homology of a two-component link consists of two $\mathbb{Q} \oplus \mathbb{Q}$ summands supported in homological gradings 0 and $2\ell k(L)$, we can see by inspection that $\ell k(L) = -3$. We can then apply equation (4) again; or rather to the same equation but where we take $\mathbb{Z}/2$ -coefficients rather than rational coefficients. Let U denote the unknot. Note that $\text{Kh}(U; \mathbb{Z}/2) \otimes W$ has support in $l := h - q$ grading 2 so that L_1 cannot be the unknot. Likewise $\text{Kh}(T(2, 3); \mathbb{Z}/2) \otimes W$ has support in $l := h - q$ grading -7 so that in fact L_2 is $T(2, -3)$.

By an application of [40, Corollary 3.2.C] and the universal coefficient theorem we have $\text{rank}(\text{Khr}(L; \mathbb{Q})) \leq 6$, so that $\text{rank}(\widehat{\text{HFK}}(L; \mathbb{Q})) \leq 12$ by the rank bound coming from Dowlin’s spectral sequence [17]. Recall that there is a spectral sequence from $\widehat{\text{HFL}}(L; \mathbb{Q})$ to $\widehat{\text{HFL}}(L_2; \mathbb{Q}) \otimes V[-\frac{3}{2}]$. Since L has linking number -3 and the second component is a copy of $T(2, -3)$ it follows that in A_2 grading $-\frac{5}{2}$ there is a $\mathbb{Q}_0 \oplus \mathbb{Q}_{-1}$ -summand, in A_2 grading $-\frac{3}{2}$ there is a $\mathbb{Q}_1 \oplus \mathbb{Q}_0$ -summand

and that in A_2 grading $-\frac{1}{2}$ there is a $\mathbb{Q}_1 \oplus \mathbb{Q}_2$ -summand. This completely determines the A_2 -graded version of the link Floer homology of L by symmetry properties and the fact that the rank is at most 12. Now, since L_1 is unknotted and the linking number of L is -3 , $\widehat{\text{HFL}}(L; \mathbb{Q})$ must have support in Alexander gradings $\pm\frac{3}{2}$. Since homogeneous summands with A_2 -grading at least 0 must all die under the spectral sequence to $\widehat{\text{HFL}}(T(2, -3); \mathbb{Q}) \otimes V[-\frac{3}{2}]$, we have that the pairs of generators in each A_2 grading must be of distinct A_1 gradings.

The span of δ -graded $\text{Kh}(L; \mathbb{Z}/2)$ is 4, so the span of δ -graded $\text{Khr}(L; \mathbb{Z}/2)$ is 2. Thus, the span of δ -graded pointed Khovanov homology, $\widetilde{\text{Kh}}(L, \mathbf{p}; \mathbb{Z}/2)$ where \mathbf{p} consists of a point on each component of L , is at most 2 by [6, Lemma 2.11] and so finally the span of δ -graded $\widehat{\text{HFL}}(L; \mathbb{Q})$ is at most 2. It follows that at most three of the homogeneous \mathbb{Q} summands with A_2 -grading at most $-\frac{1}{2}$ occur in extremal A_1 gradings. It follows in turn that $\widehat{\text{HFL}}(L; \mathbb{Q})$ is of rank two in the maximal nontrivial A_1 grading, so that U is a braid axis for $T(2, -3)$. Since the linking number is -3 , the corresponding braid is a 3-braid. Now, by Birman and Menasco's classification of 3-braids with braid-closures representing the unknot, the only two such augmented braid-closures are L9n15 and L9n16. These are distinguished by their Khovanov homology, see [28]. The result holds for oriented links by Remark 1.4. \square

3. Annular Khovanov homology

In this section we study annular Khovanov homology. In Section 3A we review structural properties of the invariant we will use in the rest of the section. In Section 3B we prove rank bounds for the annular Khovanov homology of clasp-closures and braid-closures and give some applications of the rank bounds to the study of braid-closures. In Section 3C we give two braid-closure detection results. In Section 3E we apply a rank bound from Section 3B to prove that annular Khovanov homology detects the Mazur pattern.

3A. A review. We begin with a brief review of annular Khovanov homology, which was introduced by Asaeda, Przytycki and Sikora in [1]. We work with coefficients in R , where R is either \mathbb{Z} , \mathbb{C} , \mathbb{Q} or $\mathbb{Z}/2$. Annular Khovanov homology is an R -module-valued invariant of links in the thickened annulus. The underlying chain complex for the annular Khovanov homology of an annular link L is freely generated by complete resolutions of a fixed diagram for L where each circle is decorated with a 1 or an X . The resulting homology groups carry three gradings. The first of these gradings is called the *homological grading* which we shall denote by i , the second is the *quantum grading* which we shall denote by j . These two gradings are defined just as in the Khovanov homology case. The third grading is called the *annular grading*, which we shall denote by k . This is defined as the difference between the

number of resolutions encircling the annular axis marked with a 1 and those marked with an X . The differential on annular Khovanov homology is then simply defined as the components of the differential on the Khovanov complex of the underlying link that preserve the annular grading.

We will use two exact triangles for annular Khovanov homology. Recall — say from [11, Lemma 8.2] — that annular Khovanov admits the following skein exact triangle corresponding to resolving a negative crossing:

$$(6) \quad \begin{array}{ccc} \text{AKh}(L) & \xrightarrow{\quad\quad\quad} & \text{AKh}(L_0)[n_-^0 - n_-]\{3n_- - 3n_- + 1\} \\ & \swarrow & \nwarrow \delta \\ & \text{AKh}(L_1)\{-1\} & \end{array}$$

Here n_- is the number of negative crossings in the diagram for L , n_-^0 is the number of negative crossings in the diagram for L_0 , $\{a\}$ is a shift in the quantum grading by a and $[b]$ is a shift in the homological grading by b . The map δ increases the homological grading by one. Corresponding to resolving a positive crossing we have the following exact triangle:

$$(7) \quad \begin{array}{ccc} \text{AKh}(L) & \xrightarrow{\quad\quad\quad} & \text{AKh}(L_0)\{1\} \\ & \swarrow & \nwarrow \delta \\ & \text{AKh}(L_1)[n_-^1 - n_- + 1]\{3n_-^1 - 3n_- + 2\} & \end{array}$$

Here the map δ again increases the homological grading by 1. Grigsby, Licata and Wehrli showed that for annular Khovanov homology with complex coefficients carries the structure of an $\mathfrak{sl}_2(\mathbb{C})$ representation [20]. This entails that, up to an overall grading shift in the quantum grading, $\text{AKh}(L; \mathbb{C})$ decomposes as a direct sum of vector spaces V_n^i , where V_n^i is the rank $n + 1$ vector space supported in homological grading i and quantum and annular gradings $(-n + 2p, -n + 2p)$ for all $0 \leq p \leq n$. It is not hard to see that annular Khovanov homology of an annular link L is supported in annular gradings j with $|j| \leq w(L)$, where $w(L)$ is the *wrapping number* of L ; i.e., the minimum *geometric* intersection number of L with a meridional disk for the thickened annulus. Consequently, $\text{AKh}(L; \mathbb{C})$ can only contain V_n summands with $n \leq w(L)$.

Annular Khovanov homology admits a spectral sequence to the Khovanov homology of the underlying link. The differential on annular Khovanov homology inducing this spectral sequence increases the homological grading by one, preserves the quantum grading and decreases the annular grading. Moreover, the differential forms part of an action of the $\mathfrak{sl}_2(\wedge)$ current algebra on $\text{AKh}(L; \mathbb{C})$ — a stronger structural property than being an $\mathfrak{sl}_2(\mathbb{C})$ representation. See [20, Section 6] for details.

It is not hard to see that $\text{AKh}(b(\beta), k = -n; R)$ consists of a single copy of R . This summand is generated by Plamenevskaya's *transverse invariant* [38]. This class consists of n concentric circles about the braid axis, each decorated with X . The quantum grading of this generator is the self-linking number of β — which we denote $sl(\beta)$ — and the homological grading is zero.

3B. From orderability to rank bounds. In this section we prove Theorem 3.1 and related results. The most concise version of our result is that stated in the introduction:

Theorem 3.1. *Let β be an n -braid with $n \geq 2$. Then:*

- (1) $\text{rank}(\text{AKh}(b(\beta); \mathbb{C})) \geq 2n$.
- (2) $\text{rank}(\text{AKh}(c(\beta); \mathbb{C})) \geq 4n$.

Of course, if $n = 1$, then $c(\beta)$ is undefined, while $\text{rank}(\text{AKh}(b(\beta); \mathbb{C})) = 2$. Theorem 3.1 is a direct consequence of the next — stronger — result. To state it recall that the braid group is left orderable. There are many different interpretations of the ordering on the braid group (see [16]), two of which we will use in this section. The first is the following: we write $\beta < 1$ if there is a word for β in the letters given by the standard Artin generators and their inverses which is σ -negative; i.e., if among the letters that occur in that word, the letter of the lowest index occurs only with negative powers. See [16, Chapter II, Section 1.2] for details.

Lemma 3.2. *Suppose β is a σ -negative n -braid. Then $\text{AKh}(c(\beta); \mathbb{C})$ contains a*

$$V_n^{a_1}\{a_2\} \oplus V_n^{b_1}\{b_2\} \oplus V_{n-2}^{a_1-1}\{a_2-2\} \oplus V_{n-2}^{b_1-1}\{b_2-2\}$$

summand. Moreover, the $\mathfrak{sl}_2(\wedge)$ action sends the lowest annular grading generator in $V_{n-2}^{a_1-1}\{a_2-2\}$ to the lowest annular grading generator in $V_n^{a_1}\{a_2\}$ and the lowest annular grading generator in $V_{n-2}^{b_1-1}\{b_2-2\}$ to the lowest annular grading generator in $V_n^{b_1}\{b_2\}$.

Similarly, $\text{AKh}(b(\beta); \mathbb{C})$ contains a $V_n^0\{n + sl(\beta)\} \oplus V_{n-2}^{-1}\{n - 2 + sl(\beta)\}$ summand. Moreover, the $\mathfrak{sl}_2(\wedge)$ action sends the lowest annular grading generator in $V_{n-2}^{-1}\{n + sl(\beta) - 2\}$ to the lowest annular grading generator in $V_n^0\{n + sl(\beta)\}$.

Remark 3.3. As we shall see, one can write down the values of the quantum and homological gradings of $\text{AKh}(c(\beta), k = -n; R)$ in terms of diagrammatic data for β . Comparing this to the braid closure case — where there is a single generator whose homological grading is zero and quantum grading is the self-linking number — it is natural to ask what topological or contact geometry-theoretic information these numbers contain. We do not pursue this question here.

Our strategy for the proof of Lemma 3.2 is to use properties of σ -negative words to control the annular Khovanov homology of closures of braids in the next to minimal annular grading.

Given an annular link L view $\text{CKh}(L; R)$ as a chain complex filtered with respect to the annular filtration. The differential comes in two pieces, $\partial_0 + \partial_{-2}$, where ∂_0 preserves the annular grading on $\text{CKh}(L; R)$ and ∂_{-2} decreases it by 2. $\text{AKh}(L; R)$ can be viewed as $(\text{CKh}(L; R), \partial_0)$, the first page of the corresponding spectral sequence.

Lemma 3.4. *Let β be a σ -negative braid which is not of index 1. Then there are chain maps*

$$\begin{aligned} f_c &: \text{CKh}(c(\beta); i, j, k \leq -n; R) \rightarrow \text{CAKh}(c(\beta); i-1, j, 2-n; R), \\ f_b &: \text{CKh}(b(\beta); i, j, k \leq -n; R) \rightarrow \text{CAKh}(b(\beta); i-1, j, 2-n; R). \end{aligned}$$

Moreover, ∂_{-2}^* is a left inverse to f_c or f_b on the E_1 page of the spectral sequence from $\text{CAKh}(c(\beta); R)$ or $\text{CAKh}(b(\beta); R)$ to $\text{Kh}(b(\beta); R)$.

Here by $\text{CKh}(c(\beta); i, j, k \leq -n; R)$ we mean the k -filtered part of filtration level n . It follows that $\text{AKh}(c(\beta), k = 2-n; R)$ has an $\text{AKh}(c(\beta), k = -n; R)[-1]$ summand while $\text{AKh}(b(\beta), k = 2-n; R)$ has a $\text{AKh}(b(\beta), k = -n; R)[-1]$ summand. Here, $[-1]$ indicates a shift in the homological grading by -1 .

Proof. We treat the case of clasp-closures. The proof in the braid-closure case is the same in essence and strictly easier in practice.

Since β is σ -negative β is isotopic to a braid β' that contains the inverse of an Artin generator σ_i^{-1} but not the corresponding Artin generator, σ_i . Consider the diagram D for $c(\beta')$ as in Figure 1(c). There are three complete resolutions D_1, D_2 and D_3 corresponding to $\text{CAKh}(c(\beta), k = -n; R)$; these are shown in Figure 4. There are four generators of $\text{CAKh}(c(\beta), k = -n; R)$. They can be described as follows. For each i we have a generator X_i where every circle in the resolution D_i is decorated with an X . We have a final generator $\mathbf{1}$ which corresponds to decorating the homologically essential circles in diagram D_1 with X s and the homologically inessential circle with a 1 . The nontrivial components of the differential are given by $\partial_0(X_2) = \partial_0(X_3) = X_1$ for an appropriate sign assignment.

Pick one of the crossings corresponding in D to a letter σ_i^{-1} and label it y . Consider the resolutions D'_i of D that are identical to the resolutions D_i aside from at y . Given a generator $\mathbf{x} \in \text{CAKh}(c(\beta), R; k = -n)$ define $f_c(\mathbf{x}) \in \text{CAKh}(c(\beta), R; k = 2-n)$ to be the generator which agrees with X_i on every circle in the resolution that does not involve y , and is labeled with an X on the remaining circle. Observe that if \mathbf{x} is of (i, j, k) grading $(a, b, -n)$ then $f_c(\mathbf{x})$ is of grading $(a-1, b, 2-n)$.

Since β is σ -negative, f_c is a chain map viewed as a map $(\text{CKh}(c(\beta); R), \partial_0) \rightarrow (\text{CKh}(c(\beta); R), \partial_0)$. To verify this notice that the maps corresponding to changing the resolutions in the β' part of the diagram correspond to merging circles decorated with X 's. Thus the only contributions to the differential on $\text{CAKh}(c(\beta), k = 2-n; R)$ involve the crossings contained in the part of the diagram for the clasp.




Figure 4. The resolutions for the canonical diagram, as in Figure 1(c), for a clasp-closure yielding generators of $\text{CAKh}(c(\beta), k = -n; R)$. The solid red dot is the annular axis.

One can check that ∂_{-2}^* is a left-inverse to f_c similarly; the only contributions to the differential which lower the annular filtration level correspond to changing the resolution at y . This corresponds to splitting a circle labeled with an X , resulting in two circles both labeled with X 's. \square

Proof of Lemma 3.2. We treat only the braid-closure case, since the clasp-closure case is essentially the same. Observe first that by the previous lemma, f_b induces an injection

$$f_b^* : \text{AKh}(b(\beta), k = -n; \mathbb{C}) \cong \mathbb{C}[0, sl(\beta), -n] \hookrightarrow \text{AKh}(b(\beta), (-1, sl(\beta), 2-n); \mathbb{C}).$$

Here $\mathbb{C}[0, sl(\beta), -n]$ indicates a \mathbb{C} summand supported in (i, j, k) grading $(0, sl(\beta), -n)$. Now, $\text{AKh}(b(\beta); \mathbb{C})$ carries the structure of an $\mathfrak{sl}_2(\mathbb{C})$ -representation, where each summand is supported in a single homological grading. The summand $\mathbb{C}[0, sl(\beta), -n]$ and its image under f_b^* are supported in different homological gradings and hence cannot be part of the same irreducible $\mathfrak{sl}_2(\mathbb{C})$ representation. The image $f_b^*(\mathbb{C}[0, sl(\beta), -n])$ is a minimal annular grading \mathbb{C} -summand of its $\mathfrak{sl}_2(\mathbb{C})$ representation, as else we would have $\text{rank}(\text{AKh}(b(\beta), k = -n; \mathbb{C})) \geq 2$, which is a contradiction since $b(\beta)$ is a braid-closure. It follows that the $\text{AKh}(b(\beta); \mathbb{C})$ contains the two desired representations as summands.

The structure of these summands as an $\mathfrak{sl}_2(\wedge)$ representation follow from the fact that ∂_{-2}^* is part of the $\mathfrak{sl}_2(\wedge)$ action. \square

We can now extract a rank bound for annular Khovanov homology with $\mathbb{Z}/2$ coefficients from Lemma 3.2 and the proof of Lemma 3.4.

Lemma 3.5. *Let β be a nonidentity $n > 1$ -braid. Then*

$$\begin{aligned} \text{rank}(\text{Kh}(b(\beta)); \mathbb{Z}/2) &\leq \text{rank}(\text{AKh}(b(\beta); \mathbb{Z}/2)) - 2(n - 1), \\ \text{rank}(\text{Kh}(c(\beta); \mathbb{Z}/2)) &\leq \text{rank}(\text{AKh}(c(\beta); \mathbb{Z}/2)) - 4(n - 1). \end{aligned}$$

Proof. Suppose β is as in the statement of the lemma. Since any nonidentity braid β is either σ -negative or σ -positive, we have four cases to consider. We prove the result in the σ -negative braid-closure case. The other three cases are similar.

We show that the rank of the map $\partial_{-2}^* : \text{AKh}(b(\beta); \mathbb{Z}/2) \rightarrow \text{AKh}(b(\beta); \mathbb{Z}/2)$ is at least $n - 1$; this implies the result. The universal coefficient theorem for homology is functorial, so considering the map ∂_{-2}^* we obtain the following commutative diagram, where μ is the map defined on elementary tensors by $\mu(\mathbf{x} \otimes a) \mapsto a\mathbf{x}$, and where we have suppressed $b(\beta)$ from the notation in each nontrivial entry to make room.

$$(8) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \text{AKh}_i(\mathbb{Z}) \otimes \mathbb{F} & \xrightarrow{\mu} & \text{AKh}_i(\mathbb{F}) & \longrightarrow & \text{Tor}_{\mathbb{Z}}^1(\text{AKh}_{i+1}(\mathbb{Z}), \mathbb{F}) \xrightarrow{h} 0 \\ & & \downarrow \partial_{-2}^* \otimes \mathbf{1} & & \downarrow \partial_{-2}^* & & \downarrow \\ 0 & \longrightarrow & \text{AKh}_{i+1}(\mathbb{Z}) \otimes \mathbb{F} & \xrightarrow{\mu} & \text{AKh}_{i+1}(\mathbb{F}) & \longrightarrow & \text{Tor}_{\mathbb{Z}}^1(\text{AKh}_{i+2}(\mathbb{Z}), \mathbb{F}) \xrightarrow{h} 0 \end{array}$$

Taking $\mathbb{F} = \mathbb{C}$, so that $\text{Tor}_{\mathbb{Z}}^1(\text{AKh}_j(b(\beta); \mathbb{Z}), \mathbb{F})$ vanishes for all j , and setting $i = -1$ we obtain

$$(9) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \text{AKh}_{-1}(b(\beta); \mathbb{Z}) \otimes \mathbb{C} & \xrightarrow{\mu} & \text{AKh}_{-1}(b(\beta); \mathbb{C}) & \longrightarrow & 0 \\ & & \downarrow \partial_{-2}^* \otimes \mathbf{1} & & \downarrow \partial_{-2}^* & & \\ 0 & \longrightarrow & \text{AKh}_0(b(\beta); \mathbb{Z}) \otimes \mathbb{C} & \xrightarrow{\mu} & \text{AKh}_0(b(\beta); \mathbb{C}) & \longrightarrow & 0 \end{array}$$

Now, from the proof of Lemma 3.4, $\partial_{-2}^* : \text{AKh}_{-1}(b(\beta); \mathbb{C}) \rightarrow \text{AKh}_0(b(\beta); \mathbb{C})$ has a component given by the identity map $\mathbb{C}^{\oplus(n-1)} \rightarrow \mathbb{C}^{\oplus(n-1)}$ with respect to the canonical basis for $\text{AKh}(b(\beta); \mathbb{C})$. More specifically, since ∂_{-2}^* is nontrivial on a bottom generator of $V_{n-2}^{-1}\{n-2+sl(\beta)\}$ and ∂_{-2}^* is part of the structure of $\text{AKh}(b(\beta); \mathbb{C})$ as an $\mathfrak{sl}_2(\wedge)$ -representation, we have that ∂_{-2}^* is nontrivial on the entire $V_{n-2}^{-1}\{n-2+sl(\beta)\}$ summand. It follows from diagram (9) that $\text{AKh}_{-1}(b(\beta); \mathbb{Z})$ and $\text{AKh}_0(b(\beta); \mathbb{Z})$ contain $\mathbb{Z}^{\oplus(n-1)}$ summands and that ∂_{-2}^* acts as the identity map between these two summands.

Now take $\mathbb{F} = \mathbb{Z}/2$ in diagram (8). By the commutativity of the diagram we deduce that $\partial_{-2}^* : \text{AKh}_0(b(\beta); \mathbb{Z}/2) \rightarrow \text{AKh}_0(b(\beta); \mathbb{Z}/2)$ has a component given by the identity map $(\mathbb{Z}/2)^{\oplus(n-1)} \rightarrow (\mathbb{Z}/2)^{\oplus(n-1)}$. The result follows. \square

Proof of Theorem 3.1. Suppose β is a braid. Then β is either σ -positive, σ -negative or the identity. If β is the identity then $\text{rank}(\text{AKh}(b(\beta); \mathbb{C})) = 2^n \geq 2n$.

If β is σ -negative the result follows immediately from Lemma 3.2. If β is σ -positive, the result follows from applying Lemma 3.2 to the mirror of β , which is σ -negative, and applying symmetry properties of annular Khovanov homology. \square

We can also prove an annular Khovanov homology analogue of a result of Ni from knot Floer homology [34, Theorem A.1]. To do so, we exploit a geometric

interpretation of the ordering of the braid group in terms of curve diagrams; see [16, Chapter 10]. Recall that n -braids can be viewed as mapping classes of n -punctured disks. Recall too that a braid is *right (left) veering* if it sends every *admissible* arc to the right (left). See [2, Section 3.1] for a definition of admissible. If a braid is non-right (left) veering then it is conjugate to a σ -negative (positive) braid; see the proof of [2, Proposition 3.1].

Proposition 3.6. *Suppose β is a non-right-veering and non-left-veering n -braid, with $n \geq 4$. Then $\text{rank}(\text{AKh}(b(\beta); \mathbb{C})) \geq 4n - 4$.*

Proof. Suppose β is as in the statement of the proposition. Since V is an n -braid there is a V_n^0 summand. Since β is non-right-veering, it is conjugate to a braid β' that is σ -negative. Since $b(\beta') = b(\beta)$, $\text{AKh}(b(\beta); \mathbb{C})$ has a V_{n-2}^{-1} -summand by Lemma 3.2. Similarly, since β is non-left-veering, $\text{AKh}(b(\beta); \mathbb{C})$ contains a V_{n-2}^1 summand by Lemma 3.2.

Assume towards a contradiction that there is a unique generator in (i, k) -grading $(0, 4 - n)$. Consider the generator x of (i, k) -grading $(-1, 6 - n)$ and form $\partial_{-2}^* : \text{AKh}(b(\beta); \mathbb{C}) \rightarrow \text{AKh}(b(\beta); \mathbb{C})$. Since $b(\beta)$ is isotopic to the braid-closures of σ -positive and σ -negative words, Lemma 3.2 and the symmetry properties of the spectral sequence from annular Khovanov homology to Khovanov homology under mirroring imply that $(\partial_{-2}^*)^2(x) \neq 0$, a contradiction. The result now follows from noting that $\text{AKh}(b(\beta); \mathbb{C})$ carries the structure of an $\mathfrak{sl}_2(\mathbb{C})$ -representation; specifically there must be at least one more generator in (i, k) -grading $(0, 4 - n)$, and so in turn another V_{n-4} -summand. \square

In the case that $n = 3$ we have:

Proposition 3.7. *Suppose β is a 3-braid that is non-right-veering and non-left-veering. Then $\text{rank}(\text{AKh}(b(\beta); \mathbb{C})) \geq 10$.*

Proof. Since V is a 3-braid there is a V_3^0 summand. Since β is non-right-veering there is a V_1^{-1} summand by Lemma 3.2. Since β is non-left-veering there is a V_1^1 summand by Lemma 3.2. We claim that $\text{rank}(\text{AKh}(b(\beta); \mathbb{C})) > 8$. Otherwise, with $\text{rank}(\text{AKh}(b(\beta); \mathbb{C})) \leq 8$, we would in fact have $\text{AKh}(b(\beta); \mathbb{C}) \cong V_1^{-1} \oplus V_3^0 \oplus V_1^1$. Then Lemma 3.2 would imply that all generators die under the spectral sequence from $\text{AKh}(b(\beta); \mathbb{C})$ to $\text{Kh}(b(\beta); \mathbb{C})$; this is not possible, proving the claim. The result now follows from the fact that $\text{rank}(\text{AKh}(b(\beta); \mathbb{C}))$ is even for 3-braids — which have wrapping number 3 — since it splits as a direct sum of V_3 and V_1 summands. \square

It is unclear to the author if similar results could be obtained for clasp-closures, since clasp-closures of conjugate braids are not necessarily isotopic, so the proof strategy above breaks down.

3C. Applications of the rank bound. Let β_n denote the n -braid $\sigma_1\sigma_2\dots\sigma_{n-1}$, and $\mathbb{1}_n$ denote the identity n -braid.

Proposition 3.8. *Suppose α is an n -braid, with $n > 2$. If $\text{rank}(\text{AKh}(b(\alpha); \mathbb{C})) = 2n$ then $\text{AKh}(b(\alpha); \mathbb{C}) \cong \text{AKh}(b(\beta_n^{\pm 1}); \mathbb{C})$.*

The $n = 1$ case is uninteresting, as is the $n = 2$ case since the annular Khovanov homology of all 2-braids is known [20]. Indeed, in the 2-braid case, the proposition is false; $\text{rank}(\text{AKh}(b(\mathbb{1}_2); \mathbb{Z})) = \text{rank}(\text{AKh}(b(\beta_2^{\pm 1}); \mathbb{Z})) = 4$ [20].

Proof. Suppose α is neither σ -positive nor σ -negative. Then α is the identity braid, and one can readily check that $\text{rank}(\text{AKh}(b(\alpha); \mathbb{C})) = 2^n$. It follows that $n = 1$ or $n = 2$, contradicting our assumption.

Suppose now that α is σ -negative. Then $\text{AKh}(b(\alpha); \mathbb{C})$ contains a summand of the form $V_n^0\{n + sl(\beta)\} \oplus V_{n-2}^{-1}\{sl(\beta) - 2\}$ by Lemma 3.2, so must in fact be $V_n^0\{sl(\beta)\} \oplus V_{n-2}^{-1}\{sl(\beta) - 2\}$. The σ -positive case follows by a similar argument. \square

Since annular Khovanov homology detects $\beta_1, \beta_2, \beta_3, \beta_4, \beta_5, \beta_6, \beta_8, \beta_{10}$ by [12; 13; 18], it follows from Proposition 3.8 that $\text{rank}(\text{AKh}(-; \mathbb{Z}/2))$ detects each of these braids amongst braids of the correct index. That is, we have:

Corollary 3.9. *Suppose α is an n -braid with $n \in \{3, 4, 5, 6, 8, 10\}$. If the rank of $\text{AKh}(b(\alpha); \mathbb{Z}/2)$ equals $2n$ then $\alpha = \beta_n^{\pm 1}$.*

This allows one to cut some casework from Baldwin, Hu and Sivek's proof that Khovanov homology detects $T(2, 5)$ with $\mathbb{Z}/2$ coefficients [8, Sections 4 and 5]. In fact, we can generalize part of their argument as follows:

Proposition 3.10. *Suppose K is an m -periodic link with axis of symmetry A , $\text{rank}(\text{Kh}(K; \mathbb{Z}/m)) \leq 2n$ and $\text{lk}(A, K) \geq n$. Let J denote the quotient of K viewed as an annular link about A . Then $\text{AKh}(J; \mathbb{C}) \cong \text{AKh}(b(\beta_n^{\pm 1}); \mathbb{C})$. Moreover, if $n \in \{3, 4, 5, 6, 8, 10\}$, then $J = b(\beta_n^{\pm 1})$.*

Proof. We follow closely the argument in [8]. Let J denote the quotient of K , viewed as an annular link about A . A result of Stoffregen and Zhang [41] implies that

$$\text{rank}(\text{AKh}(J; \mathbb{Z}/m)) \leq \text{rank}(\text{Kh}(K; \mathbb{Z}/m)).$$

Thus $\text{rank}(\text{AKh}(J; \mathbb{C})) \leq \text{rank}(\text{AKh}(J; \mathbb{Z}/m)) \leq 2n$, with the first inequality coming from the universal coefficient theorem. Xie defined a spectral sequence from annular Khovanov homology to an invariant called *annular instanton Floer homology* which respects the annular grading [43]. Xie and Zhang showed in [45, Theorem 1.6] that the maximum nontrivial annular grading of the annular instanton Floer homology of J is given by

$$\min\{2g(S) + |S \cap J| : S \text{ is a meridional surface}\}.$$

Here, a meridional surface in the thickened annulus is any surface which meets the boundary of the thickened annulus in a meridian, i.e., a curve that bounds a disk. Note that $\min\{2g(S) + |S \cap J|\} \geq \ell k(A, J) \geq n$. It follows that

$$\text{rank}(\text{AKh}(J; \mathbb{C}, k = n)) > 0.$$

Therefore, J is a braid by [19, Theorem 1.1], since $\text{AKh}(J; \mathbb{C})$ contains a copy of V_{n+1} and so cannot have rank greater than one in the maximum annular grading. The result then follows directly from Proposition 3.8 in general and Corollary 3.9 in the special cases. \square

3D. Braid-closures. In this section we prove the following result:

Theorem 3.11. *Annular Khovanov homology with integer coefficients detects $b(\sigma_1\sigma_2^n)$ for $-2 \leq n \leq 5$.*

We remind the reader that the $n = 1$ case was already proven in [13], so we do not discuss it here. We begin with some computations.

We compute the annular Khovanov homology of the annular links $b(\sigma_1\sigma_2^n)$. It is readily checked that

$$(10) \quad \text{AKh}(b(\sigma_1); \mathbb{C}) \cong V_3^0\{1\} \oplus V_1^0\{1\} \oplus V_1^1\{3\}$$

and that $\text{AKh}(b(\sigma_1); \mathbb{Z}/2)$ can be obtained by replacing each homogeneous \mathbb{C} summand in $\text{AKh}(b(\sigma_1); \mathbb{C})$ with a $\mathbb{Z}/2$ summand. We compute the annular Khovanov homology of the remaining links.

Lemma 3.12. *For $n \geq 1$, $\text{AKh}(b(\sigma_1\sigma_2^n); \mathbb{C})$ is given by*

$$V_3^0\{n+1\} \oplus V_1^1\{n+3\} \oplus \bigoplus_{1 \leq i \leq n-1} V_1^{1+i}\{n+1+2i\}.$$

For $n \leq -1$, $\text{AKh}(b(\sigma_1\sigma_2^n); \mathbb{C})$ is given by

$$V_3^0\{1+n\} \oplus V_1^1\{3+n\} \oplus V_1^0\{1+n\} \oplus \bigoplus_{-1 \geq i \geq n} V_1^i\{n+1+2i\}.$$

In each case $\text{AKh}(b(\sigma_1\sigma_2^n); \mathbb{Z}/2)$ is given by replacing each homogeneous \mathbb{C} -summand with a $\mathbb{Z}/2$ -summand.

Proof. Consider the standard diagram for $b(\sigma_1\sigma_2^n)$, as in Figure 1(a). We consider two cases; that in which $n \geq 1$ and that in which $n \leq -1$. We proceed by induction in both instances.

In the $n \geq 1$ case, note that $\text{AKh}(b(\sigma_1\sigma_2); \mathbb{C}) \cong V_3^0\{2\} \oplus V_1^1\{4\}$.

For the inductive step we resolve $b(\sigma_1\sigma_2^n)$ at one of the crossings corresponding to a σ_2 . Observe that the 1-resolution is the braid-closure of the identity 1-braid,

while the 0-resolution is $b(\sigma_1\sigma_2^{n-1})$. Applying the exact triangle (7) and the fact that $n_- = 0$ and $n_-^1 = n - 1$, we obtain

$$\begin{array}{ccc} \mathrm{AKh}(b(\sigma_1\sigma_2^n); \mathbb{C}) & \xrightarrow{\quad\quad\quad} & \mathrm{AKh}(b(\sigma_1\sigma_2^{n-1}); \mathbb{C})\{1\} \\ & \swarrow & \nwarrow \delta \\ & \mathrm{AKh}(b(\mathbf{1}_1); \mathbb{C})[n]\{3n-1\} & \end{array}$$

Now,

$$(11) \quad \mathrm{AKh}(\mathbf{1}_1; \mathbb{C}) \cong V_1^0.$$

For $n > 2$ this map splits by the inductive hypothesis, since there are no generators in $\mathrm{AKh}(b(\sigma_1\sigma_2^{n-1}); \mathbb{C})$ of the correct gradings to map nontrivially to $V_1^0[n]\{3n-1\}$. For $n = 2$ the result can be computed by hand or one can note that the connecting map δ , which increases the i grading by 1, must vanish as $\mathrm{Kh}(b(\sigma_1\sigma_2^2); \mathbb{C}) \cong \mathrm{Kh}(T(2, 2)); \mathbb{C})$ has two generators with i grading 2.

We now proceed to the $n \leq -1$ case. Note that

$$\mathrm{AKh}(b(\sigma_1\sigma_2^{-1}); \mathbb{C}) \cong V_3^0 \oplus V_1^1\{2\} \oplus V_1^0 \oplus V_1^{-1}\{-2\}.$$

For the inductive step we resolve $b(\sigma_1\sigma_2^n)$ at one of the crossings corresponding to a σ_2^{-1} . Applying the exact triangle (6) and noting that $n_- = -n$ and $n_-^0 = 0$ we obtain

$$\begin{array}{ccc} \mathrm{AKh}(b(\sigma_1\sigma_2^n); \mathbb{C}) & \xrightarrow{\quad\quad\quad} & \mathrm{AKh}(b(\mathbf{1}_1); \mathbb{C})[n]\{1+3n\} \\ & \swarrow & \nwarrow \\ & \mathrm{AKh}(b(\sigma_1\sigma_2^{n+1}); \mathbb{C})\{-1\} & \end{array}$$

Given equation (11) and the inductive hypothesis, the grading data implies that the exact triangle must split. The result follows.

Finally, to see that $\mathrm{AKh}(b(\sigma_1\sigma_2^n); \mathbb{Z}/2)$ is as claimed, observe that the proofs above from the case of complex coefficients carry through to the case of $\mathbb{Z}/2$ coefficients verbatim. \square

For $\mathbb{F} \in \{\mathbb{Z}/2, \mathbb{C}\}$, $\mathrm{AKh}(b(\sigma_1^{-1}\sigma_2^n); \mathbb{F})$ can be determined from Lemma 3.12 using symmetry properties of annular Khovanov homology. Since, by the Birman–Menasco classification, the 3-braid representatives of the link $T(2, n)$ with $n \neq 0$ are exactly links of the form $b(\sigma_1^{-1}\sigma_2^n)$ and $b(\sigma_1\sigma_2^n)$, this means we have computed the annular Khovanov homology of all 3-braid representatives of $T(2, n)$.

Proof of Theorem 3.11. The strategy is to use the spectral sequence from the annular Khovanov homology of an annular link to Khovanov homology of the underlying link to determine the underlying link type then to exploit the Birman–Menasco classification [15].

Suppose L is an annular link with $\text{AKh}(L; \mathbb{Z}) \cong \text{AKh}(b(\sigma_1\sigma_2^n); \mathbb{Z})$ for some n . Note that $\text{AKh}(L; R) \cong \text{AKh}(b(\sigma_1\sigma_2^n); R)$ for $R \in \{\mathbb{Q}, \mathbb{C}, \mathbb{Z}/2\}$ by the universal coefficient theorem. Since L has rank one in the maximum nontrivial k grading it follows that L is isotopic to the closure of a braid β [19]. Since the maximum nontrivial k grading is 3 it follows that β has index 3. We now split our analysis into three cases; $n = -1$, $n = -2$ and $n \geq 0$.

Case $n = -1$: We claim that $b(\beta)$ is an unknot. First, since $\text{AKh}(b(\beta); \mathbb{C})$ has support in odd quantum gradings, $b(\beta)$ has an odd number of components. Consider the spectral sequence from $\text{AKh}(b(\beta); \mathbb{Z}/2)$ to $\text{Kh}(b(\beta); \mathbb{Z}/2)$. Suppose L is neither σ -positive nor σ -negative. Then L is the identity 3-braid. This is a contradiction, since the identity 3-braid has annular Khovanov homology of rank 8. It follows that L is either σ -positive or σ -negative. Thus $\text{rank}(\text{Kh}(b(\beta); \mathbb{Z}/2)) \leq 6$ by Lemma 3.5. It follows that L has at most two components, since $\text{rank}(\text{Kh}(b(\beta); \mathbb{Z}/2)) \geq 2^m$ where m is the number of components of L . Thus, $\text{rank}(\text{Khr}(b(\beta); \mathbb{Z}/2)) \leq 3$ by [40, Corollary 3.2.C]. Thus L is either a trefoil or an unknot [3; 29]. But L cannot be a trefoil, since $\text{Kh}(T(2, \pm 3); \mathbb{Q})$ has support in quantum gradings ± 9 . It follows that L is an unknot. Since there are only three 3-braids representing the unknot up to conjugation by Murasugi's classification [33], it suffices to show that β is not $\sigma_1\sigma_2$ or $\sigma_1^{-1}\sigma_2^{-1}$. But these two braids have braid-closures with annular Khovanov homology of rank 6 over \mathbb{C} , rather than 10, completing the proof in this case.

Case $n = -2$: First, since $\text{AKh}(b(\beta); \mathbb{C})$ has support in even quantum gradings $b(\beta)$ has an even number of components. Since β is a 3-braid it follows that β has two components. Observe that $\text{rank}(\text{Kh}(b(\beta); \mathbb{Z}/2)) \leq 8$ by Lemma 3.5, so by [44, Corollary 1.4] $b(\beta)$ represents a two-component unlink, $T(2, \pm 2)$ or $T(2, \pm 4)$. By Birman–Menasco, $c(\beta)$ must be of the form $\sigma_1^{\pm 1}\sigma_2^n$ for some even $|n| \leq 4$. Annular Khovanov homology distinguishes each of these links, concluding the proof in this case.

Case $n \neq 0$: Observe that β cannot be the identity braid, since its annular Khovanov homology is not of the correct form. Moreover, β cannot be σ -negative as there are no generators of $\text{AKh}(L; \mathbb{C})$ in homological grading -1 . It follows that β is σ -positive. An application of Lemmas 3.5 and 3.12 implies that for $n \geq 1$

$$(12) \quad \text{rank}(\text{Kh}(b(\beta); \mathbb{Z}/2)) \leq \text{rank}(\text{AKh}(b(\beta); \mathbb{Z}/2)) - 4 = 2n.$$

On the other hand, for $n = 0$, equation (10) and Lemma 3.5 imply that

$$\text{rank}(\text{Kh}(b(\beta); \mathbb{Z}/2)) \leq \text{rank}(\text{AKh}(b(\beta); \mathbb{Z}/2)) - 4 = 4.$$

Thus $\text{rank}(\text{Khr}(b(\beta); \mathbb{Z}/2)) \leq \max\{n, 2\}$ by [40, Corollary 3.2.C]. We now treat three subcases:

$n = 3$. In this case, $\text{rank}(\text{Khr}(b(\beta); \mathbb{Z}/2)) \leq 3$. Since $\text{AKh}(b(\beta); \mathbb{Q})$ is supported in odd quantum gradings it has an odd number of components. Note that $b(\beta)$ can have no more than two components, since $\text{rank}(\text{Kh}(b(\beta); \mathbb{Q})) \geq 2^m$, where m is the number of components of $c(\beta)$. It follows that $b(\beta)$ is a knot. Now, $\text{rank}(\text{Khr}(b(\beta); \mathbb{Z}/2))$ is odd, so that $\text{rank}(\text{Khr}(b(\beta); \mathbb{Z}/2)) = 1$ or $\text{rank}(\text{Khr}(b(\beta); \mathbb{Z}/2)) = 3$. If $\text{rank}(\text{Khr}(b(\beta); \mathbb{Z}/2)) = 1$ then $b(\beta)$ represents the unknot by [29]. But the three braid-closures of 3-braids representing the unknot have different annular Khovanov homology from $\text{AKh}(b(\beta); \mathbb{Z})$, so $n \neq 1$. It follows that $n = 3$ and $b(\beta)$ represents a trefoil by [3]. There are four 3-braids representing trefoils by [15]. They each have distinct annular Khovanov homology by Lemma 3.12, so the result follows.

$n = 5$. Since $\text{AKh}(b(\beta); \mathbb{C})$ is supported in odd quantum gradings, $b(\beta)$ has an odd number of components. Since β is a 3-braid, $b(\beta)$ has either one or three components. If β has three components, then since $\text{rank}(\text{Kh}(b(\beta); \mathbb{Z}/2)) \leq 10$ by Table (12), the Batson–Seed link splitting spectral sequence implies that each component of $b(\beta)$ is unknotted. More specifically, for any component K of $b(\beta)$, we have $2 \text{rank}(\text{Khr}(K; \mathbb{Z}/2)) \leq \text{rank}(\text{Kh}(K; \mathbb{Z}/2)) \leq 2$ by the universal coefficient theorem and [40, Corollary 3.2.C]. Consequently, $\text{rank}(\text{Khr}(K; \mathbb{Z}/2)) = 1$ and K is the unknot by [29]. Birman and Menasco’s classification theorem [15] implies that the only 3-braid representative of the three-component unlink is the identity 3-braid. However, the identity 3-braid has distinct annular Khovanov homology from $\text{AKh}(b(\beta); \mathbb{C})$, so that L in fact does not have 3 components. It follows that $b(\beta)$ is a knot. Now, if $\text{rank}(\text{Khr}(b(\beta); \mathbb{Z}/2)) \leq 3$ we can proceed as in the $n = 3$ case and deduce that $b(\beta)$ represents a trefoil or the unknot. This is a contradiction, since the annular Khovanov homology of the corresponding braid-closures are not of the correct form. It follows that $\text{rank}(\text{Khr}(b(\beta); \mathbb{Z}/2)) = 5$. In turn it follows from [40, Corollary 3.2.C] that $\text{rank}(\text{Kh}(b(\beta); \mathbb{Z}/2)) = 10$. Consider the spectral sequence from $\text{AKh}(b(\beta); \mathbb{Z}/2)$ to $\text{Kh}(b(\beta); \mathbb{Z}/2)$. The proof of Lemma 3.5, together with the symmetry properties of the spectral sequences under mirroring, implies that the spectral sequence kills the generators in (i, j, k) gradings given by

$$(0, 9, 3), (1, 9, 1), (0, 7, 1), (1, 7, -1).$$

Since the E_∞ -page $\text{Kh}(b(\beta); \mathbb{Z}/2)$ must have rank 10, the remaining ten generators must survive. By examining their homological and quantum gradings we find that $\text{Kh}(b(\beta); \mathbb{Z}/2) \cong \text{Kh}(T(2, 5); \mathbb{Z}/2)$. Thus $b(\beta)$ is $T(2, 5)$ by [8, Theorem 1.1]. Birman and Menasco’s classification implies that $\beta = \sigma^{\pm 1} \sigma_2^5$ up to conjugation. But $b(\sigma^{-1} \sigma_2^5)$ has the wrong annular Khovanov homology, so the result follows.

$n \in \{0, 2, 4\}$. In this case $b(\beta)$ has an even number of components, because $\text{AKh}(b(\beta); \mathbb{C})$ is supported in even quantum gradings. Since $b(\beta)$ is a 3-braid it has

exactly two components. Now, $\text{rank}(\text{Kh}(b(\beta); \mathbb{Z}/2)) \leq 8$, so by [44, Corollary 1.4] $b(\beta)$ is either a two-component unlink, $T(2, \pm 2)$ or $T(2, \pm 4)$. By Birman and Menasco’s classification result [15], $c(\beta)$ must be of the form $\sigma_1^{\pm 1} \sigma_2^n$ for some even $|n| \leq 4$. Annular Khovanov homology distinguishes these links. \square

3E. Clasp-closures. We now study the annular Khovanov homology of clasp-closures of 3-braids. The results are dependent on the rank bound from Section 3B. We first state our main result, followed by a necessary lemma.

Theorem 3.13. *Annular Khovanov homology with integer coefficients detects the Mazur pattern.*

Lemma 3.14. *Suppose K is a clasp-closure of a 3-braid. If K' is an annular knot with $\text{AKh}(K; \mathbb{C}) \cong \text{AKh}(K'; \mathbb{C})$ then K' is also a clasp-closure of a 3-braid.*

Proof. Suppose K' is as in the statement of the lemma. Consider Xie’s spectral sequence from $\text{AKh}(K'; \mathbb{C})$ to $\text{AHI}(K'; \mathbb{C})$ [43]. The maximum nontrivial annular grading of $\text{AHI}(K'; \mathbb{C})$ is either 3 or 1. By [45, Theorem 1.6], if it is 1 then there is a meridional surface of Euler characteristic zero, i.e., K' is an annular link of wrapping number 1. Such links have annular Khovanov homology with maximal nontrivial annular grading one — which can be seen by viewing K' as a connect sum of the 1-braid and a wrapping number zero link and applying the Künneth formula for annular Khovanov homology — a contradiction.

It follows that $\text{AHI}(K'; \mathbb{C})$ is of rank 2 in annular grading 3, the maximum annular grading in which $\text{AHI}(K'; \mathbb{C})$ is nontrivial. It follows from [11, Proposition 8.6] that K' is a clasp-braid-closure of index 3. \square

To prove Theorem 3.13 it remains to show that annular Khovanov homology distinguishes the Mazur pattern from the other clasp-closures of 3-braids representing unknots. To that end, we give a partial computation for the annular Khovanov homology of the three types of clasp-closures representing unknots.

First, we consider the mirror of the Mazur pattern, $c(\sigma_1^{-1})$.

Lemma 3.15. *$\text{AKh}(c(\sigma^{-1}); \mathbb{C})$ is given by*

$k \rightarrow$ $i \downarrow$	-3	-1	1	3
0	\mathbb{C}_{-5}	\mathbb{C}_{-3}^2	\mathbb{C}_{-1}^2	\mathbb{C}_1
-1	\mathbb{C}_{-7}	\mathbb{C}_{-5}^2	\mathbb{C}_{-3}^2	\mathbb{C}_{-1}
-2		\mathbb{C}_{-7}	\mathbb{C}_{-5}	

$\text{AKh}(c(\sigma^{-1}); \mathbb{Z}/2)$ can be obtained by replacing every homogeneous \mathbb{C} -summand and replacing it with a $\mathbb{Z}/2$ -summand.

Proof. Consider the 0- and 1-resolutions of the crossing at the top of the diagram shown in Figure 1(c). The 0-resolution yields the $b(\sigma_1^{-1}\sigma_2^{-1})$ while the 1-resolution is $b(\sigma_1^{-1})$. Recall that $\text{AKh}(b(\sigma_1^{-1}\sigma_2^{-1}); \mathbb{C})$ is given by

$$(13) \quad \begin{array}{c|cccc} \begin{array}{c} k \rightarrow \\ i \downarrow \end{array} & -3 & -1 & 1 & 3 \\ \hline 0 & \mathbb{C}_{-5} & \mathbb{C}_{-3} & \mathbb{C}_{-1} & \mathbb{C}_1 \\ -1 & & \mathbb{C}_{-5} & \mathbb{C}_{-3} & \end{array}$$

This can be computed by hand. On the other hand, $\text{AKh}(b(\sigma_1^{-1}); \mathbb{C})$ is given by

$$(14) \quad \begin{array}{c|cccc} \begin{array}{c} k \rightarrow \\ i \downarrow \end{array} & -3 & -1 & 1 & 3 \\ \hline 0 & \mathbb{C}_{-4} & \mathbb{C}_{-2}^2 & \mathbb{C}_0^2 & \mathbb{C}_2 \\ -1 & & \mathbb{C}_{-4} & \mathbb{C}_{-2} & \end{array}$$

Now observe that $n^- = 3$, $n_0^- = 2$, and the exact triangle (6) reduces to

$$\begin{array}{ccc} \text{AKh}(c(\sigma^{-1}); \mathbb{C}) & \xrightarrow{\quad\quad\quad} & \text{AKh}(b(\sigma_1^{-1}\sigma_2^{-1}); \mathbb{C})[-1]\{-2\} \\ & \swarrow & \nwarrow \delta \\ & \text{AKh}(b(\sigma_1^{-1}); \mathbb{C})\{-1\} & \end{array}$$

Comparing the gradings listed in (13) and (14) and noting that the connecting homomorphism δ preserves the quantum grading, we see that the lower right-hand map in this triangle vanishes, yielding the desired result for $\text{AKh}(c(\sigma_1^{-1}); \mathbb{C})$. The computation for $\text{AKh}(c(\sigma_1^{-1}); \mathbb{Z}/2)$ is identical. \square

We now give a partial computation for $\text{AKh}(b(\sigma_1^{-3}\sigma_2\sigma_1^{-2}); \mathbb{C})$.

Lemma 3.16. $b(\sigma_1^{-3}\sigma_2\sigma_1^{-2})$ has annular Jones polynomial given by

$$t^{-3}(-q + q^{-1}) + t^{-1}(-q^3 + 2q) + t(-q^5 + 2q^3) + t^3(-q^7 + q^5).$$

Moreover, in annular grading 3 the annular Khovanov homology has rank two and is supported in (i, j) gradings $(3, 7)$ and $(2, 5)$.

Proof. Consider the 0 and 1-resolutions of the crossing at the top of the diagram shown in Figure 1(c) taking $\alpha = \sigma_1^{-3}\sigma_2\sigma_1^{-2}$. The 0-resolution yields the braid $b'(\sigma_1^{-3}\sigma_2\sigma_1^{-2}\sigma_2^{-1})$ but with the orientation of the component which is not a braid-closure of the 1-braid endowed with the opposite orientation — which we have indicated with the $'$. The 1-resolution is $b(\sigma_1^{-3}\sigma_2\sigma_1^{-2})$.

The annular Khovanov homology of the two braids can be computed using Hunt, Keese, Licata and Morrison's program [22]. In particular we find that

$\text{AKh}(b(\sigma_1^{-3}\sigma_2\sigma_1^{-2}\sigma_2^{-1}); \mathbb{C})$ is given by

$k \rightarrow$ $i \downarrow$	-3	-1	1	3
0	\mathbb{C}_{-8}	\mathbb{C}_{-6}	\mathbb{C}_{-4}	\mathbb{C}_{-2}
-1		\mathbb{C}_{-8}	\mathbb{C}_{-6}	
-2		\mathbb{C}_{-8}^2	\mathbb{C}_{-6}^2	
-3		\mathbb{C}_{-10}	\mathbb{C}_{-8}	
-4		\mathbb{C}_{-12}	\mathbb{C}_{-10}	
-5		\mathbb{C}_{-14}	\mathbb{C}_{-12}	

To correct for the fact that one of the components is given the nonbraid orientation we have to shift the homological grading by [2] and the quantum grading by {6}. This holds because (annular) Khovanov homology is defined by applying a cube of resolution procedure to an unoriented (annular) link diagram, and then shifting the (diagram dependent) homological and quantum gradings by quantities determined by the orientation — namely the quantum grading by $\{n_+ - 2n_-\}$ and the homological grading by $-n_-$, where n_{\pm} represents the number of positive and negative intersections — to obtain the diagram independent homological and quantum gradings. It is then straightforward to check that for the diagram at hand, reversing the orientation of the relevant component induces the shifts in homological and quantum grading as claimed above.

On the other hand, $\text{AKh}(b(\sigma_1^{-3}\sigma_2\sigma_1^{-2}); \mathbb{C})$ is given by

$k \rightarrow$ $i \downarrow$	-3	-1	1	3
1		\mathbb{C}_{-5}	\mathbb{C}_{-3}	
0	\mathbb{C}_{-7}	\mathbb{C}_{-5}^2	\mathbb{C}_{-3}^2	\mathbb{C}_{-1}
-1		\mathbb{C}_{-7}	\mathbb{C}_{-5}	
-2		\mathbb{C}_{-9}	\mathbb{C}_{-7}	
-3		\mathbb{C}_{-11}	\mathbb{C}_{-9}	
-4		\mathbb{C}_{-13}	\mathbb{C}_{-11}	
-5		\mathbb{C}_{-15}	\mathbb{C}_{-13}	

Since $n_- = 4$ and $n_-^1 = 6$, we have the exact triangle

$$\begin{array}{ccc}
 \text{AKh}(L; \mathbb{C}) & \xrightarrow{\hspace{10em}} & \text{AKh}(b'(\sigma_1^{-3}\sigma_2\sigma_1^{-2}\sigma_2^{-1}); \mathbb{C})\{1\} \\
 & \swarrow \hspace{2em} \searrow & \\
 & \text{AKh}(b(\sigma_1^{-3}\sigma_2\sigma_1^{-2}); \mathbb{C})[3]\{8\} &
 \end{array}$$

This isn't enough information to show that the exact triangle splits. However, it

does split in annular gradings ± 3 , and the decategorification of the exact triangle determines the annular Jones polynomial, as desired. \square

Let S be the annular link given by the split sum of $b(\mathbf{1}_1)$ and an unknot. Observe that the annular Jones polynomial of S is given by $J(S) = t^{-1}(1+q^{-2}) + t(1+q^2)$.

Lemma 3.17. *The annular Jones polynomial of $c(\sigma_1^n \sigma_2^{-1} \sigma_1 \sigma_2)$ is given by*

$$J(c(\sigma_1^n \sigma_2^{-1} \sigma_1 \sigma_2)) = \frac{q + (-1)^{n+1} q^{1-2n}}{1+q^2} J(S) + (-1)^n q^{-2n} J(c(\sigma_2^{-1} \sigma_1 \sigma_2)).$$

Moreover, in annular grading 3 the annular Khovanov homology has rank two and is supported in (i, j) gradings $(-n, 3-2n)$ and $(-1-n, 1-2n)$.

Proof. We first compute the annular Khovanov homology of $c(\sigma_1)$, which is isotopic to $c(\sigma_2^{-1} \sigma_1 \sigma_2)$.

Consider the 0 and 1-resolutions of the crossing at the top of the diagram shown in Figure 1(c) taking $\alpha = \sigma_1$. The 0-resolution is $b(\sigma_1 \sigma_2^{-1})$. The 1-resolution is $b(\sigma_1)$. Since $n_-^0 = 1$ and $n_- = 2$, we have

$$\begin{array}{ccc} \mathrm{AKh}(c(\sigma_1); \mathbb{C}) & \xrightarrow{\quad\quad\quad} & \mathrm{AKh}(b(\sigma_1 \sigma_2^{-1}); \mathbb{C})[-1]\{-2\} \\ & \swarrow & \nwarrow \delta \\ & \mathrm{AKh}(b(\sigma_1); \mathbb{C})\{-1\} & \end{array}$$

$\mathrm{AKh}(b(\sigma_1 \sigma_2^{-1}); \mathbb{C})[-1]\{-2\}$ and $\mathrm{AKh}(b(\sigma_1); \mathbb{C})\{-1\}$ are given respectively by

$\begin{array}{c} k \rightarrow \\ i \downarrow \end{array}$	-3	-1	1	3
0		\mathbb{C}_{-1}	\mathbb{C}_1	
-1	\mathbb{C}_{-5}	\mathbb{C}_{-3}^2	\mathbb{C}_{-1}^2	\mathbb{C}_1
-2		\mathbb{C}_{-5}	\mathbb{C}_{-3}	

and

$\begin{array}{c} k \rightarrow \\ i \downarrow \end{array}$	-3	-1	1	3
1		\mathbb{C}_1	\mathbb{C}_3	
0	\mathbb{C}_{-3}	\mathbb{C}_{-1}^2	\mathbb{C}_1^2	\mathbb{C}_3

Thus, since δ preserves the quantum grading and increases the homological grading by one, the exact triangle splits and $\mathrm{AKh}(c(\sigma_1); \mathbb{C})$ is given by

$\begin{array}{c} k \rightarrow \\ i \downarrow \end{array}$	-3	-1	1	3
1		\mathbb{C}_1	\mathbb{C}_3	
0	\mathbb{C}_{-3}	\mathbb{C}_{-1}^3	\mathbb{C}_1^3	\mathbb{C}_3
-1	\mathbb{C}_{-5}	\mathbb{C}_{-3}^2	\mathbb{C}_{-1}^2	\mathbb{C}_1
-2		\mathbb{C}_{-5}	\mathbb{C}_{-3}	

We now proceed to the general case. Remove the axis from the diagram shown in Figure 3 to obtain a diagram for the link. Observe that the 1-resolution of the

highlighted crossing is the link S , which has annular Khovanov homology given by

$k \rightarrow$ $i \downarrow$	-1	1
0	$\mathbb{C}_{-2} \oplus \mathbb{C}_0$	$\mathbb{C}_0 \oplus \mathbb{C}_2$

Now, since we can take $n_-^0 - n_- = -1$, the exact triangle (6) reduces to

$$\begin{array}{ccc}
 \text{AKh}(\sigma_1^n \sigma_2^{-1} \sigma_1 \sigma_2; \mathbb{C}) & \xrightarrow{\quad\quad\quad} & \text{AKh}(\sigma_1^{n-1} \sigma_2^{-1} \sigma_1 \sigma_2; \mathbb{C})[-1]\{-2\} \\
 & \swarrow & \nwarrow \\
 & \text{AKh}(S; \mathbb{C})\{-1\} &
 \end{array}$$

Since $\text{AKh}(S; \mathbb{C})$ is trivial in annular grading 3 this proves the second part of the result. For the first part, observe that decategorifying either of the above exact triangles we obtain

$$J(c(\sigma_1^n \sigma_2^{-1} \sigma_1 \sigma_2)) = q^{-1} J(S) - q^{-2} J(c(\sigma_1^{n-1} \sigma_2^{-1} \sigma_1 \sigma_2)).$$

The desired result follows by induction. □

Remark 3.18. One could perhaps give a complete computation of the annular Khovanov homology of the infinite family of clasp-closures using techniques of J. Wang [42]. The annular Jones polynomial was enough for our purposes, however, so we do not pursue this.

Let $r(\beta)$ denote the reverse of the braid word β written in terms of the standard Artin generators.

Lemma 3.19. *Suppose β_1 and β_2 are 3-braids such that $c(\beta_1)$ and $c(\beta_2)$ represent unknots. If $\text{AKh}(c(\beta_1); \mathbb{C}) \cong \text{AKh}(c(\beta_2); \mathbb{C})$ then $c(\beta_1) = c(\beta_2)$ or $c(\beta_1) = c(r(\beta_2))$.*

Proof. Lemmas 3.15, 3.16, and 3.17 determine the annular Khovanov homology of all of the clasp-closures up to mirroring. The annular Khovanov homology of their mirrors can be determined using formal properties of annular Khovanov homology. We can then see that no two clasp-closures of 3-braids representing unknots have the same annular Khovanov homology in annular grading 3 so the result follows. □

Proof of Theorem 3.13. Suppose that K is an annular link with $\text{AKh}(K; \mathbb{Z}) \cong \text{AKh}(c(\sigma^{-1}); \mathbb{Z})$. Since $\text{AKh}(K; \mathbb{Z})$ is supported in odd quantum gradings it follows that K has an odd number of components. K is a clasp-closure of a 3-braid by Lemma 3.14. Consider the Batson–Seed link splitting sequence for $\text{Kh}(K; \mathbb{C})$. Observe that $\text{rank}(\text{Kh}(K; \mathbb{C})) \geq 2^m$, where m is the number of components of K . Now observe that Lemma 3.2 implies that the rank of ∂_{-2}^* is at least 4 since the two V_{n-2} summands in the statement of the lemma are mapped nontrivially under ∂_{-2}^* ,

which we recall is part of the $\mathfrak{sl}_2(\mathbb{C})$ action on annular Khovanov homology. In turn it follows that $\text{rank}(\text{Kh}(K; \mathbb{C})) \leq 6$. Thus K has a single component.

We now show that K represents the unknot. An application of the universal coefficient theorem shows that $\text{AKh}(K; \mathbb{Z}/2) \cong \text{AKh}(c(\sigma^{-1}); \mathbb{Z}/2)$. Consider the spectral sequence from $\text{AKh}(K; \mathbb{Z}/2)$ to $\text{Kh}(K; \mathbb{Z}/2)$. Lemma 3.5 implies that $\text{rank}(\text{Kh}(K; \mathbb{Z}/2)) \leq 6$. It follows that $\text{rank}(\text{Khr}(K; \mathbb{Q})) \leq \text{rank}(\text{Khr}(K; \mathbb{Z}/2)) \leq 3$, so that L is either a trefoil or an unknot by [29] and [3]. However, K cannot be a trefoil because $\text{AKh}(K; \mathbb{C})$, and hence $\text{Kh}(K; \mathbb{C})$, does not contain a summand in quantum grading ± 9 .

It follows that K is a clasp-closure of one of Baldwin and Sivek’s 3-braid types. By Lemma 3.19, if two such annular links have the same annular Khovanov homology then they differ only up to reversal. But of course, $c(\sigma_1^{-1})$ and $c(r(\sigma_1^{-1}))$ are isotopic, so the result follows. \square

Acknowledgments

The author thanks Gage Martin for helpful conversations, and Subhankar Dey for further helpful conversations and careful feedback on an earlier draft of this paper. He is also grateful for [28] and [31], which he found to be very helpful throughout the course of this project. Finally, he would like to thank the referee for their careful reading of this paper.

References

- [1] M. M. Asaeda, J. H. Przytycki, and A. S. Sikora, “Categorification of the Kauffman bracket skein module of I -bundles over surfaces”, *Algebr. Geom. Topol.* **4** (2004), 1177–1210. MR
- [2] J. A. Baldwin and J. E. Grigsby, “Categorified invariants and the braid group”, *Proc. Amer. Math. Soc.* **143**:7 (2015), 2801–2814. MR
- [3] J. A. Baldwin and S. Sivek, “Khovanov homology detects the trefoils”, *Duke Math. J.* **171**:4 (2022), 885–956. MR
- [4] J. A. Baldwin and S. Sivek, “Floer homology and non-fibered knot detection”, *Forum Math. Pi* **13** (2025), art. id. e1, 65 pp. MR
- [5] J. Baldwin and D. S. Vela-Vick, “A note on the knot Floer homology of fibered knots”, *Algebr. Geom. Topol.* **18**:6 (2018), 3669–3690. MR
- [6] J. A. Baldwin, A. S. Levine, and S. Sarkar, “Khovanov homology and knot Floer homology for pointed links”, *J. Knot Theory Ramifications* **26**:2 (2017), art. id. 1740004, 49 pp. MR
- [7] J. A. Baldwin, S. Sivek, and Y. Xie, “Khovanov homology detects the Hopf links”, *Math. Res. Lett.* **26**:5 (2019), 1281–1290. MR
- [8] J. A. Baldwin, Y. Hu, and S. Sivek, “Khovanov homology and the cinquefoil”, *J. Eur. Math. Soc. (JEMS)* **27**:6 (2025), 2443–2465. MR
- [9] J. Batson and C. Seed, “A link-splitting spectral sequence in Khovanov homology”, *Duke Math. J.* **164**:5 (2015), 801–841. MR
- [10] F. Binns and S. Dey, “Cable links, annuli and sutured Floer homology”, *Math. Res. Lett.* **31**:5 (2024), 1315–1337. MR

- [11] F. Binns and S. Dey, “Floer homology, clasp-braids and detection results”, preprint, 2024. arXiv 2405.11224
- [12] F. Binns and S. Dey, “Rank bounds in link Floer homology and detection results”, *Quantum Topol.* **16**:4 (2025), 743–794. MR
- [13] F. Binns and G. Martin, “Knot Floer homology, link Floer homology and link detection”, *Algebr. Geom. Topol.* **24**:1 (2024), 159–181. MR
- [14] J. S. Birman and T. E. Brendle, “Braids: a survey”, pp. 19–103 in *Handbook of knot theory*, Elsevier, Amsterdam, 2005. MR
- [15] J. S. Birman and W. W. Menasco, “Studying links via closed braids. III. Classifying links which are closed 3-braids”, *Pacific J. Math.* **161**:1 (1993), 25–113. MR
- [16] P. Dehornoy, I. Dynnikov, D. Rolfsen, and B. Wiest, *Ordering braids*, Mathematical Surveys and Monographs **148**, American Mathematical Society, Providence, RI, 2008. MR
- [17] N. Dowlin, “A spectral sequence from Khovanov homology to knot Floer homology”, *J. Amer. Math. Soc.* **37**:4 (2024), 951–1010. MR
- [18] E. Farber, B. Reinoso, and L. Wang, “Fixed-point-free pseudo-Anosov homeomorphisms, knot Floer homology and the cinquefoil”, *Geom. Topol.* **28**:9 (2024), 4337–4381. MR
- [19] J. E. Grigsby and Y. Ni, “Sutured Khovanov homology distinguishes braids from other tangles”, *Math. Res. Lett.* **21**:6 (2014), 1263–1275. MR
- [20] J. E. Grigsby, A. M. Licata, and S. M. Wehrli, “Annular Khovanov homology and knotted Schur–Weyl representations”, *Compos. Math.* **154**:3 (2018), 459–502. MR
- [21] J. Hoste, “The first coefficient of the Conway polynomial”, *Proc. Amer. Math. Soc.* **95**:2 (1985), 299–302. MR
- [22] H. Hunt, H. Keese, A. Licata, and S. Morrison, “Computing annular Khovanov homology”, 2015. arXiv 1505.04484
- [23] A. Juhász, “Holomorphic discs and sutured manifolds”, *Algebr. Geom. Topol.* **6** (2006), 1429–1457. MR
- [24] A. Juhász, “Floer homology and surface decompositions”, *Geom. Topol.* **12**:1 (2008), 299–350. MR
- [25] A. Juhász, “The sutured Floer homology polytope”, *Geom. Topol.* **14**:3 (2010), 1303–1354. MR
- [26] M. Khovanov, “A categorification of the Jones polynomial”, *Duke Math. J.* **101**:3 (2000), 359–426. MR
- [27] “KLO: Knot-Like Objects”, software, version 0.979, available at <https://community.middlebury.edu/~mathanimations/klo>.
- [28] “The Knot Atlas”, online resource, available at https://katlas.org/wiki/Main_Page.
- [29] P. B. Kronheimer and T. S. Mrowka, “Khovanov homology is an unknot-detector”, *Publ. Math. Inst. Hautes Études Sci.* **113** (2011), 97–208. MR
- [30] E. S. Lee, “An endomorphism of the Khovanov invariant”, *Adv. Math.* **197**:2 (2005), 554–586. MR
- [31] C. Livingston and A. H. Moore, “LinkInfo: table of link invariants”, 2024, available at <http://linkinfo.math.indiana.edu>.
- [32] G. Martin, “Khovanov homology detects $T(2, 6)$ ”, *Math. Res. Lett.* **29**:3 (2022), 835–849. MR
- [33] K. Murasugi, *On closed 3-braids*, Memoirs of the American Mathematical Society **151**, American Mathematical Society, Providence, RI, 1974. MR
- [34] Y. Ni, “Exceptional surgeries on hyperbolic fibered knots”, preprint, 2020. arXiv 2007.11774

- [35] P. Ozsváth and Z. Szabó, “Holomorphic disks and knot invariants”, *Adv. Math.* **186**:1 (2004), 58–116. MR
- [36] P. Ozsváth and Z. Szabó, “Holomorphic disks, link invariants and the multi-variable Alexander polynomial”, *Algebr. Geom. Topol.* **8**:2 (2008), 615–692. MR
- [37] P. Ozsváth and Z. Szabó, “Link Floer homology and the Thurston norm”, *J. Amer. Math. Soc.* **21**:3 (2008), 671–709. MR
- [38] O. Plamenevskaya, “Transverse knots and Khovanov homology”, *Math. Res. Lett.* **13**:4 (2006), 571–586. MR
- [39] J. Rasmussen, “Floer homology and knot complements”, preprint, 2003. arXiv math/0306378
- [40] A. N. Shumakovitch, “Torsion of Khovanov homology”, *Fund. Math.* **225**:1 (2014), 343–364. MR
- [41] M. Stoffregen and M. Zhang, “Localization in Khovanov homology”, *Geom. Topol.* **28**:4 (2024), 1501–1585. MR
- [42] J. Wang, “The cosmetic crossing conjecture for split links”, *Geom. Topol.* **26**:7 (2022), 2941–3053. MR
- [43] Y. Xie, “Instantons and annular Khovanov homology”, *Adv. Math.* **388** (2021), art. id. 107864, 51 pp. MR
- [44] Y. Xie and B. Zhang, “On links with Khovanov homology of small ranks”, *Math. Res. Lett.* **29**:4 (2022), 1261–1277. MR
- [45] Y. Xie and B. Zhang, “Instanton Floer homology for sutured manifolds with tangles”, *J. Differential Geom.* **130**:3 (2025), 701–769. MR

Received August 4, 2024. Revised September 11, 2025.

FRASER BINNS
DEPARTMENT OF MATHEMATICS
PRINCETON UNIVERSITY
PRINCETON, NJ
UNITED STATES
fb1673@princeton.edu