

*Pacific  
Journal of  
Mathematics*

**POLYNOMIAL IDENTITIES AND AZUMAYA LOCI  
FOR RATIONAL QUANTUM SPHERES**

ALEXANDRU CHIRVASITU

Volume 342 No. 1

May 2026



## POLYNOMIAL IDENTITIES AND AZUMAYA LOCI FOR RATIONAL QUANTUM SPHERES

ALEXANDRU CHIRVASITU

We prove a number of structure and isomorphism results concerning the noncommutative Natsume–Olsen spheres  $\mathbb{S}_\theta^{2n-1}$  deformed along a skew-symmetric matrix  $\theta \in \mathbb{R}$ . These include (a) the fact that two  $C^*$ -algebras of the form  $\mathbb{S}_\theta^3 \otimes M_n$  are isomorphic precisely in the obvious cases; (b) the fact that  $m$  and  $n$  are recoverable from the isomorphism class of  $C(\mathbb{S}_\theta^{2m-1}) \otimes M_n$ ; (c) the PI character, PI degree and Azumaya loci of  $C(\mathbb{S}_\theta^{2m-1})$  for rational  $\theta$ , along with a realization of their centers as (function algebras of) branched cover of  $\mathbb{S}^{2n-1}$ ; and (d) for rational  $\theta$  again, the topological finite generation of  $C(\mathbb{S}_\theta^{2m-1})$  over their centers, with algebraic finite generation equivalent to being classical (equivalently, Azumaya).

### Introduction

Consider a skew-symmetric real matrix  $\theta \in M_n(\mathbb{R})$ . We will be working extensively with the following noncommutative-geometric constructs.

- The *noncommutative* (or *quantum*) *tori*  $\mathbb{T}_\theta^n$  [57, §1; 32, §12.2; 40, §1.1.5] defined as objects dual to the generator-and-relation  $C^*$ -algebras

$$A_\theta^n = C(\mathbb{T}_\theta^n) := \langle \text{unitaries } u_j, j \in [n] := \{1..n\} \mid u_k u_j = e(\theta_{jk}) u_j u_k \rangle$$

for  $e(-) := \exp(2\pi i -)$ .

- The *noncommutative spheres* [48, Definition 2.1] analogously defined by

$$C(\mathbb{S}_\theta^{2n-1}) := \langle \text{normal } t_j, j \in [n] \mid t_k t_j = e(\theta_{jk}) t_j t_k, \sum t_j^* t_j = 1 \rangle.$$

The former “glue” to produce the latter in ways reminiscent of classical topology (e.g., the *standard genus-1 (Heegaard) splitting*  $\mathbb{S}^3 = (\mathbb{T}^2 \times [0, 1]) \cup_{\mathbb{T}^2} (\mathbb{T}^2 \times [0, 1])$  of [61, Proposition 3.3]): per [48, Theorem 2.5], we have

$$(0-1) \quad C(\mathbb{S}_\theta^{2n-1}) \cong \text{Cont}_\partial(\mathbb{S}_+^{n-1} \rightarrow C(\mathbb{T}_\theta^n)),$$

*MSC2020:* 46L52, 16H05, 46M20, 55R25, 55R37, 46L85, 16R10, 55R40.

*Keywords:* Azumaya algebra, Chern class, PI algebra, bundle, classifying space, noncommutative sphere, noncommutative torus, projectively flat.

where

$$(0-2a) \quad \begin{aligned} \mathbb{S}_+^{n-1} &:= \{(s_1, \dots, s_n) \in \mathbb{S}^{n-1} \subseteq \mathbb{R}^n : s_i \geq 0\}, \\ \mathbb{S}_{+,F \subseteq [n]}^{n-1} &:= \{(s_i) \in \mathbb{S}_+^{n-1} : s_i = 0, \forall i \notin F\}, \end{aligned}$$

and ‘ $\partial$ ’ means

$$(0-2b) \quad f(\mathbb{S}_{+,F}^{n-1}) \subseteq C^*(u_i, i \in F) \subseteq A_\theta^n, \quad \forall F \subseteq [n].$$

Section 1 is concerned with reconstruction/isomorphism problems, i.e., the extent to which initial data (the deformation parameter  $\theta \in M_n(\mathbb{R})$ , or perhaps its size  $n$ ) are determined by the isomorphism or stable isomorphism classes of the quantum-sphere algebras. A paraphrased aggregate of Theorems 1.1 and 1.2 reads as follows.

**Theorem A.** (1) For  $n, n' \in \mathbb{Z}_{>0}$  and  $\theta, \theta' \in \mathbb{R}$  identified with skew-symmetric  $2 \times 2$  matrices we have

$$\begin{aligned} C(\mathbb{S}_\theta^3) \otimes M_n &\cong C(\mathbb{S}_{\theta'}^3) \otimes M_{n'} \iff C(\mathbb{T}_\theta^2) \otimes M_n \cong C(\mathbb{T}_{\theta'}^2) \otimes M_{n'} \\ &\iff n = n' \text{ and } \theta \in \pm\theta' + \mathbb{Z}. \end{aligned}$$

(2) For positive integers  $m, n$  with  $m \geq 2$  and skew-symmetric  $\theta \in M_m(\mathbb{R})$  the isomorphism class of either  $C(\mathbb{T}_\theta^m) \otimes M_n$  or  $C(\mathbb{S}_\theta^{2m-1}) \otimes M_n$  determines  $m$  and  $n$ .

In Section 2 and onward we assume the deformation parameter  $\theta \in M_n(\mathbb{Q})$  rational and examine consequent *polynomial-identity (PI)* phenomena (on which background the main text offers a brief reminder). The focus is on

- *Azumaya algebras* [16, §4.4]: one way to formalize a purely algebraic analogue of the section-space  $\Gamma(\mathcal{E} \otimes \mathcal{E}^*)$  for a vector bundle  $\mathcal{E}$  over a compact Hausdorff space; and
- the extent to which algebras fail to qualify, as measured by the *Azumaya locus* [11, §III.1.7].

Again summarizing for brevity, one rendition of Theorems 2.8 and 2.13 below is this:

**Theorem B.** Let  $\theta \in M_n(\mathbb{Q})$  be a skew-symmetric matrix for  $n \in \mathbb{Z}_{\geq 2}$ .

- (1) The center  $Z_\theta \leq C_\theta := C(\mathbb{S}_\theta^{2n-1})$  is the function algebra of a branched cover of  $\mathbb{S}^{2n-1}$  which is not, in general, a topological manifold.
- (2) The smallest  $n^2$  admitting a  $Z_\theta$ -algebra embedding  $C_\theta \leq M_n(Z_\theta)$  is

$$h_\theta := [(\mathbb{Z}^n + \text{im } \theta) : \mathbb{Z}^n],$$

so in particular  $C_\theta$  is a PI algebra of **PI-degree** (see [29, §1])  $\sqrt{h_\theta}$ .

(3) *The Azumaya locus of  $C_\theta$  consists precisely of those maximal ideals  $p \in \text{Max}(Z_\theta)$  whose restriction to  $\mathbb{S}_+^{n-1}$  along equation (0-1) belongs to some face supported by  $F \subseteq [n]$  with*

*$h_\theta > h_{\theta,F} := h$  attached to the submatrix of  $\theta$  supported on  $F$ -indexed rows/columns.*

In Section 3 we turn to the intimate connection between polynomial identities and the condition that an algebra be finitely generated as a module over its center (see e.g. [54, Chapter 6] and the motivating discussion on [3, p. 532]) in the context of studying the quantum spheres  $\mathbb{S}_\theta^{2n-1}$ . It will turn out that said finite generation (mostly) fails, but its weaker topological analogue (requiring that  $C(\mathbb{S}_\theta^{2n-1})$  contain a dense finitely generated module over its center) always holds; per Theorem 3.3:

**Theorem C.** *For  $n \in \mathbb{Z}_{\geq 2}$  and rational skew-symmetric  $\theta \in M_n(\mathbb{Q})$  the algebra  $C(\mathbb{S}_\theta^{2n-1})$  is*

- *always a topologically finitely generated module over its center  $Z_\theta$ , but*
- *algebraically finitely generated as such precisely when  $\theta$  is integral.*

Theorems B and C both link naturally to the theory of *Banach, Hilbert* and  *$C^*$  bundles* [27, §II.13; 21, pp. 7–9; 20, §1] over compact Hausdorff spaces and satellite topics: the theory of *noncommutative branched covers* initiated in [51] and phrased in the language of *finite-index conditional expectations* [28, Definition 2] is germane to the discussion below, which relies directly or indirectly on material from [7; 13; 51].

### 1. Isomorphisms of rationally deformed quantum 3-spheres

Throughout, unqualified (typically vector or algebra) *bundles* are assumed *locally trivial* [37, Definition 1.1.8]; they are to be distinguished from more general constructs termed (*F*) *Banach bundles* on [21, pp. 7–8], which will also make an appearance in Section 3.

The isomorphism problem for noncommutative 3-spheres is not difficult to resolve, given its 2-torus analogue. The following is very much in the spirit of [55, Theorem 3] for *irrational* tori. The *rational* torus version [33, Theorem 1.1] (recovered also as [56, Theorem 3.12]) typically does not involve matrix tensorands.

**Theorem 1.1.** *Consider  $n, n' \in \mathbb{Z}_{>0}$ ,  $\theta, \theta' \in \mathbb{R}$ , and set*

$$C_{\theta,n} := C(\mathbb{S}_\theta^3) \otimes M_n, \quad A_{\theta,n} := C(\mathbb{T}_\theta^2) \otimes M_n,$$

*and similarly for the primed parameters. The following conditions are equivalent.*

(a) *We have an isomorphism*

$$(1-1) \quad C_{\theta,n} \cong C_{\theta',n'}.$$

(b) *We have an isomorphism*

$$(1-2) \quad A_{\theta,n} \cong A_{\theta',n'}.$$

(c) *The parameters coincide save for trivial modifications, in the sense that*

$$n = n' \quad \text{and} \quad \theta \in \pm\theta' + \mathbb{Z}.$$

It might be worthwhile to first observe separately that the size  $n$  of the tensorand  $M_n$  can be recovered from the isomorphism class of either  $C_{\theta,n}$  or  $A_{\theta,n}$  and in fact, more generally, for noncommutative spheres/tori of arbitrary dimension.

**Theorem 1.2.** *Let  $\theta \in M_m(\mathbb{R})$  be a skew-symmetric matrix,  $m \geq 2$  and  $n \in \mathbb{Z}_{>0}$ .*

(1) *The isomorphism class of the  $C^*$ -algebra  $C(\mathbb{T}_\theta^m) \otimes M_n$  determines  $m$  and  $n$ .*

(2) *The isomorphism class of the  $C^*$ -algebra  $C(\mathbb{S}_\theta^{2m-1}) \otimes M_n$  determines  $m$  and  $n$ .*

*Proof.* We structure the argument conforming to the statement.

Statement (1) is a simple matter of unwinding some of the well-known K-theoretic results on noncommutative tori.

According to [22, Theorem 2.2],  $K_0(C(\mathbb{T}_\theta^m))$  is free abelian of rank  $2^{m-1}$  with the class of the identity as a generator. The usual isomorphism

$$K_0(A) \cong K_0(A \otimes M_n)$$

induced by the upper-corner embedding [67, Lemma 6.2.10] then makes it clear that in

$$K_0(C(\mathbb{T}_\theta^m) \otimes M_n) \cong \mathbb{Z}^{2^{m-1}}$$

the class of the identity is divisible precisely by  $n$ , hence the conclusion.

We turn to (2).

Recovering  $n$ : Equation (0-1) realizes  $C(\mathbb{S}_\theta^{2m-1})$  as a  $C(X)$ -algebra (i.e., a  $C^*$ -algebra receiving a central morphism from  $C(X)$ : see [39, Definition 1.5]) for  $X = \mathbb{S}_+^{m-1}$ . The fibers

$$C(\mathbb{S}_\theta^{2m-1})|_p := C(\mathbb{S}_\theta^{2m-1}) / (C(\mathbb{S}_\theta^{2m-1}) \cdot \{f \in C(X) : f(p) = 0\}), \quad p \in \mathbb{S}_+^{m-1},$$

are noncommutative-torus algebras  $A$  in general, and specifically to  $C(\mathbb{S}^1)$  at the  $m$  vertices of the spherical simplex  $X$ . It follows that  $n$  can be recovered from the abstract  $C^*$ -algebra  $C := C(\mathbb{S}_\theta^{2m-1}) \otimes M_n$  as the smallest dimension of an irreducible representation.

Recovering  $m$ : A small detour will first discern what can be recovered from the center

$$(1-3) \quad Z := Z(C(\mathbb{S}_\theta^{2m-1}) \otimes M_n) = Z(C(\mathbb{S}_\theta^{2m-1})) = Z(C)$$

alone. The isomorphism (0-1) specializes to  $Z \cong \text{Cont}_\partial(\mathbb{S}_+^{m-1} \rightarrow Z(C(\mathbb{T}_\theta^n)))$ , with  $\partial$  indicating the central analogue of (0-2b):

$$(1-4) \quad f(\mathbb{S}_{+,F}^{m-1}) \subseteq Z(C(\mathbb{T}_\theta^n)) \cap C^*(\{u_j \mid j \in F\}).$$

Recall [23, §2.2] that  $C(\mathbb{T}_\theta^n)$  is a cocycle twist  $C^*(\mathbb{Z}^n, \sigma)$  of the group algebra  $C^*(\mathbb{Z}^n)$  (with the generators of  $\mathbb{Z}^n$  mapping to the  $u_i$ ). The proof of [22, Lemma 2.3] then describes the center  $Z(C(\mathbb{T}_\theta^n))$  as

$$(1-5) \quad C^*(\Gamma) \subset C^*(\mathbb{Z}^n),$$

$$\Gamma := \text{kernel of the bicharacter } \mathbb{Z}^n \wedge \mathbb{Z}^n \xrightarrow{e(\theta)=\exp(2\pi i\theta)} \mathbb{S}^1,$$

with the matrix  $\theta$  regarded, as usual, as a bilinear form. The boundary condition (1-4) then reads

$$f(\mathbb{S}_{+,F}^{m-1}) \subseteq C^*(\Gamma_F) \cong C(\mathbb{T}_F), \quad \mathbb{T}_F := \widehat{\Gamma}_F,$$

where

$$(1-6) \quad \Gamma_F := \Gamma \cap \mathbb{Z}^F, \quad \mathbb{Z}^F := \text{sum of the } F\text{-indexed summands in } \mathbb{Z}^n.$$

We write  $\text{Max}(A)$  for the *maximal spectrum* [5, Exercise 1.26] of a commutative ring, typically applying the notion to commutative  $C^*$ -algebras (and occasionally omitting the qualifier “maximal”).  $X := \text{Max}(Z)$  can thus be described as follows:

- To each finite-set inclusion

$$F \subseteq F' \subseteq [m]$$

associate

$$\begin{aligned} &\text{an inclusion } \Delta_F \hookrightarrow \Delta_{F'}, \quad \Delta_F := \mathbb{S}_{+,F}^{m-1}, \\ &\text{and a quotient } \mathbb{T}_F \twoheadrightarrow \mathbb{T}_{F'} \quad \text{dual to } \Gamma_F \leq \Gamma_{F'}. \end{aligned}$$

- And then recover  $X$  as the *coend* [43, §IX.6]

$$(1-7) \quad X = \text{Max}(Z(C(\mathbb{S}_\theta^{2m-1}))) \cong \int^F \Delta_F \times \mathbb{T}_F.$$

This is a particular instance of the *geometric realization* [58, Definition 3.8.1] of a *simplicial object* [43, §VII.5]: more precisely,  $F \mapsto \mathbb{T}_F$  is a simplicial topological space (a simplicial object in the category of topological spaces), and (1-7) is its simplicial realization.

The space (1-7) does have *dimension*

$$(1-8) \quad \dim X = m - 1 + \dim \text{Max}(Z(C(\mathbb{T}_\theta^m))) = m - 1 + \text{rank ker } \theta,$$

where

- “dimension” is used in any of three senses: the *small inductive*, *large inductive* and *covering* dimensions [24, Definitions 1.1.1, 1.6.1, 1.6.7], all coincident for separable metric spaces [24, Theorem 1.7.7];
- and the *rank* of an abelian group is [35, Definition A1.59]

$$\text{rank } \Gamma := \dim_{\mathbb{Q}} \mathbb{Q} \otimes_{\mathbb{Z}} \Gamma.$$

By contrast to  $m$  (Example 1.3), this latter number (1-8) *is*, then, recoverable from the isomorphism class of  $Z = Z(C)$  alone (so  $C$  itself is not needed for this).

The fibers of  $C$  at the various points of  $X$  are of the form  $C^*(\mathbb{Z}^m / \Gamma_F, \sigma) \otimes M_n$  for subsets  $F \subseteq [m]$  with the twist induced by the original cocycle  $\sigma$  on  $\mathbb{Z}^m / \Gamma_F$  (possible, given that  $\Gamma = \Gamma_{[m]}$  is by definition the kernel of  $\theta$ ).

The abelian groups  $\mathbb{Z}^m / \Gamma_F$  have ranks

$$(1-9) \quad m - \text{rank } \Gamma_F \geq m - \text{rank } \Gamma_{[m]} = m - \text{rank } \ker \theta,$$

with equality achieved. Since the sum between (1-8) and this minimum is  $2m - 1$ , it will be enough to observe that the ranks on the left-hand side of (1-9) can be recovered from the corresponding fibers  $C^*(\mathbb{Z}^m / \Gamma_F, \sigma) \otimes M_n$ .

To see this,

- write

$$\mathbb{Z}^m / \Gamma_F \cong (\text{finite abelian group } H) \times \mathbb{Z}^r, \quad r := \text{rank } \mathbb{Z}^m / \Gamma_F$$

[10, p. VII.19, Theorem 2];

- so that  $C^*(\mathbb{Z}^m / \Gamma_F, \sigma)$  can be expressed as an iterated,  $r$ -fold crossed product

$$C^*(H) \rtimes \mathbb{Z} \rtimes \cdots \rtimes \mathbb{Z}$$

as in [32, Proposition 12.8];

- whence

$$K_0(C^*(\mathbb{Z}^m / \Gamma_F, \sigma) \otimes M_n) \cong K_0(C^*(\mathbb{Z}^m / \Gamma_F, \sigma)) \cong \mathbb{Z}^{2r-1}$$

as for noncommutative tori, via the *Pimsner–Voiculescu sequence* [6, Theorem 10.2.1].

This completes the proof. □

Note that the isomorphism class of the center (1-3) alone does *not*, in general, determine  $m$ .

**Example 1.3.** We exhibit noncommutative-sphere algebras  $C(\mathbb{S}_{\theta}^{2m-1})$  with isomorphic centers and distinct  $m$  (so also, by necessity, distinct  $\theta$ ):

(I) Take first  $m = 2$  and  $\theta = 0$ , so that

$$C(\mathbb{S}_\theta^{2m-1}) \cong C(\mathbb{S}^3)$$

and the spectrum of the center is the 3-sphere.

(II) On the other hand, take  $m = 3$  and some  $\theta$  for which the kernel  $\Gamma$  of (1-5) is of rank 1 (i.e., isomorphic to  $\mathbb{Z}$ ) and not contained in any of the subgroups

$$\mathbb{Z}^F, \quad F \subset [3] \text{ properly}$$

of (1-6). The coend (1-7) is then

$$\mathbb{S}_+^2 \times \mathbb{S}^1 / ((x, t) = (x, t'), \forall x \in \partial S_+^2, t \in \mathbb{S}^1)$$

with  $\partial$  denoting the boundary. Alternatively, in words: consider the product  $\mathbb{D}^2 \times \mathbb{S}^1$  (where  $\mathbb{D}^2$  is the closed 2-disk) and identify all copies

$$\mathbb{S}^1 \times \{t\} \subset \mathbb{S}^1 \times \mathbb{S}^1 \subset \mathbb{D}^2 \times \mathbb{S}^1$$

to each other in the obvious fashion. It is a simple exercise to see that this space is again (homeomorphic to) the 3-sphere.

Also in reference to Theorem 1.2, note that an isomorphism

$$(1-10) \quad A \otimes M_n \cong A' \otimes M_n$$

of unital  $C^*$ -algebras does not, in general, imply that  $A$  and  $A'$  themselves are isomorphic:

**Example 1.4.** Recall [45, §2] the unital  $C^*$ -algebras  $A = U_{m,n}^{nc}$  generated by the  $mn$  elements of a unitary (possibly nonsquare)  $m \times n$  matrix. As explained in loc.cit., we may as well assume that  $m \leq n$ .

This is (by definition) the universal unital  $C^*$ -algebra  $A$  whose Hilbert modules  $A^m$  and  $A^n$  are isomorphic, so their endomorphism algebras (as Hilbert modules) will of course also be isomorphic:

$$A \otimes M_m \cong M_m(A) \cong M_n(A) \cong A \otimes M_n.$$

Now fix any  $m \geq 2$ , and consider the case  $A = U_{m,m^2}^{nc}$ . We then have

$$A \otimes M_m \cong A \otimes M_{m^2} \cong (A \otimes M_m) \otimes M_m,$$

which is (1-10) with  $A' = A \otimes M_m$  and  $n = m$ . An isomorphism

$$(1-11) \quad A \cong A' \cong A \otimes M_m$$

does not exist though: according to the discussion immediately preceding [2, §6],

$$K_0(A) \cong \mathbb{Z}/(n-m) = \mathbb{Z}/(m^2-m) \text{ generated by the class of } 1 \in A,$$

so that class cannot be a multiple of  $m$ , as it would be, given an isomorphism (1-11).

The ensuing discussion freely references *Azumaya algebras* (in their various incarnations, such as those in [1, Definition 5.1.4; 46, §13.7.6; 11, §III.1.3]) and ancillary notions, including the following.

- *Polynomial identities* (or *PIs* for short), for (always unital) rings  $A$ , are noncommutative polynomials vanishing when substituting arbitrary elements of  $A$  for the variables. (See [1, Definition 2.2.1].)
- *PI algebras* are those algebras  $A$  over commutative rings (usually fields)  $\mathbb{k}$  satisfying at least one nonzero PI that is *stable* [1, §2.2.3] in the sense that all  $A \otimes_{\mathbb{k}} \mathbb{k}'$  for commutative-ring extensions  $\mathbb{k} \subseteq \mathbb{k}'$  satisfy a common nonzero identity. (See [1, Definition 2.2.41].)
- An algebra  $A$  over a commutative ring  $R$  is *Azumaya (of rank  $n^2$ )* as such [1, Definition 5.1.4] if for some faithfully flat extension  $R \rightarrow R'$  we have

$$A \otimes_R R' \cong M_n(R').$$

- Equivalently [1, Theorem 10.3.2],  $A$  is rank- $n^2$  Azumaya (unqualified, i.e., over its center) precisely when it satisfies the polynomial identities of  $n \times n$  matrices but has no quotient satisfying the identities of  $(n-1) \times (n-1)$  matrices.

We refer the reader to the sources just cited, with more specific citations accompanying the text where appropriate.

**Remark 1.5.** We will repeatedly (and henceforth tacitly) take it for granted that being Azumaya over a commutative ring is stable under scalar extension: if  $A$  is Azumaya over  $R$  then  $A_S := A \otimes_R S$  is Azumaya over  $S$  for any commutative-ring morphism  $R \rightarrow S$  [1, Proposition 5.4.28].

*Proof of Theorem 1.1.* That (c) implies the other conditions is clear enough: for tori transitioning between  $\pm\theta$  is effected by interchanging the two unitary generators, and the realization (0-1) settles the matter for spheres. We thus focus on the converse implications, seeking to deduce (c) from each of the other conditions.

(I) Rational vs. irrational parameters. Given an isomorphism (1-1), the deformation parameters  $\theta$  and  $\theta'$  are simultaneously either rational or irrational.

Indeed, if one of them ( $\theta$ , say) is irrational, then  $C_{\theta,n}$  has infinite-dimensional simple quotients isomorphic to  $A_{\theta,n}$  (simple, being a minimal tensor product of simple  $C^*$ -algebras: [32, Corollary 12.12] and [64, Corollary IV.4.21]). Furthermore, (0-1) makes it clear that *all* infinite-dimensional simple quotients are of this form.

On the other hand, if  $\theta' \in \mathbb{Q}$  then the simple quotients of  $C_{\theta',n'}$  are plainly finite-dimensional, again by (0-1).

The same argument also (more easily) handles (1-2).

(II) The irrational case. This is the simpler branch of the argument: we have already observed that if  $\theta$  and  $\theta'$  are both irrational then (1-1) implies (1-2),  $A_{\theta,n}$  and  $A_{\theta',n'}$  being the only infinite-dimensional simple quotients of  $C_{\theta,n}$  and  $C_{\theta',n'}$  respectively. We can then simply appeal to [56, Theorem 3.12].

(III) The rational case. We are now assuming that

$$\theta = \frac{p}{q} \quad \text{and} \quad \theta' = \frac{p'}{q'}$$

are rational (in lowest terms, as depicted).

We focus mostly on the implication (a)  $\implies$  (c), as it will turn out to be more elaborate, indicating along the way where and how the argument changes so as to also deliver (b)  $\implies$  (c). There is no harm in assuming  $n = n'$  throughout, as Theorem 1.2 allows.

An isomorphism (1-1) induces isomorphisms between

- the centers  $C(\mathbb{S}^3)$  of the two  $C^*$ -algebras [15, Proposition 4.5(1)],
- and hence also between the  $q \times q$  matrix bundles over the Azumaya loci

$$\mathbb{S}^3 \setminus \left( \bigsqcup \text{two circles} \right) \cong \mathbb{T}^2 \times (0, 1)$$

of  $C_{\theta,n}$  and  $C_{\theta',n'}$  [15, Proposition 4.5(3)].

Over those Azumaya loci,  $C_{\theta,n}$  and  $C_{\theta',n'}$  are isomorphic  $qn \times qn$  and  $q'n' \times q'n'$  matrix bundles over  $X = \mathbb{T}^2 \times (0, 1)$ . We thus have  $qn = q'n'$ , and because

- matrix bundles are classifiable homotopically [37, §18.2.4],
- $X$  deformation-retracts onto its slice

$$\mathbb{T}^2 \cong \mathbb{T}^2 \times \left\{ \frac{1}{2} \right\} \subset X,$$

- and over that slice the two algebras are  $A_{\theta,n}$  and  $A_{\theta',n'}$ , respectively,

the implication (b)  $\implies$  (c) reduces to (a)  $\implies$  (c), on which we henceforth focus. We have been assuming that  $n = n'$ , so that  $q = q'$  because  $qn = q'n'$ . The rest is simple conceptually, but requires a bit of unwinding of the attendant bundle-classification theory.

The  $qn \times qn$ -matrix bundles on  $X = \mathbb{T}^2 \times (0, 1)$  are classified by homotopy classes of maps  $X \rightarrow BPU(qn)$  into the *classifying space* [37, §7.2, Definition 2.7] of the projective  $qn \times qn$  unitary group [37, §18.3, Assertion 3.2 and Remark 3.3]. Writing  $[-, -]$  for homotopy classes of maps [37, §6.4, Definition 4.2], we have

$$[X, -] \cong [\mathbb{T}^2, -],$$

so we may as well assume we are working with the 2-torus.

We have an exact sequence

$$(1-12) \quad [\mathbb{T}^2, BSU(qn)] \rightarrow [\mathbb{T}^2, BPU(qn)] \rightarrow [\mathbb{T}^2, B^2\mathbb{Z}/qn] \cong H^2(\mathbb{T}^2, \mathbb{Z}/qn) \\ \cong \mathbb{Z}/qn$$

[42, Corollary 3.2.7] attached to the top row of [37, §18.3.6], where  $B^2$  denotes the iterated classifying-space construction discussed in [37, §7.4.6] for *abelian* groups (such as  $\mathbb{Z}/qn$ ). The second arrow in (1-12) is an *embedding*, given that the leftmost term  $[\mathbb{T}^2, BSU(qn)]$  is trivial: it classifies degree-0 rank- $qn$  vector bundles on the 2-torus, and those are trivial [65, p. 2, Proposition]. In other words,  $qn \times qn$ -matrix-algebra bundles over the 2-torus are classified by the characteristic class denoted by  $\beta_{qn}$  in [37, §18.3.7] (our  $qn$  being that source's  $n$ ).

Denote by  $(\widetilde{-})_q$  inverses modulo  $q$  and similarly for  $q'$ . In our case, focusing on the un-primed side, the  $qn \times qn$ -matrix bundle in question is of the form  $\mathcal{E} \otimes \mathcal{E}^*$  for a vector bundle  $\mathcal{E} \cong \mathcal{F} \otimes \mathbb{C}^n$  with rank- $q$  degree- $\widetilde{p}_q$   $\mathcal{F}$ , and the class  $\beta_{qn}$  is the image of the Chern class

$$n\widetilde{p}_q \in [\mathbb{T}^2, B^2\mathbb{Z}] \cong H^2(\mathbb{T}^2, \mathbb{Z}) \cong \mathbb{Z}$$

through the natural map  $\mathbb{Z} \rightarrow \mathbb{Z}/nq$  (arising from the vertical maps in [37, §18.3.4]) so all in all, the isomorphism of matrix bundles simply amounts to

$$\widetilde{p}_qn = \pm \widetilde{p}'_qn \pmod{qn}$$

(the  $\pm$  depending on whether the homeomorphism between the two spheres preserves or reverses orientations). This is in turn equivalent to

$$qn \mid \widetilde{p}_qn \pm \widetilde{p}'_qn \iff q \mid \widetilde{p}_q \pm \widetilde{p}'_q \iff q \mid p \pm p' \iff \theta + \mathbb{Z} = \pm\theta' + \mathbb{Z}.$$

This concludes the proof.  $\square$

**Remarks 1.6.** (1) The obvious analogue of [55, Theorem 3] does in fact go through for rational deformation parameters, giving an extension of [33, Theorem 1.1].

(2) The literature on the classification of noncommutative tori up to isomorphism or Morita equivalence is extensive: the reader can consult [57; 12; 23], for instance.

(3) The proof of Theorem 1.1 takes for granted the following discussion on Chern classes and their relation to matrix-algebra-bundle characteristic classes.

The fibration sequence associated (as in [37, §18.3.6]) to the exact sequence

$$1 \rightarrow SU(n) \rightarrow U(n) \xrightarrow{\det} \mathbb{S}^1 \rightarrow 1$$

and (part of) the top portion of the diagram in [37, §18.3.6] fit together into a

commutative diagram

$$(1-13) \quad \begin{array}{ccccc} & & B \det & \rightarrow & B\mathbb{S}^1 & \xrightarrow{\cong} & K(\mathbb{Z}, 2) \\ & \nearrow & & & \downarrow & & \downarrow \\ BSU(n) & \rightarrow & BU(n) & & & & K(\mathbb{Z}, 2) \\ & \searrow & \downarrow & & & & \downarrow \\ & & BPU(n) & \rightarrow & B^2\mathbb{Z}/n & \xrightarrow{\cong} & K(\mathbb{Z}/n, 2) \end{array}$$

where

- $K(G, m)$  denotes, as usual [37, Definition 9.6.1], the  $m$ -th Eilenberg–Mac Lane space with homotopy group  $\pi_n \cong G$  and vanishing homotopy in other degrees;
- the left-hand vertical map is the obvious one, resulting from the quotient  $U(n) \rightarrow PU(n)$  that mods out the center;
- the middle vertical arrow is obtained by applying the classifying-space (homotopy) functor  $B(-)$  [60, §3.7; 47, Proposition 8.1] to the map

$$\mathbb{S}^1 \rightarrow B\mathbb{Z}/n$$

classifying the  $\mathbb{Z}/n$ -bundle on the circle obtained via the  $n$ -fold cover

$$\mathbb{S}^1 \xrightarrow{\text{central embedding}} U(n) \xrightarrow{\det} \mathbb{S}^1;$$

- and the right-hand vertical map induced by  $\mathbb{Z} \rightarrow \mathbb{Z}/n$ , well-defined only upon choosing a generator for  $\mathbb{Z}/n$  (so a choice is involved).

For a space  $X$  the upper composition

$$(n\text{-bundles on } X) \xrightarrow[\cong]{[37, \text{Assertion 18.3.2}]} [X, BU(n)] \rightarrow [X, K(\mathbb{Z}, 2)] \cong H^2(X, \mathbb{Z})$$

simply associates the first Chern class  $c_1(\mathcal{E}) \in H^2(X, \mathbb{Z})$  to a vector bundle  $\mathcal{E}$  on  $X$ , whereas the bottom composition

$$(n^2\text{-matrix bundles on } X) \xrightarrow{\cong} [X, BPU(n)] \rightarrow [X, K(\mathbb{Z}/n, 2)] \cong H^2(X, \mathbb{Z}/n)$$

(where again the first isomorphism comes from [37, Assertion 18.3.2]) similarly associates the class  $\beta_n(\mathcal{A})$  of [37, §18.3.7] to a matrix-algebra bundle  $\mathcal{A}$ .

In conclusion, the Chern class  $c_1(\mathcal{E}) \in H^2(X, \mathbb{Z})$  of a vector bundle gets mapped to the  $H^2(X, \mathbb{Z}/n)$ -characteristic class of the corresponding matrix-algebra bundle  $\mathcal{E} \otimes \mathcal{E}^*$ .

(4) We will refer to the class  $\beta_n \in H^2(X, \mathbb{Z}/n)$  associated in [37, §18.3.7] to an  $n \times n$  matrix-algebra bundle on  $X$  as *the characteristic 2-class* of the matrix bundle, to distinguish it from the Dixmier–Douady (3-)class  $\alpha \in H^3(X, \mathbb{Z})$  also discussed in [37, §18.3.7].

## 2. Azumaya theory for higher quantum spheres

Much can be said about higher-dimensional rational tori  $C(\mathbb{T}_\theta^n) = A_\theta^n$  [32, §12.2] as well. These algebras are also Azumaya, (essentially) by [16, Proposition 7.2]. The setup there is more algebraically oriented so the discussion requires some translation, but nothing particularly problematic. The deformation-parameter data in [16, §7] consists of

- a primitive  $\ell$ -th root of unity  $q$  in the ground field (which for us is  $\mathbb{C}$ ); and
- a skew-symmetric  $n \times n$  matrix  $H = (h_{ij}) \in M_n(\mathbb{Z})$ , which will turn out to be an integer multiple of our  $\theta$ .

The  $n$  invertible generators  $x_i$  of loc. cit. are required to skew-commute in the sense that

$$(2-1) \quad x_j x_k = q^{h_{jk}} x_k x_j;$$

compare this with the usual skew-commutation relation [57, p. 193]

$$(2-2) \quad u_k u_j = e^{2\pi i \theta_{jk}} u_j u_k$$

for the unitary generators of  $A_\theta^n$ . Now, if  $\ell$  is a positive integer divisible by the denominators of all  $\theta_{jk}$ , then

$$x_\bullet = u_\bullet, \quad q = e^{-\frac{2\pi i}{\ell}}, \quad H = \ell\theta \in M_n(\mathbb{Z})$$

will effect the transition between equations (2-1) and (2-2).

Proposition 7.2 of [16] (or rather the appropriately translated version) then shows that  $A_\theta^n$  is Azumaya of rank

$$(2-3) \quad h = h_\theta := \text{cardinality of range}(\mathbb{Z}^n \xrightarrow{H=\ell\theta} (\mathbb{Z}/\ell)^n),$$

where a matrix is regarded as an operator in the usual way.

Note that although there was a choice in selecting  $\ell$ ,  $h$  itself will not depend on that choice: scaling  $\ell$  to, say,  $d\ell$  will also scale  $H$  by  $d$ , thus preserving the size of its image. In fact, the number (2-3) has the following alternative, direct description in terms of the matrix  $\theta$  alone:

**Lemma 2.1.** *For a rational skew-symmetric  $\theta \in M_n(\mathbb{Q})$  the number (2-3) can be recovered as the index*

$$(2-4) \quad h_\theta = [(\mathbb{Z}^n + \text{im } \theta) : \mathbb{Z}^n].$$

*Proof.* We can assume [9, §IX.5.1, Théorème 1] that  $\theta$  is block-diagonal, consisting of a zero block and one of the form

$$(2-5) \quad \begin{pmatrix} 0 & D \\ -D & 0 \end{pmatrix}$$

for a diagonal matrix

$$D = \text{diag}(p_1/q_1, p_2/q_2, \dots)$$

with lowest-terms nonzero entries. The quantities in (2-3) and (2-4) are now easily seen to both be equal to  $\prod_i q_i^2$ .  $\square$

**Remark 2.2.** The proof of Lemma 2.1 also makes it plain that (2-3) is indeed a square, as implicit in the discussion: it is the common dimension of the matrix-algebra fibers of an Azumaya algebra.

Still assuming  $\theta$  rational, we have:

**Notation 2.3.** Consider a skew-symmetric matrix  $\theta \in M_n(\mathbb{R})$  (usually rational).

(1) For a tuple  $\mathbf{u} := (u_i)_{i=1}^n$  of generators of  $A_\theta^n$  and an integer tuple  $\mathbf{m} = (m_i)_{i=1}^n$  we write

$$\mathbf{u}^{\mathbf{m}} := \prod_{i=1}^n u_i^{m_i}.$$

The product is to be understood as ordering the indices increasingly unless otherwise specified, but at no point will the ordering in fact matter: we will mostly be interested in  $C^*$ -subalgebras of  $A_\theta^n$  generated by such products, and a reordering simply scales by a modulus-1 complex number.

(2) The *integral kernel* of  $\theta$  is

$$\theta^\perp := \{\mathbf{m} \in \mathbb{Z}^n : \theta \mathbf{m} \in \mathbb{Z}^n\},$$

regarding  $\theta$  as a linear endomorphism on  $\mathbb{R}^n$ .

(3) For  $F \subseteq [n]$  write

- $\theta|_F$  for the  $|F| \times |F|$  matrix consisting of  $\theta$ -entries in  $F$ -labeled rows and columns;
- $h_{\theta,F} := h_{\theta|_F}$ , with  $h_\bullet$  as in (2-4).

The realization of  $A_\theta^n$  as a cocycle twist of the group algebra of  $\mathbb{Z}^n$  [23, §2.2], coupled with the proof of [22, Lemma 2.3], describing the center of such a twisted group algebra, gives

$$(2-6) \quad Z(A_\theta^n) = C^*(\mathbf{u}^{\theta^\perp}) := C^*(\mathbf{u}^{\mathbf{m}} : \mathbf{m} \in \theta^\perp) \stackrel{\theta \in M_n(\mathbb{Q})}{\cong} C^*(\mathbb{Z}^n) \cong C(\mathbb{T}^n).$$

By the Azumaya claim we can once more realize  $A_\theta^n$  as sections of a bundle of dimension- $h_\theta$  matrix algebras over  $\mathbb{T}^n$  for the  $h_\theta$  of (2-4).

**Remark 2.4.** In dealing with vector bundles over higher-dimensional tori, one additional complication not visible for  $n = 2$  (or even  $n = 3$ ) is that the cohomological classification recalled in [15, Remark 4.4(1)] no longer goes through: as observed

in the introductory remarks (p. 247) of [49] and elaborated in § 3 of the same work, there are nonflat rank-2 vector bundles on complex 2-dimensional tori (*four-dimensional* when regarded as real tori) with vanishing Chern classes.

Such pathologies thus occur as soon as one can reasonably expect them, namely whenever the torus and bundle are large enough for both first Chern classes  $c_1$  and  $c_2$  to come into play, meaning the bundle has rank  $\geq 2$  and the torus dimension  $\geq 4$ .

Incidentally, the proof of Lemma 2.1 also shows that rational noncommutative tori are decomposable into small-dimensional pieces:

**Lemma 2.5.** *For a skew-symmetric matrix  $\theta \in M_n(\mathbb{Q})$  the corresponding noncommutative torus algebra  $A_\theta^n$  decomposes as*

$$A_\theta^n \cong C(\mathbb{T}^k) \otimes \bigotimes_{i=1}^t A_{\theta_i}^2$$

for rationals  $\theta_i \in \mathbb{Q}$  and  $n = k + 2t$ .

*Proof.* Immediate from the observation made in passing in the proof of Lemma 2.1 that after a change of (integer) basis  $\theta$  is a direct sum of a  $(k \times k)$ , say zero matrix and a  $(2t \times 2t)$  matrix of the form (2-5).  $\square$

It also follows that, notwithstanding Remark 2.4 (and [15, Remark 4.4(1)]), the bundles relevant to the Azumaya structure of  $A_\theta^n$  are relatively well-behaved. To make sense of the statement, recall [41, §§I.2 and I.4] the notion of a *projectively flat* vector bundle over a space  $X$  (either plain or *Hermitian*, i.e., with structure group  $G = \mathrm{GL}(r)$  [41, §I.4] or  $G = U(r)$ ), respectively. Then:

- Consider the *principal*  $G$ -bundle  $P \rightarrow X$  associated to the vector bundle  $E$  in question, via the usual correspondence [37, §18.2, Assertion 2.3].
- The quotient  $\bar{P} := P/Z(G)$  by the center of  $G$  (scalar matrices, so either  $\mathbb{C}^\times$  when  $G = \mathrm{GL}(r)$  or  $\mathbb{S}^1$  for  $G = U(r)$ ) is then a principal bundle over the *projective* group

$$PG := G/Z(G) = \mathrm{PGL}(r) \text{ or } \mathrm{PU}(r);$$

- The original bundle is projectively flat if  $\bar{P}$  is flat in the usual sense [41, Proposition I.2.6].

**Proposition 2.6.** *For skew-symmetric  $\theta \in M_n(\mathbb{Q})$  we have*

$$A_\theta^n \cong \mathrm{End}(\mathcal{E}) = \Gamma(\mathcal{E} \otimes \mathcal{E}^*) := (\text{continuous sections of the bundle } \mathcal{E} \otimes \mathcal{E}^*)$$

for a projectively flat bundle  $\mathcal{E}$  on  $\mathbb{T}^n \cong \mathrm{Max}(Z(A_\theta^n))$  of rank  $\sqrt{h_\theta}$  for  $h_\theta = (2-4)$ . In particular,  $\mathcal{E} \otimes \mathcal{E}^*$  itself is a flat matrix-algebra bundle.

*Proof.* That the rank is as claimed has already been noted above, as a consequence of [16, Proposition 7.2]. Lemma 2.5 reduces the problem to noncommutative 2-tori. Indeed, it shows that  $\mathcal{E}$  decomposes as an *external (or exterior) tensor product* [4, §2.6], and external tensor products preserve projective flatness: the latter is definable [41, Proposition I.2.8] as the existence of a connection whose curvature takes scalar values, and there is a simple formula for the curvature of the tensor product of the natural two connections [41, §I.5, (5.15)].

The conclusion now follows from the fact that *all* complex vector bundles on a 2-torus (and indeed a compact orientable surface) are projectively flat: being completely classified by rank and Chern class ([65, Proposition, p. 2] again) a rank- $q$  bundle splits as the sum between a line bundle and a trivial rank- $(q - 1)$  bundle, etc.

As for the last claim on flatness, it follows from the general remark that  $\mathcal{E} \otimes \mathcal{E}^*$  is flat whenever  $\mathcal{E}$  is projectively flat [41, Propositions I.2.9 and/or I.4.23].  $\square$

**Remark 2.7.** The flatness of the relevant  $q \times q$  matrix bundle for  $n = 2$  and  $\theta = p/q$  is plain from its direct construction in the proof of [32, Proposition 12.2] (or [33, §2], on which the latter account is based): the bundle is obtained as the (total space of the) quotient

$$(\mathbb{T}^2 \times M_q) / (\mathbb{Z}/q)^2$$

through the diagonal action, where

$$(\mathbb{Z}/q)^2 \subset \mathbb{T}^2$$

acts on the torus via translation by the  $q$ -torsion subgroup and on  $M_q$  as described in loc. cit. (In particular, the two generators act by conjugation by two order- $q$  unitary matrices that commute up to scalars.)

A few reminders will help make sense of the statement of Theorem 2.8, a higher-dimensional generalization of [15, Proposition 4.5].

- The *PI-degree* [19, Definitions A.7.1.8 and B.4.15; 29, §1] of a PI algebra is the largest  $n$  for which the polynomial identities of  $A$  are among those of  $M_n$ . Rank- $n^2$  Azumaya algebras, for instance, have PI-degree  $n$  (as observed just after Definition B.4.15 of [19]).

The slight definition variations one typically encounters in the literature will not make a difference here, so the above will do.

- The *Azumaya locus* of an algebra  $A$  consists of those maximal ideals

$$\mathfrak{m} \in \text{Max}(Z := Z(A)) := \text{maximal spectrum of the center } Z$$

for which the *localization* [5, Example (1) post Corollary 3.2]  $A_{\mathfrak{m}}$  is Azumaya over  $Z_{\mathfrak{m}}$ .

There are again minor departures from this setup in the literature (one might,

for instance, consider all *prime* ideals with the requisite property [66, §1], impose additional finiteness/primality constraints on the algebra  $A$  [11, §III.1.7]).

Set

$$(2-7) \quad I_Y := \{f|_Y \equiv 0\} \trianglelefteq \text{Cont}_\theta(\mathbb{S}_+^{n-1} \rightarrow C(\mathbb{T}_\theta^n)) \stackrel{(0-1)}{\cong} C_\theta, \quad Y = \bar{Y} \subseteq \mathbb{S}_+^{n-1}$$

and

$$(2-8) \quad C_\theta|_Y := C_\theta/I_Y, \quad Z_\theta|_Y := Z_\theta/(I_Y \cap Z_\theta).$$

For subsets  $S \subseteq \mathbb{Z}^n$  and  $F \subseteq [n]$  we write

$$S_{\downarrow F} := \{s = (s_i)_{i=1}^n \in S : s_i = 0, \forall i \notin F\};$$

the portion of  $S$  *supported on*  $F$ , in other words. This applies to the symbol  $\theta_{\downarrow F}^\perp$  employed below.

**Theorem 2.8.** *Let  $\theta \in M_n(\mathbb{Q})$  be an  $n \times n$  skew-symmetric rational matrix for some fixed  $n \geq 2$ .*

(1) *The center of the noncommutative sphere algebra  $C_\theta := C(\mathbb{S}_\theta^{2n-1})$  is*

$$(2-9) \quad \begin{aligned} Z_\theta &:= Z(C(\mathbb{S}_\theta^{2n-1})) \\ &\cong \text{Cont}_\theta(\mathbb{S}_+^{n-1} \rightarrow Z(C(\mathbb{T}_\theta^n))) \\ &= \{\mathbb{S}_+^{n-1} \xrightarrow[\text{cont.}]{f} C^*(\mathbf{u}^{\theta^\perp}) : f(\mathbb{S}_{+,F}^{n-1}) \in C^*(\mathbf{u}^{\theta_{\downarrow F}^\perp}), \forall F \subseteq [n]\}. \end{aligned}$$

(2) *The spectrum  $X_\theta := \text{Max}(Z_\theta)$  branch-covers the classical sphere*

$$(2-10) \quad \mathbb{S}^{2n-1} \cong \text{Max}\{\mathbb{S}_+^{n-1} \xrightarrow[\text{cont.}]{f} C^*(u_i^{q_i}) : f(\mathbb{S}_{+,F}^{n-1}) \in C^*(u_i^{q_i}, i \in F), \forall F \subseteq [n]\}$$

for

$$q_i := \text{lcm}(\text{lowest-term denominators of } \theta_{ij}, j \in [n]).$$

(3)  *$C_\theta$  embeds into  $M_{\sqrt{h_\theta}}(Z_\theta)$  with  $h_\theta$  as in (2-4) and is not a  $Z_\theta$ -subalgebra of  $M_n(Z_\theta)$  for any smaller  $n$ . In particular,  $C_\theta$  is a PI algebra of PI-degree  $\sqrt{h_\theta}$ .*

*Proof.* (1) We have already recalled (2-6) that the centers  $Z(C(\mathbb{T}_\theta^n))$  of the noncommutative torus algebras are as claimed, so the conclusion follows from the identification (0-1).

(2) That the right-hand side of (2-10) is indeed a sphere is a simple topology exercise (the classical counterpart to (0-1)), given that the  $u_i^{q_i} \in A_\theta^n$  are central and hence the generators of a  $C(\mathbb{T}^n)$ . The claim is that the map

$$X_\theta \xrightarrow[\text{onto}]{\pi} \mathbb{S}^{2n-1}$$

dualizing the embedding

$$(2-11) \quad C(\mathbb{S}^{2n-1}) \cong \text{Cont}_\theta(\mathbb{S}_+^{n-1} \rightarrow Z_\downarrow(A_\theta^n) := C^*(u_i^{qi})) \\ := \{ \mathbb{S}_+^{n-1} \xrightarrow[\text{cont.}]{f} Z_\downarrow(A_\theta^n) : f(\mathbb{S}_{+,F}^{n-1}) \in C^*(u_i^{qi}, i \in F), \forall F \subseteq [n] \} \\ \longrightarrow (2-9) =: C(X_\theta)$$

is a *branched cover* in the sense of [51, §1]: an open surjection of compact Hausdorff spaces, with a finite upper bound on the cardinalities of the fibers  $\pi^{-1}(p)$ ,  $p \in \mathbb{S}^{2n-1}$ . Surjectivity is not at issue, so it is the two other requirements that require attention. Rather than attack that classical statement directly, we appeal to the theory of *noncommutative branched covers* developed in [51; 7], revolving around the notion of a  $C^*$  *conditional expectation* [6, Definition II.6.10.1].

We have embeddings

$$Z_\downarrow(A_\theta^n) \hookrightarrow Z(A_\theta^n) \hookrightarrow C^*(\mathbb{Z}^n) \cong C^*(t_i, i \in [n]), \quad t_i^{qi} = u_i^{qi}.$$

The usual [53, Proposition 8.5] (plainly of *finite index* [28, Definition 2]) expectation

$$C^*(t_i, i \in [n]) \xrightarrow{E} Z_\downarrow(A_\theta^n)$$

restricts to another such (also  $E$ ) on the intermediate  $Z(A_\theta^n)$ , compatible with the inclusions of the subgroups generated by  $t_i, i \in F$  for  $F \subseteq [n]$ .

We then have a finite-index

$$\text{Cont}(\mathbb{S}_+^{n-1} \rightarrow Z(A_\theta^n)) \xrightarrow{E \circ} \text{Cont}(\mathbb{S}_+^{n-1} \rightarrow Z_\downarrow(A_\theta^n)),$$

hence also (because of the noted  $F$ -compatibility) an analogous expectation between  $\text{Cont}_\theta$  function algebras. These, however, are exactly the two extremes of (2-11), whence the conclusion by [7, Theorem 1.4].

(3) The isomorphism (0-1) and Proposition 2.6 ensure that the irreducible  $C_\theta$ -representations are all at most  $\sqrt{h_\theta}$ -dimensional. That estimate is the best possible: in the language of [6, Definition IV.1.4.1],  $C_\theta$  is sharply  $\sqrt{h_\theta}$ -*subhomogeneous*. To verify this last sharpness assertion, quotient out the ideal  $I_p := I_{\{p\}}$  of (2-7) in (0-1) to obtain a surjection

$$C_\theta \twoheadrightarrow A_\theta^n \xrightarrow{\text{Proposition 2.6}} M_{\sqrt{h_\theta}}.$$

$C_\theta$  thus embeds [18, §2.7.3] into a product of matrix algebras  $M_n, n \leq \sqrt{h_\theta}$  with at least one  $M_{\sqrt{h_\theta}}$  quotient, and the conclusion follows.  $\square$

**Corollary 2.9.** *A noncommutative sphere algebra  $C_\theta$  is Azumaya if and only if it is commutative, i.e.,  $\theta \in M_n(\mathbb{Z})$ .*

*Proof.* This is immediate from Theorem 2.8 and its proof:  $\theta \notin M_n(\mathbb{Z})$  is equivalent to  $h_\theta > 1$ , and it remains to observe that if that condition holds then  $C_\theta$  cannot be Azumaya, for on the one hand it has an  $h_\theta$ -dimensional matrix algebra quotient, while on the other hand  $C_\theta|_p$  is abelian for any of the  $n$  vertices  $p \in \mathbb{S}_+^{n-1}$ .  $\square$

The branch-covering qualification in Theorem 2.8(2) is not redundant: as soon as  $n \geq 3$  the center  $Z_\theta$  of (2-9) need not be (the function algebra of) a sphere.

**Example 2.10.** Set

$$\theta := \begin{pmatrix} 0 & \frac{1}{2} & \frac{1}{2} \\ -\frac{1}{2} & 0 & \frac{1}{2} \\ -\frac{1}{2} & -\frac{1}{2} & 0 \end{pmatrix} \in M_3(\mathbb{Q}).$$

Denoting by  $u_1, u_2, u_3$  the generators of  $A_\theta^3$ , the sphere  $\mathbb{S}^{2n-1} = \mathbb{S}^5$  of Theorem 2.8(2) is

$$\mathbb{S}^5 \cong \text{Max Cont}_\theta(\mathbb{S}_+^2 \rightarrow C^*(u_i^2, i \in [3])),$$

while

$$X_\theta \cong \text{Max Cont}_\theta(\mathbb{S}_+^2 \rightarrow Z(A_\theta^3) = C^*(u_i^2, i \in [3], u_1 u_2 u_3)).$$

The latter space thus consists of two 5-spheres glued along the 3-dimensional complex

$$(2-12) \quad \text{Max Cont}_\theta(\partial\mathbb{S}_+^2 \rightarrow Z(A_\theta^3))$$

(one copy of which is embedded in each of the two 5-spheres). Said complex can be described as follows:

- Consider three 3-spheres  $\mathbb{S}_i^3$  indexed by  $i \in \mathbb{Z}/3$ , regarded as total spaces of the *Hopf fibration* [17, §14.1.9]  $\mathbb{S}^3 \xrightarrow{\pi} \mathbb{S}^2$ .
- In each  $\mathbb{S}_i^3$  set  $\mathbb{S}_{i,\pm}^1 := \pi^{-1}(p_\pm)$  for antipodes  $p_\pm \in \mathbb{S}^2$ .
- Glue  $\mathbb{S}_i^3$  to  $\mathbb{S}_{i+1}^3$  (indices modulo 3) by identifying  $\mathbb{S}_{i,+}^1 \cong \mathbb{S}_{i+1,-}^1$  to obtain (2-12).

$X_\theta$  is now easily seen not to be homeomorphic to  $\mathbb{S}^5$ , and indeed, not a topological manifold: points in  $\mathbb{S}_0^3$  but off the exceptional circles  $\mathbb{S}_{0,\pm}^1$  have neighborhoods homeomorphic to two copies of  $\mathbb{R}^5$  glued along a closed  $\mathbb{R}^3$ . The removal of that  $\mathbb{R}^3$  disconnects such a neighborhood, so it cannot [25, Theorem 1.8.13] be homeomorphic to a Euclidean space.

**Remark 2.11.** Example 2.10 illustrates a qualitative distinction between quantum tori and spheres: while *equivalent*  $\theta, \theta' \in M_n(\mathbb{Q})$ , in the sense [9, §IX.1.6] (appropriate for bilinear forms) that

$$\exists(T \in \text{GL}_n(\mathbb{Z})) \quad : \quad \theta' = T\theta T^t,$$

will produce isomorphic torus algebras  $A_\theta^n \cong A_{\theta'}^n$ , it may well be that  $C_\theta \not\cong C_{\theta'}$ . In fact, not even the centers of the latter two algebras need be isomorphic.

Indeed,  $Z_\theta$  will be (the function algebra of) a  $(2n - 1)$ -sphere whenever  $\theta$  is block-diagonal with blocks (2-5), for in that case, in the notation of Theorem 2.8,

$$Z(A_\theta^n) = C^*(u_i^{q_i}, i \in [n]).$$

Per Example 2.10, then, with that choice of  $\theta$  and

$$\theta' := \begin{pmatrix} 0 & \frac{1}{2} & 0 \\ -\frac{1}{2} & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} = T\theta T^t, \quad T := \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & -1 & 1 \end{pmatrix}.$$

The respective centers  $Z_{\theta, \theta'} = Z(C_{\theta, \theta'})$  are nonisomorphic.

To follow up on Corollary 2.9, determining the Azumaya locus of  $C_\theta$  is now also. We again need some vocabulary and notation preliminaries.

**Definition 2.12.** Let  $\theta \in M_n(\mathbb{Q})$  be a rational skew-symmetric matrix and recall Notation 2.3.

The *jump (sub)complex*  $\text{JMP}_\theta$  of  $\theta$  (just plain JMP when  $\theta$  is understood) is

$$\text{JMP} = \text{JMP}_\theta := \{F \in 2^{[n]} : h_{\theta, F} < h_\theta\}.$$

Identifying *simplicial complexes* and their respective *geometric realizations* [17, §8.1],  $\text{JMP}_\theta$  is a *subcomplex* [17, § 8.1; 62, §3.1] of the simplex  $\Delta^{n-1} \cong \mathbb{S}_+^{n-1}$  of (0-2a) with vertex set  $[n]$ ; this justifies the term.

**Theorem 2.13.** *The Azumaya locus of  $C_\theta$  is the complement*

$$(2-13) \quad (X_\theta = \text{Max}(Z_\theta)) \setminus \text{Max}(Z_\theta|_{\text{JMP}})$$

in the notation of (2-8).

*Proof.* (2-13) contains the Azumaya locus: Suppose  $p \in \text{Max}(Z_\theta)$  belongs precisely to the subspaces

$$X_{\theta, \hat{i}} := \text{Max}(Z_\theta|_{\mathbb{S}_{+, \hat{i}}^{n-1}}) \subseteq X_\theta, \quad i \in F \in 2^{[n]}.$$

The intersection

$$p \cap C(\mathbb{S}_+^{n-1}) \in \text{Max } C(\mathbb{S}_+^{n-1}) \cong \mathbb{S}_+^{n-1}$$

will then belong to the interior of the face  $\mathbb{S}_{+, \hat{F}}^{n-1}$ , and (0-1) specializes to

$$C_\theta / (C_\theta \cdot (p \cap C(\mathbb{S}_+^{n-1}))) \cong C^*(u_i, i \in \hat{F}) \cong A_{\theta|_F}^{\hat{F}} \subseteq A_\theta^n.$$

If  $\hat{F} \in \text{JMP}$  then (by the very definition of the jump complex) this further specializes, at every maximal ideal of the center  $Z(A_\theta^n) \cong C(\mathbb{T}^n)$ , to a *proper* inclusion

$$C_\theta / (C_\theta \cdot p) \cong (M_{\sqrt{h_{\theta, \hat{F}}}})^{[(\theta|_{\hat{F}})^\perp : \theta_{\hat{F}}^\perp]} \subsetneq M_{\sqrt{h_\theta}}$$

of a nonmatrix algebra, with  $(\theta|_{\widehat{F}})^\perp \leq \mathbb{Z}^{|\widehat{F}|}$  regarded as a subgroup of  $\mathbb{Z}^n$  by padding vectors with zero entries.  $C_\theta$ , then, cannot be Azumaya at  $p$ , for its localization  $(C_\theta)_p$  does not satisfy the polynomial identities of any  $M_n$ ,  $n < \sqrt{h_\theta}$ .

(2-13) is contained in the Azumaya locus: Every maximal ideal  $p \in (2-13)$  is contained in the interior of some

$$\text{Max}(Z_\theta|_Y) \subset \text{Max}(Z_\theta), \quad Y = \bar{Y} \subseteq \mathbb{S}_+^{n-1} \setminus \text{JMP},$$

because the intersection  $p \cap C(\mathbb{S}_+^{n-1})$  belongs to the latter open set, and it suffices to take for  $Y$  a closed neighborhood of that point still contained in that open. The conclusion now follows from the fact that for such  $Y$  the algebras  $C_\theta|_Y$  are Azumaya:

$$C_\theta|_Y \cong \text{End}(\pi^* \mathcal{E}), \quad Y \times \mathbb{T}^n \xrightarrow{\pi := 2\text{nd projection}} \mathbb{T}^n \cong \text{Max}(Z(A_\theta^n)). \quad \square$$

### 3. On and around center-finiteness

Finiteness (i.e., being a finitely generated module) over the center is well-known not to be automatic for PI algebras: [54, Example post Proposition 5.1.3], say, is of a finitely generated algebra satisfying the identities of  $M_4$  but not embeddable in a ring finite over its center. The noncommutative sphere algebras  $C_\theta$  themselves, as will become apparent, are another case in point.

Subhomogeneous  $C^*$ -algebras  $A$  (such as  $C_\theta$ ) admit (unital) embeddings

$$(3-1) \quad A \leq M_n^I \cong M_n(C(\beta I)),$$

$$\beta(-) := \text{Stone-}\check{\text{C}}\text{ech compactification [30, §6.5] of } I,$$

as noted in the proof of Theorem 2.8 (essentially by [18, §2.7.3]). But this will also not (generally) suffice to ensure center-finiteness, per Example 3.1: the latter will indeed not even be *topologically* center-finite, in the sense of containing a (norm-topology-)dense module over its center. Naturally, rings sandwiched as  $R \leq A \leq M_n(R)$  for *Noetherian* commutative  $R$  are center-finite, so Example 3.1 is in a sense a manifestation of the non-Noetherianness of the continuous-function algebras  $C(X)$  involved.

**Example 3.1.** [52, Example 3.5] (also [7, Example 3.6]) provides a  $C^*$ -algebra  $A$  equipped with a central morphism  $C(X) \rightarrow A$  (a  $C(X)$ -algebra) for

$$X := (X_0 := \bigsqcup_{n \geq 1} \mathbb{C}\mathbb{P}^n)^+ := \text{one-point compactification [68, Problem 19A] of } X_0$$

with fiber  $\mathbb{C}$  at the distinguished point  $\infty \in X$  and fibers  $M_2(\mathbb{C})$  over  $X_0$ . Because the associated  $M_2$ -bundle over  $X$  is by construction not *of finite type* (that is, [36, Definition 3.5.7] trivializable over a finite open cover of  $X_0$ ),  $A$  cannot be

topologically finitely generated as a  $C(X)$ -module by [31, Theorem 1.1] (or [14, Theorem A]) and [7, Proposition 3.7].

**Remarks 3.2.** (1) Example 3.1 is one instance of the following general setup.

- Consider compact Hausdorff spaces  $Y_n$  respectively equipped with (complex) vector  $d$ -bundles  $\mathcal{E}_n$ ; we abuse notation and conflate these [37, Remark 15.1.2] with their corresponding principal  $U(d)$ -bundles.

- Assume the attached set

$$\{\text{ind}_{U(d)}(\mathcal{E}_n)\}_n \subseteq \mathbb{Z}_{\geq 0}$$

of  $U(d)$ -indices [44, Definition 6.2.3] is unbounded: there is no finite upper bound on the cardinality of an open cover of  $Y_n$  that will trivialize  $\mathcal{E}_n$ .

- The  $\mathcal{E}_n$  glue to a vector  $d$ -bundle over  $X_0 := \bigsqcup_n Y_n$ .
- Form the bundle  $\mathcal{F} := \mathcal{E} \oplus (X_0 \times \mathbb{C})$  (i.e., add a trivial 1-dimensional summand to  $\mathcal{E} \rightarrow X_0$ ).
- Construct the corresponding endomorphism bundle  $\mathcal{F} \otimes \mathcal{F}^*$  (a  $(d + 1) \times (d + 1)$ -matrix bundle over  $X_0$ ).
- And finally, take for the  $C^*$ -algebra  $A$  (supposed to play the same role as in Example 3.1) the *unitization* [6, §II.1.2.1]  $\Gamma_0(\mathcal{F} \otimes \mathcal{F}^*)^+$ , the subscript 0 indicating sections on  $X_0$  vanishing at  $\infty$ .

(2) For path-connected (compact Hausdorff)  $Y_n$  in (1) above the unboundedness

$$\sup_n \dim Y_n, \quad \dim := \text{covering dimension} \text{ [25, Definition 1.6.7]}$$

is an essential feature of this family of examples: per [38, Proposition 2.1] (or as an immediate consequence of [50, Lemma 2.4], say) for any *paracompact* [26, §5.1] path-connected space  $Y$  there is an open cover

$$Y = \bigcup_{i=0}^{\dim Y} U_i, \quad U_i \text{ contractible in } Y.$$

The restriction of a bundle on  $Y$  will be trivializable [17, Theorem 14.3.3] over every  $U_i$ , rendering (1) inoperative.

(3) In reference to (3-1), observe that endomorphism algebras  $\text{End}(\mathcal{E}) = \Gamma(\mathcal{E} \otimes \mathcal{E}^*)$  for vector-bundles  $\mathcal{E} \rightarrow X$  over compact Hausdorff  $X$  can always be embedded unitally in some  $M_n(C(X))$  (same space  $X$ : one need not involve the typically nonmetrizable  $\beta$ (discrete  $X$ )).

To see this, consider a decomposition  $\mathcal{E} \oplus \mathcal{F} \cong \mathbf{1}^m$  of a trivial bundle over  $X$ , with  $\mathbf{1}$  denoting the trivial rank-1 bundle (one such exists by [63, Theorems 1 and 2]).

We then have

$$\mathcal{E}^d \oplus \mathcal{F}^d \cong \mathbf{1}^{md}, \quad d := \text{rank } \mathcal{E},$$

and obtain an embedding

$$\text{End}(\mathcal{E}) \xrightarrow{\iota := \text{id}^d \oplus (\text{id} \otimes \text{ev}_x)} \text{End}(\mathcal{E}^d) \times \text{End}(\mathcal{F}^d) \leq \text{End}(\mathbf{1}^{md}) \cong M_{md}(C(X)),$$

where the second component,  $\text{id} \otimes \text{ev}_x$ , means

- identifying  $\mathcal{F}^d$  with  $\mathcal{F} \otimes \mathbb{C}^d$  (focusing on the complex case to fix ideas), and
- mapping a global endomorphism  $s \in \text{End}(\mathcal{E})$  to the endomorphism of  $\mathcal{F} \otimes \mathbb{C}^d$  operating trivially on the  $\mathcal{F}$  tensorand and as the restriction  $s_x \in \text{End}(\mathbb{C}^d) \cong \text{End}(\mathcal{E}_x)$  of  $x$  to the fiber at some fixed  $x \in X$ .

Returning to the  $C_\theta$ : while (mostly) not center-finite, they nevertheless exhibit less pathology in that regard than Example 3.1 and the like.

**Theorem 3.3.** *Let  $\theta \in M_n(\mathbb{Q})$  be a skew-symmetric matrix for some  $n \in \mathbb{Z}_{\geq 2}$ .*

- (1) *The quantum sphere  $C^*$ -algebra  $C_\theta$  is center-finite precisely when it is commutative, i.e.,  $\theta \in M_n(\mathbb{Z})$ .*
- (2) *On the other hand,  $C_\theta$  is always topologically center-finite.*

*Proof.* (1) Theorem 2.8 makes  $C_\theta$  into a subhomogeneous  $Z_\theta$ -algebra, which can thus [13, Theorem A] be regarded as a  $Z_\theta$ -Hilbert module [67, Definition 15.1.5]. Finite generation would imply [67, Corollary 15.4.8] that  $C_\theta$  is also *projective* [1, Definition 5.3.1].

The Hilbert-module-to-Hilbert-bundle correspondence of [34, Scholium 6.7] and Swan's celebrated [59, Theorem 1.6.3] (originally [63, Theorems 1 and 2]) then imply that the (F) Hilbert bundle [21, p. 9] over  $X := \text{Max}(Z_\theta)$  with fibers

$$X \ni p \mapsto C_\theta|_p$$

is locally trivial, so (the sphere being connected) of constant rank. This, in turn, is equivalent (Corollary 2.9) to the commutativity of  $C_\theta$ .

(2) Once more regarding  $C_\theta$ , via Theorem 2.8(1), as the section-space  $\Gamma(\mathcal{E})$  of a subhomogeneous (F) Banach bundle  $\mathcal{E} \rightarrow \mathbb{S}^{2n-1} \cong \text{Max}(Z_\theta)$ , observe that the *strata*

$$X_d := \{p \in X : \dim \text{fiber } \mathcal{E}_p = \dim C_\theta / C_\theta \cdot p = d\}$$

are all members of the *set ring* [8, I, Definition 1.2.13] generated by the spaces

$$\text{Max}(Z_\theta|_{\mathbb{S}_{+,F}^{n-1}}) \subseteq \text{Max}(Z_\theta) = X, \quad F \in 2^{[n]}.$$

Thus they are all finite unions of path-connected paracompact spaces of finite (covering) dimension, so by [38, Proposition 2.1] they admit respective covers

$$X_d = \bigcup_{j=0}^N U_j, \quad U_j = \overset{\circ}{U}_j \text{ contractible in } \mathbb{S}_d^{2n-1} \quad (\text{some } N \in \mathbb{Z}_{\geq 0}).$$

The restrictions  $\mathcal{E}|_{X_d}$  are thus all trivializable by finite covers, hence topological finite generation via [31, Theorem 1.1].  $\square$

### Acknowledgments

The paper has benefited from fruitful exchanges with B. Badzioch, M. Khalkhali, B. Passer and I. Thompson.

### References

- [1] E. Aljadeff, A. Giambruno, C. Procesi, and A. Regev, *Rings with polynomial identities and finite dimensional representations of algebras*, AMS Colloquium Publications **66**, Amer. Math. Soc., 2020. MR
- [2] P. Ara and K. R. Goodearl, “ $C^*$ -algebras of separated graphs”, *J. Funct. Anal.* **261**:9 (2011), 2540–2568. MR
- [3] M. Artin, “On Azumaya algebras and finite dimensional representations of rings”, *J. Algebra* **11** (1969), 532–563. MR
- [4] M. F. Atiyah, *K-theory*, W. A. Benjamin, 1967. MR
- [5] M. F. Atiyah and I. G. Macdonald, *Introduction to commutative algebra*, Addison-Wesley, 1969. MR
- [6] B. Blackadar, *Operator algebras: theory of  $C^*$ -algebras and von Neumann algebras = Operator algebras and non-commutative geometry, III*, Encyclopaedia of Mathematical Sciences **122**, Springer, 2006. MR
- [7] E. Blanchard and I. Gogić, “On unital  $C(X)$ -algebras and  $C(X)$ -valued conditional expectations of finite index”, *Linear Multilinear Algebra* **64**:12 (2016), 2406–2418. MR
- [8] V. I. Bogachev, *Measure theory*, vol. 1, Springer, 2007. MR
- [9] N. Bourbaki, *Algèbre* (chapitre 9), Masson, 1959. MR
- [10] N. Bourbaki, *Algebra II* (chapters 4–7), Springer, 1990. MR
- [11] K. A. Brown and K. R. Goodearl, *Lectures on algebraic quantum groups*, Birkhäuser, 2002. MR
- [12] H. Bursztyn, “A survey on Morita equivalence of quantum tori”, preprint, 1999, available at <https://math.berkeley.edu/~alanw/242papers99/bursztyn>.
- [13] A. Chirvasitu, “Non-commutative branched covers and bundle unitarizability”, preprint, 2024. arXiv 2409.03531v1
- [14] A. Chirvasitu, “Small Banach bundles and modules”, *Proc. Amer. Math. Soc.* **153**:9 (2025), 3907–3920. MR
- [15] A. Chirvasitu and B. Passer, “Continuity and equivariant dimension”, preprint, 2025. arXiv 2403.06767v3
- [16] C. De Concini and C. Procesi, “Quantum groups”, pp. 31–140 in *D-modules, representation theory, and quantum groups* (Venice, 1992), Lecture Notes in Math. **1565**, Springer, 1993. MR

- [17] T. tom Dieck, *Algebraic topology*, European Math. Soc., Zürich, 2008. MR
- [18] J. Dixmier,  *$C^*$ -algebras*, North-Holland Mathematical Library **15**, North-Holland, Amsterdam, 1977. MR
- [19] V. Drensky and E. Formanek, *Polynomial identity rings*, Birkhäuser, Basel, 2004. MR
- [20] M. J. Dupré, *The classification and structure of  $C^*$ -algebra bundles*, Mem. Amer. Math. Soc. **222**, Amer. Math. Soc., 1979. MR
- [21] M. J. Dupré and R. M. Gillette, *Banach bundles, Banach modules and automorphisms of  $C^*$ -algebras*, Research Notes in Mathematics **92**, Pitman, 1983. MR
- [22] G. A. Elliott, “On the  $K$ -theory of the  $C^*$ -algebra generated by a projective representation of a torsion-free discrete abelian group”, pp. 157–184 in *Operator algebras and group representations* (Neptun, Romania, 1980), Monogr. Stud. Math. **17**, Pitman, 1984. MR
- [23] G. A. Elliott and H. Li, “Strong Morita equivalence of higher-dimensional noncommutative tori, II”, *Math. Ann.* **341**:4 (2008), 825–844. MR
- [24] R. Engelking, *Dimension theory*, North-Holland Mathematical Library **19**, North-Holland, Amsterdam, 1978. MR
- [25] R. Engelking, *Dimension theory*, North-Holland Mathematical Library **19**, North-Holland, Amsterdam, 1978. MR
- [26] R. Engelking, *General topology*, 2nd ed., Sigma Series in Pure Mathematics **6**, Heldermann, Berlin, 1989. MR
- [27] J. M. G. Fell and R. S. Doran, *Representations of  $*$ -algebras, locally compact groups, and Banach  $*$ -algebraic bundles, vol. 1: Basic representation theory of groups and algebras*, Pure and Applied Mathematics **125**, Academic Press, Boston, 1988. MR
- [28] M. Frank and E. Kirchberg, “On conditional expectations of finite index”, *J. Operator Theory* **40**:1 (1998), 87–111. MR
- [29] T. Gateva, “P.I. degree of tensor products of PI-algebras”, *J. Algebra* **123**:1 (1989), 64–73. MR
- [30] L. Gillman and M. Jerison, *Rings of continuous functions*, Graduate Texts in Mathematics **43**, Springer, 1960. MR
- [31] I. Gogić, “Topologically finitely generated Hilbert  $C(X)$ -modules”, *J. Math. Anal. Appl.* **395**:2 (2012), 559–568. MR
- [32] J. M. Gracia-Bondía, J. C. Várilly, and H. Figueroa, *Elements of noncommutative geometry*, Birkhäuser, 2001. MR
- [33] R. Høegh-Krohn and T. Skjelbred, “Classification of  $C^*$ -algebras admitting ergodic actions of the two-dimensional torus”, *J. Reine Angew. Math.* **328** (1981), 1–8. MR
- [34] K. H. Hofmann and K. Keimel, “Sheaf-theoretical concepts in analysis: bundles and sheaves of Banach spaces, Banach  $C(X)$ -modules”, pp. 415–441 in *Applications of sheaves* (Durham, 1977), Lecture Notes in Math. **753**, Springer, 1979. MR
- [35] K. H. Hofmann and S. A. Morris, *The structure of compact groups: a primer for the student, a handbook for the expert*, 4th ed., Studies in Mathematics **25**, De Gruyter, Berlin, 2020. MR
- [36] D. Husemoller, *Fibre bundles*, 3rd ed., Graduate Texts in Mathematics **20**, Springer, 1994. MR
- [37] D. Husemoller, M. Joachim, B. Jurčo, and M. Schottenloher, *Basic bundle theory and K-cohomology invariants*, Lecture Notes in Physics **726**, Springer, 2008. MR
- [38] I. M. James, “On category, in the sense of Lusternik–Schnirelmann”, *Topology* **17**:4 (1978), 331–348. MR

- [39] G. G. Kasparov, “Equivariant  $KK$ -theory and the Novikov conjecture”, *Invent. Math.* **91**:1 (1988), 147–201. MR
- [40] M. Khalkhali, *Basic noncommutative geometry*, 2nd ed., European Math. Soc., Zürich, 2013. MR
- [41] S. Kobayashi, *Differential geometry of complex vector bundles*, Publ. Math. Soc. Japan **15**, Princeton University Press, 1987. MR
- [42] S. O. Kochman, *Bordism, stable homotopy and Adams spectral sequences*, Fields Institute Monographs **7**, Amer. Math. Soc., 1996. MR
- [43] S. Mac Lane, *Categories for the working mathematician*, 2nd ed., Graduate Texts in Mathematics **5**, Springer, 1998. MR
- [44] J. Matoušek, *Using the Borsuk–Ulam theorem: lectures on topological methods in combinatorics and geometry*, Springer, 2003. MR
- [45] K. McClanahan, “ $K$ -theory and Ext-theory for rectangular unitary  $C^*$ -algebras”, *Rocky Mountain J. Math.* **23**:3 (1993), 1063–1080. MR
- [46] J. C. McConnell and J. C. Robson, *Noncommutative Noetherian rings*, revised ed., Graduate Studies in Mathematics **30**, Amer. Math. Soc., 2001. MR
- [47] S. A. Mitchell, “Notes on principal bundles and classifying spaces”, preprint, 2011, available at <https://sites.math.washington.edu/~mitchell/Notes/prin.pdf>.
- [48] T. Natsume and C. L. Olsen, “Toeplitz operators on noncommutative spheres and an index theorem”, *Indiana Univ. Math. J.* **46**:4 (1997), 1055–1112. MR
- [49] T. Oda, “Vector bundles on abelian surfaces”, *Invent. Math.* **13** (1971), 247–260. MR
- [50] R. S. Palais, “Homotopy theory of infinite dimensional manifolds”, *Topology* **5** (1966), 1–16. MR
- [51] A. A. Pavlov and E. V. Troitskii, “Quantization of branched coverings”, *Russ. J. Math. Phys.* **18**:3 (2011), 338–352. MR
- [52] N. C. Phillips, “Recursive subhomogeneous algebras”, *Trans. Amer. Math. Soc.* **359**:10 (2007), 4595–4623. MR
- [53] G. Pisier, *Introduction to operator space theory*, LMS Lecture Note Series **294**, Cambridge University Press, 2003. MR
- [54] C. Procesi, *Rings with polynomial identities*, Pure and Applied Mathematics **17**, Dekker, New York, 1973. MR
- [55] M. A. Rieffel, “ $C^*$ -algebras associated with irrational rotations”, *Pacific J. Math.* **93**:2 (1981), 415–429. MR
- [56] M. A. Rieffel, “The cancellation theorem for projective modules over irrational rotation  $C^*$ -algebras”, *Proc. London Math. Soc.* (3) **47**:2 (1983), 285–302. MR
- [57] M. A. Rieffel, “Noncommutative tori: a case study of noncommutative differentiable manifolds”, pp. 191–211 in *Geometric and topological invariants of elliptic operators* (Brunswick, ME, 1988), Contemp. Math. **105**, Amer. Math. Soc., 1990. MR
- [58] E. Riehl, *Categorical homotopy theory*, New Mathematical Monographs **24**, Cambridge University Press, 2014. MR
- [59] J. Rosenberg, *Algebraic  $K$ -theory and its applications*, Graduate Texts in Mathematics **147**, Springer, 1994. MR
- [60] G. Rudolph and M. Schmidt, *Differential geometry and mathematical physics, II: Fibre bundles, topology and gauge fields*, Springer, 2017. MR

- [61] M. Scharlemann, “Heegaard splittings of compact 3-manifolds”, pp. 921–953 in *Handbook of geometric topology*, North-Holland, Amsterdam, 2002. MR
- [62] E. H. Spanier, *Algebraic topology*, McGraw-Hill, 1966. Reprinted Springer, 1995. MR
- [63] R. G. Swan, “Vector bundles and projective modules”, *Trans. Amer. Math. Soc.* **105** (1962), 264–277. MR
- [64] M. Takesaki, *Theory of operator algebras I = Operator algebras and non-commutative geometry* **5**, Encyclopaedia of Mathematical Sciences **124**, Springer, 1979. MR
- [65] M. Thaddeus, “An introduction to the topology of the moduli space of stable bundles on a Riemann surface”, pp. 71–99 in *Geometry and physics* (Aarhus, 1995), Lecture Notes in Pure and Appl. Math. **184**, Dekker, New York, 1997. MR
- [66] A. Tikaradze, “On the Azumaya locus of almost commutative algebras”, *Proc. Amer. Math. Soc.* **139**:6 (2011), 1955–1960. MR
- [67] N. E. Wegge-Olsen, *K-theory and C\*-algebras: a friendly approach*, Oxford University Press, 1993. MR
- [68] S. Willard, *General topology*, Addison-Wesley, 1970. Reprinted Dover, Mineola, NY, 2004. MR

Received August 6, 2025.

ALEXANDRU CHIRVASITU  
DEPARTMENT OF MATHEMATICS  
UNIVERSITY AT BUFFALO  
UNITED STATES  
achirvas@buffalo.edu

# PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

[msp.org/pjm](http://msp.org/pjm)

## EDITORS

Don Blasius (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[blasius@math.ucla.edu](mailto:blasius@math.ucla.edu)

Matthias Aschenbrenner  
Fakultät für Mathematik  
Universität Wien  
Vienna, Austria  
[matthias.aschenbrenner@univie.ac.at](mailto:matthias.aschenbrenner@univie.ac.at)

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Atsushi Ichino  
Department of Mathematics  
Kyoto University  
Kyoto 606-8502, Japan  
[atsushi.ichino@gmail.com](mailto:atsushi.ichino@gmail.com)

Robert Lipshitz  
Department of Mathematics  
University of Oregon  
Eugene, OR 97403  
[lipshitz@uoregon.edu](mailto:lipshitz@uoregon.edu)

Kefeng Liu  
School of Sciences  
Chongqing University of Technology  
Chongqing 400054, China  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Sucharit Sarkar  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[sucharit@math.ucla.edu](mailto:sucharit@math.ucla.edu)

Dimitri Shlyakhtenko  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[shlyakht@ipam.ucla.edu](mailto:shlyakht@ipam.ucla.edu)

Ruixiang Zhang  
Department of Mathematics  
University of California  
Berkeley, CA 94720-3840  
[ruixiang@berkeley.edu](mailto:ruixiang@berkeley.edu)

## PRODUCTION

Silvio Levy, Scientific Editor, [production@msp.org](mailto:production@msp.org)

---

See inside back cover or [msp.org/pjm](http://msp.org/pjm) for submission instructions.

---

The subscription price for 2026 is US \$710/year for the electronic version, and \$965/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).


---

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004.

---

PJM peer review and production are managed by EditFLOW<sup>®</sup> from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2026 Mathematical Sciences Publishers

# PACIFIC JOURNAL OF MATHEMATICS

Volume 342 No. 1 May 2026

---

The $\bar{\partial}$ -problem on $Z(q)$ -domains	1
DEBRAJ CHAKRABARTI, PHILLIP S. HARRINGTON and ANDREW RAICH	
Polynomial identities and Azumaya loci for rational quantum spheres	37
ALEXANDRU CHIRVASITU	
From $i$ -boxes to signed words	63
ALESSANDRO CONTU, FAN QIN and QIAOLING WEI	
Near coincidences and nilpotent division fields	79
HARRIS B. DANIELS and JEREMY ROUSE	
The Cauchy problem for 1D nonlinear Schrödinger equations with repulsive delta potential for data in $L^p$ -based spaces	115
QINGQUAN DENG, PING LI and XIUHONG LONG	
Finite basis problem for involution semigroups of order four	163
MENG GAO, EDMOND W. H. LEE, YAN FENG LUO and WEN TING ZHANG	