

*Pacific
Journal of
Mathematics*

FROM i -BOXES TO SIGNED WORDS

ALESSANDRO CONTU, FAN QIN AND QIAOLING WEI

Volume 342 No. 1

May 2026

FROM i -BOXES TO SIGNED WORDS

ALESSANDRO CONTU, FAN QIN AND QIAOLING WEI

The combinatorics of i -boxes has recently been introduced by Kashiwara, Kim, Oh and Park in the study of cluster algebras arising from the representation theory of quantum affine algebras. In this article, we associate to each chain of i -boxes a signed word, which canonically determines a cluster seed, following Berenstein, Fomin and Zelevinsky. By bridging these two different languages, we are able to provide a quick solution to the problem of explicitly determining the exchange matrices associated with chains of i -boxes.

1. Introduction	63
2. Combinatorics of i -boxes	66
3. Signed words	71
4. Comparison of matrices	72
References	77

1. Introduction

1.1. Background. Cluster algebras were introduced by Fomin and Zelevinsky [FZ02]. These algebras possess distinguished elements called cluster monomials. One of the main reasons for the interest in the theory of cluster algebras is their unexpected emergence across diverse areas of mathematics. An illustrative case is the representation theory of quantum affine algebras, where cluster algebra structures are studied through the framework of monoidal categorification, starting with the inspiring work of Hernandez and Leclerc [HL10]. A monoidal category \mathcal{C} is a monoidal categorification of a cluster algebra \mathcal{A} if there is an isomorphism between the Grothendieck ring of \mathcal{C} and \mathcal{A} , such that the cluster monomials of \mathcal{A} correspond to certain simple objects of \mathcal{C} .

Let \mathfrak{g} be a simple complex finite-dimensional Lie algebra. In 2023, Kashiwara, Kim, Oh and Park [KKOP24] defined certain monoidal Serre subcategories $\mathcal{C}^{[a,b],\mathcal{D}_Q,w_0}$ of the module category of the quantum affine algebra of \mathfrak{g} , where $[a,b]$ denotes a possibly unbounded integer interval (for details; see [KKOP24, §4, §6]). To show that these categories are instances of monoidal categorification,

MSC2020: primary 13F60; secondary 17B37.

Keywords: cluster algebra, quantum affine algebra, monoidal categorification.

they introduced the combinatorics of *chains of \mathbf{i} -boxes* [KKOP24, §4, §5]. The definition of \mathbf{i} -boxes (Section 2) is based on the choice of an infinite sequence of indexes \mathbf{i} , and a chain of \mathbf{i} -boxes is a sequence of integer intervals satisfying certain technical conditions.

Let us fix a category $\mathcal{C}^{[a,b],\mathcal{D}_Q,\mathbb{w}_0}$. In this setting, the sequence \mathbf{i} is of the form $\widehat{\mathbf{w}}_0$, a particular sequence whose elements belong to the index set of a simply laced Dynkin diagram canonically associated to \mathfrak{g} . Kashiwara, Kim, Oh and Park associate an affine determinant module $M(\mathfrak{c})$ (a generalization of the Kirillov–Reshetikhin modules) to each \mathbf{i} -box \mathfrak{c} . Moreover, for each chain of \mathbf{i} -boxes $\mathfrak{C} = (\mathfrak{c}_i)$, they show the existence of a skew-symmetric exchange matrix $B(\mathfrak{C})$ such that, together with the representatives $([M(\mathfrak{c}_i)])_i$ of the modules associated to the \mathbf{i} -boxes of the chain, they give a seed for a cluster algebra structure of the Grothendieck ring of $\mathcal{C}^{[a,b],\mathcal{D}_Q,\mathbb{w}_0}$ [KKOP24, Theorem 8.1].

More precisely, when b is an integer, Kashiwara, Kim, Oh and Park start by explicitly providing the exchange matrix $B(\mathfrak{C}_{\underline{[a,b]}})$ associated to a specific chain of \mathbf{i} -boxes, denoted $\mathfrak{C}_{\underline{[a,b]}}$ (see Definition 2.5), generalizing a construction of Hernandez and Leclerc [HL11], who showed that the matrix $B(\mathfrak{C})$ can be obtained from the matrix $B(\mathfrak{C}_{\underline{[a,b]}})$ through a sequence of mutations. The case $b = \infty$ is treated through a limit procedure. At the end of [KKOP24], they state the problem of finding an explicit formula for all the matrices $B(\mathfrak{C})$. See Remark 2.11 for more details of this problem in terms of monoidal categories.

This problem is natural and fundamental for understanding the cluster structures appearing in the representation theory of quantum affine algebras. Recently, two solutions have been proposed:

- In [Con26], the first author proposed a solution translating the problem in a framework of additive categorification. Each exchange matrix $B(\mathfrak{C})$ is obtained as a submatrix of a square matrix $\overline{B}(\mathfrak{C})$, which, starting from the explicitly given matrix $\overline{B}(\mathfrak{C}_{\underline{[a,b]}})$, can be expressed via a matrix multiplication, known as Palu’s generalized mutation rule [Pal]:

$$\overline{B}(\mathfrak{C}) = P \overline{B}(\mathfrak{C}_{\underline{[a,b]}}) P^{-t},$$

where P is an invertible matrix obtained through the computation of indices of cluster-tilting objects of a suitable cluster category.

- In [KK24], Kashiwara and Kim work with sequences \mathbf{i} taking values in the index set of a generalized Cartan matrix C and with *maximal commuting families* of \mathbf{i} -boxes. Each chain of \mathbf{i} -boxes forms a maximal commuting family. For any such family \mathcal{F} , they define an $\mathcal{F} \times \mathcal{F}$ -skew-symmetrizable matrix $\widetilde{B}^{\text{KK}}(\mathcal{F})$ (see Section 4.1). When C is a symmetric Cartan matrix of

finite type and \mathbf{i} is of the form \widehat{w}_0 , each exchange matrix $B(\mathcal{C})$ can be obtained as a submatrix $B^{\text{KK}}(\mathcal{C})$ of the matrix $\widetilde{B}^{\text{KK}}(\mathcal{C})$.

The solution to the Kashiwara–Kim–Oh–Park problem in [Con26], although interesting for bridging monoidal and additive categorification of cluster algebras, is not as explicit, since it requires a multiplication of matrices. Kashiwara and Kim provide a direct formula relying on monoidal categorification and elaborate combinatorial machinery.

1.2. Main results. In this work, we propose an alternative and straightforward solution to Kashiwara, Kim, Oh and Park’s problem. The starting point is the combinatorics and the formalism of signed words. Recall that a signed word on an index set I is a sequence whose elements are of the form εh , where ε is in $\{\pm 1\}$ and h is in I . Assume that I is the index set of a generalized Cartan matrix. To each signed word \underline{h} , one can associate a seed $\mathbf{t}(\underline{h})$ following [BFZ05], which plays an important role in studying the cluster structures on double Bott–Samelson cells [SW21; Qin24b].

Let $B(\underline{h})$ be the corresponding exchange matrix. Assume that \mathbf{i} takes value in I and that b is in \mathbb{Z} . For each chain of \mathbf{i} -boxes \mathcal{C} associated to I , we define algorithmically a skew-symmetrizable matrix $B(\mathcal{C})$ in a similar fashion to [KKOP24]. Using the indices of the \mathbf{i} -boxes of the chain, we define a signed word $\underline{h}(\mathcal{C})$. Our main result states that the desired matrix $B(\mathcal{C})$ is given by the well-known matrix $B(\underline{h}(\mathcal{C}))$.

Theorem 1.1. *The matrix $B(\mathcal{C})$ equals $B(\underline{h}(\mathcal{C}))$.*

Therefore, by applying our main result to the setting where the Cartan matrix is symmetric of finite type and \mathbf{i} is of the form \widehat{w}_0 , we obtain a solution to the problem of Kashiwara Kim Oh Park, via a translation from the combinatorics of signed words to that of \mathbf{i} -boxes.

The outline of the proof of Theorem 1.1 is the following:

- (1) By direct comparison, we verify that the matrices $B(\mathcal{C}^{[a,b]})$ and $B(\underline{h}(\mathcal{C}^{[a,b]}))$ are equal.
- (2) Let \mathcal{C}' and \mathcal{C} be any chains related by a box move. We show that, if $B(\mathcal{C})$ equals $B(\underline{h}(\mathcal{C}))$, then $B(\mathcal{C}')$ also equals $B(\underline{h}(\mathcal{C}'))$.

Additionally, when I is the set of indices of a generalized Cartan matrix, we directly verify that our matrix $B(\underline{h}(\mathcal{C}))$ corresponds to Kashiwara–Kim’s matrix $B^{\text{KK}}(\mathcal{C})$ (Proposition 4.5).

Theorem 1.1 implies that the matrix $B(\mathcal{C})$ is independent of the choice of box moves from $\mathcal{C}^{[a,b]}$ to \mathcal{C} (Corollary 4.3). It also determines matrices in the case $b = +\infty$, as colimits of the matrices in the cases $b \in \mathbb{Z}$ (Corollary 4.4; see also [Qin24b]).

1.3. Notations and conventions. Choose any finite subset $K^{\text{ex}} \subset K$. For any $(K \times K^{\text{ex}})$ -matrix $Z = (Z_{ij})$ and permutation σ on K , we define the $(K \times \sigma(K^{\text{ex}}))$ -matrix σZ such that $(\sigma Z)_{\sigma i, \sigma j} := Z_{i, j}$. Let $\sigma_{j, j+1}$ denote the transposition $(j, j+1)$.

The matrix Z is called an exchange matrix if $Z_{ik} \in \mathbb{Z}$ for $(i, k) \in K \times K^{\text{ex}}$ and, moreover, it is skew-symmetrizable, i.e., there exists a diagonal matrix $D = (D_{kk})_{k \in K^{\text{ex}}}$ with diagonal entries $D_{kk} \in \mathbb{N}_{>0}$, called a skew-symmetrizer, such that $D_{ii}Z_{ik} = -D_{kk}Z_{ki}$, $\forall i, k \in K^{\text{ex}}$.

Let $[\]_+$ denote $\max(\ , 0)_+$. Let Z denote an exchange matrix. Following [FZ02], for any $k \in K^{\text{ex}}$, the mutation μ_k gives us a new exchange matrix $\mu_k Z$ such that

$$(\mu_k Z)_{ij} = \begin{cases} Z_{ij} + [Z_{ik}]_+ [Z_{kj}]_+ - [-Z_{ik}]_+ [-Z_{kj}]_+ & \text{if } i, j \neq k, \\ -Z_{ij} & \text{if } i = k \text{ or } j = k. \end{cases}$$

2. Combinatorics of i -boxes

In this section, following [KKOP24; KK24], we recall the definition and the properties of i -boxes.

For $a, b \in \mathbb{Z} \cup \{\pm\infty\}$, we write $[a, b]$ for the integer interval

$$[a, b] = \{k \in \mathbb{Z} \mid a \leq k \leq b\}.$$

The *length* l of an integer interval $[a, b]$ is defined as $l = \max(b - a + 1, 0)$.

Let I be a finite set of indices and Z be an integer interval. We write $\mathbf{i} = (i_Z)_{k \in Z}$ for a sequence of elements of I indexed by the elements of Z . For $s \in Z$ and $j \in I$, we introduce the symbols

$$(1) \quad \begin{aligned} s^+ &= \min(\{t \in Z \mid s < t, i_t = i_s\} \cup \{\infty\}), \\ s^- &= \max(\{t \in Z \mid s > t, i_t = i_s\} \cup \{-\infty\}), \\ s(j)^\oplus &= \min(\{t \in Z \mid s \leq t, i_t = j\} \cup \{\infty\}), \\ s(j)^\ominus &= \max(\{t \in Z \mid s \geq t, i_t = j\} \cup \{-\infty\}). \end{aligned}$$

Definition 2.1 [KK24, §2.1; Qin24a, §6.1]. A finite integer interval $[a, b]$ in Z is an i -box if $i_a = i_b$. For an i -box $[a, b]$, we define its *color* $i([a, b])$ as $i([a, b]) = i_a = i_b$ and its *i -cardinality* or *order* as the number of times that the index $i([a, b])$ appears in the subinterval of \mathbf{i} corresponding to $[a, b]$.

Remark 2.2. The intervals in Definition 2.1 are closely related to Kirillov–Reshetikhin modules of quantum affine algebras. They were called i -boxes in [KKOP24], which focused on the case $I = I_{\mathfrak{g}}$, $\mathbf{i} = \widehat{w}_0$ and $Z = \mathbb{Z}$, where $I_{\mathfrak{g}}$ is the set of Dynkin indices of a simply laced Lie algebra \mathfrak{g} and $\widehat{w}_0 = (i_k)_{k \in \mathbb{Z}}$ is an infinite sequence obtained from a reduced expression $\underline{w}_0 = s_{i_1} \dots s_{i_l}$ of the longest element of the

Weyl group of \mathfrak{g} by extending the sequence i_1, \dots, i_l via the rule

$$i_{k+l} = i_k^*,$$

where $(-)^*$ is the involution on the index set $I_{\mathfrak{g}}$ induced by $w_0(\alpha_i) = -\alpha_{i^*}$, for any simple root α_i , $i \in I_{\mathfrak{g}}$.

For a finite interval $[a, b]$ in Z , we define the \mathbf{i} -boxes

$$[a, b\} = [a, b(i_a)^\ominus] \text{ and } \{a, b] = [a(i_b)^\oplus, b].$$

In other terms, $[a, b\}$ and $\{a, b]$ are the largest \mathbf{i} -boxes contained in $[a, b]$ with colors i_a and i_b respectively. When we want to emphasize that an \mathbf{i} -box is of color j , we use the notation $[a, b]_j$.

Definition 2.3 [KKOP24, Definition 5.1]. Let l be in $\mathbb{N} \cup \{\infty\}$. A *chain* of \mathbf{i} -boxes of *length* l is a sequence of \mathbf{i} -boxes $\mathfrak{C} = (\mathfrak{c}_k)_{1 \leq k < l+1}$ satisfying the following conditions for any $1 \leq s < l+1$:

- (i) The union $[\tilde{a}_s, \tilde{b}_s] := \bigcup_{1 \leq k \leq s} \mathfrak{c}_k$ is an interval of length s .
- (ii) The \mathbf{i} -box \mathfrak{c}_s is the largest \mathbf{i} -box of color $i(\mathfrak{c}_s)$ contained in the interval $\bigcup_{1 \leq k \leq s} \mathfrak{c}_k$.

Condition (ii) implies $b_s \leq \tilde{b}_s < b_s^-$ and $a_s \geq \tilde{a}_s > a_s^-$. In addition, we have $[\tilde{a}_s, \tilde{b}_s] \subset [\tilde{a}_t, \tilde{b}_t]$ whenever $s < t$. We call the interval $\bigcup_{1 \leq k < l+1} \mathfrak{c}_k$ the *range* of the chain. For any $1 \leq s < l+1$, the sequence $(\mathfrak{c}_k)_{1 \leq k \leq s}$ is a chain of \mathbf{i} -boxes, called a *subchain* of \mathfrak{C} .

Remark 2.4 [KKOP24, §5]. To each chain of \mathbf{i} -boxes $\mathfrak{C} = (\mathfrak{c}_k)_{1 \leq k < l+1}$, we can bijectively associate a pair $(c, (E_k)_{1 \leq k < l})$, where $c \in Z$ and each E_k is a symbol in $\{L, R\}$, in the following way: let c be the integer such that $\{c\} = \mathfrak{c}_1$ and, for any $1 \leq k < l$, set

$$E_k = \begin{cases} L & \text{if } [\tilde{a}_{k+1}, \tilde{b}_{k+1}] = [\tilde{a}_k - 1, \tilde{b}_k], \\ R & \text{otherwise.} \end{cases}$$

In fact, given such a pair $(c, (E_k)_{1 \leq k < l})$, the associated chain of \mathbf{i} -boxes \mathfrak{C} can be recursively recovered as follows:

- $\mathfrak{c}_1 = \{c\}$;
- for any $2 \leq k < l+1$, we have

$$\mathfrak{c}_k = \begin{cases} [\tilde{a}_{k-1} - 1, \tilde{b}_{k-1}] & \text{if } E_{k-1} = L, \\ \{\tilde{a}_{k-1}, \tilde{b}_{k-1} + 1\} & \text{if } E_{k-1} = R. \end{cases}$$

We refer to $T = L$ (resp. $= R$) as a *left* (resp. *right*) *expansion operator* and to a sequence $(c, (E_k)_{1 \leq k < l})$ as a *rooted sequence of expansion operators*; see [Con26].

Definition 2.5. Let $[a, b]$ be an integer interval with $a \leq b$, $a \in \mathbb{Z} \sqcup \{-\infty\}$ and $b \in \mathbb{Z}$. Denote its length by $l = b - a + 1$. Following [KKOP24], we define $\mathfrak{C}^{[a, b]}$ as the chain of \mathbf{i} -boxes associated to the rooted sequence of expansion operators $(b, (E_k)_{1 \leq k < l})$, where $E_k = L$ for any k . Explicitly, the k -th \mathbf{i} -boxes are $c_k = [b - k + 1, b]$, $\forall k \in [1, l]$.

Definition 2.6 [KKOP24, §5]. Let $\mathfrak{C} = (c_k)$ be a chain of \mathbf{i} -boxes of length $l \leq \infty$ corresponding to a pair $(c, (E_k)_{1 \leq k < l})$.

- (i) For $1 \leq s < l$, the \mathbf{i} -box c_s is defined to be *movable* if $s = 1$ or $s \geq 2$ and $E_{s-1} \neq E_s$.
- (ii) For a movable \mathbf{i} -box c_s , the *box move* at s , denoted by ν_s , is the operation sending \mathfrak{C} to the chain $\nu_s \mathfrak{C}$, whose associated pair (c', E') is defined as follows:

$$c' = \begin{cases} c + 1 & \text{if } s = 1, E_1 = R, \\ c - 1 & \text{if } s = 1, E_1 = L, \\ c & \text{if } s > 1, \end{cases} \quad \text{and} \quad E'_k = \begin{cases} R & \text{if } E_k = L, k \in \{s-1, s\}, \\ L & \text{if } E_k = R, k \in \{s-1, s\}, \\ E_k & \text{if } k \notin \{s-1, s\}. \end{cases}$$

- (iii) We call a finite composition of box moves a *chain transformation*.

Example 2.7. Let $I = \{1, 2, 3\}$ be the set of Dynkin indices of a simple Lie algebra of type A_3 . Let $\underline{w}_0 = s_1 s_2 s_3 s_1 s_2 s_1$ be a reduced expression of the longest element Weyl group of type A and let \mathbf{i} be the sequence \widehat{w}_0 :

$$\widehat{w}_0 = \dots \underbrace{1, 3, 2, 3, 1, 2, 3, 1}_{[-3, 4]} \dots$$

The chain of \mathbf{i} -boxes $\mathfrak{C} = (c_k)_{1 \leq k \leq 8}$ of range $[-3, 4]$ associated to the rooted sequence of expansion operators $(4, (L, L, L, L, L, L, L))$ is given by

$$\begin{aligned} c_1 &= [4]_1, & c_2 &= [3]_3, & c_3 &= [2]_2, & c_4 &= [1, 4]_1, \\ c_5 &= [0, 3]_3, & c_6 &= [-1, 2]_2, & c_7 &= [-2, 3]_3, & c_8 &= [-3, 4]_1. \end{aligned}$$

Notice that the only movable \mathbf{i} -box in the chain \mathfrak{C} is c_1 . The box move at 1 sends \mathfrak{C} to the chain $\nu_1 \mathfrak{C} = (c'_k)_{1 \leq k \leq 6}$ associated to the sequence $(3, (R, L, L, L, L, L, L))$:

$$\begin{aligned} c'_1 &= [3]_3, & c'_2 &= [4]_1, & c'_3 &= [2]_2, & c'_4 &= [1, 4]_1, \\ c'_5 &= [0, 3]_3, & c'_6 &= [-1, 2]_2, & c'_7 &= [-2, 3]_3, & c'_8 &= [-3, 4]_1. \end{aligned}$$

Notice that the \mathbf{i} -box c'_2 is movable and that the associated box move sends $\nu_1 \mathfrak{C}$ to the chain associated to the sequence $(3, (L, R, L, L, L, L, L))$. Iterating this process, we see that, through a composition of box moves, the chain \mathfrak{C} is sent to

the chain $\tilde{\mathfrak{C}} = (\tilde{c}_k)_{1 \leq k \leq 8}$ associated to the sequence $(3, (L, L, L, L, L, L, R))$:

$$\begin{aligned} \tilde{c}_1 &= [3]_3, & \tilde{c}_2 &= [2]_2, & \tilde{c}_3 &= [1]_1, & \tilde{c}_4 &= [0, 3]_3, \\ \tilde{c}_5 &= [-1, 2]_2, & \tilde{c}_6 &= [-2, 3]_3, & \tilde{c}_7 &= [-3, 1]_1 & \tilde{c}_8 &= [-3, 4]_1. \end{aligned}$$

Remark 2.8 [KKOP24, Lemma 5.10; Con26, Remark 2.10]. Let $[a, b]$ be an integer interval with $b \in \mathbb{Z}$ and $a \in \mathbb{Z} \sqcup \{-\infty\}$. Then any two chains of \mathbf{i} -boxes of range $[a, b]$ are related by a chain transformation.

Remark 2.9. Let \mathfrak{C} and \mathfrak{C}' be two chain of \mathbf{i} -boxes. Assume that \mathfrak{C} and \mathfrak{C}' are related by a box move at $s \geq 1$.

- If $s \geq 2$, we have

$$i(c_{s+1}) = i(c'_s), \quad i(c_s) = i(c'_{s+1}) \quad \text{and} \quad i(c_k) = i(c'_k) \quad \text{for any } k \notin \{s, s+1\}.$$

- If $s = 1$, we have

$$i(c_2) = i(c'_1), \quad i(c'_2) = i(c_1) \quad \text{and} \quad i(c_k) = i(c'_k) \quad \text{for any } k \geq 3.$$

Definition 2.10 [KK24, Definition 2.13]. Let $\mathfrak{C} = (c_k)_k$ be a chain of \mathbf{i} -boxes with associated sequence of extension operators $(E_k)_k$. For any k , the *effective end* z of the \mathbf{i} -box $c_k = [x, y]$ is defined as

$$z = \begin{cases} y & \text{if } k = 1 \text{ or } E_{k-1} = R, \\ x & \text{if } k = 1 \text{ or } E_{k-1} = L. \end{cases}$$

2.1. The matrix associated to a chain of \mathbf{i} -boxes. From now on, let I be the set of indices of a generalized Cartan matrix $C = (c_{ij})_{i,j \in I}$. Let $[a, b]$ be an integer interval and let l be its length. Let $\mathfrak{C} = (c_k)_{k \in [1, l]}$ be a chain of \mathbf{i} -boxes of range $[a, b]$. In this subsection, following [KKOP24], under the assumption that b is an integer, we recursively associate to the chain \mathfrak{C} an exchange matrix $B(\mathfrak{C})$.

We introduce the sets

$$\begin{aligned} K(\mathfrak{C}) &= \begin{cases} \{1, \dots, l\} & \text{if } l < \infty, \\ \mathbb{N}_{\geq 1} & \text{if } l = \infty; \end{cases} \\ K(\mathfrak{C})^{\text{fr}} &= \{s \in K(\mathfrak{C}) \mid c_s = [a(i)^\oplus, b(i)^\ominus] \text{ for some } i \in I\}; \\ K(\mathfrak{C})^{\text{ex}} &= K(\mathfrak{C}) \setminus K(\mathfrak{C})^{\text{fr}}. \end{aligned}$$

For any $k \in K(\mathfrak{C})$, we define $k[1]$ as

$$k[1] = \min(\{k' \in [k+1, l] \mid c_{k'} \text{ has the same color of } c_k\} \sqcup \{+\infty\}).$$

Assume that b is in \mathbb{Z} . Following [KKOP24, §7.5], let

$$B(\mathfrak{C}_{[a, b]}^-) = (b_{jk})_{j \in K(\mathfrak{C}), k \in K(\mathfrak{C})^{\text{ex}}}$$

be the exchange matrix given by

$$b_{jk} = \begin{cases} 1 & \text{if } k = j[1], \\ -1 & \text{if } j = k[1], \\ c_{i_j, i_k} & \text{if } j < k < j[1] < k[1], \\ -c_{i_j, i_k} & \text{if } k < j < k[1] < j[1], \\ 0 & \text{otherwise.} \end{cases}$$

Although this construction was given in [KKOP24] only for Cartan matrices of ADE type (see also Remark 2.12), we apply the same formula in the more general setting considered here.

Next, for any chain of \mathbf{i} -boxes $\mathfrak{C} = (c_k)_{k \in [1, l]}$ of range $[a, b]$, fix a sequence of box moves ν_1, \dots, ν_N whose composition sends $\mathfrak{C}_{[a, b]}^{\underline{a}}$ to \mathfrak{C} . For $0 \leq s \leq N$, we write $\mathfrak{C}_s = (c_k^s)_{k \in [1, l]}$ for the chain of \mathbf{i} -boxes

$$\mathfrak{C}_s = \begin{cases} \mathfrak{C}_{[a, b]}^{\underline{a}} & \text{if } s = 0, \\ \nu_s \circ \dots \circ \nu_1 \mathfrak{C}_{[a, b]}^{\underline{a}} & \text{otherwise.} \end{cases}$$

Let B_0 be the matrix $B(\mathfrak{C}_{[a, b]}^{\underline{a}})$ and, for any $1 \leq s \leq N$, recursively define the exchange matrix B_s as follows:

- (i) If the \mathbf{i} -box c_{s+1}^{s-1} has the same color as c_s^{s-1} , then

$$B_s = \mu_s(B_{s-1}).$$

- (ii) If the \mathbf{i} -boxes c_{s+1}^{s-1} and c_s^{s-1} have different colors, then

$$B_s = \sigma_{s, s+1}(B_{s-1}).$$

Finally, we set $B(\mathfrak{C}) = B_N$. In the next section, we will show that the matrix $B(\mathfrak{C})$ does not depend on the choice of the composition of box-moves sending $\mathfrak{C}_{[a, b]}^{\underline{a}}$ to \mathfrak{C} .

Remark 2.11. Let \mathfrak{g} be a finite-dimensional simple complex Lie algebra and let \mathfrak{g} be the Lie algebra of simply laced type associated to the unfolding of the Dynkin diagram of \mathfrak{g} [FO21]. Assume that b is in $\mathbb{Z} \cup \{\infty\}$ and that $\mathbf{i} = (i_k)_{k \in [a, b]}$ is a subsequence of the sequence $\widehat{w}_0 = (i_k)_{k \in \mathbb{Z}}$, for a certain reduced expression \underline{w}_0 of the longest element of the Weyl group of \mathfrak{g} . In [KKOP24], Kashiwara, Kim, Oh and Park associate a *monoidal seed* (see [KKOP24, Definition 7.2] for the terminology) to each chain of \mathbf{i} -boxes \mathfrak{C} , defining a family of *commuting real prime* modules $M(\mathfrak{C})$ and showing the existence of a companion exchange matrix $B(\mathfrak{C})$. To do this, when b is an integer they use the above procedure [KKOP24, Proposition 8.11], and when $b = +\infty$ they define $B(\mathfrak{C})$ as a colimit of the matrices associated to the subchain of \mathfrak{C} . By [KKOP24, Proposition 7.14, Lemma 7.16], this operation is well defined. It follows from this last proposition that the exchange matrix $B(\mathfrak{C})$ does not depend on the choice of the sequence of box moves sending $\mathfrak{C}_{[a, b]}^{\underline{a}}$ to \mathfrak{C} .

Nevertheless, [KKOP24] does not provide an explicit description of the coefficients of $B(\mathfrak{C})$, whose determination is stated as an open problem.

3. Signed words

Recall that $C = (c_{ij})_{i,j \in I}$ is a generalized Cartan matrix. Let l be in $\mathbb{N} \sqcup \{+\infty\}$.

Definition 3.1. A signed word on the index set I is a sequence $\underline{\mathbf{h}} = (\varepsilon_k h_k)_{1 \leq k < l+1}$ such that, for any k , $\varepsilon_k \in \{\pm 1\}$ and $h_k \in I$. Denote $\mathbf{h}_k = \varepsilon_k h_k$ and $|\mathbf{h}_k| = h_k$.

We introduce the sets

$$\begin{aligned} K(\underline{\mathbf{h}}) &= [1, l]; \\ K(\underline{\mathbf{h}})^{\text{ex}} &= \{s \in K(\underline{\mathbf{h}}) \mid \exists t \in [s+1, l], h_t = h_s\}; \\ K(\underline{\mathbf{h}})^{\text{fr}} &= K(\mathfrak{C}) \setminus K(\underline{\mathbf{h}})^{\text{ex}}. \end{aligned}$$

For any $k \in K(\underline{\mathbf{h}})$, we define $k[1]$ as

$$(2) \quad k[1] = \min(\{k' \in [k+1, l] \mid |\mathbf{h}_{k'}| = |\mathbf{h}_k|\} \sqcup \{+\infty\}).$$

Note that $k[1]$ in (2) and s^+ in (1) should not be confused.

Following [BFZ05; SW21; Qin24a, (6.1)], for any signed word $\underline{\mathbf{h}}$, we define $\tilde{B}(\underline{\mathbf{h}}) = (\tilde{b}_{jk})_{j,k \in K(\underline{\mathbf{h}})}$ by

$$(3) \quad \tilde{b}_{jk} = \begin{cases} \varepsilon_k & \text{if } k = j[1], \\ -\varepsilon_j & \text{if } j = k[1], \\ \varepsilon_k c_{|\mathbf{h}_j|, |\mathbf{h}_k|} & \text{if } \varepsilon_{j[1]} = \varepsilon_k, j < k < j[1] < k[1], \\ \varepsilon_k c_{|\mathbf{h}_j|, |\mathbf{h}_k|} & \text{if } \varepsilon_k = -\varepsilon_{k[1]}, j < k < k[1] < j[1], \\ -\varepsilon_j c_{|\mathbf{h}_j|, |\mathbf{h}_k|} & \text{if } \varepsilon_{k[1]} = \varepsilon_j, k < j < k[1] < j[1], \\ -\varepsilon_j c_{|\mathbf{h}_j|, |\mathbf{h}_k|} & \text{if } \varepsilon_j = -\varepsilon_{j[1]}, k < j < j[1] < k[1], \\ 0 & \text{otherwise.} \end{cases}$$

It has a skew-symmetrizer $\tilde{D}(\underline{\mathbf{h}})$, which is a diagonal matrix with diagonal entries $\tilde{D}_{ii} = h_i$. Let $B(\underline{\mathbf{h}})$ denote its $K(\underline{\mathbf{h}}) \times K(\underline{\mathbf{h}})^{\text{ex}}$ -submatrix.

From now on, we will always assume that $B(\underline{\mathbf{h}})$ is a locally finite matrix, i.e., for any j , only finitely many b_{jk} are nonzero and, for any k , only finitely many b_{jk} are nonzero. This assumption allows us to extend results in [SW21; Qin24a] for $l \in \mathbb{N}$ to the case $l = +\infty$ as in [Qin24b].

Let $\underline{\mathbf{h}}_{[j,k]}$ denote the sequence $(\mathbf{h}_s)_{s \in [j,k]}$.

Definition 3.2 [SW21, Section 2.3; Qin24b, Section 3.2]. Let $\underline{\mathbf{h}} = (\mathbf{h}_k)_{k \in [1,l]}$ be a signed word.

- The *left reflection* of $\underline{\mathbf{h}}$ is the operation sending $\underline{\mathbf{h}}$ to the signed word $\underline{\mathbf{h}}' = (-\mathbf{h}_1, \underline{\mathbf{h}}_{[2,l]})$.

- Let $j \in [1, l-1]$ be such that \underline{h}_j and \underline{h}_{j+1} have different signs. Then the *flip* of \underline{h} at j is the operation sending \underline{h} to the signed word $\underline{h}' = (\underline{h}_{[1, j-1]}, \underline{h}_{j+1}, \underline{h}_j, \underline{h}_{[j+2, l]})$.

Proposition 3.3 [SW21, Proposition 3.7; Qin24b, Section 3.2]. *Let $\underline{h} = (\underline{h}_k)_{k \in [1, l]}$ and $\underline{h}' = (\underline{h}'_k)_{k \in [1, l]}$ be two signed words.*

- (1) *If \underline{h}' is obtained from \underline{h} via a left reflection, then $B(\underline{h}') = B(\underline{h})$.*
- (2) *If \underline{h}' is obtained from \underline{h} via a flip at j , then*

$$B(\underline{h}') = \begin{cases} \sigma_{j, j+1} B(\underline{h}) & \text{if } h_j \neq h'_j, \\ \mu_j(B(\underline{h})) & \text{if } h_j = h'_j. \end{cases}$$

4. Comparison of matrices

Recall that I denotes the index set of a generalized Cartan matrix C . Let $[a, b]$ be an interval and $\mathbf{i} = (i_k)_{k \in [a, b]}$ a sequence of elements of I . In the following, we will consider chains of \mathbf{i} -boxes defined with respect to the sequence \mathbf{i} . Let $\mathcal{C} = (c_k)_{k \in [1, l]}$ be a chain of \mathbf{i} -boxes on $[a, b]$ and let $(E_k)_{1 \leq k < l}$ be the associated sequence of expansion operators. For any $1 \leq k < l + 1$, we write $c_k = [a_k, b_k]$. We associate to \mathcal{C} a signed word $\underline{h}(\mathcal{C}) = (\mathbf{h}_k)_{1 \leq k < l+1}$ as follows:

- We set \mathbf{h}_1 equal to the color h_1 of the \mathbf{i} -box c_1 .
- For any $2 \leq k < l + 1$, we set $\mathbf{h}_k = \varepsilon_k h_k$ where h_k is the color of the \mathbf{i} -box c_k and the sign ε_k is defined by

$$\begin{cases} 1 & \text{if } E_{k-1} = L, \\ -1 & \text{if } E_{k-1} = R. \end{cases}$$

It follows from our construction that $K(\mathcal{C}) = K(\underline{h}(\mathcal{C}))$, $K(\mathcal{C})^{\text{ex}} = K(\underline{h}(\mathcal{C}))^{\text{ex}}$ and $K(\mathcal{C})^{\text{fr}} = K(\underline{h}(\mathcal{C}))^{\text{fr}}$, and the definition of $k[1]$ becomes identical. We will simply denote $K(\mathcal{C})$ by K below. Note that different chains of \mathbf{i} -boxes correspond to different signed words, but not every signed word comes from a chain of \mathbf{i} -boxes.

Remark 4.1. Assume $k'[1] = k$ for some $k', k \in K = [1, \ell]$, and set $k' = k[-1]$. By our constructions, the following properties (1) to (6) are equivalent in each case:

- | | |
|------------------------------------------------|------------------------------------------------|
| (1) $\varepsilon_k = 1$ | (1) $\varepsilon_k = -1$ |
| (2) $E_{k-1} = L$ | (2) $E_{k-1} = R$ |
| (3) a_k is the effective end of $[a_k, b_k]$ | (3) b_k is the effective end of $[a_k, b_k]$ |
| (4) $\tilde{a}_k = a_k$ | (4) $\tilde{b}_k = b_k$ |
| (5) $a_k = a_{k[-1]}^-$ | (5) $b_k = b_{k[-1]}^+$ |
| (6) $b_k = b_{k[-1]}$ | (6) $a_k = a_{k[-1]}$ |

Assume that b is an integer. In the following, we want to show that the matrices $B(\mathcal{C})$ and $B(\underline{\mathbf{h}}(\mathcal{C}))$ are equal, thus providing a solution to the Kashiwara–Kim–Oh–Park problem.

To start with, consider the chain of \mathbf{i} -boxes $\mathcal{C}_{-}^{[a,b]}$. Then the signed word $\underline{\mathbf{h}}(\mathcal{C}_{-}^{[a,b]}) = (\mathbf{h}_k)_{k \in [1,l]}$ is given by

$$\mathbf{h}_k = i_{b-k+1}.$$

Then the matrix $B(\underline{\mathbf{h}}(\mathcal{C}_{-}^{[a,b]})) = (\tilde{b}_{jk})_{j \in K, k \in K^{\text{ex}}}$ simplifies to

$$\tilde{b}_{jk} = \begin{cases} 1 & \text{if } k = j[1], \\ -1 & \text{if } j = k[1], \\ c_{|\mathbf{h}_j|, |\mathbf{h}_k|} & \text{if } j < k < j[1] < k[1], \\ -c_{|\mathbf{h}_j|, |\mathbf{h}_k|} & \text{if } k < j < k[1] < j[1]. \end{cases}$$

Therefore, $B(\underline{\mathbf{h}}(\mathcal{C}_{-}^{[a,b]}))$ coincides with the matrix $B(\mathcal{C}_{-}^{[a,b]})$.

Proposition 4.2. *Let $\mathcal{C} = (c_k)_k$ be any chain of \mathbf{i} -boxes of range $[a, b]$. Suppose that the matrix $B(\mathcal{C})$ equals $B(\underline{\mathbf{h}}(\mathcal{C}))$. Then for any movable \mathbf{i} -box c_s of \mathcal{C} , we also have $B(v_s(\mathcal{C})) = B(\underline{\mathbf{h}}(v_s(\mathcal{C})))$.*

Proof. Recall that we write $\underline{\mathbf{h}}(\mathcal{C}) = (\mathbf{h}_k)_k = (\varepsilon_k h_k)_k$. Denote the sequence of expansion operators associated to \mathcal{C} by $(E_k)_k$.

Suppose that s is greater than 1. Since c_s is movable, we have $E_{s-1} \neq E_s$. Then, at the level of the associated signed word, we have $\varepsilon_s \neq \varepsilon_{s+1}$. Moreover, by Remark 2.9, the signed word associated to $v_s(\mathcal{C})$ is $\underline{\mathbf{h}}(v_s(\mathcal{C})) = (\mathbf{h}'_k)_k$, where

$$\mathbf{h}'_k = \begin{cases} \mathbf{h}_k & \text{if } k \notin \{s, s+1\}, \\ \mathbf{h}_{s+1} & \text{if } k = s, \\ \mathbf{h}_s & \text{if } k = s+1. \end{cases}$$

In particular, the signed word $\underline{\mathbf{h}}(v_s(\mathcal{C}))$ is obtained from $\underline{\mathbf{h}}(\mathcal{C})$ through a flip move at s . We have two cases:

- (1) If the \mathbf{i} -boxes c_s and c_{s+1} have the same color, then $|\mathbf{h}_s| = |\mathbf{h}_{s+1}|$. Therefore, the matrices $B(\underline{\mathbf{h}}(v_s(\mathcal{C})))$ and $B(v_s(\mathcal{C}))$ are equal, since they are the mutation of the matrix $B(\underline{\mathbf{h}}(\mathcal{C})) = B(\mathcal{C})$ at s .
- (2) If the \mathbf{i} -boxes c_s and c_{s+1} have different color, then $|\mathbf{h}_s| \neq |\mathbf{h}_{s+1}|$. Therefore, the matrices $B(\underline{\mathbf{h}}(v_s(\mathcal{C})))$ and $B(v_s(\mathcal{C}))$ are equal, since they are obtained from the matrix $B(\underline{\mathbf{h}}(\mathcal{C})) = B(\mathcal{C})$ by applying σ_s .

If $s = 1$, by Remark 2.9, the signed word $\underline{h}(v_s(\mathfrak{C})) = (\mathbf{h}'_k)_k$ associated to $v_s(\mathfrak{C})$ is given by

$$\mathbf{h}'_k = \begin{cases} \mathbf{h}_k & \text{if } k \geq 3, \\ |\mathbf{h}_2| & \text{if } k = 1, \\ -\varepsilon_2 \mathbf{h}_1 & \text{if } k = 2. \end{cases}$$

Therefore, up to a left reflection which does not change the B -matrix, the signed word $\underline{h}(v_s(\mathfrak{C}))$ is obtained from $\underline{h}(\mathfrak{C})$ through a flip move, and we can conclude as above that $B(v_s(\mathfrak{C})) = B(\underline{h}(v_s(\mathfrak{C})))$. \square

Proof of Theorem 1.1. Recall that \mathfrak{C} could be obtained from $\mathfrak{C}^{[a,b]}$ by finite many box moves, and $B(\mathfrak{C}^{[a,b]}) = B(\underline{h}(\mathfrak{C}^{[a,b]}))$. Applying Proposition 4.2 repeatedly, we obtain $B(\mathfrak{C}) = B(\underline{h}(\mathfrak{C}))$. \square

Corollary 4.3. *The matrix $B(\mathfrak{C})$ does not depend on the choice of the chain transformation sending $\mathfrak{C}^{[a,b]}$ to \mathfrak{C} .*

Finally, assume that $b = +\infty$. Consider, for $s \geq 1$, the subchains $\mathfrak{C}_s = (c_k)_{1 \leq k \leq s}$ of \mathfrak{C} . As a corollary of Proposition 4.2, the coefficients of the exchange matrices $B(\mathfrak{C}_s)$ stabilize:

Corollary 4.4. *For any $s \geq 1$, if $(i, j) \in K(\mathfrak{C}_s) \times K^{\text{ex}}(\mathfrak{C}_s)$, then $B(\mathfrak{C}_t)_{ij} = B(\mathfrak{C}_s)_{ij}$ for any $t \geq s$.*

Therefore, we can define the exchange matrix $B(\mathfrak{C})$ as the colimit of the matrices $B(\mathfrak{C}_s)$. In other words, we have

$$B(\mathfrak{C})|_{K(\mathfrak{C}_s) \times K^{\text{ex}}(\mathfrak{C}_s)} = B(\mathfrak{C}_s) \quad \text{for any } s \geq 1.$$

4.1. Comparison of $B(\underline{h}(\mathfrak{C}))$ with Kashiwara–Kim’s matrix. We still write $[a, b]$ for an interval and $\mathfrak{C} = (c_k)_{k \in [1, l]}$ for a chain of \mathbf{i} -boxes $c_k = [a_k, b_k]$ of range $[a, b]$. For $1 \leq k \leq l$, h_k is the color of the \mathbf{i} -box c_k . Let \tilde{D} be the diagonal $l \times l$ -matrix with diagonal entries $d_1 = d_{h_1}, \dots, d_l = d_{h_l}$, where the $(d_i)_{i \in I}$ are the diagonal entries of the minimal symmetrizer of the generalized Cartan matrix C . Following [KK24], we define $\tilde{B}^{\text{KK}}(\mathfrak{C}) = (b_{kk'}^{\text{KK}})_{k, k' \in K(\mathfrak{C})}$ as the skew-symmetrizable $l \times l$ -matrix with skew-symmetrizer \tilde{D} and whose positive entries are given as follows:

- (4) $b_{jk}^{\text{KK}} = \begin{cases} 1 & \text{if } (a_j = a_k \text{ and } b_k = b_j^-) \text{ or } (b_j = b_k \text{ and } a_k = a_j^-), \\ -c_{h_j, h_k} & \text{if } c_{h_j, h_k} < 0 \text{ and one of the following conditions holds:} \end{cases}$
- (a) $[a_j, b_j^+] \in \mathfrak{C}$, a_j is the effective end of $[a_j, b_j]$, $a_k^- < a_j < a_k < b_k < b_j^+ < b_k^+$,
 - (b) $[a_j, b_j^+] \in \mathfrak{C}$, b_k is the effective end of $[a_k, b_k]$, $a_k^- < a_j < b_j < b_k < b_j^+ < b_k^+$,
 - (c) $[a_k^-, b_k] \in \mathfrak{C}$, b_k is the effective end of $[a_k, b_k]$, $a_j^- < a_k^- < a_j < b_j < b_k < b_j^+$,
 - (d) $[a_k^-, b_k] \in \mathfrak{C}$, a_j is the effective end of $[a_j, b_j]$, $a_j^- < a_k^- < a_j < a_k < b_k < b_j^+$.

Let $B^{\text{KK}}(\mathfrak{C})$ denote the restriction of the matrix $\tilde{B}(\mathfrak{C})$ to the indexes $K(\mathfrak{C}) \times K^{\text{ex}}(\mathfrak{C})$. When the Cartan matrix is symmetric of finite type and \underline{i} is of the form \hat{w}_0 , it is proved in [KK24] that $B^{\text{KK}}(\mathfrak{C}) = B(\mathfrak{C})$. Then Theorem 1.1 implies that $B^{\text{KK}}(\mathfrak{C}) = B(\underline{h}(\mathfrak{C}))$. In the next proposition, in the general case where C is a generalized Cartan matrix, we compare them directly without using Theorem 1.1.

Proposition 4.5. *The matrix $B^{\text{KK}}(\mathfrak{C})$ is equal to the matrix $B(\underline{h}(\mathfrak{C}))$.*

Proof. $\tilde{B}^{\text{KK}}(\mathfrak{C})$ and $\tilde{B}(\underline{h}(\mathfrak{C}))$ have the same skew-symmetrizer. It suffices to prove that the positive (j, k) -entries are the same, where at least one of j, k belongs to K^{ex} .

Let $j, k \in [1, l]$ be such that $c_{h_j, h_k} < 0$. Recall from (4) the conditions (a), (b), (c) and (d) which characterize such entries. Define another four conditions:

- (i) $\varepsilon_{j[1]} = \varepsilon_k = -1$, $j < k < j[1] < k[1]$;
- (ii) $\varepsilon_k = -\varepsilon_{k[1]} = -1$, $j < k < k[1] < j[1]$;
- (iii) $\varepsilon_{k[1]} = \varepsilon_j = 1$, $k < j < k[1] < j[1]$;
- (iv) $\varepsilon_j = -\varepsilon_{j[1]} = 1$, $k < j < j[1] < k[1]$.

If any of these conditions is true, then $b_{jk} = -c_{h_j, h_k}$. Moreover, conditions (i), (ii), (iii) and (iv) are mutually exclusive, while (a) (resp. (c)) and (b) (resp. (d)) can hold at the same time.

Step 1. First, assume that (i) holds. Using Remark 4.1, we have:

- $\varepsilon_k = -1$ is equivalent to b_k being the effective end of $[a_k, b_k]$.
- $\varepsilon_{j[1]} = -1$ is equivalent to $[a_j, b_j^+]$ belonging to \mathfrak{C} , i.e., $\mathfrak{c}_{j[1]} = [a_j, b_j^+]$.
- $j < k$ implies $[a_j, b_j] \subset [\tilde{a}_k, \tilde{a}_k]$. Then we deduce $a_k^- < \tilde{a}_k < a_j$.
- $j < k$ and $\varepsilon_k = -1$ imply that $b_j < \tilde{b}_k = b_k$.
- $k < j[1]$ and $\varepsilon_{j[1]} = -1$ imply that $b_k < \tilde{b}_{j[1]} = b_j^+$.
- $j[1] < k[1]$ and $\varepsilon_{j[1]} = -1$ implies $b_j^+ < b_k^+$. This claim follows from the fact that $b_j^+ = b_{j[1]} \leq \tilde{b}_{k[1]}$ and from the construction of $\mathfrak{c}_{k[1]}$, which implies that $\tilde{b}_{k[1]} \leq b_k^+$.

Therefore, (i) implies (b). Now, assume that (b) holds. Then we have the effective ends $b_k = \tilde{b}_k$ and $b_{j[1]} = \tilde{b}_{j[1]} = b_j^+$. Moreover, $\mathfrak{c}_{j[1]} = [a_j, b_j^+]$. We claim that (b) implies one of (i)–(iv).

First, we cannot have $j[1] < k$ since $b_j^+ = \tilde{b}_{j[1]} > \tilde{b}_k = b_k$. In addition, we cannot have $k[1] < j$. Otherwise, it would follow that $\mathfrak{c}_{k[1]} \subset [\tilde{a}_j, \tilde{b}_j]$, so that $b_{k[1]} \leq \tilde{b}_j < b_j^+$ and $a_{k[1]} \geq \tilde{a}_j > a_j^-$. But since we have either $b_{k[1]} = b_k^+$ or $a_{k[1]} = a_k^-$, both cases lead to contradiction with (b).

If $j < k < j[1] < k[1]$, we are in case (i).

If $j < k < k[1] < j[1]$, we claim that $\varepsilon_{k[1]} = 1$, i.e, we are in case (ii). To see this, assume $\varepsilon_{k[1]} = -1$. Then $\tilde{b}_{k[1]} = b_k^+ > b_j^+ = \tilde{b}_{j[1]}$, which contradicts $k[1] < j[1]$.

If $k < j < k[1] < j[1]$, as before, $k[1] < j[1]$ and (b) imply $\varepsilon_{k[1]} = 1$. We claim that $\varepsilon_j = 1$, i.e, we are in case (iii). To see this, note that $b_j < b_k$ but $j > k$, so b_j is not the effective end of \mathfrak{c}_j , i.e, $\varepsilon_j = 1$. The claim follows.

If $k < j < j[1] < k[1]$, as before, $j > k$ and $b_j < b_k$ imply $\varepsilon_j = 1$. So we are in case (iv).

Step 2. Similarly, assume that (ii) holds. We have:

- $\varepsilon_k = -1$ is equivalent to b_k being the effective end of $[a_k, b_k]$.
- $\varepsilon_{k[1]} = 1$ is equivalent to $[a_k^-, b_k]$ belonging to \mathfrak{C} , i.e., $\mathfrak{c}_{k[1]} = [a_k^-, b_k]$.
- $j < k$ implies $a_k^- < \tilde{a}_k < a_j$.
- $j < k$ and $\varepsilon_k = -1$ imply that $b_j < \tilde{b}_k = b_k$.
- $k[1] < j[1]$ and $\varepsilon_{k[1]} = 1$ imply $b_k < b_j^+$ and $a_j^- < a_k^-$. This follows from the fact that $a_k^- = a_{k[1]} \geq \tilde{a}_{j[1]} \geq a_j^-$ and $b_k = b_{k[1]} \leq \tilde{b}_{j[1]} \leq b_j^+$.

Therefore, (ii) implies (c).

Now assume that (c) holds. Then we have the effective ends $b_k = \tilde{b}_k$ and $a_{k[1]} = \tilde{a}_{k[1]} = a_k^-$. Moreover, $\mathfrak{c}_{k[1]} = [a_k^-, b_k]$. We claim that (c) implies one of (i)–(iv).

First, we cannot have $k[1] < j$. In fact, since $\mathfrak{c}_j = [a_j, b_j] \subset [a_k^-, b_k] = \mathfrak{c}_{k[1]}$, if $j > k[1]$, none of a_j, b_j could be an effective end. In addition, we cannot have $j[1] < k$. Otherwise, it would follow that $\mathfrak{c}_{j[1]} \subset [\tilde{a}_k, \tilde{b}_k]$, thus $b_{j[1]} \leq \tilde{b}_k = b_k$ and $a_{j[1]} \geq \tilde{a}_k > a_k^-$. Since either $b_{j[1]} = b_j^+$ or $a_{j[1]} = a_j^-$ holds, and both contradict (c), we are led to a contradiction.

If $j < k < k[1] < j[1]$, we are in case (ii).

If $j < k < j[1] < k[1]$, we claim that $\varepsilon_{j[1]} = -1$, i.e, we are in case (i). To see this, assume $\varepsilon_{j[1]} = 1$. Then $\tilde{a}_{j[1]} = a_j^- < a_k^- = \tilde{a}_{k[1]}$, which contradicts $j[1] < k[1]$.

If $k < j < j[1] < k[1]$, as before, $j[1] < k[1]$ and (c) imply $\varepsilon_{j[1]} = -1$. Moreover, $b_j < b_k$ and $j > k$ imply that b_j is not the effective end of \mathfrak{c}_j , i.e., $\varepsilon_j = 1$. So we are in case (iv).

If $k < j < k[1] < j[1]$, as before, $j > k$ and $b_j < b_k$ imply $\varepsilon_j = 1$. So we are in case (iii).

Step 3. Let us deduce the remaining cases from the previous steps. Let σ denote the order reversing automorphism on \mathbb{Z} such that $\sigma(x) = -x$. Consider the word $\mathbf{i}' = (i'_k) := (i_{\sigma k})$ and the chain of \mathbf{i}' -boxes $\mathfrak{C}' = (\mathfrak{c}'_s)_{s \in [1, l]}$ such that $\mathfrak{c}'_s := [a'_s, b'_s] := [\sigma b_s, \sigma a_s]$. Then, for any j, k in K , the signed word $\underline{\mathbf{h}}(\mathfrak{C})$ satisfies conditions (i), (ii), (iii), or (iv) if and only if, respectively, the signed word $\underline{\mathbf{h}}(\mathfrak{C}') = (\varepsilon'_i h_i)$ satisfies conditions (iii), (iv), (i), or (ii) with j and k swapped:

- (iii) $\varepsilon_{j[1]} = \varepsilon_k = 1$, $j < k < j[1] < k[1]$,
- (iv) $\varepsilon_k = -\varepsilon_{k[1]} = 1$, $j < k < k[1] < j[1]$,
 - (i) $\varepsilon_{k[1]} = \varepsilon_j = -1$, $k < j < k[1] < j[1]$,
 - (ii) $\varepsilon_j = -\varepsilon_{j[1]} = -1$, $k < j < j[1] < k[1]$.

Similarly, for any j, k in K , the chain of \mathbf{i} -boxes \mathfrak{C} satisfies the conditions (a), (b), (c), or (d) if and only if, respectively, the chain of \mathbf{i} -boxes \mathfrak{C}' satisfies the conditions (c), (d), (a), or (b) with j and k swapped:

- (c) $[(a')_j^-, (b')_j] \in \mathfrak{C}'$, $(b')_j$ is the effective end of $[(a')_j, (b')_j]$, $(b')_k^+ > (b')_j > (b')_k > (a')_k > (a')_j^- > (a')_k^-$;
- (d) $[(a')_j^-, (b')_j] \in \mathfrak{C}'$, $(a')_k$ is the effective end of $[(a')_k, (b')_k]$, $(b')_k^+ > (b')_j > (a')_j > (a')_k > (a')_j^- > (a')_k^-$;
- (a) $[(a')_k, b_k^+] \in \mathfrak{C}'$, $(a')_k$ is the effective end of $[(a')_k, (b')_k]$, $(b')_j^+ > (b')_k^+ > (b')_j > (a')_j > (a')_k > (a')_j^-$;
- (b) $[(a')_k, (b')_k^+] \in \mathfrak{C}'$, $(b')_j$ is the effective end of $[(a')_j, (b')_j]$, $(b')_j^+ > (b')_k^+ > (b')_j > (b')_k > (a')_k > (a')_j^-$.

Combining with the results in Steps 1 and 2, we obtain that (iii) implies (d), that (iv) implies (a) and that, if any among (a), and (d) holds, then we are in one of the cases (i)–(iv).

Finally, we obtain the desired claim by comparing the explicit formula for the entries of the matrices, using the results of the previous steps. \square

Funding. This work was supported by the National Natural Science Foundation of China [grant number 12422102 to Fan Qin].

References

- [BFZ05] A. Berenstein, S. Fomin, and A. Zelevinsky, “Cluster algebras, III: Upper bounds and double Bruhat cells”, *Duke Math. J.* **126**:1 (2005), 1–52. MR
- [Con26] A. Contu, “Solution of a problem in monoidal categorification by additive categorification”, *J. Algebra* **691** (2026), 128–185. MR
- [FO21] R. Fujita and S.-j. Oh, “Q-data and representation theory of untwisted quantum affine algebras”, *Comm. Math. Phys.* **384**:2 (2021), 1351–1407.
- [FZ02] S. Fomin and A. Zelevinsky, “Cluster algebras, I: Foundations”, *J. Amer. Math. Soc.* **15**:2 (2002), 497–529. MR
- [HL10] D. Hernandez and B. Leclerc, “Cluster algebras and quantum affine algebras”, *Duke Math. J.* **154**:2 (2010), 265–341. MR
- [HL11] D. Hernandez and B. Leclerc, “Quantum Grothendieck rings and derived Hall algebras”, 2011. arXiv 1109.0862v1
- [KK24] M. Kashiwara and M. Kim, “Exchange matrices of I-boxes”, *Adv. Math.* **485** (2026), art. id. 110711, 67 pp. MR

- [KKOP24] M. Kashiwara, M. Kim, S.-j. Oh, and E. Park, “Monoidal categorification and quantum affine algebras, II”, *Invent. Math.* **236**:2 (2024), 837–924.
- [Pal] Y. Palu, “Grothendieck group and generalized mutation rule for 2-Calabi–Yau triangulated categories”, *J. Pure Appl. Algebra* **213**:7 (2009), 1438–1449. MR
- [Qin24a] F. Qin, “Analogues of the dual canonical bases for cluster algebras from Lie theory”, preprint, 2024. arXiv 2407.02480
- [Qin24b] F. Qin, “Based cluster algebras of infinite rank”, preprint, 2024. arXiv 2409.02881
- [SW21] L. Shen and D. Weng, “Cluster structures on double Bott–Samelson cells”, *Forum Math. Sigma* **9** (2021), art. id. e66, 89 pp. MR

Received August 17, 2025.

ALESSANDRO CONTU
RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES
KYOTO UNIVERSITY
KYOTO
JAPAN
alessandro20.contu@gmail.com

FAN QIN
SCHOOL OF MATHEMATICAL SCIENCES
BEIJING NORMAL UNIVERSITY
BEIJING, 100875
CHINA
qin.fan.math@gmail.com

QIAOLING WEI
CAPITAL NORMAL UNIVERSITY
BEIJING
CHINA
wql03ster@gmail.com

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Fakultät für Mathematik
Universität Wien
Vienna, Austria
matthias.aschenbrenner@univie.ac.at

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Atsushi Ichino
Department of Mathematics
Kyoto University
Kyoto 606-8502, Japan
atsushi.ichino@gmail.com

Robert Lipshitz
Department of Mathematics
University of Oregon
Eugene, OR 97403
lipshitz@uoregon.edu

Kefeng Liu
School of Sciences
Chongqing University of Technology
Chongqing 400054, China
liu@math.ucla.edu

Sucharit Sarkar
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
sucharit@math.ucla.edu

Dimitri Shlyakhtenko
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
shlyakht@ipam.ucla.edu

Ruixiang Zhang
Department of Mathematics
University of California
Berkeley, CA 94720-3840
ruixiang@berkeley.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2026 is US \$710/year for the electronic version, and \$965/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2026 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 342 No. 1 May 2026

The $\bar{\partial}$ -problem on $Z(q)$ -domains	1
DEBRAJ CHAKRABARTI, PHILLIP S. HARRINGTON and ANDREW RAICH	
Polynomial identities and Azumaya loci for rational quantum spheres	37
ALEXANDRU CHIRVASITU	
From i -boxes to signed words	63
ALESSANDRO CONTU, FAN QIN and QIAOLING WEI	
Near coincidences and nilpotent division fields	79
HARRIS B. DANIELS and JEREMY ROUSE	
The Cauchy problem for 1D nonlinear Schrödinger equations with repulsive delta potential for data in L^p -based spaces	115
QINGQUAN DENG, PING LI and XIUHONG LONG	
Finite basis problem for involution semigroups of order four	163
MENG GAO, EDMOND W. H. LEE, YAN FENG LUO and WEN TING ZHANG	