

*Pacific
Journal of
Mathematics*

**ON THE TEST PROPERTIES OF THE FROBENIUS
ENDOMORPHISM**

OLGUR CELIKBAS, ARASH SADEGHI AND YONGWEI YAO

ON THE TEST PROPERTIES OF THE FROBENIUS ENDOMORPHISM

OLGUR CELIKBAS, ARASH SADEGHI AND YONGWEI YAO

We prove two theorems concerning the test properties of the Frobenius endomorphism over commutative Noetherian local rings of prime characteristic p . Our first theorem generalizes a result of Funk and Marley on the vanishing of Ext and Tor modules, while our second theorem generalizes one of our previous results on maximal Cohen–Macaulay tensor products. In these earlier results, we replace ${}^e R$ with a more general module ${}^e M$, where R is a Cohen–Macaulay ring, M is a Cohen–Macaulay R -module with full support, and ${}^e M$ is the module viewed as an R -module via the e -th iteration of the Frobenius endomorphism. We also provide examples and present applications of our results, yielding new characterizations of the regularity of local rings.

1. Introduction

Throughout the paper, all rings are assumed to be commutative and Noetherian. By (R, \mathfrak{m}, k) , we mean R is a local ring with a unique maximal ideal \mathfrak{m} and residue field k .

Let R be a ring of prime characteristic p , $F : R \rightarrow R$ be the Frobenius endomorphism, and let M be an R -module. Each iteration F^e of F defines a new R -module structure on M , denoted by ${}^e M$, whose scalar multiplication is given as follows: For $r \in R$ and $x \in {}^e M$, we have that $r \cdot x = r^{p^e} x$. We say that R is F -finite if, for some $e \geq 1$ (or equivalently, for all $e \geq 1$), the module ${}^e R$ is finitely generated over R ; see, for example, [24]. We denote by $F_R^e(-)$ the scalar extension along the e -th iteration $F_R^e : R \rightarrow R$ of F . Thus, if $\sum_i n_i \otimes s_i \in F_R^e(M)$, where $F_R^e(M) = M \otimes_R {}^e R$, $n_i \in M$ and $r_i \in R$, then $r \cdot (\sum_i n_i \otimes s_i) = \sum_i n_i \otimes (r s_i)$, with $r s_i$ being the product of r and s_i in R . Note that $F_R^e(M)$ is the S -module $M \otimes_R S$ obtained via the base change $F^e : R \rightarrow S = R$.

The module structure of ${}^e R$ (as an R -module) contains important information about the homological properties of the ring R . For example, a remarkable result

MSC2020: primary 13A35, 13C14, 13D07; secondary 13C10, 13C11.

Keywords: Cohen–Macaulay module, depth and torsion properties of tensor products of modules, Frobenius endomorphism, ring of prime characteristic, Ext and Tor.

of Kunz [17] shows that R is regular if and only if eR is a flat R -module for some (or equivalently, for all) $e \geq 1$; see also [4] and [25] for extensions of this result. Motivated by Kunz's result, the test properties of the Frobenius endomorphism have been extensively studied.

If N is a finitely generated R -module, it follows from the work of Herzog [12] and Peskine and Szpiro [23; 24] that $\text{pd}_R(N) < \infty$ if and only if $\text{Tor}_i^R({}^eR, N) = 0$ for infinitely many e and for all $i \geq 1$. Avramov and Miller [3] showed that, if R is a complete intersection, the vanishing of a single $\text{Tor}_n^R({}^eR, N)$ for some $e \geq 1$ and $n \geq 1$ suffices to conclude that $\text{pd}_R(N) < \infty$. Koh and Lee [16] developed ideas rooted in techniques of Burch [7], Herzog [12], and Hochster [13], and showed that eR detects the finiteness of N even when finitely many Tor modules vanish. Specifically, Koh and Lee proved that, given integers $e \gg 0$ and $t \geq 1$, if $\text{Tor}_i^R({}^eR, N) = 0$ for $i = t, \dots, t + \text{depth}(R)$, then $\text{pd}_R(N) < \infty$. They further showed that in the case where R is Cohen–Macaulay, the number of vanishing Tor modules can be reduced by one. We refer the reader to the expository work [21] of Miller for further details.

In this paper, we focus on the following result of Funk and Marley [11; 10], which examines the vanishing of $\text{Tor}_i^R({}^eR, N)$ for the case where R is Cohen–Macaulay and N is possibly an infinitely generated R -module.

1.1 (Funk and Marley [11, 3.1 and 3.2]). Let (R, \mathfrak{m}, k) be a d -dimensional Cohen–Macaulay local ring of prime characteristic p , with $d \geq 1$, and let N be an R -module. Given integers $e \gg 0$ and $t \geq 1$, the following hold:

- (i) If $\text{Tor}_i^R({}^eR, N) = 0$ for all $i = t, \dots, t + d - 1$, then $\text{fd}_R(N) \leq d$.
- (ii) If R is F -finite and $\text{Ext}_R^i({}^eR, N) = 0$ for all $i = t, \dots, t + d - 1$, then $\text{id}_R(N) \leq d$.

One of the main goals of this paper is to generalize the result of Funk and Marley stated in 1.1. In fact, we prove more and establish the following theorem:

Theorem 1.2. *Let (R, \mathfrak{m}, k) be a d -dimensional local ring of prime characteristic p , M be a finitely generated Cohen–Macaulay R -module such that $\text{Supp}_R(M) = \text{Spec}(R)$, and let N be an R -module. Assume $\text{depth}(R) \geq 1$. Given integers $t \geq 1$ and $e \gg 0$, we have the following:*

- (i) *If $\text{Tor}_i^R({}^eM, N) = 0$ for all $i = t, \dots, t + d - 1$, then $\text{fd}_R(N) \leq d$.*
- (ii) *If R is F -finite and $\text{Ext}_R^i({}^eM, N) = 0$ for all $i = t, \dots, t + d - 1$, then $\text{id}_R(N) \leq d$.*
- (iii) *If N is finitely generated and $\text{Ext}_R^i(N, {}^eM) = 0$ for all $i = t, \dots, t + d - 1$, then $\text{pd}_R(N) \leq t - 1$.*

Parts (i) and (ii) of Theorem 1.2 recover the Funk–Marley result for the case where $M = R$. Note that, in each part of Theorem 1.2, only d consecutive vanishings

of Ext or Tor modules are needed to conclude the homological property the module of N .

We first record several preliminary results in Section 2 and then prove the first two parts of [Theorem 1.2](#) as [Theorem 3.2](#) in Section 3. The third part of [Theorem 1.2](#) is established as [Theorem 4.3](#) in Section 4. Additionally, in [Example 3.4](#), we show that the conclusion of [Theorem 1.2](#) may fail if the ring R in question has zero depth.

Li [18] proved that, if (R, \mathfrak{m}, k) is a Cohen–Macaulay local ring, N is a finitely generated R -module with rank, and $F_R^e(N)$ is maximal Cohen–Macaulay for some $e \gg 0$, then N is free. In [8], the authors of the present paper replaced the rank hypothesis on N with the weaker assumption that N is generically free, and proved the following result:

1.3 (Celikbas, Sadeghi, Yao [8, 1.3]). Let (R, \mathfrak{m}, k) be a Cohen–Macaulay local ring of prime characteristic p , and let M and N be finitely generated R -modules. Assume N is generically free, that is, $N_{\mathfrak{p}}$ is free over $R_{\mathfrak{p}}$ for all $\mathfrak{p} \in \text{Ass}(R)$. If $F_R^e(N)$ is maximal Cohen–Macaulay for some $e \gg 0$, then N is free.

In [Section 4](#), as a byproduct of [Theorem 1.2](#)(iii), we generalize [1.3](#) and prove the following result; see [Corollary 4.5](#). Note that $F_R^e(N) \otimes_R M$ is the S -module $(N \otimes_R S) \otimes_S M$, where $R \rightarrow S = R$ is the e -th iteration of the Frobenius endomorphism.

Theorem 1.4. *Let (R, \mathfrak{m}, k) be a local ring of prime characteristic p , and let M and N be finitely generated R -modules. Assume the following conditions hold:*

- (i) M is Cohen–Macaulay and $\text{Supp}_R(M) = \text{Spec}(R)$.
- (ii) N is generically free, that is, $N_{\mathfrak{p}}$ is a free $R_{\mathfrak{p}}$ -module for all $\mathfrak{p} \in \text{Ass}(R)$.

If $F_R^e(N) \otimes_R M$ is maximal Cohen–Macaulay for some $e \gg 0$, then N is free.

[Examples 4.8](#) and [4.9](#) showcase the necessity of the hypotheses $\text{Supp}_R(M) = \text{Spec}(R)$ and N is generically free in [Theorem 1.4](#). An immediate consequence of [Theorem 1.4](#) over one-dimensional rings can be stated as follows:

Corollary 1.5. *Let (R, \mathfrak{m}, k) be a one-dimensional reduced local ring of prime characteristic p and let $0 \neq I$ be an ideal of R . If $F_R^e(N) \otimes_R I$ is torsion-free for some finitely generated R -module N and $e \gg 0$, then N is free.*

[Theorem 1.2](#)(iii), in addition to [Theorem 1.4](#), has other applications, namely [Corollaries 4.4](#), [4.6](#), and [4.7](#). Moreover, in [Corollary 4.10](#), we obtain new characterizations of the regularity in terms of the vanishing of Ext and Tor.

2. Preliminaries

In this section, we record several preliminary results and observations that are necessary for our arguments in the subsequent sections. For the main results of this paper, one can skip this section and proceed to [Sections 3](#) and [4](#).

2.1. Let (R, \mathfrak{m}, k) be a local ring and let M be an R -module.

- (i) We set $M^\vee = \text{Hom}_R(M, E_R(k))$, where $E_R(k)$ is the injective hull of k . Note that $\text{Hom}_R(-, E_R(k))$ is a faithful exact functor.
- (ii) Assume that M is finitely generated over R . Given $n \geq 1$, we denote by $\Omega_R^n M$ the n -th syzygy of M , namely, the image of the n -th differential map in a minimal free resolution of M . By convention, $\Omega_R^0 M = M$.
- (iii) If $M \neq 0$ is finitely generated over R and $\dim_R(M) = t$, we define the *Hilbert–Samuel multiplicity* of M as

$$e_R(M) = t! \lim_{n \rightarrow \infty} \frac{\text{length}_R(M/\mathfrak{m}^n M)}{n^t},$$

which is a positive integer; see, for example, [20, p. 107].

2.2. Let (R, \mathfrak{m}, k) be a local ring and let N and X be R -modules.

- (1) Assume $\text{Soc}_R(X) \not\subseteq \mathfrak{m}X$ and let $x \in \text{Soc}_R(X) - \mathfrak{m}X$. This implies that $Rx \cong k$ and the nonzero map $Rx \hookrightarrow X \twoheadrightarrow X/\mathfrak{m}X$, $x \mapsto \bar{x}$, splits. Therefore, $Rx \hookrightarrow X$ splits and thus k is a direct summand of X as an R -module.
- (2) Assume R has prime characteristic p and $X = {}^e M$ for some R -module $0 \neq M$ and $e \geq 0$.
 - (a) If $\text{Soc}_R(X) \not\subseteq \mathfrak{m}X$, that is, ${}^e(0 :_M \mathfrak{m}^{[p^e]}) \not\subseteq {}^e(\mathfrak{m}^{[p^e]} M)$, then part (1) shows that k is a direct summand of ${}^e M$ as an R -module.
 - (b) Assume M is finitely generated over R with $\text{depth}_R(M) = 0$. For all $e \gg 0$, we have ${}^e(0 :_M \mathfrak{m}^{[p^e]}) = \text{Soc}_R({}^e M) \supseteq {}^e(\text{Soc}_R(M)) \not\subseteq {}^e(\mathfrak{m}^{[p^e]} M)$. Hence, part (i) shows that k is a direct summand of ${}^e M$ for every $e \gg 0$.
 - (c) Assume (R, \mathfrak{m}, k) is Artinian. Then ${}^e(0 :_M \mathfrak{m}^{[p^e]}) \not\subseteq {}^e(\mathfrak{m}^{[p^e]} M)$ for all $e \gg 0$. By part (i), k is a direct summand of ${}^e M$ for all $e \gg 0$. Thus, given $i \geq 1$, if $\text{Ext}_R^i({}^e M, N) = 0$ for some $e \gg 0$, then $\text{Ext}_R^i(k, N) = 0$ and hence N is injective; see [6, 3.1.12] and also [9, 2.0.10].

2.3. Let (R, \mathfrak{m}, k) be a local ring of prime characteristic p . There exists a local flat ring homomorphism $(R, \mathfrak{m}, k) \rightarrow (S, \mathfrak{n}, \ell)$ such that S is F-finite, $\mathfrak{m}S = \mathfrak{n}$ and $|\ell| = \infty$; hence S is faithfully flat over R , $\dim(S) = \dim(R)$ and $e(R) = e(S)$; it also follows that R is Cohen–Macaulay (respectively, regular) if and only if S is Cohen–Macaulay (respectively, regular). (For example, with $\hat{R} \cong k[[x_1, \dots, x_m]]/I$, we can pick $S = \bar{k}[[x_1, \dots, x_m]]/I\bar{k}[[x_1, \dots, x_m]]$, where \bar{k} is the algebraic closure of k .) In the case where S is such an extension of R and M is a finitely generated R -module, it follows that M is a Cohen–Macaulay R -module with $\text{Supp}_R(M) = \text{Spec}(R)$ if and only if $M \otimes_R S$ is a Cohen–Macaulay S -module with $\text{Supp}_S(M \otimes_R S) = \text{Spec}(S)$; also, M is free over R if and only if $M \otimes_R S$ is free over S .

2.4. Let (R, \mathfrak{m}, k) be a local ring of prime characteristic p and let $M \neq 0$ be a finitely generated R -module. We set:

$$\text{crs}_p(M) = \min\{e \geq 0 \mid (0 :_{M/(\underline{x})M} \mathfrak{m}^{[p^e]}) \not\subseteq \mathfrak{m}^{[p^e]}(M/(\underline{x})M) \text{ for an } M\text{-regular sequence } \underline{x}\},$$

$$\text{drs}_p(M) = \min\{e \geq 0 \mid \mathfrak{m}^{[p^e]}M \subseteq (\underline{x})M \text{ for a system of parameters } \underline{x} \text{ on } M\}.$$

It seems unknown whether or not $\sup\{\text{crs}_p(M_{\mathfrak{p}}) \mid \mathfrak{p} \in \text{Supp}_R(M)\}$ is finite in general.

2.5. Let (R, \mathfrak{m}, k) be a local ring of prime characteristic p and let $M \neq 0$ be a finitely generated Cohen–Macaulay R -module. It follows that $\text{crs}_p(M) \leq \text{drs}_p(M)$. If $|k| = \infty$, then $\text{drs}_p(M) \leq \lceil \log_p e_R(M) \rceil$; see, for example, [8, 2.4]. Therefore, we have

$$\begin{aligned} \sup\{\text{crs}_p(M_{\mathfrak{p}}) \mid \mathfrak{p} \in \text{Supp}_R(M)\} &\leq \begin{cases} \max\{\lceil \log_p e_R(M) \rceil, \text{crs}_p(M)\}, \\ \lceil \log_p e_R(M) \rceil & \text{if } |k| = \infty, \end{cases} \\ \sup\{\text{drs}_p(M_{\mathfrak{p}}) \mid \mathfrak{p} \in \text{Supp}_R(M)\} &\leq \begin{cases} \max\{\lceil \log_p e_R(M) \rceil, \text{drs}_p(M)\}, \\ \lceil \log_p e_R(M) \rceil & \text{if } |k| = \infty. \end{cases} \end{aligned}$$

2.6. Let $R \rightarrow S$ be a ring homomorphism, M be an S -module, and let X be an R -module.

- (i) $\text{Hom}_R(M, X)$ has an S -module structure defined as follows: For $s \in S$ and $\alpha \in \text{Hom}_R(M, X)$, we set $s \cdot \alpha \in \text{Hom}_R(M, X)$ as $(s \cdot \alpha)(m) = \alpha(sm)$ for all $m \in M$.
- (ii) $\text{Hom}_R(X, M)$ has an S -module structure defined as follows: For $s \in S$ and $\alpha \in \text{Hom}_R(X, M)$, we set $s \cdot \alpha \in \text{Hom}_R(X, M)$ as $(s \cdot \alpha)(x) = s\alpha(x)$ for all $x \in X$.
- (iii) $M \otimes_R X$ has an S -module structure as follows: For $s \in S$ and $\sum_i m_i \otimes x_i \in M \otimes_R X$, with $m_i \in M$ and $x_i \in X$, we define

$$s \cdot \left(\sum_i m_i \otimes x_i\right) = \sum_i (sm_i) \otimes x_i \in M \otimes_R X.$$

The following observation is used in several proofs in the sequel. Note that, over a Noetherian ring, every finitely generated module is finitely presented.

2.7. Let $f : R \rightarrow S$ be a ring homomorphism, A be an R -module, and let B be an S -module. Assume E is an injective S -module. Given $n \geq 0$, the following hold:

- (i) $\text{Hom}_S(\text{Tor}_n^R(A, B), E) \cong \text{Ext}_R^n(A, \text{Hom}_S(B, E))$; see [26, 10.63].
- (ii) If A is finitely presented over R , then

$$\text{Hom}_S(\text{Ext}_R^n(A, B), E) \cong \text{Tor}_n^R(A, \text{Hom}_S(B, E)).$$

(iii) If (R, \mathfrak{m}, k) is local, $R = S$, and $f = \text{id}$, then parts (i) and (ii) imply that

$$\text{fd}_R(B) = \text{id}_R(B^\vee) \quad \text{and} \quad \text{id}_R(B) = \text{fd}_R(B^\vee).$$

2.8 (Auslander–Buchsbaum [2]). Let (R, \mathfrak{m}, k) be a d -dimensional local ring and let N be an R -module. If $\text{id}_R(N) < \infty$, then $\text{id}_R(N) \leq d$. Thus, if $\text{fd}_R(N) < \infty$, then $\text{fd}_R(N) \leq d$; see 2.7(iii).

The next observation is used in the proofs of Lemma 3.8 and Proposition 3.11.

2.9. Let (R, \mathfrak{m}, k) be a d -dimensional local ring and let N be an R -module. Consider a minimal injective resolution of N

$$I = (0 \rightarrow I^0 \xrightarrow{h_1} I^1 \xrightarrow{h_2} \dots \xrightarrow{h_n} I^n \xrightarrow{h_{n+1}} I^{n+1} \rightarrow \dots),$$

where

$$I^j = \bigoplus_{\mathfrak{p} \in \text{Spec}(R)} E_R(R/\mathfrak{p})^{\oplus \mu_j(\mathfrak{p}, N)} \quad \text{and} \quad \mu_j(\mathfrak{p}, N) = \text{rank}_{k(\mathfrak{p})} (\text{Ext}_{R_{\mathfrak{p}}}^j(k(\mathfrak{p}), N))$$

for $j \geq 0$. Here, $\mu_j(\mathfrak{p}, N)$ is not necessarily finite. Note that $\text{Hom}_{R_{\mathfrak{p}}}(k(\mathfrak{p}), (h_j)_{\mathfrak{p}}) = 0$ for $j \geq 0$. If $\mathfrak{q} \in \text{Spec}(R)$ and we localize I at \mathfrak{q} , then the resulting $I_{\mathfrak{q}}$ is a minimal injective resolution of $N_{\mathfrak{q}}$ over $R_{\mathfrak{q}}$ and $I_{\mathfrak{q}}^j = \bigoplus_{\mathfrak{p} \subseteq \mathfrak{q}} E_R(R/\mathfrak{p})^{\oplus \mu_j(\mathfrak{p}, N)}$ for all $j \geq 0$; see, for example, [5] and [15, 3.15 and Appendix 20–24].

Assume $\text{id}_{R_{\mathfrak{p}}}(N_{\mathfrak{p}}) < \infty$ for all $\mathfrak{p} \in \text{Spec}(R) - \{\mathfrak{m}\}$. Then $\text{id}_{R_{\mathfrak{p}}}(N_{\mathfrak{p}}) \leq \dim(R_{\mathfrak{p}}) \leq d - 1$ for all $\mathfrak{p} \in \text{Spec}(R) - \{\mathfrak{m}\}$; see 2.8. This implies that $I^i = E_R(k)^{\oplus \mu_i(\mathfrak{m}, N)}$ for all $i \geq d$. Consequently, I has the form

$$I = (0 \rightarrow I^0 \rightarrow \dots \rightarrow I^{d-1} \rightarrow E_R(k)^{\oplus \mu_d(\mathfrak{m}, N)} \xrightarrow{h_{d+1}} E_R(k)^{\oplus \mu_{d+1}(\mathfrak{m}, N)} \rightarrow \dots).$$

2.10. Let (R, \mathfrak{m}, k) be a local ring, I be an ideal of R such that $\sqrt{I} = \sqrt{\text{Ann}_R(M)}$, M be a finitely generated R -module, and let N be an R -module. Then:

(i) $\text{Hom}_R(M, N) = 0 \iff \text{grade}_R(I, N) \geq 1 \iff \text{Hom}_R(R/I, N) = 0$; see [6, 1.2.3 or 1.2.10(e)].

(ii) It follows from part (i) and 2.7(i) that

$$\begin{aligned} M \otimes_R N = 0 &\iff \text{Hom}_R(M, N^\vee) = 0 \iff \text{Hom}_R(R/I, N^\vee) = 0 \\ &\iff (R/I \otimes_R N)^\vee = 0 \iff R/I \otimes_R N = 0 \iff N = IN. \end{aligned}$$

(iii) If $\text{Supp}_R(M) = \text{Spec}(R)$ (or equivalently, $I = 0$), it follows from parts (i) and (ii) that

$$\text{Hom}_R(M, N) = 0 \iff M \otimes_R N = 0 \iff N = 0.$$

(iv) $\text{Ass}_R(\text{Hom}_R(M, N)) = \text{Supp}_R(M) \cap \text{Ass}_R(N)$; see, for example, [6, 1.2.28].

Remark 2.11. Assume I , M , and N are as in 2.10, but R is (Noetherian as always) not necessarily local. Then the implications considered in 2.10 still hold since they can all be verified locally. More precisely, it follows that

$$\mathrm{Hom}_R(M, N) = 0 \iff \mathrm{Hom}_R(R/I, N) = 0 \quad \text{and} \quad M \otimes_R N = 0 \iff N = IN.$$

In particular, if $\mathrm{Supp}_R(M) = \mathrm{Spec}(R)$, we have

$$\mathrm{Hom}_R(M, N) = 0 \iff N = 0 \iff M \otimes_R N = 0.$$

These implications establish the fact $\mathrm{Ass}_R(\mathrm{Hom}_R(M, N)) = \mathrm{Supp}_R(M) \cap \mathrm{Ass}_R(N)$, namely the equality stated in 2.10(iv), still holds.

2.12. Let (R, \mathfrak{m}, k) be a local ring and let M be a finitely generated R -module. Consider a minimal free representation $P_1 \xrightarrow{\partial_1} P_0 \rightarrow M \rightarrow 0$. The *transpose* $\mathrm{Tr}_R M$ of M is defined as the cokernel of the R -dual map $\partial_1^* = \mathrm{Hom}_R(\partial_1, R)$. We refer the reader to [1] for the details of the following:

- (i) There is an exact sequence of R -modules $0 \rightarrow M^* \rightarrow P_0^* \rightarrow P_1^* \rightarrow \mathrm{Tr}_R M \rightarrow 0$.
- (ii) It follows that, up to isomorphism, $\mathrm{Tr}_R M$ is uniquely determined.
- (iii) It follows that $\mathrm{Tr}_R(\mathrm{Tr}_R M) \cong M$.
- (iv) M is free if and only if $\mathrm{Tr}_R M$ is free.

The following result from [8] is necessary for our proof of Theorem 1.4.

2.13 [8, 2.2]. Let (R, \mathfrak{m}, k) be a local ring and let M and N be finitely generated R -modules. Assume $n \geq 1$ is an integer. Assume further the following conditions hold:

- (i) $N_{\mathfrak{p}}$ is free for all $\mathfrak{p} \in \mathrm{Spec}(R) - \{\mathfrak{m}\}$.
- (ii) $\mathrm{depth}_R(M \otimes_R N) \geq n$.
- (iii) $\mathrm{depth}_R(M) \geq n - 1$.

Then $\mathrm{Ext}_R^i(\mathrm{Tr}_R N, M) = 0$ for all $i = 1, \dots, n$.

3. A generalization of a result of Funk and Marley

The main results of this section are captured in Theorem 3.2 below, stated earlier as parts (i) and (ii) of Theorem 1.2. They generalize this result already quoted in 1.1:

3.1 (Funk and Marley [11, 3.1 and 3.2]). Let (R, \mathfrak{m}, k) be a d -dimensional Cohen–Macaulay local ring of prime characteristic p , with $d \geq 1$, and let N be an R -module. Given integers $t \geq 1$ and $e \gg 0$, the following hold:

- (i) If $\mathrm{Tor}_i^R({}^e R, N) = 0$ for all $i = t, \dots, t + d - 1$, then $\mathrm{fd}_R(N) \leq d$.
- (ii) If R is F -finite and $\mathrm{Ext}_R^i({}^e R, N) = 0$ for all $i = t, \dots, t + d - 1$, then $\mathrm{id}_R(N) \leq d$.

The original statement of 3.1 in [11] includes the case where $d = 0$; in fact this case follows from the techniques of Koh and Lee used in the proof of [16, 2.6]; see also [11, 2.8] and [21, 2.2.8]. Note also that [11, 3.1 and 3.2], namely 3.1, is stated in terms of complexes of R -modules, but its proof naturally reduces to the case of modules. For this reason, we consider only modules when generalizing 3.1 in Theorems 3.2 and 4.3, which can also be extended to the complex case in a similar manner, as explained in the proof of [11, 3.1].

Theorem 3.2. *Let (R, \mathfrak{m}, k) be a d -dimensional local ring of prime characteristic p , M be a finitely generated Cohen–Macaulay R -module such that $\text{Supp}_R(M) = \text{Spec}(R)$, and let N be an R -module. Assume $\text{depth}(R) \geq 1$. Given $t \geq 1$ and $e \gg 0$, the following hold:*

- (i) *If $\text{Tor}_i^R({}^eM, N) = 0$ for all $i = t, \dots, t + d - 1$, then $\text{fd}_R(N) \leq d$.*
- (ii) *If R is F-finite and $\text{Ext}_R^i({}^eM, N) = 0$ for all $i = t, \dots, t + d - 1$, then $\text{id}_R(N) \leq d$.*

Remark 3.3. Let R be a ring of prime characteristic p , M be a finitely generated R -module such that $\text{Supp}_R(M) = \text{Spec}(R)$, and let N be an R -module. Let $e \geq 0$.

- (i) Assume R is F-finite. Then $\text{Hom}_R({}^eM, N) = 0 \implies N = 0$; see 2.10(iii).
- (ii) Without the F-finite assumption, we have ${}^eM \otimes N = 0 \implies N = 0$, as locally this reduces to the F-finite case and then follows from 2.10(iii).

Thus, Theorem 3.2 still holds when $t = 0$. This relies on $\text{Supp}_R({}^eM) = \text{Spec}(R)$ and does not depend on the choice of e or M being Cohen–Macaulay.

Theorem 3.2 generalizes 3.1 in the case where $\text{depth}(R) \geq 1$. For a generalization of 3.1 in the case where $\text{depth}(R) = 0$ (that is, the $d = 0$ case of 3.1), see Proposition 3.10. Before presenting our proof of Theorem 3.2 at the end of the section, we would like to discuss the sharpness of the result and list some corollaries of the theorem. We will also prove Propositions 3.10 and 3.11, which the proof of Theorem 3.2 relies on.

The following example shows that the positive depth assumption on the ring is necessary for Theorem 3.2.

Example 3.4. Let $R = \mathbb{F}_p[[x, y]]/(x^2, xy)$ and let $M = R/(x)$. Then R is an F-finite ring with $\text{depth}(R) = 0$ and $\text{dim}(R) = 1$, and M is a Cohen–Macaulay R -module such that $\text{Supp}_R(M) = \text{Spec}(R)$. Moreover, ${}^eM \cong M^{\oplus p^e}$ for all $e \geq 0$.

Let $N = R/(y)$. Then $\text{Tor}_1^R(M, N) = 0$ so that $\text{Tor}_1^R({}^eM, N) = 0$ for all $e \geq 0$. If $\text{pd}_R(N) < \infty$, then N is free since $\text{depth}(R) = 0$. Hence $\text{pd}_R(N) = \text{fd}_R(N) = \infty$. This shows that the positive depth assumption is needed for Theorem 3.2(i).

Next let $N = M$. Then $\text{Ext}_R^1(M, N) = 0$ so that $\text{Ext}_R^1({}^eM, M) = 0$ for all $e \geq 0$. If $\text{id}_R(N) < \infty$, then $\text{id}_R(N) = \text{depth}(R) = 0$, that is, N is injective. Hence, $\text{id}_R(N) = \infty$ (one can also conclude that $\text{id}_R(N) = \infty$ since R is not Cohen–Macaulay). Thus the positive depth assumption is needed for Theorem 3.2(ii).

We give several corollaries of [Theorem 3.2](#). The next corollary covers the particular case where $S = \hat{R}$, the \mathfrak{m} -adic completion of (R, \mathfrak{m}) .

Corollary 3.5. *Let $f : (R, \mathfrak{m}, k) \rightarrow (S, \mathfrak{n}, \ell)$ be a flat local ring homomorphism, where R is a d -dimensional local ring of prime characteristic p and $\mathfrak{m}S = \mathfrak{n}$, M be a finitely generated Cohen–Macaulay R -module such that $\text{Supp}_R(M) = \text{Spec}(R)$, and let N be a finitely generated S -module. Assume $\text{depth}(R) \geq 1$. Given $t \geq 1$ and $e \gg 0$, if $\text{Tor}_i^R({}^eM, N) = 0$ for all $i = t, \dots, t + d - 1$, then $\text{pd}_S(N) \leq d$.*

Proof. Observe $\text{Tor}_i^R(k, N) \cong \text{Tor}_i^S(\ell, N)$ for all $i \geq 0$. By [Theorem 3.2](#), we have that $\text{fd}_R(N) \leq d$. Set $r = \text{fd}_R(N)$. It follows that $\text{Tor}_r^R(k, N) \neq 0 = \text{Tor}_{r+1}^R(k, N)$. Therefore, $\text{pd}_S(N) = r \leq d$. \square

Corollary 3.6. *Let (R, \mathfrak{m}, k) be a d -dimensional local ring of prime characteristic p , M be a finitely generated Cohen–Macaulay R -module such that $\text{Supp}_R(M) = \text{Spec}(R)$, and let N be an R -module. Assume $\text{depth}(R) \geq 1$. Given $t \geq 1$ and $e \gg 0$, if $\text{Ext}_R^i({}^eM, N^\vee) = 0$ for all $i = t, \dots, t + d - 1$, then $\text{id}_R(N^\vee) \leq d$.*

Proof. The vanishing of $\text{Ext}_R^i({}^eM, N^\vee)$ yields the vanishing of $\text{Tor}_i^R({}^eM, N)$; see [2.7\(i\)](#). Thus $\text{id}_R(N^\vee) = \text{fd}_R(N) \leq d$ by [Theorems 3.2\(i\)](#) and [2.7\(iii\)](#). \square

Corollary 3.7. *Let (R, \mathfrak{m}, k) be a d -dimensional local ring of prime characteristic p , M be a finitely generated Cohen–Macaulay R -module such that $\text{Supp}_R(M) = \text{Spec}(R)$, and let N be an R -module. Assume $\text{depth}(R) \geq 1$. Given $t \geq 1$ and $e \gg 0$, assume at least one of the following conditions holds:*

- (i) $\text{Ext}_R^i(N, ({}^eM)^\vee) = 0$ for all $i = t, \dots, t + d - 1$.
- (ii) $\text{Ext}_R^i(N, {}^e(M^\vee)) = 0$ for all $i = t, \dots, t + d - 1$.

Then $\text{fd}_R(N) \leq d$.

Proof. The vanishing of $\text{Ext}_R^i(N, ({}^eM)^\vee)$ yields the vanishing of $\text{Tor}_i^R(N, {}^eM)$; see [2.7\(i\)](#). Thus, case (i) follows from [Theorem 3.2\(i\)](#).

Similarly, the vanishing of $\text{Ext}_R^i(N, {}^e(M^\vee))$ yields the vanishing of $\text{Tor}_i^R(N, {}^eM)$ by [2.7\(i\)](#), where $R \rightarrow S$ is the Frobenius F^e with $(S, \mathfrak{n}, \ell) = (R, \mathfrak{m}, k)$, $A = N$ over R , $B = M$ over S , and $E = E_S(\ell)$ is the injective hull of ℓ over S . Therefore, case (ii) also follows from [Theorem 3.2\(i\)](#). \square

Next, we prepare some auxiliary results for the [proof of Theorem 3.2](#). To begin with, we present [Lemma 3.8](#) and [Corollary 3.9](#) which are akin to [[11](#), 4.5 and 4.6].

Lemma 3.8. *Let (R, \mathfrak{m}, k) be a d -dimensional F -finite local ring of prime characteristic p , M be a finitely generated R -module such that $\text{Supp}_R(M) = \text{Spec}(R)$, and let N be an R -module. Set $\delta = \max\{\text{depth}_{R_{\mathfrak{p}}}(M_{\mathfrak{p}}) \mid \mathfrak{p} \in \text{Spec}(R)\}$ and let $t \geq 1$ be an integer. If*

$$\sup\{e \mid \text{Ext}_R^i({}^eM, N) = 0 \text{ for all } i = t, \dots, t + \delta\} \geq \sup\{\text{crs}_p(M_{\mathfrak{p}}) \mid \mathfrak{p} \in \text{Spec}(R)\},$$

then $\text{id}_R(N) \leq \min\{d, t + \delta - 1\}$. Therefore, if $\text{Ext}_R^i({}^eM, N) = 0$ for all $i = t, \dots, t + \delta$ and for infinitely many e , then $\text{id}_R(N) \leq \min\{d, t + \delta - 1\}$.

Proof. By 2.8, it suffices to show $\text{id}_R(N) \leq t + \delta - 1$, and we proceed by induction on d . The case where $d = 0$ follows from 2.2(2)(iii). Hence, we assume $d \geq 1$.

As R is F-finite, eM is a finitely presented R -module. Thus, for all $\mathfrak{p} \in \text{Spec}(R)$, we have the isomorphisms

$$\text{Ext}_R^i({}^eM, N)_{\mathfrak{p}} \cong \text{Ext}_{R_{\mathfrak{p}}}^i({}^eM_{\mathfrak{p}}, N_{\mathfrak{p}}).$$

So, for all $\mathfrak{p} \in \text{Spec}(R) - \{\mathfrak{m}\}$, the induction hypothesis dictates $\text{id}_{R_{\mathfrak{p}}}(N_{\mathfrak{p}}) < t + \delta$. By 2.9, N has a minimal injective resolution of the form

$$I = (0 \rightarrow I^0 \rightarrow \dots \rightarrow I^{t+\delta-1} \rightarrow E_R(k)^{\oplus \mu_{t+\delta}} \rightarrow E_R(k)^{\oplus \mu_{t+\delta+1}} \rightarrow \dots),$$

in which $\mu_i = \mu_i(\mathfrak{m}, N)$ for $i \geq t + \delta$. Set $\text{depth}_R(M) = v$. By assumption, there exists $e \geq \text{crs}_{\mathfrak{p}}(M)$ such that $\text{Ext}_R^i({}^eM, N) = 0$ for $i = t + \delta - v, \dots, t + \delta$. Since $e \geq \text{crs}_{\mathfrak{p}}(M)$, 2.4 yields a maximal M -regular sequence $\underline{x} = \{x_1, \dots, x_v\}$ such that

$$(0 :_{M/(\underline{x})M} \mathfrak{m}^{[p^e]}) \not\subseteq \mathfrak{m}^{[p^e]}(M/(\underline{x})M).$$

Thus, k is a direct summand of ${}^e(M/(\underline{x})M)$; see 2.2. As x_1 is M -regular, there is a short exact sequence $0 \rightarrow {}^eM \rightarrow {}^eM \rightarrow {}^e(M/x_1M) \rightarrow 0$. This, together with $\text{Ext}_R^i({}^eM, N) = 0$ for $i = t + \delta - v, \dots, t + \delta$, implies that

$$\text{Ext}_R^i({}^e(M/x_1M), N) = 0 \quad \text{for all } i = t + \delta - v + 1, \dots, t + \delta.$$

As $\underline{x} = \{x_1, \dots, x_v\}$ is M -regular, inductively we get $\text{Ext}_R^{t+\delta}({}^e(M/(\underline{x})M), N) = 0$. Therefore, $\text{Ext}_R^{t+\delta}(k, N) = 0$ since k is a direct summand of ${}^e(M/(\underline{x})M)$. In view of 2.9, we deduce

$$\mu_{t+\delta} = \text{rank}_k(\text{Ext}_R^{t+\delta}(k, N)) = 0.$$

Therefore, $I^{t+\delta} = E_R(k)^{\oplus \mu_{t+\delta}} = 0$, which concludes that $\text{id}_R(N) \leq t + \delta - 1$. \square

Corollary 3.9. *Let (R, \mathfrak{m}, k) be a d -dimensional local ring of prime characteristic p , M be a finitely generated R -module such that $\text{Supp}_R(M) = \text{Spec}(R)$, and let N be an R -module. Let S be a ring extension of R as in 2.3. Set*

$$\delta = \max\{\text{depth}_{S_{\mathfrak{p}}}(M \otimes_R S_{\mathfrak{p}}) \mid \mathfrak{p} \in \text{Spec}(S)\}$$

and let $t \geq 1$ be an integer. If

$$\sup\{e \mid \text{Tor}_i^R({}^eM, N) = 0 \text{ for all } i = t, \dots, t + \delta\} \geq \sup\{\text{crs}_{\mathfrak{p}}(M \otimes_R S_{\mathfrak{p}}) \mid \mathfrak{p} \in \text{Spec}(S)\},$$

then $\text{fd}_R(N) \leq d$. In particular, if $\text{Tor}_i^R({}^eM, N) = 0$ for all $i = t, \dots, t + \delta$ and for infinitely many e , then $\text{fd}_R(N) \leq d$.

Proof. It suffices to show $\text{fd}_S(N \otimes_R S) \leq \dim(S)$. Thus, given the assumption, we may assume that $R = S$ is already F-finite. By 2.7(i), the vanishing of $\text{Tor}_i^R({}^eM, N)$ yields vanishing of $\text{Ext}_R^i({}^eM, N^\vee)$. Now the claim follows from Lemma 3.8; see 2.7(iii). \square

As an application of Lemma 3.8 and Corollary 3.9, we obtain the following proposition that generalizes a result of Takahashi and Yoshino [27, 5.3].

Proposition 3.10. *Let (R, \mathfrak{m}, k) be a d -dimensional local ring of prime characteristic p , M be a finitely generated Cohen–Macaulay R -module such that $\text{Supp}_R(M) = \text{Spec}(R)$, and let N be an R -module. Then, given $t \geq 1$ and $e \gg 0$, the following hold:*

- (i) *If $\text{Tor}_i^R({}^eM, N) = 0$ for all $i = t, \dots, t + d$, then $\text{fd}_R(N) \leq d$.*
- (ii) *If R is F-finite and $\text{Ext}_R^i({}^eM, N) = 0$ for all $i = t, \dots, t + d$, then $\text{id}_R(N) \leq d$.*

Proof. For (i), we may assume that R is F-finite with $|k| = \infty$; see 2.3. In this case, we have $\sup\{\text{crs}_p(M_{\mathfrak{p}}) \mid \mathfrak{p} \in \text{Spec}(R)\} \leq \lceil \log_p e_R(M) \rceil < \infty$. The rest follows from Corollary 3.9.

For (ii), without the assumption that $|k| = \infty$, we have

$$\sup\{\text{crs}_p(M_{\mathfrak{p}}) \mid \mathfrak{p} \in \text{Spec}(R)\} \leq \max\{\lceil \log_p e_R(M) \rceil, \text{crs}_p(M)\} < \infty.$$

The rest follows from Lemma 3.8. \square

Our proof of Proposition 3.11 is inspired by some of the techniques employed by Funk and Marley in the proof of [11, 3.2]. To distinguish various module structures in the proof, we present 3.11 in the context of a general ring homomorphism $f : R \rightarrow S$. Subsequently, in the proof of Theorem 3.2, we apply Proposition 3.11 to $F^e : R \rightarrow R$, the iterated Frobenius endomorphism. Recall that, over a local ring (R, \mathfrak{m}, k) , we set $(-)^{\vee} = \text{Hom}_R(-, E_R(k))$.

Proposition 3.11. *Let $f : (R, \mathfrak{m}, k) \rightarrow (S, \mathfrak{n}, \ell)$ be a module-finite local homomorphism of local rings, with $d = \dim(R) \geq 1$, M be an S -module, and let N be an R -module. Assume the following:*

- (i) *$\text{Ext}_R^i(M, N) = 0$ for all $i = t, \dots, t + d - 1$ for some $t \geq 1$.*
- (ii) *M is a finitely generated S -module such that $\text{Supp}_R(M) = \text{Spec}(R)$.*
- (iii) *There exists $\underline{x} = \{x_1, \dots, x_d\} \subseteq \mathfrak{n}$ such that \underline{x} is M -regular and $\mathfrak{m}M \subseteq (\underline{x})M$.*
- (iv) *$\text{id}_{R_{\mathfrak{p}}}(N_{\mathfrak{p}}) < \infty$ for all $\mathfrak{p} \in \text{Spec}(R) - \{\mathfrak{m}\}$.*
- (v) *$\text{fd}_{R_{\mathfrak{p}}}((N^{\vee})_{\mathfrak{p}}) < \infty$ for all $\mathfrak{p} \in \text{Ass}(R)$.*

Then $\text{id}_R(N) \leq d$.

Proof. As $\text{id}_{R_{\mathfrak{p}}}(N_{\mathfrak{p}}) < \infty$ for each $\mathfrak{p} \in \text{Spec}(R) - \{\mathfrak{m}\}$, 2.9 dictates that N has a minimal injective resolution of the form

$$\mathbf{I} = (0 \rightarrow I^0 \xrightarrow{h_1} \dots \rightarrow I^{d-1} \xrightarrow{h_d} E_R(k)^{\oplus \mu_d(m, N)} \rightarrow \dots \xrightarrow{h_{t+d}} E_R(k)^{\oplus \mu_{t+d}(m, N)} \rightarrow \dots).$$

We apply $\text{Hom}_R(M, -)$ to \mathbf{I} and use our assumption (i) to obtain an exact sequence

$$(3.11.1) \quad M_{t-1} \xrightarrow{g_t} \dots \xrightarrow{g_{t+d-2}} M_{t+d-2} \xrightarrow{g_{t+d-1}} M_{t+d-1} \xrightarrow{g_{t+d}} M_{t+d} \xrightarrow{\rho} C \rightarrow 0,$$

where $M_i = \text{Hom}_R(M, I^i)$, $g_i = \text{Hom}_R(M, h_i)$ and $C = \text{coker}(g_{t+d})$. Note that each M_i is an S -module; see 2.6(i). Hence, C is an S -module as well.

Claim 1. *The induced sequence $\overline{M_{t+d-1}} \xrightarrow{\overline{g_{t+d}}} \overline{M_{t+d}} \xrightarrow{\overline{\rho}} \overline{C} \rightarrow 0$, in which*

$$\overline{M}_i = \text{Hom}_R(M/(\underline{\mathbf{x}})M, I^i), \quad \overline{g}_{t+d} = \text{Hom}_R(M/(\underline{\mathbf{x}})M, h_{t+d})$$

and $\overline{C} = \text{Hom}_S(S/(\underline{\mathbf{x}}), C)$ is exact.

Proof. Since I^i is injective over R and x_1 is regular on M , an application of $\text{Hom}_R(-, I^i)$ to the exact sequence $0 \rightarrow M \xrightarrow{x_1} M \rightarrow M/x_1M \rightarrow 0$ induces a short exact sequence

$$(\Delta_i) \quad 0 \leftarrow M_i \xleftarrow{x_1} M_i \leftarrow \text{Hom}_R(M/x_1M, I^i) \leftarrow 0.$$

We combine the short exact sequences (Δ_i) , for all $i = t-1, \dots, t+d$, with the exact sequence (3.11.1), and obtain the exact sequence of S -modules

$$\text{Hom}_R(M/x_1M, I^t) \rightarrow \dots \rightarrow \text{Hom}_R(M/x_1M, I^{t+d}) \rightarrow \text{Hom}_S(S/x_1S, C) \rightarrow 0.$$

Inductively, as $\underline{\mathbf{x}} = \{x_1, \dots, x_d\}$ is an M -regular sequence, we realize the following exact sequence of S -modules that is naturally induced from (3.11.1), as claimed:

$$\text{Hom}_R(M/(\underline{\mathbf{x}})M, I^{t+d-1}) \xrightarrow{\overline{g}_{t+d}} \text{Hom}_R(M/(\underline{\mathbf{x}})M, I^{t+d}) \xrightarrow{\overline{\rho}} \text{Hom}_S(S/(\underline{\mathbf{x}})S, C) \rightarrow 0.$$

Claim 2. $\ker(\rho) = 0$.

Proof. The assumption $\mathfrak{m}M \subseteq (\underline{\mathbf{x}})M$ says that, as an R -module, $M/(\underline{\mathbf{x}})M$ is a direct sum of copies of k . Since \mathbf{I} is a minimal injective resolution of N , we see that

$$\overline{g}_{t+d} = \text{Hom}_R(M/(\underline{\mathbf{x}})M, h_{t+d}) = 0.$$

In view of the exact sequence in Claim 1, we get $\ker(\overline{\rho}) = 0$. Moreover, $\overline{\rho}$ can be identified as $\text{Hom}_S(S/(\underline{\mathbf{x}}), \rho)$ up to the natural isomorphism $\text{Hom}_S(S/(\underline{\mathbf{x}}), M_i) \cong \text{Hom}_R(M/(\underline{\mathbf{x}})M, I^i)$. Hence $(0 :_{\ker(\rho)}(\underline{\mathbf{x}})) \cong \ker(\overline{\rho}) = 0$. Given that M is finitely generated over R , we have

$$\text{Ass}_R(\ker(\rho)) \subseteq \text{Ass}_R(M_{t+d}) \subseteq \text{Ass}_R(E_R(k)^{\oplus \mu_{t+d}(m, N)}) \subseteq \{\mathfrak{m}\}.$$

Thus $\text{Ass}_S(\ker(\rho)) \subseteq \{\mathfrak{n}\}$, as \mathfrak{n} is the only prime ideal of S lying over \mathfrak{m} . So, if $\ker(\rho) \neq 0$, then $\mathfrak{n} \in \text{Ass}_S(\ker(\rho))$ and hence $0 \neq (0 :_{\ker(\rho)} \mathfrak{n}) \subseteq (0 :_{\ker(\rho)} \underline{\mathbf{x}})$, which contradicts the conclusion $(0 :_{\ker(\rho)} \underline{\mathbf{x}}) = 0$ above. This completes the proof of the claim.

Now that we know $\ker(\rho) = 0$, the exact sequence (3.11.1) forces $g_{t+d} = 0$, which gives rise to an exact sequence

$$\text{Hom}_R(M, I^{t+d-2}) \xrightarrow{g_{t+d-1}} \text{Hom}_R(M, I^{t+d-1}) \longrightarrow 0.$$

Since M is finitely presented over R , we apply $(-)^{\vee}$ to the exact sequence above and obtain the following exact sequence in light of 2.7(ii):

$$(3.11.2) \quad M \otimes_R (I^{t+d-2})^{\vee} \xleftarrow{1 \otimes h_{t+d-1}^{\vee}} M \otimes_R (I^{t+d-1})^{\vee} \longleftarrow 0.$$

Next, let us return to the injective resolution I of N . Consider the exact sequence

$$0 \longrightarrow N \longrightarrow I^0 \xrightarrow{h_1} \dots \longrightarrow I^{t+d-2} \xrightarrow{h_{t+d-1}} I^{t+d-1} \xrightarrow{\theta} D \longrightarrow 0,$$

where D is the cokernel of the map h_{t+d-1} . Applying $(-)^{\vee}$ to the exact sequence above, we get an exact sequence

$$(3.11.3) \quad 0 \longleftarrow N^{\vee} \longleftarrow (I^0)^{\vee} \longleftarrow \dots \longleftarrow (I^{t+d-2})^{\vee} \xleftarrow{h_{t+d-1}^{\vee}} (I^{t+d-1})^{\vee} \xleftarrow{\theta^{\vee}} D^{\vee} \longleftarrow 0$$

with each $(I^i)^{\vee}$ flat over R ; see 2.7(iii). It follows that

$$\text{Ass}_R(D^{\vee}) \subseteq \text{Ass}_R((I^{t+d-1})^{\vee}) \subseteq \text{Ass}(R).$$

Claim 3. $(D^{\vee})_{\mathfrak{p}} = 0$ for all $\mathfrak{p} \in \text{Ass}(R)$.

Proof. Fix any $\mathfrak{p} \in \text{Ass}(R)$. From assumption (v) and 2.8, we see that

$$\text{fd}_{R_{\mathfrak{p}}}((N^{\vee})_{\mathfrak{p}}) \leq \dim(R_{\mathfrak{p}}) \leq d.$$

Thus, (3.11.3) localized at \mathfrak{p} gives rise to a flat resolution of $N_{\mathfrak{p}}$ over $R_{\mathfrak{p}}$, which can be used to compute $\text{Tor}_i^{R_{\mathfrak{p}}}(-, (N^{\vee})_{\mathfrak{p}})$. By 2.7(ii), we have

$$\text{Tor}_{t+d-1}^{R_{\mathfrak{p}}}(M_{\mathfrak{p}}, (N^{\vee})_{\mathfrak{p}}) = (\text{Tor}_{t+d-1}^R(M, N^{\vee}))_{\mathfrak{p}} \cong (\text{Ext}_R^{t+d-1}(M, N)^{\vee})_{\mathfrak{p}} = 0.$$

Moreover, we have $\text{Tor}_{t+d}^{R_{\mathfrak{p}}}(M_{\mathfrak{p}}, (N^{\vee})_{\mathfrak{p}}) = 0$ since $t+d > \text{fd}_{R_{\mathfrak{p}}}((N^{\vee})_{\mathfrak{p}})$. Next, we apply $M_{\mathfrak{p}} \otimes_{R_{\mathfrak{p}}} -$ to (3.11.3) localized at \mathfrak{p} . The vanishing of $\text{Tor}_i^{R_{\mathfrak{p}}}(M_{\mathfrak{p}}, (N^{\vee})_{\mathfrak{p}})$, for $i = t+d-1$ and $i = t+d$, forces the exact sequence

$$M_{\mathfrak{p}} \otimes_{R_{\mathfrak{p}}} ((I^{t+d-2})^{\vee})_{\mathfrak{p}} \xleftarrow{1 \otimes (h_{t+d-1}^{\vee})_{\mathfrak{p}}} M_{\mathfrak{p}} \otimes_{R_{\mathfrak{p}}} ((I^{t+d-1})^{\vee})_{\mathfrak{p}} \xleftarrow{1 \otimes (\theta^{\vee})_{\mathfrak{p}}} M_{\mathfrak{p}} \otimes_{R_{\mathfrak{p}}} (D^{\vee})_{\mathfrak{p}} \longleftarrow 0.$$

Comparing this with (3.11.2) localized at \mathfrak{p} , we see $M_{\mathfrak{p}} \otimes_{R_{\mathfrak{p}}} (D^{\vee})_{\mathfrak{p}} = 0$. Hence $(D^{\vee})_{\mathfrak{p}} = 0$ since $\text{Supp}_{R_{\mathfrak{p}}}(M_{\mathfrak{p}}) = \text{Spec}(R_{\mathfrak{p}})$; see 2.10. This completes the proof of the claim. \square

Overall, we have $\text{Ass}_R(D^{\vee}) \subseteq \text{Ass}(R)$ and $(D^{\vee})_{\mathfrak{p}} = 0$ for all $\mathfrak{p} \in \text{Ass}(R)$. This forces $D^{\vee} = 0$, which implies $D = 0$. Therefore, $\text{id}_R(N) < t + d$, so $\text{id}_R(N) \leq d$ by 2.8. \square

We are now ready to prove Theorem 3.2, a consequence of Propositions 3.10 and 3.11. Recall that $F : R \rightarrow R$ denotes the Frobenius endomorphism.

Proof of Theorem 3.2. It suffices to prove part (i) for the case where R is F-finite with $|k| = \infty$; see 2.3. Thus, we can obtain part (i) from part (ii) via duality; see 2.7(iii).

To prove part (ii), note that eM is a finitely generated (and hence a finitely presented) R -module with $\text{Supp}_R({}^eM) = \text{Supp}_R(M) = \text{Spec}(R)$. Since e is sufficiently large, there exists an M -regular sequence $\underline{x} = \{x_1, \dots, x_d\} \subseteq \mathfrak{m}$ such that $\mathfrak{m}({}^eM) \subseteq {}^e(\underline{x})M$; see 2.4. In light of 2.7(ii), the vanishing of $\text{Ext}_R^i({}^eM, N)$ implies the vanishing of $\text{Tor}_i^R({}^eM, N^{\vee})$ for all $i = t, \dots, t + d - 1$. Therefore, $\text{id}_{R_{\mathfrak{p}}}(N_{\mathfrak{p}}) < \infty$ and $\text{fd}_{R_{\mathfrak{p}}}((N^{\vee})_{\mathfrak{p}}) < \infty$ for all $\mathfrak{p} \in \text{Spec}(R) - \{\mathfrak{m}\}$; see Proposition 3.10. Note that $\text{Ass}(R) \subseteq \text{Spec}(R) - \{\mathfrak{m}\}$ since $\text{depth}(R) \geq 1$. Now we apply Proposition 3.11, with $R = S$ and $f = F^e$, and deduce that $\text{id}_R(N) \leq d$. \square

Remark 3.12. We conclude this section by pointing out some lower bounds for the integer e that ensure the validity of the proofs of certain previously stated results.

- (a) It is enough to assume $e \geq \lceil \log_p e_R(M) \rceil$ in part (i) of both Theorem 3.2 and Proposition 3.10.
- (b) When $|k| = \infty$, it is enough to assume $e \geq \lceil \log_p e_R(M) \rceil$ in part (ii) of both Theorem 3.2 and Proposition 3.10. When $|k| < \infty$, it is enough to assume $e \geq \max \{ \lceil \log_p e_R(M) \rceil, \text{drs}_p(M) \}$ in Theorem 3.2. When $|k| < \infty$, it is enough to assume $e \geq \max \{ \lceil \log_p e_R(M) \rceil, \text{crs}_p(M) \}$ in Proposition 3.10.
- (c) It is enough to assume $e \geq \lceil \log_p e_R(M) \rceil$ in Corollaries 3.5, 3.6, and 3.7.

It is proved in [14, 2.17] that, if R is an excellent ring and M is a finitely generated R -module, then the set $\sup \{ e_{R_{\mathfrak{m}}}(M_{\mathfrak{m}}) : \mathfrak{m} \in \text{Max}(R) \}$ is finite; see [14, 2.17]. We use this fact and state a global version of Theorem 3.2.

Theorem 3.13. *Let R be a d -dimensional ring of prime characteristic p , with $d \geq 1$, M be a finitely generated Cohen–Macaulay R -module such that $\text{Supp}_R(M) = \text{Spec}(R)$, and let N be an R -module. Assume $\text{depth}(R_{\mathfrak{m}}) \geq \min \{ 1, \dim(R_{\mathfrak{m}}) \}$ for each maximal ideal \mathfrak{m} of R .*

- (i) Assume R is excellent, $s = \sup \{e_{R_{\mathfrak{m}}}(M_{\mathfrak{m}}) : \mathfrak{m} \in \text{Max}(R)\}$, and let e be an integer such that $e \geq \lceil \log_p s \rceil$. Given $t \geq 1$, if $\text{Tor}_i^R({}^e M, N) = 0$ for all $i = t, \dots, t + d - 1$, then $\text{fd}_R(N) \leq d$.
- (ii) Assume R is F -finite and the residue field of $R_{\mathfrak{m}}$ is infinite for each maximal ideal \mathfrak{m} of R . Let e be an integer such that

$$e \geq \sup \{ \lceil \log_p e_{R_{\mathfrak{m}}}(M_{\mathfrak{m}}) \rceil, \text{drs}_p(M_{\mathfrak{m}}) : \mathfrak{m} \in \text{Max}(R) \}.$$

Given $t \geq 1$, if $\text{Ext}_R^i({}^e M, N) = 0$ for all $i = t, \dots, t + d - 1$, then $\text{id}_R(N) \leq d$.

Proof. Note that $s < \infty$ due to [14, 2.17]. In proving the first part, we have that

$$\text{Tor}_i^{R_{\mathfrak{m}}}({}^e M_{\mathfrak{m}}, N_{\mathfrak{m}}) = 0 \quad \text{for all } i = t, \dots, t + d - 1.$$

If $\dim(R_{\mathfrak{m}}) = 0$, then the residue field of $R_{\mathfrak{m}}$ is a direct summand of ${}^e M_{\mathfrak{m}}$, and hence $\text{fd}_{R_{\mathfrak{m}}}(N_{\mathfrak{m}}) \leq \dim(R_{\mathfrak{m}}) \leq d$; see 2.2(2)(i). On the other hand, if $\dim(R_{\mathfrak{m}}) \geq 1$, then part (i) of Theorem 3.2 yields $\text{fd}_{R_{\mathfrak{m}}}(N_{\mathfrak{m}}) \leq d$; see Remark 3.12(a). This implies that $\text{fd}_R(N) \leq d$ as flat dimension can be computed locally. We can prove the second part similarly since $\sup \{ \lceil \log_p e_{R_{\mathfrak{m}}}(M_{\mathfrak{m}}) \rceil, \text{drs}_p(M_{\mathfrak{m}}) : \mathfrak{m} \in \text{Max}(R) \}$ is finite under our setting; see Remark 3.12(a). \square

4. On the homological properties of the Frobenius endomorphism

The aim of this section is to prove Theorem 1.2(iii) and to establish Theorem 1.4. Let us point out that Theorem 1.4 generalizes [8, 1.3]; see 1.3.

The layout of this section is as follows: We begin by preparing some auxiliary results. Then we establish Theorem 1.2(iii) in Theorem 4.3. Finally, making use of 2.13 and Theorem 4.3, we produce a proof of Theorem 1.4. Along the way, we also discuss the sharpness of our results; see Examples 4.8 and 4.9.

The first auxiliary result, namely Proposition 4.1, is akin to [16, 2.6] and [22, Theorem A]. Even though it suffices to apply the proposition to the identity map $1_R : R \rightarrow R$ in the sequel, we present it more generally in terms of a ring homomorphism $f : R \rightarrow S$.

Proposition 4.1. *Let $f : (R, \mathfrak{m}, k) \rightarrow (S, \mathfrak{n}, \ell)$ be a local homomorphism of local rings of prime characteristic p , $M \neq 0$ be a finitely generated R -module, and let N be a finitely generated S -module. Set $d = \dim(R)$ and $v = \text{depth}_R(M)$. Given $t \geq 1$ and $e \geq \text{crs}_p(M)$, if $\text{Ext}_R^i(N, {}^e M) = 0$ for all $i = t, \dots, t + v$, then $\text{fd}_R(N) \leq t - 1$, and thus $\text{pd}_R(N) \leq t - 1 + d$.*

Proof. By our choice of e , there exists a maximal M -regular sequence $\underline{x} = \{x_1, \dots, x_v\}$ such that k is a direct summand of ${}^e(M/(\underline{x})M)$ over R ; see 2.2 and 2.4.

As x_1 is M -regular, there is a short exact sequence $0 \rightarrow {}^e M \rightarrow {}^e M \rightarrow {}^e(M/x_1 M) \rightarrow 0$. This, together with $\text{Ext}_R^i(N, {}^e M) = 0$ for $i = t, \dots, t + v$, implies that

$$\text{Ext}_R^i(N, {}^e(M/x_1M)) = 0 \quad \text{for all } i = t, \dots, t + v - 1.$$

Inductively, as $\underline{x} = \{x_1, \dots, x_v\}$ is M -regular, we get $\text{Ext}_R^t(N, {}^e(M/(\underline{x})M)) = 0$. Since k is a direct summand of ${}^e(M/(\underline{x})M)$, we see that $\text{Ext}_R^t(N, k) = 0$, which implies $\text{fd}_R(N) \leq t - 1$; see [22, 2.1]. Note that every flat R -module has projective dimension at most d ; see [28, 4.2.8]. Therefore, we conclude that $\text{pd}_R(N) \leq t - 1 + d$. \square

As in Propositions 3.11 and 4.1, we present Proposition 4.2 in the context of a general ring homomorphism $f : R \rightarrow S$, allowing us to distinguish various module structures in the proof. When Proposition 4.2 is applied in the proof of Theorem 4.3, the homomorphism will be $F^e : R \rightarrow R$, the e -th iteration of the Frobenius endomorphism.

Proposition 4.2. *Let (R, \mathfrak{m}, k) be a local ring, $f : R \rightarrow S$ be a ring homomorphism, N be a finitely generated R -module, and let M be a finitely generated S -module. Assume the following hold:*

- (i) $\text{Ext}_R^i(N, M) = 0$ for all $i = t, \dots, t + d - 1$ for some $d \geq 1$ and $t \geq 1$.
- (ii) There exists $\underline{x} = \{x_1, \dots, x_d\} \subseteq \text{Jac}(S)$ such that \underline{x} is M -regular and $\mathfrak{m}M \subseteq (\underline{x})M$.

Then $\text{Hom}_R(\Omega_R^t(N), M) = 0$.

Proof. Consider a minimal free resolution of N over R :

$$F = (\dots \rightarrow F_{t+d} \xrightarrow{h_{t+d}} F_{t+d-1} \xrightarrow{h_{t+d-1}} \dots \xrightarrow{h_{t+1}} F_t \xrightarrow{h_t} F_{t-1} \rightarrow \dots \rightarrow 0).$$

Applying $\text{Hom}_R(-, M)$ to F and using assumption (i), we get an induced exact sequence

$$(4.2.1) \quad M_{t+d} \xleftarrow{g_{t+d}} M_{t+d-1} \xleftarrow{g_{t+d-1}} \dots \xleftarrow{g_{t+1}} M_t \xleftarrow{g_t} M_{t-1} \xleftarrow{g_{t-1}} G \leftarrow 0,$$

in which $M_i = \text{Hom}_R(F_i, M)$, $g_i = \text{Hom}_R(h_i, M)$, and $G = \ker(g_t)$. All M_i , and hence G , are S -modules; see 2.6(ii). In fact, each M_i is isomorphic to a finite direct sum of M . Thus all M_i are finitely generated S -modules.

As F_i is free over R and x_1 is regular on M , an application of $\text{Hom}_R(F_i, -)$ to the exact sequence $0 \rightarrow M \xrightarrow{x_1} M \rightarrow M/x_1M \rightarrow 0$ induces a short exact sequence

$$(\Gamma_i) \quad 0 \longrightarrow M_i \xrightarrow{x_1} M_i \longrightarrow \text{Hom}_R(F_i, M/x_1M) \longrightarrow 0.$$

We combine the exact sequences (Γ_i) , for all $i = t - 1, \dots, t + d$, with the exact sequence (4.2.1) and obtain the following exact sequence of S -modules:

$$\text{Hom}_R(F_{t+d-1}, M/x_1M) \leftarrow \dots \leftarrow \text{Hom}_R(F_{t-1}, M/x_1M) \leftarrow G/x_1G \leftarrow 0.$$

Inductively, as $\underline{x} = \{x_1, \dots, x_d\}$ is an M -regular sequence, we realize the following exact sequence of S -modules that is naturally induced from (4.2.1):

$$(4.2.2) \quad \overline{M}_t \xleftarrow{\overline{g}_t} \overline{M}_{t-1} \xleftarrow{i} \overline{G} \longleftarrow 0,$$

in which $\overline{M}_i = \text{Hom}_R(F_i, M/(\underline{x})M)$, $\overline{g}_i = \text{Hom}_R(h_i, M/(\underline{x})M)$ and $\overline{G} = G/(\underline{x})G$. Up to isomorphism, we may write

$$\overline{M}_i = M_i/(\underline{x})M_i = M_i \otimes_S S/(\underline{x})S, \quad \overline{G} = G \otimes_S S/(\underline{x})S$$

and thus $i = \iota \otimes 1_{S/(\underline{x})S}$.

The assumption $\mathfrak{m}M \subseteq (\underline{x})M$ implies that $M/(\underline{x})M$ is annihilated by \mathfrak{m} . Since F is a minimal free resolution of N , we conclude $\overline{g}_i = \text{Hom}_R(h_i, M/(\underline{x})M) = 0$. Thus, the exactness of (4.2.2) forces $\text{im}(i) = \ker(\overline{g}_t) = \overline{M}_{t-1}$, meaning that $\text{im}(\iota) + (\underline{x})M_{t-1} = M_{t-1}$. As M_{t-1} is finitely generated over S and $(\underline{x})S \subseteq \text{Jac}(S)$, we obtain $\text{im}(\iota) = M_{t-1}$, thanks to Nakayama's lemma. This forces $\ker(g_{t+1}) = \text{im}(g_t) = 0$ due to the exactness of (4.2.1). Finally, as $\Omega_R^t(N) \cong \text{coker}(h_{t+1})$, we see that

$$\text{Hom}_R(\Omega_R^t(N), M) \cong \ker(\text{Hom}_R(h_{t+1}, M)) = \ker(g_{t+1}) = 0. \quad \square$$

Equipped with Propositions 4.1 and 4.2, we are now ready to prove the result stated in Theorem 1.2(iii).

Theorem 4.3. *Let (R, \mathfrak{m}, k) be a d -dimensional local ring of prime characteristic p , M be a finitely generated Cohen–Macaulay R -module such that $\text{Supp}_R(M) = \text{Spec}(R)$, and let N be a finitely generated R -module. Assume $\text{depth}(R) \geq 1$. Given $t \geq 1$ and $e \gg 0$, if $\text{Ext}_R^i(N, {}^eM) = 0$ for all $i = t, \dots, t + d - 1$, then $\text{pd}_R(N) \leq \min\{t - 1, \text{depth}(R)\}$.*

Proof. Since e is sufficiently large, there exists $\underline{x} = \{x_1, \dots, x_d\} \subseteq \mathfrak{m}$ such that \underline{x} is M -regular and $\mathfrak{m}({}^eM) \subseteq {}^e((\underline{x})M)$; see 2.4. Upon an application of Proposition 4.2 to the map $F^e : R \rightarrow R$, we see that the vanishing of $\text{Ext}_R^i(N, {}^eM)$ for $i = t, \dots, t + d - 1$ implies $\text{Hom}_R(\Omega_R^t(N), {}^eM) = 0$.

To prove the claim by contradiction, suppose that $\Omega_R^t(N) \neq 0$ and select \mathfrak{p} in $\text{Ass}_R(\Omega_R^t(N)) \subseteq \text{Ass}(R)$; then $\mathfrak{p} \neq \mathfrak{m}$ since $\text{depth}(R) \geq 1$. It follows from Proposition 4.1 that $\text{pd}_{R_{\mathfrak{p}}}(N_{\mathfrak{p}}) < \infty$. Hence, $N_{\mathfrak{p}}$ is free by the Auslander–Buchsbaum formula. So, $\Omega_R^t(N)_{\mathfrak{p}} \neq 0$ is free over $R_{\mathfrak{p}}$. Also, since $\text{Supp}_R({}^eM) = \text{Supp}_R(M) = \text{Spec}(R)$, we see that $({}^eM)_{\mathfrak{p}} \neq 0$. Hence $\text{Hom}_{R_{\mathfrak{p}}}(\Omega_R^t(N)_{\mathfrak{p}}, ({}^eM)_{\mathfrak{p}}) \neq 0$, which contradicts the conclusion $\text{Hom}_R(\Omega_R^t(N), {}^eM) = 0$. Thus $\Omega_R^t(N) = 0$, so $\text{pd}_R(N) \leq t - 1$. This proves that $\text{pd}_R(N) \leq \min\{t - 1, \text{depth}(R)\}$. \square

We now record several corollaries of Theorem 4.3. It is worth noting that the positive depth assumption in the theorem is necessary; see Example 3.4.

Corollary 4.4. *Let (R, \mathfrak{m}, k) be a d -dimensional local ring of prime characteristic p , M be a finitely generated Cohen–Macaulay R -module such that $\text{Supp}_R(M) = \text{Spec}(R)$, and let N be a finitely generated R -module. Assume $\text{depth}(R) \geq 1$. Given $t \geq 1$ and $e \gg 0$, assume that at least one of the following conditions holds:*

- (i) $\text{Tor}_i^R(N, ({}^e M)^\vee) = 0$ for all $i = t, \dots, t + d - 1$.
- (ii) $\text{Tor}_i^R(N, {}^e(M^\vee)) = 0$ for all $i = t, \dots, t + d - 1$.

Then $\text{pd}_R(N) \leq \min\{t - 1, \text{depth}(R)\}$.

Proof. The vanishing of $\text{Tor}_i^R(N, ({}^e M)^\vee)$ yields the vanishing of $\text{Ext}_R^i(N, {}^e M)$; see 2.7(ii). Thus, case (i) follows from Theorem 4.3. Similarly, the vanishing of $\text{Tor}_i^R(N, {}^e(M^\vee))$ yields the vanishing of $\text{Ext}_R^i(N, {}^e M)$ by 2.7(ii) where $R \rightarrow S$ is the Frobenius F^e with $(S, \mathfrak{n}, \ell) = (R, \mathfrak{m}, k)$, $A = N$ over R , $B = M$ over S , and $E = E_S(\ell)$ is the injective hull of ℓ over S . Therefore, case (ii) also follows from Theorem 4.3. \square

Next, we provide a proof of Theorem 1.4, restated here as Corollary 4.5 for convenience. Before doing so, we present a few facts needed for the argument.

Recall that $F_R^e(-)$ denotes the scalar extension along $F^e : R \rightarrow R$, the e -th iteration of the Frobenius endomorphism. Given R -modules X and Y , it follows that

$$X \otimes_R {}^e Y \cong (F_R^e(X) \otimes_R Y) \quad \text{and} \quad F_{R_p}^e(X_p) \cong (F_R^e(X))_p \quad \text{for all } p \in \text{Spec}(R).$$

Moreover, we have $\text{Supp}_R(X) = \text{Supp}_R(F_R^e(X))$. Although the finitely generated case suffices for our argument in Corollary 4.5, we briefly discuss the general situation where X may not be finitely generated. To establish the equality $\text{Supp}_R(X) = \text{Supp}_R(F_R^e(X))$, it is enough to assume R is local. If $X \neq 0$ is finitely generated, then $F_R^e(X) \neq 0$ via a surjection $X \twoheadrightarrow k$. In the general case, pick a flat local ring homomorphism $\varphi : R \rightarrow A$, where A is F-finite; such a ring map always exists by 2.3. Setting $S = R$ and $B = A$, and letting $F_A^e : A \rightarrow B$ denote the e -th iteration of the Frobenius of A , we see that the composition

$$\varphi \circ F_R^e : R \xrightarrow{F_R^e} S \xrightarrow{\varphi} B$$

agrees with $F_A^e \circ \varphi : R \xrightarrow{\varphi} A \xrightarrow{F_A^e} B$. Thus

$$(X \otimes_R S) = 0 \implies (X \otimes_R S) \otimes_S B = 0 \implies (X \otimes_R A) \otimes_A B = 0 \implies X \otimes_R A = 0 \implies X = 0.$$

Here, the second implication follows since $\varphi \circ F_R^e = F_A^e \circ \varphi$, the third implication follows from 2.10(iii) as A is F-finite, while the fourth one holds because φ is faithfully flat. This establishes what we want to show. We also refer the reader to Marley’s result in [19, 2.1] for an alternative proof of $\text{Supp}_R(X) = \text{Supp}_R(F_R^e(X))$.

Corollary 4.5. *Let (R, \mathfrak{m}, k) be a local ring of prime characteristic p , and let M and N be finitely generated R -modules. Assume the following conditions hold:*

- (i) M is Cohen–Macaulay and $\text{Supp}_R(M) = \text{Spec}(R)$.
- (ii) N is generically free, that is, $N_{\mathfrak{p}}$ is a free $R_{\mathfrak{p}}$ -module for all $\mathfrak{p} \in \text{Ass}(R)$.

If $F_R^e(N) \otimes_R M$ is a maximal Cohen–Macaulay R -module for some $e \gg 0$, then N is free.

Proof. Without loss of generality we may assume R is F-finite (and also $|k| = \infty$); see 2.3. Set $d = \dim(R)$ and proceed by induction on d .

As R is F-finite, the assumption that $F_R^e(N) \otimes_R M$ is maximal Cohen–Macaulay can be interpreted as that the module $N \otimes_R {}^eM$ is maximal Cohen–Macaulay. If $\text{depth}(R) = 0$, then N is free since we assume it is generically free. Hence, we assume $\text{depth}(R) \geq 1$. Note that, by the induction hypothesis, we may assume $N_{\mathfrak{p}}$ is a free $R_{\mathfrak{p}}$ -module for all $\mathfrak{p} \in \text{Spec}(R) - \{\mathfrak{m}\}$. Note also that $\text{depth}_R({}^eM) = d = \text{depth}_R(N \otimes_R {}^eM)$. Therefore, we use 2.13 for the case where $n = d$ and conclude that $\text{Ext}_R^i(\text{Tr}_R N, {}^eM) = 0$ for all $i = 1, \dots, d$. Now, Theorem 4.3 implies that $\text{Tr}_R N$ is free. Consequently, N is free; see 2.12(iv). \square

Corollary 4.6. *Let (R, \mathfrak{m}, k) be a local ring of prime characteristic p , and let M and N be finitely generated R -modules. Assume the following conditions hold:*

- (i) M is Cohen–Macaulay.
- (ii) M has constant rank on $\text{Min}(R)$ and N is generically free.

If $F_R^e(N) \otimes_R M$ is maximal Cohen–Macaulay for some $e \gg 0$, then N is free.

Proof. Suppose $\text{Min}(R) \not\subseteq \text{Supp}_R(M)$. Then $\text{Supp}_R(M) \cap \text{Min}(R) = \emptyset$ since M has constant rank on $\text{Min}(R)$. This implies that $\dim_R(M) < \dim(R)$. However, this is not possible as $\dim(R) = \dim_R(F_R^e(N) \otimes_R M) \leq \dim_R(M)$. So, $\text{Min}(R) \subseteq \text{Supp}_R(M)$ and hence $\text{Supp}_R(M) = \text{Spec}(R)$. Now, the claim follows from Corollary 4.5. \square

Corollary 4.7. *Let (R, \mathfrak{m}, k) be a local ring of prime characteristic p , M be a finitely generated Cohen–Macaulay R -module, and let N be a finitely generated R -module. Assume:*

- (i) For each $\mathfrak{p} \in \text{Min}(R)$, there is an integer $r_{\mathfrak{p}} \geq 1$ such that $M_{\mathfrak{p}} \cong R_{\mathfrak{p}}^{\oplus r_{\mathfrak{p}}}$.
- (ii) N is generically free and $\dim_R(N) = \dim(R)$.

If $F_R^e(N) \otimes_R M$ is Cohen–Macaulay for some $e \gg 0$, then N is free.

Proof. It follows that $\dim_R(F_R^e(N) \otimes_R M) = \dim_R(F_R^e(N)) = \dim_R(N) = \dim(R)$ because $\text{Supp}_R(M) = \text{Spec}(R)$. Now, the claim follows from Corollary 4.5. \square

We illustrate the sharpness of [Theorem 1.4](#) with some examples. Specifically, [Example 4.8](#) demonstrates that the theorem’s conclusion may fail if the module M in question does not have full support. Similarly, [Example 4.9](#) highlights the necessity of the assumption that M is generically free for the theorem to hold.

Example 4.8. Let $R = \mathbb{F}_p[[x, y]]/(xy)$, $M = R/(x)$, and $N = M$. Note that M is maximal Cohen–Macaulay, and $N_{\mathfrak{p}}$ is a free $R_{\mathfrak{p}}$ -module for all $\mathfrak{p} \in \text{Ass}(R)$. It follows that $N \otimes_R {}^e M$ is maximal Cohen–Macaulay for all $e \geq 0$ because $N \otimes_R {}^e M \cong N \otimes_R M^{\oplus p^e} \cong M^{\oplus p^e}$. However, N is not free. Here $\text{Spec}(R) \ni (y) \notin \text{Supp}_R(M)$. This example also shows that the constant rank assumption in [Corollary 4.6](#) is necessary. As a separate observation, $\text{Ext}_R^1(M, M) = 0$. Thus, the full support assumption is necessary in [Theorem 4.3](#).

Example 4.9. Let $R = \mathbb{F}_p[[x, y]]/(x^2, xy)$, $M = R/(x)$, and $N = M$. Note that M is Cohen–Macaulay, $\text{Supp}_R(M) = \text{Spec}(R)$, and $N_{\mathfrak{p}}$ is a free $R_{\mathfrak{p}}$ -module for all $\mathfrak{p} \in \text{Min}(R)$. It follows that $N \otimes_R {}^e M$ is maximal Cohen–Macaulay for all $e \geq 0$ since $N \otimes_R {}^e M \cong M^{\oplus p^e}$. However, N is not free. Here we have that $\text{Ass}(R) \ni \mathfrak{m} \notin \text{Min}(R)$.

It is proved in [\[4, 1.1\]](#) that, if there is a finitely generated module $N \neq 0$ over a local ring R of prime characteristic p such that $\text{fd}_R({}^r N) < \infty$ or $\text{id}_R({}^r N) < \infty$ for some $r \geq 1$, then R is regular. We use this fact and obtain the following consequences of [Theorem 1.2](#); cf. [\[29, 6.8 and 6.10\]](#).

Corollary 4.10. *Let (R, \mathfrak{m}, k) be a d -dimensional local ring of prime characteristic p , M be a finitely generated Cohen–Macaulay R -module such that $\text{Supp}_R(M) = \text{Spec}(R)$, and let $N \neq 0$ be a finitely generated R -module. Assume $\text{depth}(R) \geq 1$. Given $t \geq 1, r \geq 1$ and $e \gg 0$, we further assume that at least one of the following conditions holds:*

- (i) $\text{Tor}_i^R({}^e M, {}^r N) = 0$ for all $i = t, \dots, t + d - 1$.
- (ii) R is F -finite and $\text{Ext}_R^i({}^e M, {}^r N) = 0$ for all $i = t, \dots, t + d - 1$.
- (iii) ${}^r N$ is finitely generated over R and $\text{Ext}_R^i({}^r N, {}^e M) = 0$ for all $i = t, \dots, t + d - 1$.

Then R is regular.

Proof. In view of [Theorem 1.2](#), the claims follow from [\[4, 1.1\]](#). □

Remark 4.11. We point out some lower bounds for the integer e that ensure the validity of the proofs of certain previously stated results.

- (a) It is enough to assume $e \geq \lceil \log_p e_R(M) \rceil$ in [Corollaries 1.5 and 4.5](#).
- (b) It is enough to assume $e \geq \lceil \log_p e_R(M) \rceil$ in both [Theorem 4.3](#) and [Corollary 4.4](#). This is because we may use [2.3](#) and assume $|k| = \infty$ in the proofs of these results.

- (c) It is enough to assume $e \geq \max \{ \lceil \log_p e_R(M) \rceil, \text{drs}_p(M) \}$ in [Corollary 4.10](#) in general. If $|k| = \infty$, however, it suffices to assume $e \geq \lceil \log_p e_R(M) \rceil$ in [Corollary 4.10](#).

We do not know whether the Cohen–Macaulay assumption on M is necessary in [Theorem 1.4](#). This raises the following question:

Question 4.12. Let (R, \mathfrak{m}, k) be a local ring of prime characteristic p and let M and N be finitely generated R -modules. Assume $\text{Supp}_R(M) = \text{Spec}(R)$ and N is generically free. If $F_R^e(N) \otimes_R M$ is maximal Cohen–Macaulay for some $e \gg 0$, then must N be free? What if R is Cohen–Macaulay, or $N = M$?

We conclude this section with a global version of [Theorem 4.3](#); we skip its proof as it is similar to that of [Theorem 3.13](#).

Theorem 4.13. *Let R be a d -dimensional excellent ring of prime characteristic p , M be a finitely generated Cohen–Macaulay R -module such that $\text{Supp}_R(M) = \text{Spec}(R)$, and let N be a finitely generated R -module. Assume $\text{depth}(R_{\mathfrak{m}}) \geq \min\{1, \dim(R_{\mathfrak{m}})\}$ for each maximal ideal \mathfrak{m} of R . Set*

$$s = \sup \{ e_{R_{\mathfrak{m}}}(M_{\mathfrak{m}}) : \mathfrak{m} \in \text{Max}(R) \},$$

and let e be an integer such that $e \geq \lceil \log_p s \rceil$. Given $t \geq 1$, if $\text{Ext}_R^i(N, {}^e M) = 0$ for all $i = t, \dots, t + d - 1$, then $\text{pd}_R(N) \leq \min\{t - 1, \text{depth}(R)\}$.

Acknowledgements

Part of this work was completed during Celikbas’s visit to the Department of Mathematics and Statistics at Georgia State University in February 2018, and during the visit of Celikbas, Sadeghi, and Yao to the Nesin Mathematics Village in İzmir, Türkiye, in May 2019.

The authors thank these two institutions for their hospitality. They also thank Mohsen Asgarzadeh for his comments on a previous version of the manuscript. Finally, the authors thank the referee for a careful reading of the manuscript and for helpful comments.

References

- [1] M. Auslander and M. Bridger, *Stable module theory*, Memoirs of the American Mathematical Society **94**, American Mathematical Society, 1969. [MR](#)
- [2] M. Auslander and D. A. Buchsbaum, “Homological dimension in noetherian rings, II”, *Trans. Amer. Math. Soc.* **88** (1958), 194–206. [MR](#)
- [3] L. L. Avramov and C. Miller, “Frobenius powers of complete intersections”, *Math. Res. Lett.* **8**:1-2 (2001), 225–232. [MR](#)
- [4] L. L. Avramov, M. Hochster, S. B. Iyengar, and Y. Yao, “Homological invariants of modules over contracting endomorphisms”, *Math. Ann.* **353**:2 (2012), 275–291. [MR](#)

- [5] H. Bass, “On the ubiquity of Gorenstein rings”, *Math. Z.* **82** (1963), 8–28. [MR](#)
- [6] W. Bruns and J. Herzog, *Cohen–Macaulay rings*, Cambridge Studies in Advanced Mathematics **39**, Cambridge University Press, 1993. [MR](#)
- [7] L. Burch, “On ideals of finite homological dimension in local rings”, *Proc. Cambridge Philos. Soc.* **64** (1968), 941–948. [MR](#)
- [8] O. Celikbas, A. Sadeghi, and Y. Yao, “Tensoring with the Frobenius endomorphism”, *Homology Homotopy Appl.* **20**:1 (2018), 251–257. [MR](#)
- [9] T. H. Funk, *Frobenius and homological dimensions of complexes*, PhD thesis, University of Nebraska – Lincoln, 2021, available at <https://digitalcommons.unl.edu/mathstudent/107>.
- [10] T. Funk and T. Marley, “Correction to: Frobenius and homological dimensions of complexes”, *Collect. Math.* **71**:2 (2020), 299–300. [MR](#)
- [11] T. Funk and T. Marley, “Frobenius and homological dimensions of complexes”, *Collect. Math.* **71**:2 (2020), 287–297. [MR](#)
- [12] J. Herzog, “Ringe der Charakteristik p und Frobeniusfunktoren”, *Math. Z.* **140** (1974), 67–78. [MR](#)
- [13] M. Hochster, “Cyclic purity versus purity in excellent Noetherian rings”, *Trans. Amer. Math. Soc.* **231**:2 (1977), 463–488. [MR](#)
- [14] M. Hochster and Y. Yao, “Frobenius splitting, strong F-regularity, and small Cohen–Macaulay modules”, *Trans. Amer. Math. Soc.* **376**:9 (2023), 6729–6765. [MR](#)
- [15] S. B. Iyengar, G. J. Leuschke, A. Leykin, C. Miller, E. Miller, A. K. Singh, and U. Walther, *Twenty-four hours of local cohomology*, Graduate Studies in Mathematics **87**, American Mathematical Society, 2007. [MR](#)
- [16] J. Koh and K. Lee, “Some restrictions on the maps in minimal resolutions”, *J. Algebra* **202**:2 (1998), 671–689. [MR](#)
- [17] E. Kunz, “Characterizations of regular local rings of characteristic p ”, *Amer. J. Math.* **91** (1969), 772–784. [MR](#)
- [18] J. Li, “Frobenius criteria of freeness and Gorensteinness”, *Arch. Math. (Basel)* **98**:6 (2012), 499–506. [MR](#)
- [19] T. Marley, “The Frobenius functor and injective modules”, *Proc. Amer. Math. Soc.* **142**:6 (2014), 1911–1923. [MR](#)
- [20] H. Matsumura, *Commutative ring theory*, 2nd ed., Cambridge Studies in Advanced Mathematics **8**, Cambridge University Press, 1989. [MR](#)
- [21] C. Miller, “The Frobenius endomorphism and homological dimensions”, pp. 207–234 in *Commutative algebra* (Grenoble and Lyon, 2001), *Contemp. Math.* **331**, Amer. Math. Soc., 2003. [MR](#)
- [22] S. Nasseh, M. Tousei, and S. Yassemi, “Characterization of modules of finite projective dimension via Frobenius functors”, *Manuscripta Math.* **130**:4 (2009), 425–431. [MR](#)
- [23] C. Peskine and L. Szpiro, “Sur la topologie des sous-schémas fermés d’un schéma localement noethérien, définis comme support d’un faisceau cohérent localement de dimension projective finie”, *C. R. Acad. Sci. Paris Sér. A-B* **269** (1969), A49–A51. [MR](#)
- [24] C. Peskine and L. Szpiro, “Dimension projective finie et cohomologie locale. Applications à la démonstration de conjectures de M. Auslander, H. Bass et A. Grothendieck”, *Inst. Hautes Études Sci. Publ. Math.* **42** (1973), 47–119. [MR](#)
- [25] A. G. Rodicio, “On a result of Avramov”, *Manuscripta Math.* **62**:2 (1988), 181–185. [MR](#)

- [26] J. J. Rotman, *An introduction to homological algebra*, 2nd ed., Springer, 2009. [MR](#)
- [27] R. Takahashi and Y. Yoshino, “Characterizing Cohen-Macaulay local rings by Frobenius maps”, *Proc. Amer. Math. Soc.* **132**:11 (2004), 3177–3187. [MR](#)
- [28] J. Xu, *Flat covers of modules*, Lecture Notes in Mathematics **1634**, Springer, 1996. [MR](#)
- [29] M. R. Zargar, O. Celikbas, M. Gheibi, and A. Sadeghi, “Homological dimensions of rigid modules”, *Kyoto J. Math.* **58**:3 (2018), 639–669. [MR](#)

Received March 13, 2025. Revised January 31, 2026.

OLGUR CELIKBAS
SCHOOL OF MATHEMATICAL AND DATA SCIENCES
WEST VIRGINIA UNIVERSITY
MORGANTOWN, WV
UNITED STATES
olgur.celikbas@math.wvu.edu

ARASH SADEGHI
DOKHANIAT 49179-66686
GORGAN
IRAN
sadeghiarash61@gmail.com

YONGWEI YAO
DEPARTMENT OF MATHEMATICS AND STATISTICS
GEORGIA STATE UNIVERSITY
ATLANTA, GA
UNITED STATES
yyao@gsu.edu

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Fakultät für Mathematik
Universität Wien
Vienna, Austria
matthias.aschenbrenner@univie.ac.at

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Atsushi Ichino
Department of Mathematics
Kyoto University
Kyoto 606-8502, Japan
atsushi.ichino@gmail.com

Kefeng Liu
School of Sciences
Chongqing University of Technology
Chongqing 400054, China
liu@math.ucla.edu

Sucharit Sarkar
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
sucharit@math.ucla.edu

Dimitri Shlyakhtenko
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
shlyakht@ipam.ucla.edu

Ruixiang Zhang
Department of Mathematics
University of California
Berkeley, CA 94720-3840
ruixiang@berkeley.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2026 is US \$710/year for the electronic version, and \$965/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2026 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 342 No. 2 June 2026

On multilinear maximal operators along homogeneous curves	207
LARS BECKER and BEN KRAUSE	
The plain sphere number of a link	217
RYAN BLAIR, ALEXANDRA KJUCHUKOVA and ELLA PFAFF	
On the test properties of the Frobenius endomorphism	235
OLGUR CELIKBAS, ARASH SADEGHI and YONGWEI YAO	
Symplectic automorphisms of a surface with genus two fibration and their action on CH_0	259
JIABIN DU and WENFEI LIU	
Non-braid-positive hyperbolic L -space knots	275
KEISUKE HIMENO	
The Rankin–Selberg integral on GSp_2 for square-free levels	299
SEIJI KUGA and MASAO TSUZUKI	
Sum of the squares of the p' -character degrees	351
NGUYEN N. HUNG, J. MIQUEL MARTÍNEZ and GABRIEL NAVARRO	
On the first eigenvalue of the Hodge Laplacian of submanifolds	381
CHRISTOS-RAENT ONTI	
Howe duality for the dual pair $\text{SL}_2(\mathbb{R}) \times F_{4,1}$: a ping pong of K -types	387
GORDAN SAVIN	
Liouville theorems and new gradient estimates for positive solutions to $\Delta_p u + au^q = 0$ on a complete manifold	395
YOUDE WANG and LIQIN ZHANG	
p -nuclearity of reduced group L^p -operator algebras	427
ZHEN WANG	
A higher-rank analog of the strong openness property	441
JINGCAO WU	