

*Pacific
Journal of
Mathematics*

**A HIGHER-RANK ANALOG OF THE
STRONG OPENNESS PROPERTY**

JINGCAO WU

A HIGHER-RANK ANALOG OF THE STRONG OPENNESS PROPERTY

JINGCAO WU

We study a strong openness property for singular Hermitian vector bundles (E, h) that are Griffiths-semipositive.

1. Introduction

The strong openness conjecture is an important subject in complex geometry. It was completely solved by Q. Guan and X. Zhou as follows:

Theorem 1.1 [11, Theorem 1.1]. *Let Δ^n be the unit polydisc, and let (L, φ) be a singular Hermitian line bundle on Δ^n . Assume that the associated curvature satisfies $i\Theta_{L, \varphi} \geq 0$, and that F is a holomorphic section of L which satisfies*

$$\int_{\Delta^n} |F|^2 e^{-\varphi} d\lambda < \infty,$$

where $d\lambda$ is the Lebesgue measure. After shrinking Δ^n if necessary, there exists a positive ε such that

$$\int_{\Delta^n} |F|^2 e^{-(1+\varepsilon)\varphi} d\lambda < \infty.$$

Equivalently, the following equality concerning multiplier ideal sheaves holds:

$$(1) \quad \bigcup_{\varepsilon > 0} \mathcal{I}((1+\varepsilon)\varphi) = \mathcal{I}(\varphi).$$

It has led to fruitful developments, such as [3; 1; 2; 13; 9; 12]. Among them, we mention the following variant for later use.

Corollary 1.1 [3, Corollary B.2]. *Let φ, ψ be plurisubharmonic functions on a domain $U \subseteq \mathbb{C}^n$. Assume that ψ has analytic singularities. Then for any compact subset $K \Subset U$, there exists a positive ε such that*

$$e^{\psi-\varphi} \in L_{\text{loc}}^1(U, K) \quad \text{if and only if} \quad e^{\psi-(1+\varepsilon)\varphi} \in L_{\text{loc}}^1(U, K).$$

This research was supported by NSFC (grant 12201381) and Shanghai Scientific Sailing Program (grant 22YF1412500).

MSC2020: [32L10](#), [32U05](#), [32D15](#).

Keywords: strong openness, singular metric, vector bundle.

Here $L^1_{\text{loc}}(U, K)$ denotes the set of Lebesgue measurable functions f defined in a neighborhood of K in U such that $|f|$ is locally integrable in a neighborhood of every point $x \in K$.

It is natural to ask if the *strong openness property* holds for the higher-rank vector bundles. There are some discussions in [16; 18; 19], and we should make a brief review first. For an arbitrary holomorphic vector bundle E and a singular Hermitian metric h , they considered the multiplier submodule sheaf $\mathcal{E}(h)$ defined as

$$\mathcal{E}(h)_x := \{F_x \in E_x \mid |F_x|_h^2 \text{ is integrable around } x\}.$$

Suppose that $\{h_j\}$ is a sequence of singular metrics decreasing to h . Then, under a certain Nakano-type positivity (on $\{h_j\}$ or h), they concluded that

$$(2) \quad \bigcup_j \mathcal{E}(h_j) = \mathcal{E}(h).$$

In this paper, we investigate this problem from another point of view. More precisely, we apply the deep relationship between the Hermitian metrics on E and its tautological line bundle $\mathcal{O}_E(1)$, to reduce everything to the rank one case. This theory is known as the Finsler geometry, which will be recalled later. Then we obtain that

Theorem 1.2. *Let E be a holomorphic vector bundle of rank r over Δ^n , and let h be a singular Hermitian metric on E such that (E, h) is Griffiths semipositive. Assume that $\det h$ has analytic singularities. If F is a holomorphic section of E satisfying*

$$\int_{\Delta^n} |F|_h^2 d\lambda < \infty,$$

then after shrinking Δ^n if necessary, there exists a positive ε (independent of F) such that

$$\int_{\Delta^n} |F|_h^{2(1+\varepsilon)} d\lambda < \infty.$$

In principle, the Griffiths-type positivity is strictly weaker than the Nakano-type positivity. But we should also notice that [16; 18; 19] made no restrictions on the singularity of $\det h$. Hence, there is no direct relationship between [16; 18; 19] and **Theorem 1.2**.

Next, let us consider general singular metrics. Remember that in the line bundle case, if we furthermore suppose that $\mathcal{S}(\varphi) = \mathcal{O}_{\Delta^n}$, the equality (1) is then called the *openness property*. It was proved by [4], and certainly can be seen as a special case of **Theorem 1.1**. We generalize it to the higher-rank vector bundles as follows:

Theorem 1.3. *Let E be a holomorphic vector bundle of rank r over Δ^n_R , where Δ^n_R is a polydisc with radius R slightly larger than 1. Let h be a singular Hermitian*

metric on E such that (E, h) is Griffiths semipositive, and let φ be the metric on $\mathcal{O}_E(1)$ induced by h . Assume that

$$(3) \quad \mathcal{I}((r + 1)\varphi) = \mathcal{O}_{\mathbb{P}(E^*)}.$$

Then there exist positive numbers \hat{R} and ε such that

$$\int_{\Delta_{\hat{R}}^n} |F|_h^{2(1+\varepsilon)} d\lambda < \infty$$

for any holomorphic section F of E .

As we will see in Section 4, (3) implies that

$$\int_{\Delta_{\hat{R}}^n} |F|_h^2 d\lambda < \infty$$

for any F , namely

$$\mathcal{E}(h) = E.$$

But the converse seems not obvious.

This paper is organized as follows: in Section 2 we will review the basic materials of singular metrics and Finsler geometry, and in Section 3 we give the proof of Theorem 1.2. In the end we prove Theorem 1.3.

2. Preliminary

Let X be a complex manifold of dimension n , and let $p : E \rightarrow X$ be a holomorphic vector bundle over X of rank r .

2.1. Singular Hermitian metrics.

Definition 2.1 [5; 21; 22]. A singular Hermitian metric on E is a map h that associates to every point $x \in X$ a singular Hermitian inner product $|\cdot|_{h,x} : E_x \rightarrow [0, +\infty]$ on the complex vector space E_x , subject to the following two conditions:

1. h is finite and positive definite almost everywhere, meaning that for all x outside a set of Lebesgue measure zero, $|\cdot|_{h,x}$ is a Hermitian inner product on E_x ;
2. h is measurable, meaning that the function

$$|F|_h : U \rightarrow [0, +\infty], \quad x \mapsto |F(x)|_{h,x},$$

is measurable for any open $U \subseteq X$ and $F \in \Gamma(U, E)$.

Definition 2.2 [22]. Let h be a singular Hermitian metric on E , which canonically induces a singular metric h^* on the dual bundle E^* .

- (1) (E, h) is called Griffiths-seminegative (or negatively curved) if, for any (local) holomorphic section F of E , the function $\log |F|_h^2$ is plurisubharmonic.

(2) (E, h) is called Griffiths-semipositive (or positively curved) if (E^*, h^*) is Griffiths-seminegative.

Proposition 2.1 [22, Proposition 1.3]. *Let E be a holomorphic vector bundle over Δ^n , equipped with a singular Hermitian metric h .*

- (1) (See also [5], Proposition 3.1.) *If (E, h) is Griffiths-semipositive, then over any smaller polydisk there is a sequence of smooth, Griffiths positive metrics $\{h_\nu\}$ increasing pointwise to h .*
- (2) *If (E, h) is Griffiths-seminegative, then $\log \det h$ is a plurisubharmonic function.*

Definition 2.3 [6]. A plurisubharmonic function φ is said to have analytic singularities if φ can be written locally as

$$\varphi = \alpha \log(|f_1|^2 + \dots + |f_N|^2) + v,$$

where $\alpha \in \mathbb{R}_+$, v is a bounded function and the f_j are holomorphic functions.

When (E, h) is a Griffiths-semipositive singular Hermitian vector bundle, the function $-\log \det h$ is plurisubharmonic by Proposition 2.1(2). Then we can ask it has analytic singularities, and briefly say that $\det h$ has analytic singularities. Obviously, it not necessarily implies that h itself has analytic singularities.

2.2. Finsler geometry revisited. We only collect the necessary materials. One could refer to [8; 15] for a sophisticated comprehension.

Let $x = (x^1, \dots, x^n)$ be a local coordinate system in X and let $w = (w^0, \dots, w^{r-1})$ be the fiber coordinate system defined by a local holomorphic frame

$$W = \{W^0, \dots, W^{r-1}\}$$

of E . Let h be a smooth Hermitian metric on E . We write

$$h_i = \frac{\partial h}{\partial x^i}, \quad h_{\bar{j}} = \frac{\partial h}{\partial \bar{x}^j}, \quad h_{i\bar{j}} = \frac{\partial^2 h}{\partial x^i \partial \bar{x}^j}, \quad h_{i\alpha} = \frac{\partial^2 h}{\partial x^i \partial w^\alpha},$$

and so on, to denote the differentiation with respect to x^i, \bar{x}^j ($1 \leq i, j \leq n$) and w^α, \bar{w}^β ($0 \leq \alpha, \beta \leq r - 1$).

Denote by $q : \mathbb{P}(E) \rightarrow X$ the natural projection from the projectivized bundle to the ambient space, and h induces a Hermitian metric q^*h on q^*E . Then as a subbundle,

$$\mathcal{O}_{\mathbb{P}(E)}(-1) := \{((x, [w]), Z) \in q^*E \mid Z = \lambda w, \lambda \in \mathbb{C}\}$$

inherits a metric from q^*E , whose weight function is denoted by $-\psi$. Accordingly, we can define the metrics $\psi, -\varphi$ and φ on $\mathcal{O}_{\mathbb{P}(E)}(1), \mathcal{O}_{\mathbb{P}(E^*)}(-1)$ and $\mathcal{O}_{\mathbb{P}(E^*)}(1)$ respectively in an obvious way. By definition, one easily verifies: when (E, h) is

Griffiths-semipositive, $(\mathcal{O}_{\mathbb{P}(E^*)}(1), \varphi)$ is semipositive. For this reason, we usually denote $\mathcal{O}_{\mathbb{P}(E^*)}(1)$ by $\mathcal{O}_E(1)$, and call it the tautological line bundle of E .

Remark 2.1. Following the same procedure, a singular Hermitian metric h on E induces singular metrics on $\mathcal{O}_{\mathbb{P}(E)}(-1)$, $\mathcal{O}_{\mathbb{P}(E)}(1)$, $\mathcal{O}_E(-1)$ and $\mathcal{O}_E(1)$ respectively.

Moreover, if (E, h) is Griffiths-semipositive, $\mathcal{O}_E(1)$ is pseudo-effective equipped with the corresponding metric as is shown in [21], Proposition 2.3.5.

Next let us recall the celebrated curvature formula in [15]. Remember that ψ is the weight function of the metric on $\mathcal{O}_{\mathbb{P}(E)}(1)$ induced by h . We first expand $i\partial\bar{\partial}\psi$ on $\mathbb{P}(E)$ as follows:

$$i\partial\bar{\partial}\psi = i(g_{i\bar{j}}dx^i \wedge d\bar{x}^j + g_{i\bar{\beta}}dx^i \wedge d\bar{e}^\beta + g_{\alpha\bar{j}}de^\alpha \wedge d\bar{x}^j + g_{\alpha\bar{\beta}}de^\alpha \wedge d\bar{e}^\beta).$$

Here (w^0, \dots, w^{r-1}) lifts as (e^1, \dots, e^{r-1}) , and we adopt the summation convention of Einstein. Note if we furthermore ask the holomorphic frame W to be normal with respect h at x , $ig_{\alpha\bar{\beta}}de^\alpha \wedge d\bar{e}^\beta$ is just the Fubini–Study metric on $\mathbb{P}(E)_x$. Thus the matrix $[g_{\alpha\bar{\beta}}]$ is invertible everywhere. Denote by $[g^{\beta\alpha}]$ the inverse matrix. Then we can define the conformal basis by

$$\{\delta e^\alpha = de^\alpha + g^{\beta\alpha}g_{i\bar{\beta}}dx^i, dx^i\}.$$

It is shown in [15] that on this basis we can rewrite $i\partial\bar{\partial}\psi$ as

$$(4) \quad i\partial\bar{\partial}\psi = -\Psi + \omega_{FS},$$

where

$$\Psi = i\Theta_{\alpha\bar{\beta}i\bar{j}} \frac{e^\alpha \bar{e}^\beta}{h_{\sigma\bar{\tau}} e^\sigma \bar{e}^\tau} dx^i \wedge d\bar{x}^j \quad \text{and} \quad \omega_{FS} = ig_{\alpha\bar{\beta}} \delta e^\alpha \wedge \delta \bar{e}^\beta.$$

Restricted on each $\mathbb{P}(E)_x$, ω_{FS} is just the Fubini–Study metric up to a coordinate transform.

We should also make a brief explanation of Ψ . $\Theta_{\alpha\bar{\beta}i\bar{j}}$ is the curvature tensor of E associated with h . Moreover, remember that (w^0, \dots, w^{r-1}) lifts to (e^1, \dots, e^{r-1}) . Locally, say on $\{w^0 \neq 0\}$, we have $e^\alpha = w^\alpha/w^0$ for $\alpha = 1, \dots, r-1$. In view of this relationship, Ψ is interpreted as

$$\Psi = i\Theta_{\alpha\bar{\beta}i\bar{j}} \frac{w^\alpha \bar{w}^\beta}{h_{\sigma\bar{\tau}} w^\sigma \bar{w}^\tau} dx^i \wedge d\bar{x}^j.$$

Since both $\Theta_{\alpha\bar{\beta}i\bar{j}}$ and $h_{\sigma\bar{\tau}}$ are independent of w^α and $w^\alpha \bar{w}^\beta / (h_{\sigma\bar{\tau}} w^\sigma \bar{w}^\tau)$ is homogeneous of degree 0, this expression is indeed a function of (e^1, \dots, e^{r-1}) . Therefore Ψ is well-defined.

In practice, we will apply (4) to the tautological bundle $(\mathcal{O}_E(1), \varphi)$ on $\mathbb{P}(E^*)$. It enables us to establish a crucial isometry:

$$K_{\mathbb{P}(E^*)/X}^{-1} \simeq \mathcal{O}_E(r) \otimes \pi^* \det E^*.$$

3. Strong openness

We briefly recall the L^2 -representation in [20; 17] for a (singular) Hermitian metric of a vector bundle, which helps us to reduce everything to the line bundle case; then we apply Corollary 1.1 to obtain our desired estimate.

We will work with the following setup: let X be a complex manifold, and let E be a holomorphic vector bundle of rank r over X . Let $\pi : \mathbb{P}(E^*) \rightarrow X$ be the natural projection. For a Hermitian metric h on E , we denote by φ the weight function of the induced metric h_L on $\mathcal{O}_E(1)$. Suppose that h is smooth, then $i\partial\bar{\partial}\varphi$ is a Fubini–Study-type metric along each fiber $\mathbb{P}(E^*)_x$. We briefly denote it by $\omega_x := (i\partial\bar{\partial}\varphi)|_{\mathbb{P}(E^*)_x}$, then the volume $\text{Vol}(\mathbb{P}(E^*)_x)$ against ω_x equals 1 for every x . This is standard in Finsler geometry, and one could refer to [7, Lemma 2.1] for a beautiful explanation.

3.1. The L^2 -representation. We have a canonical isomorphism $\pi_*\mathcal{O}_E(1) \simeq E$ [14, Chapter II, Proposition 7.11]). Hence, for any local holomorphic section F of E , $|F|^2e^{-\varphi}$ is understood in an obvious way. Combining with [20; 17], we obtain:

Proposition 3.1. *Up to a multiplication of a constant, for every $x \in X$ and $F \in E_x$ we have*

$$(5) \quad |F|_{h,x}^2 = \int_{\mathbb{P}(E^*)_x} |F|^2 e^{-\varphi} \omega_x^{r-1}.$$

Furthermore, φ induces an isometry between the canonical isomorphism:

$$K_{\mathbb{P}(E^*)/X}^{-1} \simeq \mathcal{O}_E(r) \otimes \pi^* \det E^*.$$

Namely, if (e^1, \dots, e^{r-1}) is a fiber coordinate system of $\mathbb{P}(E^*)$, then

$$(6) \quad \omega_x^{r-1} = i^{r-1} \pi^* \det h^* \cdot e^{-r\varphi} de^1 \wedge d\bar{e}^1 \wedge \dots \wedge de^{r-1} \wedge d\bar{e}^{r-1}$$

up to a constant.

Moreover, when (E, h) is a Griffiths-semipositive singular Hermitian vector bundle, the integral of the right hand side of (5) is still well-defined, and (5) holds almost everywhere.

Proof. The smooth version of the L^2 -representation (5) is nothing but a reformulation of [17], Theorem 7.1. In fact, if we take $k = 1$ and $F = \mathcal{O}_X$, (3.3) in [17] defines a metric f on E via $\pi_*\mathcal{O}_E(1) \simeq E$, which is exactly

$$\frac{1}{(r-1)!} \int_{\mathbb{P}(E^*)_x} |F|^2 e^{-\varphi} \omega_x^{r-1}.$$

Then (7.3) there implies that $f = h$ up to multiplication by $\frac{(r+1)^{r-1}}{r!}$, which is the desired conclusion.

As for (6), it is enough to show that the associated curvature forms of both sides are equal. For this purpose, we apply (4). Keep the notation there. Without loss of generality, we can make $\{W^0, \dots, W^{r-1}\}$ normal for h^* at a fixed point x . Then at $(x; e^1, \dots, e^{r-1})$, ω_x is just the standard Fubini–Study metric on \mathbb{P}^{r-1} and

$$\Psi = i\Theta_{\alpha\bar{\beta}i\bar{j}} \frac{w^\alpha \bar{w}^\beta}{|w|^2} dx^i \wedge d\bar{x}^j.$$

Taking the integral against ω_x^{r-1} , we obtain

$$\begin{aligned} \pi_* \Psi &= i\Theta_{\alpha\bar{\beta}i\bar{j}} \left(\int_{\mathbb{P}(E^*)_x} \frac{w^\alpha \bar{w}^\beta}{|w|^2} \omega_x^{r-1} \right) dx^i \wedge d\bar{x}^j = i\Theta_{\alpha\bar{\beta}i\bar{j}} \frac{\delta_{\alpha\bar{\beta}}}{r} dx^i \wedge d\bar{x}^j \\ &= \frac{i}{r} \sum_{\alpha} \Theta_{\alpha\bar{\alpha}i\bar{j}} dx^i \wedge d\bar{x}^j = \frac{i}{r} \Theta_{\det E^*, \det h^*} \end{aligned}$$

at x , hence everywhere. On the other hand, remember that $\int_{\mathbb{P}(E^*)_x} \omega_x^{r-1} = 1$,

$$\int_{\mathbb{P}(E^*)/X} i\pi^* \Theta_{\det E^*, \det h^*} \wedge \omega_x^{r-1} = i\Theta_{\det E^*, \det h^*}.$$

Here $\int_{\mathbb{P}(E^*)/X} \cdot \wedge \omega_x^{r-1}$ refers to taking the integral along fibers. It exactly implies that there exists an element γ with

$$(7) \quad \Psi = \frac{i}{r} \pi^* \Theta_{\det E^*, \det h^*} + \gamma \quad \text{and} \quad \int_{\mathbb{P}(E^*)/X} \gamma \wedge \omega_x^{r-1} = 0.$$

Combining with the fact that ω_{FS} is the Fubini–Study metric along each fiber of π and the canonical isomorphism

$$K_{\mathbb{P}(E^*)/X}^{-1} \simeq \mathcal{O}_E(r) \otimes \pi^* \det E^*,$$

we conclude from (7) that

$$i\Theta_{\mathcal{O}_E(r), r\varphi} = ir\partial\bar{\partial}\varphi = -r\Psi + r\omega_{FS} = -i\pi^* \Theta_{\det E^*, \det h^*} + i\Theta_{K_{\mathbb{P}(E^*)/X}^{-1}, \omega_x^{r-1}}.$$

The proof of (6) is then finished. Observe that by setting

$$(G, h_G) := (\mathcal{O}_E(r+1) \otimes \pi^* \det E^*, h_L^{r+1} \otimes \pi^* \det h^*),$$

we can interpret (5) as

$$(8) \quad |F|_{h,x}^2 = \int_{\mathbb{P}(E^*)_x} |F|_{h_G}^2$$

in view of (6). The definition of $\int_{\mathbb{P}(E^*)_x} |F|_{h_G}^2$ is based on

$$\mathcal{O}_E(1) \simeq K_{\mathbb{P}(E^*)/X} \otimes \mathcal{O}_E(r+1) \otimes \pi^* \det E^*,$$

and is carefully explained in [20], Sect. 2.2.

Whereas the singular version of (5) is nothing but a reformulation of [20], Proposition 3.1. Indeed, by [20] the integral of the right hand side of (8) is still well-defined when h is singular and (E, h) is Griffiths-semipositive. Moreover, (8) holds outside $V := \{\det h = \infty\}$. Note V must be a set of measure zero since h is finite almost everywhere, which concludes the last assertion. \square

We next restate and prove **Theorem 1.2**:

Theorem 3.1. *Let E be a holomorphic vector bundle of rank r over Δ^n , and let h be a singular Hermitian metric on E such that (E, h) is Griffiths-semipositive. Assume that $\det h$ has analytic singularities. If F is a holomorphic section of E satisfying*

$$\int_{\Delta^n} |F|_h^2 d\lambda < \infty,$$

then after shrinking Δ^n if necessary, there exists a positive ε (independent of F) such that

$$\int_{\Delta^n} |F|_h^{2(1+\varepsilon)} d\lambda < \infty.$$

Proof. Keep the notations of **Proposition 3.1**. Due to (the singular version of) (8) and Fubini’s theorem, it is enough to show that there exists a positive ε such that

$$\int_{\Delta^n} \left(\int_{\mathbb{P}(E^*)_x} |F|_{h_G}^2 \right)^{1+\varepsilon} d\lambda < \infty.$$

By hypothesis, $\pi^* \det h^* = e^\psi$, where ψ is a plurisubharmonic function with analytic singularities. So we have

$$\begin{aligned} \int_{\Delta^n} \left(\int_{\mathbb{P}(E^*)_x} |F|_{h_G}^2 \right)^{1+\varepsilon} d\lambda &= \int_{\Delta^n} (\det h^*)^{1+\varepsilon} \left(\int_{\mathbb{P}(E^*)_x} |F|^2 e^{-(r+1)\varphi} \right)^{1+\varepsilon} d\lambda \\ &\leq C \int_{\Delta^n} (\det h^*)^{1+\varepsilon} \left(\int_{\mathbb{P}(E^*)_x} (|F|^2 e^{-(r+1)\varphi})^{1+\varepsilon} \right) d\lambda \\ &= C \int_{\Delta^n} \int_{\mathbb{P}(E^*)_x} e^{(1+\varepsilon)(\log |F|^2 + \psi) - (1+\varepsilon)(r+1)\varphi} d\lambda. \end{aligned}$$

The inequality is a simple application of Hölder’s inequality, and C is a universal positive constant.

On the other hand, by assumption F a priori satisfies

$$\int_{\Delta^n} \int_{\mathbb{P}(E^*)_x} e^{\log |F|^2 + \psi - (r+1)\varphi} d\lambda < \infty.$$

So after shrinking Δ^n if necessary, by **Corollary 1.1** there exists a positive ε_0 such that

$$(9) \quad \int_{\Delta^n} \int_{\mathbb{P}(E^*)_x} e^{\log |F|^2 + \psi - (1+\varepsilon_0)(r+1)\varphi} d\lambda < \infty.$$

Since $e^{\varepsilon_0(\log |F|^2 + \psi)}$ is bounded, (9) implies that

$$\int_{\Delta^n} \left(\int_{\mathbb{P}(E^*)_x} |F|_{h_G}^2 \right)^{1+\varepsilon_0} d\lambda < \infty.$$

In the end, notice that [23], Theorem 1.2 indicates that $\mathcal{E}(h)$ is coherent hence is locally finitely generated in our situation. So we can even obtain a uniform constant ε_0 for all holomorphic sections of $\mathcal{E}(h)$. The proof is complete. \square

4. Openness

Now we would like to remove the restriction in Theorem 3.1 that $\det h$ has analytic singularities. A natural idea is to approximate h by a sequence of smooth metrics $\{h_\nu\}$, whose existence is due to Proposition 2.1. Then we apply Theorem 3.1 on each h_ν (which gives a corresponding ε_ν), and take the limit to obtain the desired estimate. For this purpose, we need a universal lower bound for these ε_ν . The related topic is called the effectiveness of the *strong openness property*. It is much involved even in the line bundle case (see [10], Theorem 1.3). As a compromise, we will instead apply the following effective version of the *openness property* in [4].

The line bundle case revisited.

Theorem 4.1 [4, Theorem 1.1]. *Let φ be a plurisubharmonic function in B^n with $\varphi \leq 0$. Assume that*

$$\int_{B^n} e^{-\varphi} d\lambda < \infty.$$

Then there is a number $\varepsilon > 0$ such that

$$\int_{B^{n/2}} e^{-(1+\varepsilon)\varphi} d\lambda < \infty.$$

Here B^n and $B^{n/2}$ refer to the n -dimensional balls of radii 1 and 1/2 respectively. Moreover, ε can be taken so that

$$\varepsilon \geq \frac{\delta_n}{\int_{B^n} e^{-\varphi} d\lambda},$$

where δ_n depends only on the dimension.

Let us briefly recall some crucial steps in the proof of Theorem 4.1 for readers' benefit. For any $s \geq 0$ and holomorphic function h , let

$$\varphi_s := \max(\varphi + s, 0)$$

and

$$\|h\|_s^2 := \int_{B^{n/2}} |h|^2 e^{-2\varphi_s} d\lambda.$$

The factor 2 in the exponent is necessary, and we define the norm $\|\cdot\|_s$ here over $B^n/2$ (rather than B^n in [4]) for our purposes. Then a simple variant of Proposition 2.1 in [4] indicates that for $0 < \varepsilon < 1$,

$$(10) \quad \int_{B^n/2} e^{-(1+\varepsilon)\varphi} d\lambda = a_\varepsilon \int_0^\infty e^{(1+\varepsilon)s} \|1\|_s^2 ds + b_\varepsilon \text{Vol}(B^n).$$

Here

$$a_\varepsilon = \frac{1 - \varepsilon^2}{2} \quad \text{and} \quad b_\varepsilon = \frac{1 - \varepsilon}{2^{n+1}}.$$

Pick a universal constant δ_n so that if w is holomorphic and

$$\int_{B^n} |w|^2 d\lambda \leq 4\delta_n,$$

then

$$\sup_{B^n/2} |w| \leq \frac{1}{10}.$$

The existence of such a δ_n is due to the mean value property of holomorphic functions. In particular, δ_n depends only on the dimension.

Now $\int_{B^n} e^{-\varphi} d\lambda < \infty$ by hypothesis,

$$\int_0^\infty e^s \left(\int_{B^n} e^{-2\varphi_s} \right) ds \leq \int_{B^n} e^{-\varphi} d\lambda < \infty$$

due to (the original version of) Proposition 2.1 in [4]. Then by Theorem 3.3 of [4], for any $\varepsilon > 0$ and $s > \frac{1}{2\varepsilon}$ there is a holomorphic function h_s such that

$$(11) \quad \int_{B^n} |1 - h_s|^2 d\lambda \leq 4\varepsilon \int_{B^n} e^{-\varphi} d\lambda,$$

and

$$(12) \quad \int_{B^n} |h_s|^2 e^{-2\varphi_s} d\lambda \leq e^{-(1+2\varepsilon)s} \text{Vol}(B^n).$$

Take

$$\varepsilon = \frac{\delta_n}{\int_{B^n} e^{-\varphi} d\lambda}.$$

Then (11) implies that the holomorphic function $1 - h_s$ satisfies

$$\int_{B^n} |1 - h_s|^2 d\lambda \leq 4\delta_n,$$

hence $\sup_{B^n/2} |1 - h_s| \leq \frac{1}{10}$. Therefore h_s satisfies

$$1 < 2|h_s|^2$$

on $B^n/2$. Multiply by $e^{-2\varphi_s}$ and integrate on $B^n/2$, we obtain from this inequality

and (12) that

$$(13) \quad \|1\|_s^2 \leq 2 \int_{B^n/2} |h_s|^2 e^{-2\varphi_s} d\lambda \leq 2 \int_{B^n} |h_s|^2 e^{-2\varphi_s} d\lambda \leq 2e^{-(1+2\varepsilon)s} \text{Vol}(B^n).$$

Now multiply the both sides of (13) by $e^{(1+\varepsilon)s}$ and integrate from 0 to infinity. Combining with (10) we get

$$\begin{aligned} (14) \quad & \int_{B^n/2} e^{-(1+\varepsilon)\varphi} d\lambda \\ & \leq a_\varepsilon \int_0^{\frac{1}{2\varepsilon}} e^{(1+\varepsilon)s} \|1\|_s^2 ds + a_\varepsilon \int_{\frac{1}{2\varepsilon}}^\infty e^{(1+\varepsilon)s} 2e^{-(1+2\varepsilon)s} \text{Vol}(B^n) ds + b_\varepsilon \text{Vol}(B^n) \\ & \leq a_\varepsilon \text{Vol}(B^n/2) \int_0^{\frac{1}{2\varepsilon}} e^{(1+\varepsilon)s} ds + a_\varepsilon \text{Vol}(B^n) \int_{\frac{1}{2\varepsilon}}^\infty e^{-\varepsilon s} ds + b_\varepsilon \text{Vol}(B^n) \\ & = \frac{a_\varepsilon}{2^n} \text{Vol}(B^n) \frac{e^{\frac{1+\varepsilon}{2\varepsilon}} - 1}{1 + \varepsilon} + a_\varepsilon \text{Vol}(B^n) \frac{1}{\varepsilon \sqrt{e}} + b_\varepsilon \text{Vol}(B^n) \\ & = \left(\frac{a_\varepsilon (e^{\frac{1+\varepsilon}{2\varepsilon}} - 1)}{2^n (1 + \varepsilon)} + \frac{a_\varepsilon}{\varepsilon \sqrt{e}} + b_\varepsilon \right) \text{Vol}(B^n) \\ & =: c_\varepsilon \text{Vol}(B^n). \end{aligned}$$

Since (13) is valid when $s > \frac{1}{2\varepsilon}$, we have to divide the integral into two parts. The integral from 0 to $\frac{1}{2\varepsilon}$ is standard, since $0 \leq \varphi_s \leq \frac{1}{2\varepsilon}$; the integral from $\frac{1}{2\varepsilon}$ to infinity is estimated via (13), and it leads to the second inequality. The rest is routine.

Proof of Theorem 1.3. Now we are ready to prove the openness for the higher-rank case. Let E be a holomorphic vector bundle of rank r over Δ_R^n , where Δ_R^n is a polydisc with radius R slightly larger than 1. Let (e^1, \dots, e^{r-1}) be a fiber coordinate system of $\mathbb{P}(E^*)$, and let $de \wedge d\bar{e}$ be short for

$$de^1 \wedge d\bar{e}^1 \wedge \dots \wedge de^{r-1} \wedge d\bar{e}^{r-1}.$$

Take a finite coordinate chart $\{B_j^{n+r-1}\}$ of $\mathbb{P}(E^*)$. For an arbitrary singular Hermitian metric h on E , denote by φ the induced metric on $\mathcal{O}_E(1)$, and

$$\int_{B^{n+r-1}} e^{-(r+1)\varphi} d\lambda := \max_j \left\{ \int_{B_j^{n+r-1}} e^{-(r+1)\varphi} d\lambda \right\}$$

with the convention that

$$\int_{B^{n+r-1}} e^{-(r+1)\varphi} d\lambda = +\infty \quad \text{if some} \quad \int_{B_j^{n+r-1}} e^{-(r+1)\varphi} d\lambda = +\infty.$$

We restate and prove Theorem 1.3.

Theorem 4.2. *Assume that (E, h) is Griffiths-semipositive, and*

$$\mathcal{I}((r + 1)\varphi) = \mathcal{O}_{\mathbb{P}(E^*)}.$$

Then there exist positive numbers \hat{R} and ε such that

$$\int_{\Delta_{\hat{R}}^n} |F|_h^{2(1+\varepsilon)} d\lambda < \infty$$

for any holomorphic section F of E . Moreover, ε can be taken so that

$$\varepsilon \geq \frac{\delta_{n+r-1}}{\int_{B^{n+r-1}} e^{-(r+1)\varphi} d\lambda},$$

where δ_{n+r-1} depends only on the dimension.

Proof. Pick an $R' \in (1, R)$. By Proposition 2.1(1), there exists a sequence of smooth, Griffiths positive metrics $\{h_\nu\}$ over $\Delta_{R'}^n$, increasing pointwise to h . Let φ_ν be the metric on $\mathcal{O}_E(1)$ induced by h_ν , and let $\omega_{x,\nu} := (i\partial\bar{\partial}\varphi_\nu)|_{\mathbb{P}(E^*)_x}$. Consequently, φ_ν decreases to φ . Since both φ and φ_ν are plurisubharmonic, we can assume $\varphi, \varphi_\nu \leq 0$ without loss of generality.

Now by assumption we have

$$\int_{B^{n+r-1}} e^{-(r+1)\varphi_\nu} d\lambda \leq \int_{B^{n+r-1}} e^{-(r+1)\varphi} d\lambda < \infty.$$

Then apply Theorem 4.1 ((14), more precisely), we obtain

$$(15) \quad \int_{B^{n+r-1/2}} e^{-(1+\varepsilon)(r+1)\varphi_\nu} d\lambda \leq c_\varepsilon \text{Vol}(B^{n+r-1}),$$

where ε can be taken so that

$$\varepsilon \geq \frac{\delta_{n+r-1}}{\int_{B^{n+r-1}} e^{-(r+1)\varphi_\nu} d\lambda}.$$

Since φ_ν decreases to φ ,

$$\frac{\delta_{n+r-1}}{\int_{B^{n+r-1}} e^{-(r+1)\varphi_\nu} d\lambda} \searrow \frac{\delta_{n+r-1}}{\int_{B^{n+r-1}} e^{-(r+1)\varphi} d\lambda}.$$

Let

$$\varepsilon = \frac{\delta_{n+r-1}}{\int_{B^{n+r-1}} e^{-(r+1)\varphi} d\lambda},$$

and take the limit of (15) with respect to ν . We finally obtain

$$\int_{B^{n+r-1/2}} e^{-(1+\varepsilon)(r+1)\varphi} d\lambda \leq c_\varepsilon \text{Vol}(B^{n+r-1}) < \infty.$$

In the end, observe that we can take a positive number $\hat{R} < R'$ such that the total

space of $\mathbb{P}(E^*)|_{\Delta_{\hat{R}}^n}$ is finitely covered by $B_j^{n+r-1}/2$. It exactly implies that

$$\begin{aligned} \int_{\Delta_{\hat{R}}^n} |F|_{h_\nu}^{2(1+\varepsilon)} d\lambda &= \int_{\Delta_{\hat{R}}^n} \left(\int_{\mathbb{P}(E^*)_x} |F|^2 e^{-\varphi_\nu} \omega_{x,\nu}^{r-1} \right)^{1+\varepsilon} d\lambda \\ &\leq \int_{\Delta_{\hat{R}}^n} \left(\int_{\mathbb{P}(E^*)_x} (|F|^2 e^{-\varphi_\nu})^{1+\varepsilon} \omega_{x,\nu}^{r-1} \right) \left(\int_{\mathbb{P}(E^*)_x} \omega_{x,\nu}^{r-1} \right)^{\frac{\varepsilon}{1+\varepsilon}} d\lambda \\ &= \int_{\Delta_{\hat{R}}^n} \left(\int_{\mathbb{P}(E^*)_x} (|F|^2 e^{-(\frac{r}{1+\varepsilon}+1)\varphi_\nu})^{1+\varepsilon} \cdot \pi^* \det h_\nu^* \cdot i^{r-1} de \wedge d\bar{e} \right) d\lambda \\ &\leq \sum_j \int_{B_j^{n+r-1}/2} |F|^{2(1+\varepsilon)} e^{-(r+1+\varepsilon)\varphi_\nu} \cdot \pi^* \det h_\nu^* d\lambda \\ &\leq C \sum_j \int_{B_j^{n+r-1}/2} e^{-(r+1+\varepsilon)\varphi_\nu} d\lambda \\ &\leq C \sum_j \int_{B_j^{n+r-1}/2} e^{-(r+1+\varepsilon)\varphi} d\lambda < \infty \end{aligned}$$

for any F . In the forth inequality we abuse the notation that $d\lambda$ refers to the Lebesgue measure on both $\Delta_{\hat{R}}^n$ and $B_j^{n+r-1}/2$. The fifth inequality is due to the fact that $\{\det h_\nu^*\}$ is a decreasing sequence of smooth functions. So $|F|^{2(1+\varepsilon)} \pi^* \det h_\nu^*$ is bounded by some positive constant C only depends on F . As ν tends to zero, we obtain

$$\int_{\Delta_{\hat{R}}^n} |F|_h^{2(1+\varepsilon)} d\lambda < \infty$$

for any F . The proof is complete. □

Let $\{U_j\}$ be a finite coordinate covering of $\mathbb{P}(E^*)$, and let $\{\rho_j\}$ be the associated partition of unity. When $\mathcal{S}((r+1)\varphi) = \mathcal{O}_{\mathbb{P}(E^*)}$, for any holomorphic section F we have

$$\begin{aligned} \int_{\Delta^n} |F|_{h_\nu}^2 d\lambda &= \int_{\Delta^n} \left(\int_{\mathbb{P}(E^*)_x} |F|^2 e^{-\varphi_\nu} \omega_{x,\nu}^{r-1} \right) d\lambda \\ &= \int_{\Delta^n} \left(\int_{\mathbb{P}(E^*)_x} |F|^2 e^{-(r+1)\varphi_\nu} \cdot \pi^* \det h_\nu^* \cdot i^{r-1} de \wedge d\bar{e} \right) d\lambda \\ &= \sum_j \int_{U_j} \rho_j |F|^2 e^{-(r+1)\varphi_\nu} \cdot \pi^* \det h_\nu^* d\lambda \\ &\leq C \sum_j \int_{U_j} e^{-(r+1)\varphi_\nu} d\lambda \\ &\leq C \sum_j \int_{U_j} e^{-(r+1)\varphi} d\lambda < \infty. \end{aligned}$$

As ν tends to zero, we obtain

$$\int_{\Delta^n} |F|_h^2 d\lambda < \infty$$

for any F . Hence

$$(16) \quad \mathcal{E}(h) = E.$$

However, if one attempts to deduce $\mathcal{S}((r+1)\varphi) = \mathcal{O}_{\mathbb{P}(E^*)}$ from (16), the influence of the zero locus of $\det h^*$ cannot be ignored. Hence the converse is not clear.

Acknowledgement

The author thanks Prof. Jixiang Fu for discussion and continuing encouragement. Thanks are also due to the referees for pointing out [20], which helped improve this paper.

References

- [1] S. J. Bao and Q. A. Guan, “ L^2 extension and effectiveness of strong openness property”, *Acta Math. Sin. (Engl. Ser.)* **38**:11 (2022), 1949–1964. [MR](#)
- [2] S. J. Bao and Q. A. Guan, “ L^2 extension and effectiveness of L^p strong openness property”, *Acta Math. Sin. (Engl. Ser.)* **39**:5 (2023), 814–826. [MR](#)
- [3] R. J. Berman, S. Boucksom, and M. Jonsson, “A variational approach to the Yau-Tian-Donaldson conjecture”, *J. Amer. Math. Soc.* **34**:3 (2021), 605–652. [MR](#)
- [4] B. Berndtsson, “The openness conjecture for plurisubharmonic functions”, preprint, 2013. [arXiv 1305.5781](#)
- [5] B. Berndtsson and M. Păun, “Bergman kernels and the pseudoeffectivity of relative canonical bundles”, *Duke Math. J.* **145**:2 (2008), 341–378. [MR](#)
- [6] J.-P. Demailly, *Analytic methods in algebraic geometry*, *Surveys of Modern Mathematics* **1**, International Press, Somerville, MA, 2012. [MR](#)
- [7] H. Feng, K. Liu, and X. Wan, “Chern forms of holomorphic Finsler vector bundles and some applications”, *Internat. J. Math.* **27**:4 (2016), art. id. 1650030, 22 pp. [MR](#)
- [8] P. A. Griffiths, “Hermitian differential geometry, Chern classes, and positive vector bundles”, pp. 185–251 in *Global analysis: (papers in honor of K. Kodaira)*, Princeton Univ. Press, 1969. [MR](#)
- [9] Q. Guan, “A sharp effectiveness result of Demailly’s strong openness conjecture”, *Adv. Math.* **348** (2019), 51–80. [MR](#)
- [10] Q. Guan and X. Zhou, “Effectiveness of Demailly’s strong openness conjecture and related problems”, *Invent. Math.* **202**:2 (2015), 635–676. [MR](#)
- [11] Q. Guan and X. Zhou, “A proof of Demailly’s strong openness conjecture”, *Ann. of Math. (2)* **182**:2 (2015), 605–616. [MR](#)
- [12] Q. Guan and X. Zhou, “Strong openness of multiplier ideal sheaves and optimal L^2 extension”, *Sci. China Math.* **60**:6 (2017), 967–976. [MR](#)
- [13] Q. Guan, Z. Li, and X. Zhou, “Stability of multiplier ideal sheaves”, *Chinese Ann. Math. Ser. B* **43**:5 (2022), 819–832. [MR](#)
- [14] R. Hartshorne, *Algebraic geometry*, *Graduate Texts in Mathematics* **52**, Springer, 1977. [MR](#)
- [15] S. Kobayashi, “Negative vector bundles and complex Finsler structures”, *Nagoya Math. J.* **57** (1975), 153–166. [MR](#)

- [16] L. Lempert, “Modules of square integrable holomorphic germs”, pp. 311–333 in *Analysis meets geometry*, Birkhäuser, 2017. [MR](#)
- [17] K. Liu, X. Sun, and X. Yang, “Positivity and vanishing theorems for ample vector bundles”, *J. Algebraic Geom.* **22**:2 (2013), 303–331. [MR](#)
- [18] Z. Liu, B. Xiao, H. Yang, and X. Zhou, “Multiplier submodule sheaves and a problem of Lempert”, preprint, 2021. [arXiv 2111.13452](#)
- [19] Z. Liu, H. Yang, and X. Zhou, “New properties of multiplier submodule sheaves”, *C. R. Math. Acad. Sci. Paris* **360** (2022), 1205–1212. [MR](#)
- [20] Y. Liu, Z. Liu, H. Yang, and X. Zhou, “A Le Potier-type isomorphism twisted with multiplier submodule sheaves”, *J. Math. Pures Appl.* (9) **202** (2025), art. id. 103760, 38 pp. [MR](#)
- [21] M. Păun and S. Takayama, “Positivity of twisted relative pluricanonical bundles and their direct images”, *J. Algebraic Geom.* **27**:2 (2018), 211–272. [MR](#)
- [22] H. Raufi, “Singular hermitian metrics on holomorphic vector bundles”, *Ark. Mat.* **53**:2 (2015), 359–382. [MR](#)
- [23] Y. Zou, “Note on the singular Hermitian metrics with analytic singularities”, preprint, 2022. [arXiv 2209.05053](#)

Received March 19, 2025. Revised June 26, 2025.

JINGCAO WU
SCHOOL OF MATHEMATICS
SHANGHAI UNIVERSITY OF FINANCE AND ECONOMICS
SHANGHAI
CHINA
wujincao@shufe.edu.cn

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Fakultät für Mathematik
Universität Wien
Vienna, Austria
matthias.aschenbrenner@univie.ac.at

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Atsushi Ichino
Department of Mathematics
Kyoto University
Kyoto 606-8502, Japan
atsushi.ichino@gmail.com

Kefeng Liu
School of Sciences
Chongqing University of Technology
Chongqing 400054, China
liu@math.ucla.edu

Sucharit Sarkar
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
sucharit@math.ucla.edu

Dimitri Shlyakhtenko
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
shlyakht@ipam.ucla.edu

Ruixiang Zhang
Department of Mathematics
University of California
Berkeley, CA 94720-3840
ruixiang@berkeley.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2026 is US \$710/year for the electronic version, and \$965/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and [Web of Knowledge \(Science Citation Index\)](#).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2026 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 342 No. 2 June 2026

On multilinear maximal operators along homogeneous curves	207
LARS BECKER and BEN KRAUSE	
The plain sphere number of a link	217
RYAN BLAIR, ALEXANDRA KJUCHUKOVA and ELLA PFAFF	
On the test properties of the Frobenius endomorphism	235
OLGUR CELIKBAS, ARASH SADEGHI and YONGWEI YAO	
Symplectic automorphisms of a surface with genus two fibration and their action on CH_0	259
JIABIN DU and WENFEI LIU	
Non-braid-positive hyperbolic L -space knots	275
KEISUKE HIMENO	
The Rankin–Selberg integral on GSp_2 for square-free levels	299
SEIJI KUGA and MASAO TSUZUKI	
Sum of the squares of the p' -character degrees	351
NGUYEN N. HUNG, J. MIQUEL MARTÍNEZ and GABRIEL NAVARRO	
On the first eigenvalue of the Hodge Laplacian of submanifolds	381
CHRISTOS-RAENT ONTI	
Howe duality for the dual pair $\text{SL}_2(\mathbb{R}) \times F_{4,1}$: a ping pong of K -types	387
GORDAN SAVIN	
Liouville theorems and new gradient estimates for positive solutions to $\Delta_p u + au^q = 0$ on a complete manifold	395
YOUDE WANG and LIQIN ZHANG	
p -nuclearity of reduced group L^p -operator algebras	427
ZHEN WANG	
A higher-rank analog of the strong openness property	441
JINGCAO WU	