

*Pacific  
Journal of  
Mathematics*

Volume 343 No. 1

July 2026

# PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

[msp.org/pjm](http://msp.org/pjm)

## EDITORS

Don Blasius (Managing Editor)  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[blasius@math.ucla.edu](mailto:blasius@math.ucla.edu)

Matthias Aschenbrenner  
Fakultät für Mathematik  
Universität Wien  
Vienna, Austria  
[matthias.aschenbrenner@univie.ac.at](mailto:matthias.aschenbrenner@univie.ac.at)

Vyjayanthi Chari  
Department of Mathematics  
University of California  
Riverside, CA 92521-0135  
[chari@math.ucr.edu](mailto:chari@math.ucr.edu)

Atsushi Ichino  
Department of Mathematics  
Kyoto University  
Kyoto 606-8502, Japan  
[atsushi.ichino@gmail.com](mailto:atsushi.ichino@gmail.com)

Kefeng Liu  
School of Sciences  
Chongqing University of Technology  
Chongqing 400054, China  
[liu@math.ucla.edu](mailto:liu@math.ucla.edu)

Sucharit Sarkar  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[sucharit@math.ucla.edu](mailto:sucharit@math.ucla.edu)

Dimitri Shlyakhtenko  
Department of Mathematics  
University of California  
Los Angeles, CA 90095-1555  
[shlyakht@ipam.ucla.edu](mailto:shlyakht@ipam.ucla.edu)

Ruixiang Zhang  
Department of Mathematics  
University of California  
Berkeley, CA 94720-3840  
[ruixiang@berkeley.edu](mailto:ruixiang@berkeley.edu)

## PRODUCTION

Silvio Levy, Scientific Editor, [production@msp.org](mailto:production@msp.org)

---

See inside back cover or [msp.org/pjm](http://msp.org/pjm) for submission instructions.

---

The subscription price for 2026 is US \$710/year for the electronic version, and \$965/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).


---

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004.

---

PJM peer review and production are managed by EditFLOW<sup>®</sup> from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2026 Mathematical Sciences Publishers

## THE GENERIC EXTENSION MAP AND MODULAR STANDARD MODULES

JOHANNES DROSCHL

**We study two classes of  $\ell$ -modular standard modules of the general linear group. The first class is obtained by reducing existing standard modules over  $\overline{\mathbb{Q}}_\ell$  to  $\overline{\mathbb{F}}_\ell$  with respect to their natural integral structure. The second class is obtained by studying the generic extension map of the cyclical quiver, which was motivated by the construction of certain monomial bases of quantum algebras. In the latter case we also manage to prove a modular version of the Langlands classification, similar to the work of Langlands and Zelevinsky over  $\mathbb{C}$ . We also compute the corresponding  $\ell$ -modular Rankin–Selberg  $L$ -functions and check that they agree with the  $L$ -functions of their  $\mathbb{C}$ -parameters constructed by Kurinczuk and Matringe.**

### 1. Introduction

Let  $F$  be a nonarchimedean local field with ring of integers  $\mathfrak{O}_F$  and residue field  $k$ , whose cardinality we denote by  $q$ , and  $G$  be a reductive group over  $F$ . We also fix a prime  $\ell$  not dividing  $q$  together with a field of coefficients  $R \in \{\overline{\mathbb{F}}_\ell, \overline{\mathbb{Q}}_\ell, \mathbb{C}\}$ . The classification of irreducible smooth representations of  $G(F)$  over  $R$  is of great importance in establishing the local Langlands correspondence. If  $R = \mathbb{C}$ , the classification was first achieved by Langlands in what is now known as the *Langlands classification*. His classification associates to each irreducible smooth representation  $\pi$  of  $G(F)$  a parabolic subgroup  $P \subseteq G$ , a tempered representation  $\sigma$  of the  $F$ -points of the Levi component  $M$  of  $P$  and a character  $\eta$  of the center of  $M$  satisfying certain inequalities. Then  $\pi$  can be realized as the unique quotient of the parabolically induced representation  $\mathcal{S}(\pi) = \text{ind}_P^{G(F)}(\sigma \otimes \eta)$ , known as the standard module associated to  $\pi$ , and the triple  $(P, \sigma, \eta)$  is unique up to certain natural symmetries.

The analytical nature of the Langlands classification as sketched above makes it clear that it is a nontrivial task to extend it to fields of nonzero characteristic, as soon as one ventures beyond the banal setting of [22]. Therefore the construction of modular standard modules remained an open problem. The proposed definitions

---

*MSC2020:* 11F70, 22E50.

*Keywords:* Rankin–Selberg  $L$ -function, modular representation theory, local Langlands correspondence.

given in this paper avoid the above issue by considering standard modules which are, in the modular case, not necessarily induced representations, but rather certain subrepresentations of the space of Whittaker functions.

To motivate this approach we return for a moment to the problem of the classification of irreducible modular representations. We recall that in the case  $G = \mathrm{GL}_n$  Bernstein and Zelevinsky gave in [32; 2; 3] a second classification of irreducible complex representation in terms of combinatorial objects called *multisegments*. To this date this complete second classification is a feature only present in the representation theory of  $\mathrm{GL}_n$  and its inner forms, although certain extensions of these ideas to more general groups have been pursued [10]. Furthermore, the Langlands and Zelevinsky classifications are related by the Aubert–Zelevinsky involution, an involution on the set of isomorphism classes of the irreducible representations of  $\mathrm{GL}_n(\mathbb{F})$ ; see [32; 1]. It thus fell upon the classification of Bernstein and Zelevinsky in the case  $G = \mathrm{GL}_n$  to offer itself to generalization in the modular case.

In the works of Vignéras [30] and Mínguez and Sécherre [23], this feat has been accomplished by extending their ideas. In general, the study of irreducible representations of  $\mathrm{GL}_n$  over  $\overline{\mathbb{F}}_\ell$  shows many similarities to the study of complex representations, but with several key differences as we will see later. For example, no longer is every cuspidal representation supercuspidal.

Note that the usefulness of standard modules in the complex case goes much further than the simple classification of irreducible representations. A prominent example, and the main motivator of this paper, are the local *Rankin–Selberg  $L$ -factors*  $L(s, \rho, \rho')$  of [11], which are defined for pairs of complex representations of Whittaker type  $(\rho, \rho')$  and they are rational functions in  $q^s$ . In particular, if  $G = \mathrm{GL}_n$ , every standard module  $\mathcal{S}(\pi)$  is of Whittaker type, which in turn allows one to define the invariant  $L(s, \mathcal{S}(\pi), \mathcal{S}(\pi'))$  for any pair of irreducible representations of  $\mathrm{GL}_n$ . It comes as no surprise that the  $L$ -factor  $L(s, \mathcal{S}(\pi), \mathcal{S}(\pi'))$  can be explicitly computed in terms of the Langlands parameters of  $\pi$  and  $\pi'$ .

Passing from the complex to the modular setting, the definition of local Rankin–Selberg  $L$ -factors has been extended in [14] to the more general fields  $R$  we are going to consider in this paper. The authors again associate to a pair of representations of Whittaker type an  $L$ -factor  $L(X, \pi, \pi')$ , which is a rational function in  $X$  (with  $X = q^{-s}$  in the  $\mathbb{C}$ -case this recovers the original construction of [12]). The Rankin–Selberg  $L$ -factors so obtained satisfy a functional equation, which in turn gives rise to  $\epsilon$ - and  $\gamma$ -factors associated to the pair  $(\pi, \pi')$ . In [16] the authors associate to each irreducible representation of  $\mathrm{GL}_n(\mathbb{F})$  a so-called  $C$ -parameter  $C(\pi)$ , which is a modular version of the Langlands parameter. Moreover, they equip  $C$ -parameters with a tensor product  $\otimes_{s,s}$  and define their  $L$ -,  $\epsilon$ - and  $\gamma$ -factors.

In this paper we hope to initiate the investigation of possible candidates for standard modules in the modular setting and their properties. We do this by

giving two natural possible candidates for standard modules, one obtained by reducing the ones defined over  $\overline{\mathbb{Q}}_\ell$  to  $\overline{\mathbb{F}}_\ell$ , the other one by considering the so-called *generic extension map*, which is rooted in the analysis of certain (quantum)-algebras associated to certain Dynkin or affine Dynkin quivers. We will prove that the former always contain the latter and note that at the moment the evidence points towards them being equal, although we do not treat this question in this paper.

The properties discussed above give us natural constraints our potential standard modules should satisfy. Firstly, they should admit a modular formulation of the Langlands classification and secondly, they should allow us to define Rankin–Selberg  $L$ -functions, which should be explicitly computable using the C-parameters of the underlying irreducible representations.

The contents of this paper can thus be roughly divided into three parts: the construction of standard modules, establishing certain representation-theoretic properties and finally the computation of their Rankin–Selberg  $L$ -factors.

To state our results more precisely, we need to introduce some notation. Let  $\psi : F \rightarrow R$  be a smooth, additive character and recall the space of Whittaker functions,

$$\mathcal{W}(\psi) := \left\{ f : G_n \rightarrow R : f \left( \begin{pmatrix} 1 & u_{1,2} & \cdots & \cdots \\ 0 & \ddots & \cdots & \cdots \\ 0 & \cdots & 1 & u_{n-1,n} \\ 0 & \cdots & 0 & 1 \end{pmatrix} g \right) = \psi \left( \sum_{i=1}^{n-1} u_{i,i+1} \right) f(g), f \text{ locally constant} \right\}.$$

We also recall that a smooth representation  $\pi$  of  $G_n$  of finite length is called of Whittaker type if

$$\dim_R \operatorname{Hom}_{G_n}(\pi, \mathcal{W}(\psi)) = 1,$$

and in this case we denote its image by  $\mathcal{W}(\pi, \psi)$ . We let

$$\mathfrak{Rep}_{\mathcal{W}, \psi, n} = \{\mathcal{W}(\pi, \psi) : \pi \text{ a representation of } G_n \text{ of Whittaker type}\}.$$

Let us recall that to a cuspidal representation of  $G_m$  we can associate  $o(\rho) \in \mathbb{N} \cup \{\infty\}$ , with  $o(\rho) = \infty$  if and only if  $\operatorname{char}(R) = 0$ , and and that if  $o(\rho)$  is finite, it is the minimal  $k \in \mathbb{Z}_{>0}$  such that  $\rho \cong \rho|^{-k}$ . We let moreover be  $e(\rho) = \ell$  if  $o(\rho) = 1$  and  $e(\rho) = o(\rho)$  otherwise. To two integers  $a \leq b$  and  $\rho$  cuspidal, we associate a segment (over  $R$ )  $[a, b]_\rho = (\rho|^{-a}, \dots, \rho|^{-b})$  and we identify  $[a, b]_\rho$  with  $[a + o(\rho), b + o(\rho)]_\rho$ . We also write  $[a, b]_\rho^\vee = [-b, -a]_{\rho^\vee}$ . A finite formal sum of segments is called a multisegment and we denote the set of multisegments over  $R$  by  $\operatorname{Mult}_R$ , to which we extend the operation  $(-)^\vee$ . A multisegment is called aperiodic if it does not contain a submultisegment of the form

$$[a, b]_\rho + [a+1, b+1]_\rho + \cdots + [a+e(\rho)-1, b+e(\rho)-1]_\rho,$$

and we denote the set of aperiodic multisegments by  $\text{Mult}_R^{\text{ap}}$ . We decompose any multisegment  $\mathfrak{m}$  as  $\mathfrak{m}_b + \mathfrak{m}_{nb}$ , where  $\mathfrak{m}_{nb}$  consists of those  $[a, b]_\rho$  with  $o(\rho) = 1$ . We let  $\text{Mult}_{R, \square}^{\text{ap}}$  be the set of aperiodic multisegments such that  $\mathfrak{m} = \mathfrak{m}_b$ . Note that if  $\text{char}(R) = 0$  all multisegments are aperiodic and  $\text{Mult}_{\overline{\mathbb{Q}}_\ell} = \text{Mult}_{\overline{\mathbb{Q}}_\ell, \square}^{\text{ap}}$ . Then there exists a bijection

$$\langle - \rangle : \text{Mult}_R^{\text{ap}} \rightarrow \mathfrak{Irr}(R);$$

see [25] and [23]. The definition of  $\text{Mult}_{R, \square}^{\text{ap}}$  is motivated by the following facts. If, as usual, we denote by  $\times$  the normalized parabolic induction, then  $\langle \mathfrak{m} \rangle \cong \langle \mathfrak{m}_b \rangle \times \langle \mathfrak{m}_{nb} \rangle$  and

$$L(X, C(\langle \mathfrak{m} \rangle) \otimes_{s,s} C(\langle \mathfrak{n} \rangle)) = L(X, C(\langle \mathfrak{m}_b \rangle) \otimes_{s,s} C(\langle \mathfrak{n}_b \rangle)).$$

In particular, representations with nontrivial  $L$ -functions are precisely those coming from  $\text{Mult}_{R, \square}^{\text{ap}}$ . The set  $\text{Mult}_R^{\text{ap}}$  moreover admits an order, denoted by  $\preceq$ , which comes from the *degeneration order* on certain quiver-varieties, which we will discuss in a moment. Our goal is then to construct a map

$$\mathcal{S}_\psi : \text{Mult}_R^{\text{ap}} \rightarrow \mathfrak{Rep}_{W, \psi} = \bigcup_{n \in \mathbb{N}} \mathfrak{Rep}_{W, \psi, n}$$

satisfying, among others, the following properties.

- (1)  $L(X, \mathcal{S}_\psi(\mathfrak{m}) \otimes \mathcal{S}_{\psi^{-1}}(\mathfrak{n})) = L(X, C(\langle \mathfrak{m} \rangle) \otimes C(\langle \mathfrak{n} \rangle))$ .
- (2)  $\mathcal{S}_\psi(\mathfrak{m})$  admits  $\langle \mathfrak{m} \rangle$  as a quotient and  $\langle \mathfrak{m} \rangle$  appears in the Jordan–Hölder decomposition of  $\mathcal{S}_\psi(\mathfrak{m})$  with multiplicity one.
- (3)  $\mathcal{S}_\psi(\mathfrak{m})$  admits  $\langle \mathfrak{m} \rangle$  as its unique irreducible quotient.

Properties (2) and (3) would imply that  $\dim_R \text{Hom}_{G_n}(\mathcal{S}_\psi(\mathfrak{m}), \mathcal{S}_{\psi^{-1}}(\mathfrak{m}^\vee)^\vee) = 1$  and every morphism in this space would factor through  $\langle \mathfrak{m} \rangle$ . The description of irreducible representations in the style of properties (2) and (3) is also known as the *Langlands classification*, and has not yet been achieved over  $\overline{\mathbb{F}}_\ell$ , unlike the case  $\text{char}(R) = 0$ , where the relevant statements were first proved in [32] (see p. 2).

In this paper we construct such a map  $\mathcal{S}_\psi$  satisfying (1) and (2), as well as a map

$$\mathcal{S}_{\text{gen}, \psi} : \text{Mult}_{R, \square}^{\text{ap}} \rightarrow \mathfrak{Rep}_{W, \psi}$$

satisfying the analogues of properties (2) and (3) for multisegments in  $\text{Mult}_{R, \square}^{\text{ap}}$ . We also show that  $\mathcal{S}_{\text{gen}, \psi}(\mathfrak{m}) \subseteq \mathcal{S}_\psi(\mathfrak{m})$  for  $\mathfrak{m} \in \text{Mult}_{R, \square}$ . If  $\text{char}(R) = 0$ , these two constructions agree.

We start with a sketch of the construction of  $\mathcal{S}_\psi$ . If  $\tilde{\rho}$  is a cuspidal representation over  $\overline{\mathbb{Q}}_\ell$  admitting an integral structure, we denote its reduction mod  $\ell$  by  $r_\ell(\tilde{\rho})$ . If  $r_\ell(\tilde{\rho})$  is again a cuspidal representation  $\rho$ , we say  $\tilde{\rho}$  is a lift of  $\rho$ . It is then


straightforward to define a lift  $\tilde{m} \in \text{Mult}_{\overline{\mathbb{Q}}_\ell}$  of  $m \in \text{Mult}_{\overline{\mathbb{F}}_\ell}$  by first saying that a segment  $[a', b']_{\tilde{\rho}}$  is a lift of  $[a, b]_\rho$  if  $\tilde{\rho}$  is a lift of  $\rho$ ,  $a = a' \bmod o(\rho)$  and  $b - a = b' - a'$ , and then extending this notion linearly to all multisegments. In this case one can equip  $\mathcal{S}_\psi(\tilde{m})$  with its natural integral structure given by those Whittaker functions that take values in  $\overline{\mathbb{Z}}_\ell$ , and we denote the reduction mod  $\ell$  of this specific integral structure by  $\overline{\mathcal{S}_\psi(\tilde{m})}$ . For  $m$  an aperiodic multisegment over  $\overline{\mathbb{F}}_\ell$ , we write

$$\mathcal{S}_\psi(m) := \bigcap_{\tilde{m} \text{ lift of } m} \overline{\mathcal{S}_\psi(\tilde{m})},$$

where the intersection was taken in  $\mathcal{W}(\psi)$ .

**Theorem 1.** *The map  $\mathcal{S}_\psi: \text{Mult}_{\overline{\mathbb{F}}_\ell}^{\text{ap}} \rightarrow \mathfrak{Xep}_{\mathcal{W}, \psi}$  satisfies property (1). If  $m \in \text{Mult}_{\overline{\mathbb{F}}_\ell, \square}^{\text{ap}}$ , then  $\langle m \rangle$  is a quotient of  $\mathcal{S}_\psi(m)$  and appears with multiplicity 1 in it.*

The proof of this theorem uses our second construction of standard modules, denoted by  $\mathcal{S}_{\text{gen}, \psi}$ , to which we come now. We will focus on the case  $R = \overline{\mathbb{F}}_\ell$ , but the construction works similarly for  $\text{char}(R) = 0$ . In the introduction, for the sake of clarity, we will focus on multisegments of the form  $[a_1, b_1]_\rho + \cdots + [a_k, b_k]_\rho$  for a fixed cuspidal  $\rho$  with  $o(\rho) > 1$ . We denote this set by  $\text{Mult}_{\overline{\mathbb{F}}_\ell}(\rho)$  and the aperiodic multisegments in it by  $\text{Mult}_{\overline{\mathbb{F}}_\ell}(\rho)^{\text{ap}}$ . Let  $Q$  be the cyclical quiver with  $o(\rho)$  vertices and oriented counterclockwise.



Then the isomorphism classes of finite-dimensional, nilpotent  $\mathbb{C}$ -representations of  $Q$ , denoted by  $[Q]$ , are in bijection with  $\text{Mult}_{\overline{\mathbb{F}}_\ell}(\rho)$ . The order  $\leq$  on multisegments mentioned above is nothing but the degeneration order on  $[Q]$  transported to  $\text{Mult}_{\overline{\mathbb{F}}_\ell}(\rho)$ . In [28; 4; 27] the authors investigate the so-called *generic extension map*  $*$ :  $[Q] \times [Q] \rightarrow [Q]$ , which sends the tuple  $([M], [N])$  to  $[X]$ , where  $X \in \text{Ext}_Q^1(M, N)$  is such that  $\dim_{\mathbb{C}} \text{Hom}_Q(X, X)$  is minimal. The product so constructed,

$$*: \text{Mult}_{\overline{\mathbb{F}}_\ell}(\rho) \times \text{Mult}_{\overline{\mathbb{F}}_\ell}(\rho) \rightarrow \text{Mult}_{\overline{\mathbb{F}}_\ell}(\rho),$$

is associative. The subset of  $\text{Mult}_{\overline{\mathbb{F}}_\ell}(\rho)$  generated by multisegments of the form  $[i, i]_\rho$  is then precisely the set of aperiodic multisegments  $\text{Mult}_{\overline{\mathbb{F}}_\ell}(\rho)^{\text{ap}}$ .

The next result was motivated by a similar consideration of monomial bases of quantum and Hall algebras; see for example [7] or [28].

**Theorem 2.** *Let  $\mathfrak{m} = [i_1, i_1]_\rho * \cdots * [i_k, i_k]_\rho \in \text{Mult}_{\overline{\mathbb{F}}_\ell}^{\text{ap}}(\rho)$ . Then*

$$\mathcal{S}_{\text{gen}, \psi}(\mathfrak{m}) := \mathcal{W}(\rho|^{-|i_1} \times \cdots \times \rho|^{-|i_k}, \psi)$$

*only depends on  $\mathfrak{m}$  and satisfies properties (1) and (2).*

This construction allows us to define the desired map:

**Theorem 3.** *Let  $\mathfrak{m}, \mathfrak{m}' \in \text{Mult}_{R, \square}^{\text{ap}}$ . Then  $\mathcal{S}_{\text{gen}, \psi} : \text{Mult}_{R, \square}^{\text{ap}} \rightarrow \mathfrak{Rep}_{W, \psi}$  is defined by*

$$\mathcal{S}_{\text{gen}, \psi}(\mathfrak{m} * \mathfrak{m}') = \mathcal{W}(\mathcal{S}_{\text{gen}, \psi}(\mathfrak{m}) \times \mathcal{S}_{\text{gen}, \psi}(\mathfrak{m}'), \psi).$$

*Moreover, if  $\text{char}(R) = 0$ ,  $\mathcal{S}_\psi = \mathcal{S}_{\text{gen}, \psi}$  and if  $R = \overline{\mathbb{F}}_\ell$ ,  $\mathcal{S}_{\text{gen}, \psi}(\mathfrak{m}) \subseteq \mathcal{S}_\psi(\mathfrak{m})$ .*

Although the following seems at the moment out of reach, examples of small rank seem to suggest the equality  $\mathcal{S}_\psi = \mathcal{S}_{\text{gen}, \psi}$  on  $\text{Mult}_{R, \square}^{\text{ap}}$  as well as the equality  $\mathcal{S}_\psi(\mathfrak{m}) = \overline{\mathcal{S}_\psi(\tilde{\mathfrak{m}})}$  for any lift  $\tilde{\mathfrak{m}}$  of  $\mathfrak{m}$ .

## 2. Notation

Let  $F$  be a nonarchimedean local field with ring of integers  $\mathfrak{O}_F$  and residue field  $k$ , whose cardinality we denote by  $q$ . We also choose a uniformizer  $\varpi_F$  of the maximal ideal of  $\mathfrak{O}_F$  and fix a prime  $\ell$  not dividing  $q$ . We let  $|\cdot|$  be the absolute value of  $F$  such that  $|\varpi_F| = q^{-1}$ .

From now on  $R$  is one of the algebraically closed fields  $\overline{\mathbb{F}}_\ell$  or  $\overline{\mathbb{Q}}_\ell$ . We let  $\Lambda$  be the maximal ideal of  $\overline{\mathbb{Z}}_\ell \subseteq \overline{\mathbb{Q}}_\ell$  and fix a square root  $q^{\frac{1}{2}}$  of  $q$  in  $\overline{\mathbb{Z}}_\ell$ . By abuse of notation, we are also going to denote its image in  $\overline{\mathbb{F}}_\ell$  by  $q^{\frac{1}{2}}$ .

For  $n \in \mathbb{Z}_{\geq 0}$  we let  $G_n := \text{GL}_n(F)$ . We denote by  $1_n$  the identity in  $G_n$  and by

$$w_n = \begin{pmatrix} & & & 1 \\ & & & \\ & & \ddots & \\ & & & \\ 1 & & & \end{pmatrix}$$

the antidiagonal. For  $m, n \in \mathbb{N}$ , we denote by  $w_{n,m}$  the block-diagonal embedding of  $(1_m, w_n)$  into  $G_{n+m}$  and the space  $M_{n,m}$  of  $n \times m$  matrices with entries in  $F$ .

Inside  $G_n$  we consider for  $k \in \mathbb{Z}$  the closed subsets

$$G_n^k := \{g \in G_n : \text{val}_F(\det(g)) = k\},$$

where  $\text{val}_F$  denotes the valuation of  $F$  with  $\text{val}_F(\varpi) = 1$ . We let  $B_n$  be the Borel subgroup of  $G_n$  of upper diagonal matrices and  $N_n$  its unipotent subgroup. More generally, if  $\alpha = (\alpha_1, \dots, \alpha_k)$  is a partition of  $n$ , we denote by  $P_\alpha$  the parabolic subgroup containing  $B_n$  with Levi component  $G_\alpha := G_{\alpha_1} \times \cdots \times G_{\alpha_k}$ . Its opposite

parabolic subgroup  $\bar{P}_\alpha$  is conjugate to  $P_{(\alpha_k, \dots, \alpha_1)}$ . We denote by

$$H_{n,m} := \left\{ \begin{pmatrix} g & x \\ 0 & n \end{pmatrix} : g \in G_n, x \in M_{n,m}, n \in N_m \right\} \subseteq G_{n+m}$$

and let  $P_n := H_{n-1,1}$  be the mirabolic subgroup of  $G_n$ . Let  $C_c^\infty(\mathbb{F}^n)$  be the space of  $R$ -valued locally constant and compactly supported functions on  $\mathbb{F}^n$  and we set  $\epsilon_n := (0, \dots, 0, 1) \in \mathbb{F}^n$ .

**2.1. Representations.** In the next sections we recall the usual setup for  $\ell$ -adic and  $\ell$ -modular representation theory. For details, see [29]. Let  $G \subseteq G_n$  be a closed subgroup. We denote by  $\mathfrak{Rep}(G, R)$  the category of smooth, admissible representations of finite length of  $G$ . Whenever possible we will omit the field of coefficients  $R$ . Let  $\alpha = (\alpha_1, \dots, \alpha_k)$  be a partition of  $n$ . We let  $\mathfrak{Rep}_\alpha = \mathfrak{Rep}(G_\alpha)$ , let  $\mathfrak{Irr}_\alpha$  be the set of isomorphism classes of irreducible representations in  $\mathfrak{Rep}_\alpha$  and set

$$\mathfrak{Rep} := \bigcup_{n \in \mathbb{N}} \mathfrak{Rep}_n, \quad \mathfrak{Irr} := \bigcup_{n \in \mathbb{N}} \mathfrak{Irr}_n.$$

If  $G \subseteq H \subseteq G_n$  are closed subgroups, we denote the functors of normalized induction by  $\text{Ind}_H^G$  and their compactly supported version by  $\text{ind}_H^G$ . We recall the normalized Jacquet functor and parabolic induction corresponding to parabolic subgroups  $P = MN$  of  $G_\alpha$ , which give rise to the exact functors

$$r_P : \mathfrak{Rep}_\alpha \rightarrow \mathfrak{Rep}(M), \quad \text{ind}_P^{G_\alpha} : \mathfrak{Rep}(M) \rightarrow \mathfrak{Rep}_\alpha.$$

We write  $r_\alpha := r_{P_\alpha}$  and  $\bar{r}_\alpha := r_{\bar{P}_\alpha}$ . Recall that  $r_\alpha$  and  $\bar{r}_\alpha$  are the left and right adjoints, respectively, of  $\text{ind}_{P_\alpha}^{G_n}$ , by Frobenius and Bernstein reciprocities (respectively). By abuse of notation we will also notate the maps they induce between the respective Grothendieck groups by the same letters. As is conventional, we will write

$$\pi_1 \times \cdots \times \pi_k := \text{ind}_{P_\alpha}^{G_n}(\pi_1 \otimes \cdots \otimes \pi_k).$$

If  $\chi$  is a smooth character of  $\mathbb{F}$  and  $\pi \in \mathfrak{Rep}$ , we write  $\chi\pi = \chi(\det)\pi$ . If  $\pi \in \mathfrak{Rep}_n$ , we denote the corresponding element in the Grothendieck of  $\mathfrak{Rep}_n$  by  $[\pi]$  and define the length of  $\pi$  as the number of its irreducible subquotients counted with multiplicity. If  $\pi$  is a representation of  $G_n$ , we denote by  $\pi^\epsilon$  the representation obtained by twisting  $\pi$  by  $g \mapsto w_n(g^{-1})^t w_n$ . We recall that if  $\pi$  is irreducible and  $\ell > 2$ , then  $\pi^\epsilon \cong \pi^\vee$  (see for example [29] or [23, Remark 2.7]), and

$$(\pi_1 \times \pi_2)^\epsilon = \pi_2^\epsilon \times \pi_1^\epsilon.$$

Finally, we recall that parabolic induction on  $G_n$  is commutative on the level of Grothendieck groups, in particular for  $\pi, \pi' \in \mathfrak{Irr}$  such that  $\pi \times \pi'$  is irreducible,

$\pi \times \pi' \cong \pi' \times \pi$ ; see [29, 1.16] for  $\ell > 2$ , [32, Theorem 1.9] for the case  $R = \mathbb{C}$  and [23, Proposition 2.6] in general.

**2.1.1. Cuspidal representations.** A representation  $\rho \in \mathfrak{Irr}_n$  is called cuspidal if for all nontrivial partitions  $\alpha$  of  $n$ ,  $r_\alpha(\rho) = 0$ . It is called supercuspidal if there exists no nontrivial partition  $\alpha$  and  $\pi \in \mathfrak{Irr}_\alpha$  such that  $\rho$  appears as a subquotient of  $\text{ind}_{P_\alpha}^{G_n} \pi$ . In general, supercuspidal implies cuspidal and if  $R = \overline{\mathbb{Q}}_\ell$ , cuspidal implies supercuspidal. We denote the subset of  $\mathfrak{Irr}_n$  consisting of cuspidal respectively supercuspidal representations by  $\mathfrak{C}_n$  respectively  $\mathfrak{SC}_n$  and define

$$\mathfrak{C} := \bigcup_{n \in \mathbb{N}} \mathfrak{C}_n, \quad \mathfrak{SC} := \bigcup_{n \in \mathbb{N}} \mathfrak{SC}_n.$$

We also recall the notation of cuspidal support, meaning that if  $\pi \in \mathfrak{Irr}$  there exist up to possible permutation and isomorphism unique  $\rho_1, \dots, \rho_k \in \mathfrak{C}$  such that  $\pi \hookrightarrow \rho_1 \times \dots \times \rho_k$ . We denote by  $\text{cusp}(\pi) := [\rho_1] + \dots + [\rho_k]$  the cuspidal support of  $\pi$ . Weakening in the above definition the condition of being a subrepresentation to being a subquotient and cuspidal representations to supercuspidal representations gives rise to the supercuspidal support  $\text{scusp}(\pi)$ . If at any point the field of coefficients becomes important, we will add it in parentheses to the respective category or set, e.g.,  $\mathfrak{Irr}(\overline{\mathbb{F}}_\ell)$  versus  $\mathfrak{Irr}(\overline{\mathbb{Q}}_\ell)$ .

We recall the following notions; see for example [24, §3.4, §4.5]. Let  $\rho \in \mathfrak{C}$  and recall that  $\rho \times \chi\rho$  is reducible if and only if  $\chi \cong |\cdot|^\pm$ . We recall the cuspidal line

$$\mathbb{Z}[\rho] := \{[\rho|\cdot|^k] : k \in \mathbb{Z}\}$$

and denote the cardinality of  $\mathbb{Z}[\rho]$  by  $o(\rho)$ . Note that  $o(\rho)$  is finite if and only if  $R = \overline{\mathbb{F}}_\ell$ . Set

$$e(\rho) := \begin{cases} o(\rho) & \text{if } o(\rho) > 1, \\ \ell & \text{otherwise.} \end{cases}$$

One can associate to  $\rho$  an integer  $f(\rho)$  via type theory. If  $o(\rho) > 1$ ,  $o(\rho)$  is the order of  $q^{f(\rho)}$  in  $R$ . We let  $\mathfrak{C}_n^\square$  be the subset of  $\mathfrak{C}_n$  consisting of  $\rho$  such that  $o(\rho) > 1$ ; i.e., those  $\rho$  which make  $\rho \times \rho$  irreducible. Finally, we say  $\rho$  and  $\rho'$  are in different cuspidal lines if  $[\rho'] \notin \mathbb{Z}[\rho]$  and we set  $\mathfrak{C}^\square := \bigcup_{n \in \mathbb{N}} \mathfrak{C}_n^\square$ .

**2.2. Multisegments.** We now recall the combinatorics of multisegment (for details, see [32; 23]). Let  $\rho \in \mathfrak{C}_m$  and  $a \leq b \in \mathbb{Z}$ . A segment is a sequence

$$[a, b]_\rho = ([\rho|\cdot|^a], \dots, [\rho|\cdot|^b]).$$

Two segments  $[a, b]_\rho$  and  $[a', b']_{\rho'}$  are equal if and only if  $\rho'|-|^a \cong \rho|-|^a$  and  $b - a = b' - a'$ . We also let  $[a, b]_\rho^\vee = [-b, -a]_{\rho^\vee}$ . Define

$$\begin{aligned} {}^-[a, b]_\rho &= [a + 1, b]_\rho, & [a, b]_\rho^- &= [a, b - 1]_\rho, \\ {}^+[a, b]_\rho &= [a - 1, b]_\rho, & [a, b]_\rho^+ &= [a, b + 1]_\rho, \\ a_\rho([a, b]_\rho) &= a \in \begin{cases} \mathbb{Z}/(o(\rho)\mathbb{Z}) & \text{if } o(\rho) < \infty, \\ \mathbb{Z} & \text{if } o(\rho) = \infty, \end{cases} \\ b_\rho([a, b]_\rho) &= b \in \begin{cases} \mathbb{Z}/(o(\rho)\mathbb{Z}) & \text{if } o(\rho) < \infty, \\ \mathbb{Z} & \text{if } o(\rho) = \infty. \end{cases} \end{aligned}$$

We define the length and degree of a segment as  $l([a, b]_\rho) := b - a + 1 \in \mathbb{Z}$  and  $\deg([a, b]_\rho) = (b - a + 1)m \in \mathbb{Z}$ . The cuspidal support of  $[a, b]_\rho$  is defined as  $\text{cusp}([a, b]_\rho) := [\rho|-|^a] + \cdots + [\rho|-|^b]$ . By abuse of notation we will often write  $\rho$  for the segment  $[0, 0]_\rho$ .

A multisegment is a formal finite sum and we extend the length  $l$ , the degree  $\deg$ , the notion of cuspidal support, and  $(-)^{\vee}$  linearly. We let  $\text{Mult}_R$  be the set of multisegments and  $\text{Mult}_R(\rho)$  be the set of multisegments consisting only of segments of the form  $[a, b]_\rho$ . A multisegment is called aperiodic if it does not contain a submultisegment of the form

$$[a, b]_\rho + [a + 1, b + 1]_\rho + \cdots + [a + e(\rho) - 1, b + e(\rho) - 1]_\rho.$$

For any subset  $\mathcal{N} \subseteq \text{Mult}_R$ , we denote by  $\mathcal{N}^{\text{ap}}$  the aperiodic multisegments in  $\mathcal{N}$ . For any  $\mathfrak{m} \in \text{Mult}_R$ , we decompose  $\mathfrak{m} = \mathfrak{m}_b + \mathfrak{m}_{nb}$  with  $\mathfrak{m}_b$  consisting of all segments  $[a, b]_\rho$  in  $\mathfrak{m}$  with  $\rho \in \mathfrak{C}^\square$ . We let  $\text{Mult}_{R, \square}$  be the set of multisegments such that  $\mathfrak{m} = \mathfrak{m}_b$ .

**Theorem 2.2.1** [23, §9; 32, §6]. *There exists a surjective map*

$$Z : \text{Mult}_R \rightarrow \mathfrak{Irr}$$

*satisfying the following.*

- (1)  $Z(\mathfrak{m}) \in \mathfrak{Irr}_{\deg(\mathfrak{m})}$ .
- (2)  $Z$  restricted to aperiodic multisegments is a bijection.
- (3)  $Z(\mathfrak{m})^\vee \cong Z(\mathfrak{m}^\vee)$ .
- (4) For  $\mathfrak{m} \in \text{Mult}_R^{\text{ap}}$ ,  $\text{cusp}(Z(\mathfrak{m})) = \text{cusp}(\mathfrak{m})$ . If  $\mathfrak{m} \in \text{Mult}_R(\rho)^{\text{ap}}$  with  $\rho \in \mathfrak{S}\mathfrak{C}$ , then  $\text{scusp}(Z(\mathfrak{m})) = \text{cusp}(Z(\mathfrak{m}))$ .
- (5) If  $\mathfrak{m} = \mathfrak{m}_1 + \mathfrak{m}_2$  then  $Z(\mathfrak{m})$  is a subquotient of  $Z(\mathfrak{m}_1) \times Z(\mathfrak{m}_2)$  and appears with multiplicity one in its Jordan–Hölder decomposition.
- (6)  $Z([a, b]_\rho)$  is a subrepresentation of  $\rho|-|^a \times \cdots \times \rho|-|^b$ .

(7) If  $\rho_1, \dots, \rho_l \in \mathcal{C}$  lie in pairwise different cuspidal lines and  $\mathfrak{m} = \mathfrak{m}_1 + \dots + \mathfrak{m}_k$  with  $\mathfrak{m}_i \in \text{Mult}_R(\rho_i)$ ,

$$Z(\mathfrak{m}) \cong Z(\mathfrak{m}_1) \times \dots \times Z(\mathfrak{m}_k).$$

If  $\mathfrak{m}$  consists only of segments of length 1, we call  $Z(\mathfrak{m})$  nondegenerate. Finally, we recall the Aubert–Zelevinsky involution  $(-)^* : \mathcal{Irr} \rightarrow \mathcal{Irr}$  (see [1; 32; 25]), and denote

$$\langle - \rangle := (Z)^* : \text{Mult}_R^{\text{ap}} \rightarrow \mathcal{Irr}.$$

This map preserves the cuspidal support and if  $\mathfrak{m}, \mathfrak{m}_1, \dots, \mathfrak{m}_k \in \text{Mult}_R^{\text{ap}}$ , the multiplicity of  $\langle \mathfrak{m} \rangle$  in  $\langle \mathfrak{m}_1 \rangle \times \dots \times \langle \mathfrak{m}_k \rangle$  is the multiplicity of  $Z(\mathfrak{m})$  in  $Z(\mathfrak{m}_1) \times \dots \times Z(\mathfrak{m}_k)$ .

**Lemma 2.2.1.** *Let  $\rho, \rho' \in \mathcal{C}^\square$  be in different cuspidal lines. Moreover, let  $\pi$  be an irreducible subquotient of  $\rho_1 \times \dots \times \rho_k$ ,  $\rho_i \in \mathbb{Z}[\rho]$ , and  $\pi'$  be an irreducible subquotient of  $\rho'_1 \times \dots \times \rho'_l$ ,  $\rho'_i \in \mathbb{Z}[\rho]$ . Then  $\pi \times \pi'$  is irreducible.*

*Proof.* By the classification of cuspidal representations in [21, §6], the cuspidal lines of the cuspidal supports of  $\pi$  and  $\pi'$  do not intersect. Indeed, if  $\sigma$  is a cuspidal representation appearing in the cuspidal support of  $\pi$ , its supercuspidal support consists of representations in  $\mathbb{Z}[\rho]$ ; see [23, Theorem 9.36(3)]. By the uniqueness of the cuspidal and supercuspidal supports, it follows that the cuspidal representations appearing in  $\pi$  and  $\pi'$  lie in different cuspidal lines. The claim follows from property (7) in Theorem 2.2.1.  $\square$

Let  $\mathfrak{m} \in \text{Mult}_R$  and  $\Delta = [a, b]_\rho$ ,  $\Delta' = [a', b']_\rho$  be two segments in  $\mathfrak{m}$  with

$$a + 1 \leq a' \leq b + 1 \leq b'.$$

An *elementary operation* on  $\mathfrak{m}$  refers to changing the segment in the following way:

$$\mathfrak{m} \mapsto \mathfrak{m} - \Delta - \Delta' + [a, b']_\rho + [a', b]_\rho.$$

We will say that the elementary operation is of the form  $[a, b]_\rho + [a', b']_\rho \mapsto [a, b']_\rho + [a', b]_\rho$ . We write  $\mathfrak{n} \leq \mathfrak{m}$  if  $\mathfrak{n}$  can be obtained by repeated applications of elementary operations to  $\mathfrak{m}$  and  $\mathfrak{n} < \mathfrak{m}$  if  $\mathfrak{n} \leq \mathfrak{m}$  and  $\mathfrak{m} \neq \mathfrak{n}$ . We call two segments  $\Delta, \Delta'$  linked if there exists a presentation  $\Delta = [a, b]_\rho$ ,  $\Delta' = [a', b']_\rho$  such that one of the inequalities

$$a + 1 \leq a' \leq b + 1 \leq b' \quad \text{or} \quad a' + 1 \leq a \leq b' + 1 \leq b$$

is satisfied, i.e., there exists a multisegment  $\mathfrak{n}$  with  $\mathfrak{n} < \Delta + \Delta'$ . The multisegment  $\mathfrak{m} = \Delta_1 + \dots + \Delta_k$  is called *unlinked*, if for all  $i \neq j \in \{1, \dots, k\}$   $\Delta_i$  and  $\Delta_j$  are unlinked.

**Lemma 2.2.2** [32, Theorem 7.1; 8, Theorem 6.4.1]. *Let  $\mathfrak{m} = \Delta_1 + \cdots + \Delta_k \in \mathcal{M}ult_R$  and  $\mathfrak{n} \in \mathcal{M}ult_R^{\text{ap}}$ . Then  $\langle \Delta_1 \rangle \times \cdots \times \langle \Delta_k \rangle$  contains  $\langle \mathfrak{n} \rangle$  as an irreducible subquotient if and only if  $\mathfrak{n} \leq \mathfrak{m}$ . In particular,  $\langle \mathfrak{m} \rangle \cong \langle \Delta_1 \rangle \times \cdots \times \langle \Delta_k \rangle$  if and only if  $\mathfrak{m}$  is unlinked.*

*Proof.* The case in  $R = \overline{\mathbb{Q}}_\ell$  was treated in [32, Theorem 7.1] and the case  $R = \overline{\mathbb{F}}_\ell$  in [8, Theorem 6.4.1]. (In [8], the claim was proven for  $Z(-)$  instead of  $\langle - \rangle$ , however, applying the Aubert–Zelevinsky yields the equivalence.)

To see the claim regarding the irreducibility, note that the induced representation is irreducible if and only if it contains no irreducible subquotient  $\langle \mathfrak{n}' \rangle$  for  $\mathfrak{m} \neq \mathfrak{n}' \in \mathcal{M}ult_R$ . If  $\mathfrak{n}'$  is not aperiodic, it is easy to construct  $\mathfrak{n}'' < \mathfrak{n}'$  via one elementary operation, and hence  $\mathfrak{n}'' < \mathfrak{n}' \leq \mathfrak{m}$ . Repeating this process, we arrive at  $\mathfrak{n} < \mathfrak{n}' \leq \mathfrak{m}$  with  $\mathfrak{n}$  aperiodic, and hence by the above lemma  $\langle \mathfrak{n} \rangle$  is a subquotient of the induced representation. Thus the induced representation is irreducible if and only if it contains no  $\langle \mathfrak{n} \rangle$  with  $\mathfrak{n} \in \mathcal{M}ult_R^{\text{ap}}$ , which by the above can only happen if and only if  $\mathfrak{n} \leq \mathfrak{m}$ . □

Assume for a moment that  $R = \overline{\mathbb{Q}}_\ell$  and let  $\Delta, \Delta'$  be two segments. We say  $\Delta$  precedes  $\Delta'$  if  $\Delta = [a, b]_\rho$ ,  $\Delta' = [a', b']_\rho$  and

$$a + 1 \leq a' \leq b + 1 \leq b'.$$

Thus,  $\Delta$  and  $\Delta'$  are unlinked if and only if  $\Delta$  does not precede  $\Delta'$  and vice versa. Let  $\mathfrak{m} = \Delta_1 + \cdots + \Delta_k \in \mathcal{M}ult_{\overline{\mathbb{Q}}_\ell}$ . We say  $(\Delta_1, \dots, \Delta_k)$  is in *arranged form* if for all  $i, j \in \{1, \dots, k\}$ ,  $i < j$   $\Delta_i$  does not precede  $\Delta_j$ . Any multisegment over  $\overline{\mathbb{Q}}_\ell$  admits an arranged form and any two arranged forms can be obtained from each other by repeatedly changing the order of two neighboring unlinked segments, i.e., replacing  $(\dots, \Delta_i, \Delta_{i+1}, \dots) \mapsto (\dots, \Delta_{i+1}, \Delta_i, \dots)$  if  $\Delta_i$  and  $\Delta_{i+1}$  are unlinked.

**2.3. Integral structures.** We recall the following results of [29, I.9] on integral structures. Let  $\alpha$  be a partition of  $n \in \mathbb{N}$  and  $(\tilde{\pi}, V) \in \mathfrak{R}ep_\alpha(\overline{\mathbb{Q}}_\ell)$ . We recall that an integral structure of  $\tilde{\pi}$  is a  $\overline{\mathbb{Z}}_\ell$ -lattice  $\mathfrak{l}$  in  $V$  which generates  $V$ , i.e., the natural map induces an isomorphism  $\mathfrak{l} \otimes_{\overline{\mathbb{Z}}_\ell} \overline{\mathbb{Q}}_\ell \cong V$ . If  $\tilde{\pi}$  admits an integral structure it is called *integral*. For  $\tilde{\pi} \in \mathfrak{R}ep_\alpha$  integral and  $\mathfrak{l}$  an integral structure,  $\mathfrak{l} \otimes_{\overline{\mathbb{Z}}_\ell} \overline{\mathbb{F}}_\ell$  is an object in  $\mathfrak{R}ep_\alpha(\overline{\mathbb{F}}_\ell)$ . We denote by

$$r_\ell(\tilde{\pi}) := [\mathfrak{l} \otimes_{\overline{\mathbb{Z}}_\ell} \overline{\mathbb{F}}_\ell],$$

which is an element in the Grothendieck group of  $\mathfrak{R}ep_\alpha(\overline{\mathbb{F}}_\ell)$ . It is  $r_\ell(\tilde{\pi})$  which is independent of the chosen integral structure; the representation  $\mathfrak{l} \otimes_{\overline{\mathbb{Z}}_\ell} \overline{\mathbb{F}}_\ell$  is highly dependent on the specific  $\mathfrak{l}$  and in general not irreducible. Moreover, the representation  $\text{ind}_{P_\alpha}^{G_n} \tilde{\pi}$  is integral, with the natural integral structure  $\text{ind}_{P_\alpha}^{G_n} \mathfrak{l}$ , i.e., the  $\mathfrak{l}$ -valued functions. Then

$$(\text{ind}_{P_\alpha}^{G_n} \mathfrak{l}) \otimes_{\overline{\mathbb{Z}}_\ell} \overline{\mathbb{F}}_\ell \cong \text{ind}_{P_\alpha}^{G_n} (\mathfrak{l} \otimes_{\overline{\mathbb{Z}}_\ell} \overline{\mathbb{F}}_\ell).$$

Moreover, if  $\tilde{\tau}$  is a subrepresentation (resp. quotient) of  $\tilde{\pi}$ , then  $\mathfrak{l} \cap \tilde{\tau}$  (resp. the image of  $\mathfrak{l}$  in  $\tilde{\tau}$ ) is an integral structure of  $\tilde{\tau}$ .

Finally, the functors  $r_\alpha$  and  $r_\ell$  commute on the level of Grothendieck groups by [6, Proposition 1.4(i)], i.e.,

$$r_\alpha(r_\ell(\tilde{\pi})) = r_\ell(r_\alpha(\tilde{\pi})).$$

An integral representation  $\tilde{\pi} \in \mathfrak{Irr}(\overline{\mathbb{Q}}_\ell)$  is called a *lift* of  $\pi \in \mathfrak{Irr}(\overline{\mathbb{F}}_\ell)$  if  $r_\ell(\tilde{\pi}) = [\pi]$ . If  $\Delta = [a, b]_\rho$  is a segment over  $\overline{\mathbb{F}}_\ell$ , a segment  $\tilde{\Delta} = [a', b']_{\tilde{\rho}}$  over  $\overline{\mathbb{Q}}_\ell$  is called a lift of  $\Delta$  if  $\tilde{\rho}$  is a lift of  $\rho$ ,  $b' - a' = b - a$  and  $a = a' \bmod o(\rho)$ . Finally, we extend this notation of a lift to segments  $\text{Mult}_{\overline{\mathbb{F}}_\ell}$  linearly.

In [29, III] the Bushnell–Kutzko construction for  $\ell$ -adic and modular representations was carried out. It was shown that every cuspidal representation over  $\overline{\mathbb{F}}_\ell$  admits a cuspidal lift  $\tilde{\rho}$  to  $\overline{\mathbb{Q}}_\ell$ , and that if  $\tilde{\rho}$  is such a lift of  $\rho$ ,  $f(\rho) = f(\tilde{\rho})$ .

**Lemma 2.3.1** [23, Theorem 9.39]. *Let  $\mathfrak{m} \in \text{Mult}_{\overline{\mathbb{F}}_\ell}^{\text{ap}}$  and let  $\tilde{\mathfrak{m}}$  be a lift of  $\mathfrak{m}$ . Then  $Z(\mathfrak{m})$  appears with multiplicity 1 in  $r_\ell(Z(\tilde{\mathfrak{m}}))$  and  $\langle \mathfrak{m} \rangle$  appears with multiplicity 1 in  $r_\ell(\langle \tilde{\mathfrak{m}} \rangle)$ .*

*Proof.* The claim for  $Z$  can be found in [23, Theorem 9.39]. The one for  $\langle - \rangle$  is derived from the definition of  $\langle - \rangle$  in [1] as follows. Recall the definition of  $(-)^*$ : For  $[\pi]$  an element in the Grothendieck group of  $\mathfrak{Rep}_n$ , one first defines a nonzero element  $D([\pi])$  in the same Grothendieck group, which is represented by the cohomology of a complex obtained from  $[\pi]$  by applying repeatedly the Jacquet functor and parabolic induction. In particular, if  $[\pi] \leq [\tau]$ , then  $D([\pi]) \leq D([\tau])$ . Moreover, if  $\pi$  is irreducible, there exists a unique irreducible summand  $[\pi^*]$  in  $D(\pi)$  with the same cuspidal support as  $\pi$  and if  $R = \overline{\mathbb{Q}}_\ell$ , then  $D(\pi) = [\pi^*]$ . The above construction implies that  $D$  commutes with  $r_\ell$ . Thus,  $r_\ell(\langle \tilde{\mathfrak{m}} \rangle) = r_\ell(D(Z(\tilde{\mathfrak{m}}))) = D(r_\ell(Z(\tilde{\mathfrak{m}})))$ , where the latter contains  $D(Z(\mathfrak{m})) \geq [\langle \mathfrak{m} \rangle]$  by the first claim.  $\square$

**2.4. Generic extensions.** In this section we recall the composition algebra of the cyclic quiver; compare [28]. We fix  $1 < n \in \mathbb{N} \cup \{\infty\}$  and consider the cyclic quiver  $Q$  with vertices

$$I = \begin{cases} \mathbb{Z}/(n\mathbb{Z}) & \text{if } n < \infty, \\ \mathbb{Z} & \text{if } n = \infty, \end{cases}$$

and an arrow from  $i$  to  $j$  if  $j = i + 1 \bmod n$  if  $n$  is finite and  $j = i + 1$  if  $n = \infty$ . We recall that a representation of  $Q$  is nothing but a finite-dimensional  $I$ -graded  $\mathbb{C}$ -vector  $V$  space together with a linear map  $T : V \mapsto V$  of weight 1. We call the representation nilpotent if  $T^N = 0$  for large enough  $N \in \mathbb{N}$ , and associate to  $V$  the dimension vector  $\text{grdim} V \in \mathbb{N}^I$ , whose  $i$ -th entry equals  $\dim_{\mathbb{C}} V_i$ . Note that if  $n = \infty$ , for all but finitely many  $i \in I$ , the dimension of  $\dim_{\mathbb{C}} V_i$  vanishes. We call a vector  $d \in \mathbb{N}^I$  a dimension vector if it is 0 for all but finitely many  $i \in I$ . We

let  $E_{\mathbf{d}} = \{(V_{\mathbf{d}}, T)\}$  be the set of finite-dimensional, nilpotent representations of  $Q$  over  $\mathbb{C}$  with underlying graded vector space

$$V_{\mathbf{d}} := \bigoplus_{i \in I} \mathbb{C}^{d_i},$$

where we write for  $\mathbb{C}^0$  the trivial vector space. We let  $G_{\mathbf{d}} := \prod_{i \in I} \mathrm{GL}_{d_i}(\mathbb{C})$ , where  $\mathrm{GL}_0(\mathbb{C})$  denotes the trivial group, and note that  $G_{\mathbf{d}}$  acts on  $E_{\mathbf{d}}$  by conjugation. The orbits of this action are naturally parametrized as follows. We let  $\mathcal{M}ult(Q)$  be the set of multisegments, i.e., the formal finite sums of segments  $[a, b]$ ,  $a \leq b \in \mathbb{Z}$  up to the equivalence

$$[a, b] \sim [a', b'] \quad \text{if} \quad \begin{cases} a = a' \bmod n, b - a = b' - a' & \text{if } n < \infty, \\ a = a', b' = b & \text{if } n = \infty. \end{cases}$$

We write  $[a, b]^\vee = [-b, -a]$ ,  $l([a, b]) = b - a + 1$  and extend these operations linearly to  $\mathcal{M}ult(Q)$ . To  $[a, b] \in \mathcal{M}ult(Q)$  we associate the indecomposable representation  $\lambda([a, b])$  whose underlying vector space has as a basis  $b - a + 1$  vectors  $e_1, \dots, e_{b-a+1}$ , with  $e_i$  in degree  $i \bmod n$  if  $n < \infty$  and in degree  $i$  if  $n = \infty$  and  $T(e_i) = e_{i+1}$  for  $i \leq b - a$  and  $T(e_{b-a+1}) = 0$ . The dimension vector of this representation, denoted by  $\mathrm{grdim}[a, b]$ , has as its  $i$ -th entry

$$(\mathrm{grdim}[a, b])_i = \begin{cases} \#\{x \in \mathbb{Z} : a \leq x \leq b, x = i \bmod n\} & \text{if } n < \infty, \\ 1 & \text{if } n = \infty, a \leq i \leq b, \\ 0 & \text{otherwise.} \end{cases}$$

More generally, we associate to a multisegment  $\mathfrak{m} = [a_1, b_1] + \dots + [a_k, b_k]$  the representation  $\lambda(\mathfrak{m}) := \lambda([a_1, b_1]) \oplus \dots \oplus \lambda([a_k, b_k])$ . We denote its dimension vector  $\mathrm{grdim} \mathfrak{m} = \mathrm{grdim}[a_1, b_1] + \dots + \mathrm{grdim}[a_k, b_k]$ . We call a multisegment aperiodic if either  $n = \infty$  or  $n < \infty$  and it does not contain a multisegment of the form  $[a, b] + \dots + [a+n-1, b+n+1]$ . We denote the set of aperiodic multisegments by  $\mathcal{M}ult(Q)^{\mathrm{ap}}$ .

Via this construction, the  $G_{\mathbf{d}}$ -orbits  $[E_{\mathbf{d}}]$  of  $E_{\mathbf{d}}$  are then in bijection with multisegments  $\mathfrak{m}$  such that  $\mathbf{d} = \mathrm{grdim} \mathfrak{m}$ . For  $M$  a representation in  $E_{\mathbf{d}}$ , we denote by  $[M] = G_{\mathbf{d}} \cdot M$  its orbit.

We write  $[M] \leq [N]$  for two orbits in  $E_{\mathbf{d}}$  if  $[N]$  is in the closure of  $[M]$  with respect to the analytic topology. This relation gives rise to the so-called *degeneration order*. We also recall the associative product

$$* : [E_{\mathbf{d}}] \times [E_{\mathbf{d}'}] \rightarrow [E_{\mathbf{d}+\mathbf{d}'}]$$

given by sending  $([M], [N]) \mapsto [M] * [N]$  to the orbit of their *generic extension*; see [7, §3] for cyclic quivers and [27, §2] for Dynkin quivers. The generic extension of two representations  $M$  and  $N$  is defined as the set of  $X \in \mathrm{Ext}_Q^1(M, N)$  for which

the dimension of the complex algebraic variety  $\overline{[X]}$  is maximal, or equivalently, for which  $\dim_{\mathbb{C}} \text{Hom}_Q(X, X)$  is minimal. Any  $X, X'$  in the generic extension of  $M$  and  $N$  lie in the same equivalence class, thus we can define  $[M] * [N]$  as  $[X]$  for some  $X$  in the generic extension of  $M$  and  $N$ .

**Lemma 2.4.1** ([27, Proposition 2.4]; see also [7, Proposition 3.4]). *Let  $M \in E_{\mathbf{d}}, N \in E_{\mathbf{d}'}, X \in E_{\mathbf{d}+\mathbf{d}'}$ . Then  $[M] * [N] \leq [X]$  if and only if there exist  $[M] \leq [M'], [N] \leq [N']$  such that there exists a short exact sequence  $0 \rightarrow M' \rightarrow X \rightarrow N' \rightarrow 0$ . In particular, if  $[M] \leq [M'], [N] \leq [N']$  then  $[M] * [N] \leq [M'] * [N']$ .*

For a word  $w = i_1 \dots i_k$  of indices in  $I$ , we write

$$\mathfrak{m}_{\text{gen}}(w) = [i_1, i_1] * \dots * [i_k, i_k],$$

and we denote the set of words in  $I$  by  $\Omega$ . We can describe  $\mathfrak{m}_{\text{gen}}(w)$  recursively as follows; see for example [7, p. 285 and Proposition 3.7].

For  $\mathfrak{m} \in \text{Mult}(Q)$  and  $i \in I$  we let

$$i + \mathfrak{m} := \begin{cases} \mathfrak{m} + [i, i] & \text{if there does not exist a segment of the form } [i+1, b] \text{ in } \mathfrak{m}, \\ \mathfrak{m} + {}^+\Delta - \Delta & \text{where } \Delta \text{ is the longest segment of the form } [i+1, b] \text{ in } \mathfrak{m}. \end{cases}$$

Similarly, we let

$$\mathfrak{m} + i := \begin{cases} \mathfrak{m} + [i, i] & \text{if there does not exist a segment of the form } [a, i-1] \text{ in } \mathfrak{m}, \\ \mathfrak{m} + \Delta^+ - \Delta & \text{where } \Delta \text{ is the longest segment of the form } [a, i-1] \text{ in } \mathfrak{m}. \end{cases}$$

**Lemma 2.4.2** [7, Proposition 3.7]. *In this notation, we have  $\mathfrak{m}_{\text{gen}}(iw) = i + \mathfrak{m}_{\text{gen}}(w)$  and  $\mathfrak{m}_{\text{gen}}(wi) = \mathfrak{m}_{\text{gen}}(w) + i$ .*

From now on we implicitly identify the isomorphism classes of representations of  $Q$  with  $\text{Mult}(Q)^{\text{ap}}$ .

**Theorem 2.4.1.** *The map  $\mathfrak{m}_{\text{gen}} : \Omega \rightarrow \text{Mult}(Q)$  has image  $\text{Mult}(Q)^{\text{ap}}$  and two words  $w$  and  $w'$  give rise to the same multisegment if and only if they are related by the degenerate Serre relations.*

- (1)  $ij = ji$  if  $i$  and  $j$  are not neighbors.
- (2)  $i(i+1)i = ii(i+1)$  and  $i(i+1)(i+1) = (i+1)i(i+1)$  if  $n > 2$ .
- (3)  $i(i+1)ii = ii(i+1)i$  if  $n = 2$ .

*Proof.* For a proof see [28, Theorem A, Theorem B], where the author considers the algebras  $H_t(Q)$  we consider below specialised to the case that  $t$  is a prime number. We sketch the argument for the sake of the reader; see also [27, Theorem 4.2]. Let  $t$  be an indeterminate and consider the  $\mathbb{Q}[t]$ -algebra  $H_t(Q)$ , which is generated by variables  $x_1, \dots, x_n$  satisfying the following relations:

- (1)  $x_i x_j = x_j x_i$  if  $i, j$  are not neighbors in  $Q$ ;

- (2)  $x_i^2 x_{i+1} - t x_{i+1} x_i^2 = (t+1)x_i x_{i+1} x_i$  and  $x_i x_{i+1}^2 - t x_{i+1}^2 x_i = (t+1)x_{i+1} x_i x_{i+1}$  if  $n > 2$ ;
- (3)  $t x_2 x_1^3 - (t^2 + t + 1)(x_1 x_2 x_1^2 + x_1^2 x_2 x_1) = t x_1^3$  and  $t x_1 x_2^3 - (t^2 + t + 1)(x_2 x_1 x_2^2 + x_2^2 x_1 x_2) = t x_2^3$  if  $n = 2$ .

Given a dimension vector  $\mathbf{d}$ , we can ask for the rank of the free  $\mathbb{Q}[t]$ -submodule  $H_t^{\mathbf{d}}(Q)$  of  $H_t(Q)$  spanned by the monomials containing  $x_i$  with multiplicity  $\mathbf{d}_i$ . Specializing at  $t = 1$ , we have that  $H_1(Q)$  is the universal enveloping algebra of a certain upper-triangular part of a Lie algebra, and obtain by [28, Proposition 7.2] that the rank equals the number of aperiodic elements in  $\text{Mult}(Q)$  with dimension vector  $\mathbf{d}$ . The author achieves this by constructing an explicit PBW basis of the latter space. But on the other hand, we obtain by specializing at  $t = 0$  the associative algebra on  $\Omega$  subject to the Serre relations. We have a morphism of algebras

$$H_0(Q) \rightarrow \mathbb{Q}[\text{Mult}(Q)^{\text{ap}}],$$

where the right side is equipped with the generic extension product and we send  $i$  to  $[i, i]$ . Indeed, one just needs to check that the expressions involved in the Serre relations give rise to the same multisegments. For example, if  $n > 2$ ,

$$i(i+1)i = [i, i+1] + [i, i] = ii(i+1).$$

The other relations can be checked similarly. This map is surjective by [28, §4], and since for  $\mathbf{d}$  a cuspidal support,  $H_0^{\mathbf{d}}(Q)$  has image in the multisegments with cuspidal support  $\mathbf{d}$ , the claim follows, because the dimensions agree.  $\square$

**Remark.** There exists a second map  $\mathfrak{m}_{\text{crys}} : \Omega \rightarrow \text{Mult}(Q)$  linked to  $\rho$ -derivatives and certain crystal bases of quantum groups; see for example [8] or [17]. Even though  $\mathfrak{m}_{\text{crys}}$  admits a similar, although slightly more involved, recursive description as  $\mathfrak{m}_{\text{gen}}$ , these two maps differ in general.

If  $w = i_1 \dots i_k \in \Omega$ , we let  $w^\vee = (-i_k) \dots (-i_1)$ . It follows from the definitions that  $(i + \mathfrak{m})^\vee = \mathfrak{m}^\vee + (-i)$  and hence

$$\mathfrak{m}_{\text{gen}}(w^\vee) = \mathfrak{m}_{\text{gen}}(w)^\vee.$$

**2.4.1. Degeneration order and Serre relations.** In this subsection we describe how, given two words  $v, w \in \Omega$ , one can decide whether  $[\mathfrak{m}_{\text{gen}}(v)] \preceq [\mathfrak{m}_{\text{gen}}(w)]$ . Elementary operations on  $\text{Mult}(Q)$  can be defined in a manner completely analogous to those on  $\text{Mult}_R$  and the first step in answering the above question is the following.

**Proposition 2.1** [33, Theorem 2.2; 13; 26, Theorem 3.12]. *The degeneration order and the order by elementary operations are equivalent, i.e.,*

$$[\lambda(\mathfrak{m})] \preceq [\lambda(\mathfrak{n})]$$

*if and only if  $\mathfrak{m}$  can be obtained by  $\mathfrak{n}$  via finitely many elementary operations.*

Let  $v, w \in \Omega$ . We write  $v \leq w$  if  $v$  can be obtained from  $w$  by applying a finite sequence of the following moves.

- (1)  $ij \mapsto ji$  if  $i$  and  $j$  are not neighbors in  $I$ .
- (2) If  $n > 2$ ,  $(i+1)i \mapsto i(i+1)$  and  $ii(i+1) \mapsto i(i+1)i$ .
- (3) If  $n = 2$ ,  $i(i+1)(i+1) \mapsto (i+1)i(i+1)$ .

The Serre relations are invariant under these moves.

The goal of this section is to prove the following proposition.

**Proposition 2.2.** *Let  $v, w \in \Omega$ . If  $v \geq w$  then  $\mathfrak{m}_{\text{gen}}(v) \leq \mathfrak{m}_{\text{gen}}(w)$ . If  $n = \infty$ , then  $\mathfrak{m}_{\text{gen}}(v) \leq \mathfrak{m}_{\text{gen}}(w)$  implies  $v \geq w$ .*

It is possible that there exists a more straightforward proof than the direct, purely combinatorial proof we offer, namely by relating the questions to properties of PBW bases as in the proof of Theorem 2.4.1. Note that the second implication in the proposition is wrong if  $n$  is not infinite. For example, if  $n = 3$ , we can set  $v = 002211102$  and  $w = 022110021$ . Then  $w$  is maximal with respect to the above order but

$$\begin{aligned} \mathfrak{m}_{\text{gen}}(v) &= [0, 2] + [0, 1] + [2, 3] + [2, 2] + [1, 1] \\ &> \mathfrak{m}_{\text{gen}}(w) = [0, 2] + [-1, 1] + [2, 3] + [1, 1]. \end{aligned}$$

*Proof of Proposition 2.2.* Assume first that  $v \leq w$ . Then it suffices by Lemma 2.4.1 to show that  $[i, i] + [i+1, i+1] = (i+1)+i \geq i+(i+1) = [i, i+1]$  and  $i+i+(i+1) = i+(i+1)+i$  if  $n > 2$ . The first is seen to be a simple elementary operation whereas the second is a Serre relation. The case  $n = 2$  can be checked analogously.

For the other direction (in the case  $n$  being infinite) write  $\mathfrak{n} = \mathfrak{m}_{\text{gen}}(v)$  and  $\mathfrak{m} = \mathfrak{m}_{\text{gen}}(w)$ ,  $\mathfrak{n} < \mathfrak{m}$ . Using the Serre relations, it suffices to find some words  $v' \leq w'$  with  $\mathfrak{n} = \mathfrak{m}_{\text{gen}}(v')$  and  $\mathfrak{m} = \mathfrak{m}_{\text{gen}}(w')$ . We will thus construct for any aperiodic  $\mathfrak{n} < \mathfrak{m}$  two such words  $v' \leq w'$  via induction on the length of  $\mathfrak{m}$ .

It clearly suffices to treat the case where  $\mathfrak{n}$  is obtained from  $\mathfrak{m}$  via one elementary operation  $[a, b] + [a', b'] \mapsto [a, b'] + [a', b]$ ,  $a \leq a' + 1 \leq b \leq b' + 1$  and  $\mathfrak{n} < \mathfrak{m}$  is minimal, i.e., there exists no  $\mathfrak{k}$  with  $\mathfrak{n} < \mathfrak{k} < \mathfrak{m}$ . This implies in particular the following. If there exists  $[c, d] \in \mathfrak{m}$  with  $a \leq c \leq a' \leq b \leq d \leq b'$ , then  $[c, d]$  is one of  $[a, b]$ ,  $[a', b']$ ,  $[a, b']$  or  $[a', b]$ . Indeed, if there would exist a  $[c, d]$  different from these four segments we could decompose the above elementary operation in the following way. Assume that  $a < c < a'$ ; the other case follows analogously. Then we can first apply  $[a', b'] + [c, d] \mapsto [a', d] + [c, b']$  to  $\mathfrak{m}$ , then  $[a, b] + [a', d] \mapsto [a', b] + [a, d]$  and finally  $[a, d] + [c, b'] \mapsto [a, b'] + [c, d]$  to obtain  $\mathfrak{n}$ , contradicting the minimality of the elementary operation.

Note that it makes no difference whether we prove the claim for  $\mathfrak{m}$  or  $\mathfrak{m}^\vee$ ; thus we can assume without loss of generality that  $b - a \geq b' - a'$ . We now let  $[c, d]$

be the longest segment in  $m$  with  $a \leq c$  such that  $[c + 1, d + 1]$  does not appear in  $m$ . Set  $m' = m - [c, d] + [c + 1, d]$  and by construction  $c + m' = m$ . If  $[c, d]$  is not the precise copy of  $[a', b']$  involved in the elementary operation, it follows straightforwardly that the elementary operation descends to an elementary operation on  $m'$  yielding a multisegment  $n' \preceq m'$  with  $c + n' = n$ . In this case the claim follows by the induction hypothesis. On the other hand, if  $[c, d] = [a', b']$ , it follows by construction and the assumption  $b - a \geq b' - a'$  that  $b - a = b' - a'$  and for all  $i \in \{0, \dots, a' - a\}$ ,  $[a + i, b + i]$  appears in  $m$ . By the minimality of the elementary operation this implies that  $[a', b'] = [a - 1, b - 1]$ . Moreover,  $[a, b]$  is a longest segment in  $m$ .

It follows that for  $m'' = m - [a, b] - [a', b'] + [a + 1, b] + [a' + 1, b']$  we have  $a + (a + 1) + m'' = n$  and  $(a + 1) + a + m'' = m$ , finishing the claim.  $\square$

Let  $w = i_1 \dots i_l \in \Omega$  and  $m \in \text{Mult}(Q)$ . We call  $w$  a *descendant* of  $m$  if it is obtained from  $m$  in the following, recursive, way. Write  $m = [a_1, b_1] + \dots + [a_k, b_k]$  and choose  $i \in \{1, \dots, k\}$ . Then choose a descendant  $w'$  of  $m' = [a_1, b_1] + \dots + [a_i + 1, b_i] + \dots + [a_k, b_k]$ . If  $m'$  is empty,  $w'$  is the empty word. Finally,  $w$  is a descendant of  $m$  if it is of the form  $w = a_i w'$  for some  $a_i$  and  $w'$  as above.

**Lemma 2.4.3.** *Let  $w \in \Omega$  and  $m \in \text{Mult}(Q)$ . Then  $w$  is a descendant of  $m$  if and only if  $m_{\text{gen}}(w) \preceq m$ .*

*Proof.* We argue by induction on the length of  $m$ , the case of length 1 being trivially true, and we use the notation employed above the lemma. First assume that  $w$  is a descendant of  $m$  and write  $w = a_i w'$ ,  $w'$  a descendant of  $m'$ , as above the lemma. By the induction hypothesis we have that  $m_{\text{gen}}(w') \preceq m'$  and hence  $m_{\text{gen}}(w) \preceq a_i + m'$ . It thus suffices to show that  $a_i + m' \preceq m$ , which follows quickly. If  $[a_i + 1, b_i]$  is the longest segment in  $m'$  starting in  $a_i + 1 \pmod n$ , we have in fact equality. Otherwise, let  $[a_i + 1, d]$  be the respective longest segment. Then  $a_i + m' = m - [a_i + 1, d] + [a_i, d]$ . Note that we can perform the elementary operation  $[a_i, b_i] + [a_i + 1, d] \mapsto [a_i, d] + [a_i + 1, b_i]$  on  $m$ , yielding  $a_i + m'$ , and thus proving the claim.

For the other direction, assume that  $m_{\text{gen}}(w) \preceq m$  and write  $w = cw'$ ,  $c \in I$ . Then it follows from [8, Lemma 6.3.3] that  $c \in \{a_1, \dots, a_k\}$  and  $m_{\text{gen}}(w') \preceq m'$ , where  $m'$  is of the form  $[a_1, b_1] + \dots + [a_j + 1, b_j] + \dots + [a_k, b_k]$  for some  $j$  with  $a_j = c \pmod n$ . The claim follows immediately.  $\square$

### 3. Whittaker models

We follow the setup of [2] (see [14] for the case of nonzero characteristic). Let  $\psi_R$  be an additive character  $\psi_R : F \rightarrow R$ . Moreover, we demand that  $\psi_{\overline{\mathbb{Q}}_\ell} \otimes_{\overline{\mathbb{Z}}_\ell} \overline{\mathbb{F}}_\ell = \psi_{\overline{\mathbb{F}}_\ell}$ . By abuse of notation, we will from now on write  $\psi = \psi_R$ . We extend  $\psi$  to  $N_n$  by

defining

$$\psi(u) = \psi \left( \sum_{i=1}^{n-1} u_{i,i+1} \right).$$

We define the space of Whittaker functions of  $G_n$  with respect to  $\psi$  as

$$\mathcal{W}(\psi) = \mathcal{W}_n(\psi) := \text{Ind}_{N_n}^{G_n} \psi = \left\{ W : G_n \rightarrow R \text{ locally constant} : \right. \\ \left. W(ug) = \psi(u)W(g) \text{ for all } u \in N_n, g \in G_n \right\},$$

on which  $G_n$  acts by right-translation. We call  $\pi \in \mathfrak{Rep}_n$  of Whittaker type if

$$\dim_R \text{Hom}_{G_n}(\pi, \mathcal{W}(\psi)) = 1,$$

in which case we denote the image of  $\pi$  in  $\mathcal{W}(\psi)$  by  $\mathcal{W}(\pi, \psi)$ . Thus,  $\mathcal{W}(\pi, \psi)$  is socle-irreducible and its unique irreducible subrepresentation appears with multiplicity 1 in it. Moreover, if

$$\mathcal{W}(\pi, \psi) \rightarrow \mathcal{W}(\pi', \psi)$$

is a nonzero map for a second representation  $\pi'$  of Whittaker type, it is injective. If  $\pi$  is on top of that irreducible we call it *generic*.

Let  $W$  be a Whittaker function. We define the map  $\widetilde{W}$  by

$$g \mapsto \widetilde{W}(g) := W(w_n(g^{-1})^t).$$

Finally, if  $\pi$  is of Whittaker type, we set

$$\mathcal{W}(\pi, \psi)^c = \{ \widetilde{W}(g) : W \in \mathcal{W}(\pi, \psi) \} = \mathcal{W}(\pi^c, \psi^{-1}).$$

We recall that if  $\pi_1$  and  $\pi_2$  are of Whittaker type, so is  $\pi_1 \times \pi_2$  (see [2; 3], and also [29, III.1.10]), and then  $\mathcal{W}(\pi_1 \times \pi_2, \psi) = \mathcal{W}(\mathcal{W}(\pi_1, \psi) \times \mathcal{W}(\pi_2, \psi), \psi)$ . We denote by  $\mathfrak{Rep}_{W, \psi, n}$  the set of finite-length subrepresentations of  $\mathcal{W}_n(\psi)$  which are of Whittaker type and we set

$$\mathfrak{Rep}_{W, \psi} := \bigcup_{n \in \mathbb{N}} \mathfrak{Rep}_{W, \psi, n}.$$

Then  $\mathfrak{Rep}_{W, \psi}$  can be equipped with an associative product

$$* : \mathfrak{Rep}_{W, \psi} \times \mathfrak{Rep}_{W, \psi} \rightarrow \mathfrak{Rep}_{W, \psi}, \quad (\pi, \pi') \mapsto \mathcal{W}(\pi \times \pi', \psi).$$

Associativity is an easy consequence of the uniqueness of the equality  $\mathcal{W}(\pi_1 \times \pi_2, \psi) = \mathcal{W}(\mathcal{W}(\pi_1, \psi) \times \mathcal{W}(\pi_2, \psi), \psi)$ .

If  $\Delta = [a, b]_\rho$  is a segment, the representation  $\langle \Delta \rangle$  is of Whittaker type if  $l(\Delta) < e(\rho)$ , by [23, Remark 8.14]. Every cuspidal representation is of Whittaker type.

**3.1. Derivatives.** We recall the four exact functors in the definition of the Bernstein–Zelevinsky derivatives; see [2; 3], and compare [29, III.1]. Recall also the groups  $P_n$  from Section 2 and write the unipotent subgroups as

$$\{U_{n-1,1} := \begin{pmatrix} 1 & x \\ & 1 \end{pmatrix} \in G_n : x \in M_{n-1,1}\}.$$

The four functors are

- (1)  $\Psi^+ : \mathfrak{R}\text{ep}_{n-1} \rightarrow \mathfrak{R}\text{ep}(P_n)$ , the extension by the trivial representation of  $U_{n-1,1}$  and twisted by  $|\cdot|^{-\frac{1}{2}}$ ;
- (2)  $\Psi^- : \mathfrak{R}\text{ep}(P_n) \rightarrow \mathfrak{R}\text{ep}_{n-1}$ , the  $U_{n-1,1}$ -coinvariants twisted by  $|\cdot|^{-\frac{1}{2}}$ ;
- (3)  $\Phi^+ = \text{ind}_{P_{n-1}U_{n-1,1}}^{P_n}(- \otimes \psi) : \mathfrak{R}\text{ep}(P_{n-1}) \rightarrow \mathfrak{R}\text{ep}(P_n)$ ;
- (4)  $\Phi^- : \mathfrak{R}\text{ep}(P_n) \rightarrow \mathfrak{R}\text{ep}(P_{n-1})$ , the  $(U_{n-1,1}, \psi)$ -coinvariants twisted by  $|\cdot|^{-\frac{1}{2}}$ .

Let  $\tau \in \mathfrak{R}\text{ep}(P_n)$ ,  $k \in \{1, \dots, n\}$  and set  $\tau_{(k)} := (\Phi^-)^{k-1}(\tau)$  and  $\tau^{(k)} := \Psi^-(\tau_{(k)})$ . For  $\pi \in \mathfrak{R}\text{ep}_n$  we set  $\pi_{(k)} := (\pi|_{P_n})_{(k)}$ ,  $\pi^{(k)} := (\pi|_{P_n})^{(k)}$  and  $\pi^{(0)} := \pi$ .

**3.2. Derivatives and Whittaker models.** We recall also the Kirillov model of a representation  $\pi$  of Whittaker type. It is defined as the  $P_n$ -representation given by

$$\mathcal{K}(\pi, \psi) := \{W|_{P_n} : W \in \mathcal{W}(\pi, \psi)\} \subseteq \text{Ind}_{N_n}^{P_n} \psi.$$

**Theorem 3.2.1** [9, 4.3; 20; 15]. *Let  $\pi \in \mathfrak{R}\text{ep}_n$  be of Whittaker type. The map  $W \mapsto W|_{P_n}$  is injective on  $\mathcal{W}(\pi, \psi)$ .*

We have the following description of  $(\Phi^-)^k$ .

**Lemma 3.2.1** [5, Proposition 1.3]. *Let  $\pi$  be of Whittaker type. Then we can identify  $(\Phi^-)^k(\mathcal{K}(\pi, \psi))$  with the space*

$$\left\{ p \mapsto |\det(p)|^{-\frac{k}{2}} W \begin{pmatrix} p & \\ & 1_k \end{pmatrix} : p \in P_{n-k}, W \in \mathcal{W}(\pi, \psi) \right\}.$$

The authors only prove the claim for  $R = \mathbb{C}$ , but the same method works for arbitrary base fields.

The description of the space  $\Psi^-(\Phi^-)^k(\mathcal{K}(\pi, \psi))$  is trickier. Let  $\tau$  be a subrepresentation of  $\mathcal{K}(\pi, \psi)^{(k)}$  with central character  $\chi$ . Let  $\sigma$  be the inverse image of  $\tau$  in  $\mathcal{K}(\pi, \psi)$ . For  $W \in \sigma$  and  $g \in G_{n-k}$  define the map

$$(1) \quad S(W)(g) = \lim_{z \rightarrow 0} |z|^{\frac{k-n}{2}} |\det(g)|^{\frac{k}{2}} \chi^{-1}(z) W \begin{pmatrix} zg & \\ & 1_k \end{pmatrix}.$$

Here  $z \in \mathbb{F}^\times$  is seen as an element of  $Z_{n-k}$  and the limit becomes stationary for  $z$  small enough.

**Proposition 3.1** [5, Proposition 1.7; 18, Corollary 2.1]. *The map  $S : \sigma \mapsto \mathcal{W}(\psi)$  induces the nonzero, injective map*

$$\bar{S} : \tau \mapsto \mathcal{W}(\psi).$$

Whittaker models are a useful place to look for integral structures, as the next theorem shows.

**Theorem 3.2.2** [31, Theorem 2]. *For  $\pi$  an integral representation of Whittaker type over  $\overline{\mathbb{Q}}_\ell$  of  $G_n$ , set*

$$\mathcal{W}^{\text{en}}(\pi, \psi) = \{W \in \mathcal{W}(\pi, \psi) : W(G_n) \subseteq \overline{\mathbb{Z}}_\ell\}.$$

*If  $\pi \in \mathfrak{Rep}$  is integral and of Whittaker type, then  $\mathcal{W}^{\text{en}}(\pi, \psi)$  is an integral structure of  $\mathcal{W}(\pi, \psi)$ .*

In this case we denote

$$\overline{\mathcal{W}(\pi, \psi)} := \mathcal{W}^{\text{en}}(\pi, \psi) \otimes_{\overline{\mathbb{Z}}_\ell} \overline{\mathbb{F}}_\ell.$$

**Lemma 3.2.2.** *Let  $\pi_1, \dots, \pi_k$  be integral representations of Whittaker type. Then the nonzero map*

$$\mathcal{W}(\pi_1, \psi) \times \dots \times \mathcal{W}(\pi_n, \psi) \rightarrow \mathcal{W}(\pi_1 \times \dots \times \pi_k, \psi)$$

*respects the integral structures and hence induces a nonzero map*

$$\overline{\mathcal{W}(\pi_1, \psi)} \times \dots \times \overline{\mathcal{W}(\pi_n, \psi)} \rightarrow \overline{\mathcal{W}(\pi_1 \times \dots \times \pi_k, \psi)}.$$

*Proof.* See for example the proof of [14, Theorem 2.26]. □

We recall some useful properties of Whittaker models.

**Proposition 3.2** [15, Proposition 3.7]. *Let  $\pi_1 \in \mathfrak{Rep}_{n_1}$ ,  $\pi_2 \in \mathfrak{Rep}_{n_2}$*

$$\mathcal{W}(\pi_2, \psi) \subseteq \mathcal{W}(\pi_1 \times \pi_2, \psi)^{(n_1)}.$$

**Theorem 3.2.3** [15, Theorem 3.10]. *Let  $n \geq 2$  and  $\tau$  be an a submodule of  $\text{Ind}_{N_n}^{P_n} \psi$ . If for  $k \in \{1, \dots, n-1\}$ ,  $\tau^{(k)}$  admits a central character, for any  $W_0 \in \mathcal{W}(\tau^{(k)}, \psi)$  and  $\phi \in C_c^\infty(\mathbb{F}^{n-k})$  there exists  $W \in \tau$  such that for all  $g \in G_{n-k}$*

$$W \begin{pmatrix} g & \\ & 1_k \end{pmatrix} = W_0(g) \phi(\epsilon_{n-k} g) |\det(g)|^{\frac{k-1}{2}}.$$

The proofs of these two results in [15] apply to  $R = \mathbb{C}$ , but the methods readily generalize to the more general settings presented here.

**3.3. Rankin–Selberg  $L$ -factors.** We now recall the construction of Rankin–Selberg  $L$ -factors as presented in [14]. For complex representations this goes back to the classical text [12]. Let  $\pi \in \mathfrak{Rep}_n$  and  $\pi' \in \mathfrak{Rep}_m$  be representations of Whittaker type.

**Definition.** Let  $W \in \mathcal{W}(\pi, \psi)$ ,  $W' \in \mathcal{W}(\pi', \psi^{-1})$  and  $k \in \mathbb{Z}$ .

(1) The case  $n = m$ : Let  $\phi \in C_c^\infty(\mathbb{F}^n)$  and define

$$c_k(W, W', \phi) := \int_{N_n \backslash G_n^k} W(g)W'(g)\phi(\epsilon_k g) \, d_l g,$$

$$I(X, W, W, \phi) = \sum_{k \in \mathbb{Z}} c_k(W, W', \phi)X^k \in R(X).$$

(2) The case  $n < m$ : Let  $j \in \{0, \dots, n - m - 1\}$  and define

$$c_k(W, W', j) := \int_{M_{j,m}} \int_{N_n \backslash G_n^k} W(g)W' \begin{pmatrix} g & & & \\ & x & & \\ & & 1_j & \\ & & & 1_{n-m-j} \end{pmatrix} d_l g \, d_l x,$$

$$I(X, W, W, j) = \sum_{k \in \mathbb{Z}} c_k(W, W', j)X^k \in R(X).$$

(3) The case  $n > m$ : Analogous to the case  $n < m$ .

Having defined this, we can now recall the definition of the  $L$ -factors.

**Theorem 3.3.1** [14, Theorem 3.5]. *Let  $\pi \in \mathfrak{Rep}_n$  and  $\pi' \in \mathfrak{Rep}_m$  be representations of Whittaker type.*

- (1) *The case  $n = m$ . The ideal spanned by  $I(X, W, W', \phi)$ , where we vary over  $W \in \mathcal{W}(\pi, \psi)$ ,  $W' \in \mathcal{W}(\pi', \psi^{-1})$  and  $\phi \in C_c^\infty(\mathbb{F}^n)$  is fractional and admits a generator  $L(X, \pi, \pi')$  with  $L(X, \pi, \pi')^{-1} \in R[X]$ , which is normalized by demanding  $L(0, \pi, \pi') = 1$ .*
- (2) *The case  $n \neq m$ . Fix  $j \in \{0, \dots, |n-m|-1\}$ . The ideal spanned by  $I(X, W, W', j)$ , where we vary over  $W \in \mathcal{W}(\pi, \psi)$  and  $W' \in \mathcal{W}(\pi', \psi^{-1})$  is fractional, independent of  $j$  and admits a generator  $L(X, \pi, \pi')$  with  $L(X, \pi, \pi')^{-1} \in R[X]$ , which is normalized by demanding  $L(0, \pi, \pi') = 1$ .*

In particular  $L(X, \pi, \pi') = L(X, \pi', \pi)$ . As usual, these  $L$ -factors satisfy a functional equation, giving rise to an  $\epsilon$ -factor. For  $\phi \in C_c^\infty(\mathbb{F}^n)$ , we denote by  $\hat{\phi}$  its Fourier transform with respect to the character  $\psi$ .

**Theorem 3.3.2** [14, Corollary 3.11; 14, Lemma 3.12]. *Let  $\pi \in \mathfrak{Rep}_n$ ,  $\pi' \in \mathfrak{Rep}_m$  be representations of Whittaker type with central characters  $c_\pi$  and  $c_{\pi'}$ , and let  $n \leq m$ . Let  $W \in \mathcal{W}(\pi, \psi)$ ,  $W' \in \mathcal{W}(\pi', \psi^{-1})$ ,  $\phi \in C_c^\infty(\mathbb{F}^n)$  and  $j \in \{0, \dots, m - n - 1\}$ . Then there exists  $\epsilon(X, \pi, \pi', \psi) \in R[X, X^{-1}]^\times$  such that the following holds.*

(1) *The case  $n = m$ .*

$$\frac{I(q^{-1}X^{-1}, \tilde{W}, \tilde{W}', \hat{\phi})}{L(q^{-1}, \pi^c, \pi'^c)} = c_{\pi'}(-1)^{n-1} \epsilon(X, \pi, \pi', \psi) \frac{I(X, W, W', \phi)}{L(X, \pi, \pi')}.$$

(2) *The case  $n < m$ .*

$$\frac{I(q^{-1}X^{-1}, \rho(w_{n,n-m})\tilde{W}, \tilde{W}', m-n-j-1)}{L(q^{-1}, \pi^c, \pi'^c)} = c_{\pi'}(-1)^{n-1} \epsilon(X, \pi, \pi', \psi) \frac{I(X, W, W', j)}{L(X, \pi, \pi')}.$$

The last local factor, the  $\gamma$ -factor, is finally defined as follows.

**Definition.** Let  $\pi, \pi' \in \mathfrak{R}\mathfrak{e}\mathfrak{p}$  be representations of Whittaker type. Then we define

$$\gamma(X, \pi, \pi', \psi) := \epsilon(X, \pi, \pi', \psi) \frac{L(q^{-1}X^{-1}, \pi^c, \pi'^c)}{L(X, \pi, \pi')} \in R(X).$$

We now recall the most important properties of these local factors.

**Lemma 3.3.1** [14, Theorem 3.13]. *Let  $\pi, \pi', \pi'' \in \mathfrak{R}\mathfrak{e}\mathfrak{p}$  be representations of Whittaker type and  $\tau$  a subrepresentation of  $\pi$  of Whittaker type. Then*

$$\gamma(X, \pi, \pi', \psi) = \gamma(X, \tau, \pi', \psi) \quad \text{and} \quad L(X, \tau, \pi')^{-1} | L(X, \pi, \pi')^{-1}.$$

Moreover, we have the so-called inductivity relation

$$\gamma(X, \pi \times \pi'', \pi', \psi) = \gamma(X, \pi, \pi', \psi) \gamma(X, \pi'', \pi', \psi).$$

Finally, let us remark how these factors interact with respect to reduction mod  $\ell$ .

**Lemma 3.3.2** [14, Theorem 4.1, §4.1]. *Let  $\pi, \pi' \in \mathfrak{R}\mathfrak{e}\mathfrak{p}$  be two integral representations of Whittaker type over  $\overline{\mathbb{Q}}_\ell$ . Then  $L(X, \pi, \pi')$ ,  $\epsilon(X, \pi, \pi', \psi)$ ,  $\gamma(X, \pi, \pi', \psi)$  lie in  $\overline{\mathbb{Z}}_\ell(X)$  and*

$$\gamma(X, \overline{\mathcal{W}(\pi, \psi)}, \overline{\mathcal{W}(\pi', \psi)}, \psi) = r_\ell(\gamma(X, \pi, \pi', \psi)), \\ L(X, \overline{\mathcal{W}(\pi, \psi)}, \overline{\mathcal{W}(\pi', \psi)})^{-1} | r_\ell(L(X, \pi, \pi')^{-1}).$$

**Lemma 3.3.3.** *Let  $\pi \in \mathfrak{R}\mathfrak{e}\mathfrak{p}_n, \pi' \in \mathfrak{R}\mathfrak{e}\mathfrak{p}_m$  be two representation of Whittaker type and  $k \in \{0, \dots, n\}$ . Let  $\tau$  be a subrepresentation of  $\pi^{(k)}$  admitting a central character. Then*

$$L(X, \tau, \pi')^{-1} | L(X, \pi, \pi')^{-1}.$$

*Proof.* As explained in the proof of [14, Lemma 4.6, Proposition 4.7] (see also [12, Lemma 9.2]), the claim holds true if one shows the following. For any  $W_0 \in \mathcal{W}(\tau, \psi)$  and  $\phi \in C_c^\infty(\mathbb{F}^{n-k})$  there exists  $W \in \mathcal{W}(\pi, \psi)$  such that for all  $g \in G_{n-k}$

$$W \begin{pmatrix} g & & & \\ & 1_{n-k} & & \\ & & & \end{pmatrix} = W_0(g) \phi(\epsilon_{n-k} g) |\det(g)|^{\frac{k-1}{2}}.$$

But by assumption on  $\tau^{(k)}$ ,  $\mathcal{W}(\tau^{(k)}, \psi)$  admits a central character; hence this follows from Theorem 3.2.3.  $\square$

We finish by recalling the  $L$ -factors of  $C$ -parameters of [16]. For the moment we slightly extend this notation to aperiodic multisegments and to avoid confusion we will denote our parameters by  $C'$ .

**Definition.** Let  $\rho, \rho' \in \mathfrak{C}$ . We set

$$L(X, C'(\rho), C'(\rho')) := \begin{cases} (1 - (\chi(\varpi_F)X)^{f(\rho)})^{-1} & \text{if } \rho' \cong \chi\rho^\vee, \rho \in \mathfrak{C}^\square, \chi \text{ an unramified character,} \\ 1 & \text{otherwise.} \end{cases}$$

Let  $\Delta = [a, b]_\rho, \Delta' = [a', b']_{\rho'}$  be two segments. Then we set

$$L(X, C'(\Delta), C'(\Delta')) := \begin{cases} \prod_{i=a}^b L(X, C'(\rho|-^i), C'(\rho'|-^b)) & \text{if } l(\Delta) \leq l(\Delta'), \\ \prod_{i=a'}^{b'} L(X, C'(\rho|-^b), C'(\rho'|-^i)) & \text{if } l(\Delta) \geq l(\Delta'). \end{cases}$$

Finally, if  $\mathfrak{m}, \mathfrak{n} \in \text{Mult}_R$  of the form  $\mathfrak{m} = \Delta_1 + \dots + \Delta_k, \mathfrak{n} = \Delta'_1 + \dots + \Delta'_l$  we set

$$L(X, C'(\mathfrak{m}), C'(\mathfrak{n})) := \prod_{\substack{1 \leq i \leq k \\ 1 \leq j \leq l}} L(X, C'(\Delta_i), C'(\Delta'_j)).$$

**3.4. Associative products and the map  $\mathcal{S}_{\text{gen}, \psi}$ .** For the rest of this section, we fix an additive character  $\psi$  of  $F$ . We also fix  $\rho \in \mathfrak{C}^\square$  and let  $Q$  be the quiver of Section 2.4 with  $n = o(\rho) > 1$  and set of indices  $I$ . Fix a cuspidal support

$$\mathbf{d} = \sum_{i \in I} d_i [\rho|-^i]$$

and note that  $d_i = 0$  for all but finitely many  $i \in I$  and hence  $\mathbf{d}$  gives rise to a dimension vector denoted by the same letter  $\mathbf{d}$  for  $Q$ . We let  $\text{Mult}_R(\rho)_{\mathbf{d}}$  be the set of multisegments with support  $\mathbf{d}$ . We then recall the natural bijection

$$\text{Mult}_R(\rho)_{\mathbf{d}} \leftrightarrow \{G_{\mathbf{d}}\text{-orbits in } E_{\mathbf{d}}\},$$

which respects the orders on each side, by Proposition 2.1. This bijection allows us to import the generic extension product to  $\text{Mult}_R(\rho)$ , which by abuse of notation we also denote by  $*$ . This product has the following representation-theoretic interpretation. By Theorem 2.4.1, there exists for  $\mathfrak{m} \in \text{Mult}_R^{\text{ap}}(\rho)$  representations  $\rho_1, \dots, \rho_k$  in  $\mathbb{Z}[\rho]$  such that  $\mathfrak{m} = \rho_1 * \dots * \rho_k$ . We then set

$$\mathcal{S}_{\text{gen}, \psi}(\mathfrak{m}) := \mathcal{W}(\rho_1 \times \dots \times \rho_k, \psi).$$

This representation is independent of our choices; to prove this, we first note the following analogue of the degenerate Serre relations.

**Lemma 3.4.1.** *Let  $\rho' \in \mathbb{Z}[\rho]$  be such that  $\rho' \not\cong \rho|\cdot|^{\pm 1}$ . Then*

$$\mathcal{W}(\rho \times \rho', \psi) = \mathcal{W}(\rho' \times \rho, \psi).$$

If  $o(\rho) > 2$ ,

$$\begin{aligned} \mathcal{W}(\rho \times \rho \times \rho|\cdot|, \psi) &= \mathcal{W}(\rho \times \rho|\cdot| \times \rho, \psi), \\ \mathcal{W}(\rho \times \rho|\cdot| \times \rho|\cdot|, \psi) &= \mathcal{W}(\rho|\cdot| \times \rho \times \rho|\cdot|, \psi). \end{aligned}$$

If  $o(\rho) = 2$ ,

$$\mathcal{W}(\rho \times \rho|\cdot| \times \rho \times \rho, \psi) = \mathcal{W}(\rho \times \rho \times \rho|\cdot| \times \rho, \psi).$$

*Proof.* Under the hypothesis,  $\rho \times \rho'$  is irreducible, and

$$\mathcal{W}(\rho \times \rho', \psi) \cong \rho \times \rho' \cong \rho' \times \rho \cong \mathcal{W}(\rho' \times \rho, \psi).$$

Hence they are equal. We only show the remaining claims for  $o(\rho) > 2$ ; the case  $o(\rho) = 2$  follows analogously. We start with the equality  $\mathcal{W}(\rho \times \rho \times \rho|\cdot|, \psi) = \mathcal{W}(\rho \times \rho|\cdot| \times \rho, \psi)$ . By [23, Proposition 7.17],

$$\rho \times \rho \times \rho|\cdot| \twoheadrightarrow \rho \times \langle [0, 1]_\rho \rangle,$$

which is irreducible by Lemma 2.2.2 and of Whittaker type. Thus the uniqueness of the Whittaker model forces

$$\mathcal{W}(\rho \times \rho \times \rho|\cdot|, \psi) \cong \rho \times \langle [0, 1]_\rho \rangle.$$

Similarly,

$$\mathcal{W}(\rho \times \rho|\cdot| \times \rho, \psi) \cong \langle [0, 1]_\rho \rangle \times \rho.$$

By the commutativity of parabolic induction on the Grothendieck group we have  $\langle [0, 1]_\rho \rangle \times \rho \cong \rho \times \langle [0, 1]_\rho \rangle$ ; the claim follows. The equality  $\mathcal{W}(\rho \times \rho|\cdot| \times \rho|\cdot|, \psi) = \mathcal{W}(\rho|\cdot| \times \rho \times \rho|\cdot|, \psi)$  follows by an analogous argument.  $\square$

From the lemma and Theorem 2.4.1 we obtain:

**Corollary 3.4.1.** *The representation  $\mathcal{S}_{\text{gen}, \psi}(\mathfrak{m})$  is independent of the sequence  $\rho_1, \dots, \rho_k$  with  $\rho_1 * \dots * \rho_k = \mathfrak{m}$ . Moreover, the map  $\mathcal{S}_{\text{gen}, \psi} : \text{Mult}_R(\rho)^{\text{ap}} \rightarrow \mathfrak{Rep}_W$  respects the respective products, i.e., for  $\mathfrak{m}_1, \mathfrak{m}_2 \in \text{Mult}_R(\rho)^{\text{ap}}$ , we have*

$$\mathcal{W}(\mathcal{S}_{\text{gen}, \psi}(\mathfrak{m}_1) \times \mathcal{S}_{\text{gen}, \psi}(\mathfrak{m}_2), \psi) = \mathcal{S}_{\text{gen}, \psi}(\mathfrak{m}_1 * \mathfrak{m}_2)$$

Next we define the map  $\mathcal{S}_{\text{gen}, \psi}^{\cup}$  by setting for  $\mathfrak{m} \in \text{Mult}_R(\rho)$  not necessarily aperiodic

$$\mathcal{S}_{\text{gen}, \psi}^{\cup}(\mathfrak{m}) := \bigcup_{\substack{\mathfrak{n} \leq \mathfrak{m}, \\ \mathfrak{n} \text{ aperiodic}}} \mathcal{S}_{\text{gen}, \psi}(\mathfrak{n}).$$

Here the union is taken in the space of Whittaker functions.

**Proposition 3.3.** *Let  $\mathfrak{n}, \mathfrak{m} \in \text{Mult}_R(\rho)$  with  $\mathfrak{n} \leq \mathfrak{m}$ . Then*

$$\mathcal{S}_{\text{gen}, \psi}^{\cup}(\mathfrak{n}) \subseteq \mathcal{S}_{\text{gen}, \psi}^{\cup}(\mathfrak{m}).$$

*Moreover, let  $\mathfrak{m}' \in \text{Mult}_R(\rho)$  such that  $\mathfrak{m}' * \rho = \mathfrak{m}$ . Then*

$$\mathcal{W}(\mathcal{S}_{\text{gen}, \psi}^{\cup}(\mathfrak{m}') \times \rho, \psi) \subseteq \mathcal{S}_{\text{gen}, \psi}^{\cup}(\mathfrak{m}).$$

*Similarly, if  $\mathfrak{m}' \in \text{Mult}_R(\rho)$  such that  $\rho * \mathfrak{m}' = \mathfrak{m}$ , then*

$$\mathcal{W}(\rho \times \mathcal{S}_{\text{gen}, \psi}^{\cup}(\mathfrak{m}'), \psi) \subseteq \mathcal{S}_{\text{gen}, \psi}^{\cup}(\mathfrak{m}).$$

*Proof.* The first claim follows from the definition. For the second claim we argue as follows. We only treat the case  $\mathfrak{m}' * \rho = \mathfrak{m}$ , as the other one follows similarly. Then for  $\mathfrak{n}' \leq \mathfrak{m}'$  we have by Lemma 2.4.1 that  $\mathfrak{n}' * \rho \leq \mathfrak{m}$  and hence  $\mathcal{W}(\mathcal{S}_{\text{gen}, \psi}^{\cup}(\mathfrak{m}') \times \rho, \psi) \subseteq \mathcal{S}_{\text{gen}, \psi}^{\cup}(\mathfrak{m})$  by the first claim.  $\square$

Finally, for  $\mathfrak{m} \in \text{Mult}_{R, \square}$ , write  $\mathfrak{m} = \mathfrak{m}_1 + \cdots + \mathfrak{m}_k$  with  $\mathfrak{m}_i \in \text{Mult}_{R, \square}(\rho_i)$ , where the  $\rho_i \in \mathcal{C}^{\square}$  are in pairwise different cuspidal lines. Then we set

$$\mathcal{S}_{\text{gen}, \psi}(\mathfrak{m}) := \mathcal{W}(\mathcal{S}_{\text{gen}, \psi}(\mathfrak{m}_1) \times \cdots \times \mathcal{S}_{\text{gen}, \psi}(\mathfrak{m}_k), \psi), \quad \mathcal{S}_{\text{gen}, \psi}^{\cup}(\mathfrak{m}) := \bigcup_{\substack{\mathfrak{n} \leq \mathfrak{m}, \\ \mathfrak{n} \text{ aperiodic}}} \mathcal{S}_{\text{gen}, \psi}(\mathfrak{n}),$$

and if  $\mathfrak{m}' = \mathfrak{m}'_1 + \cdots + \mathfrak{m}'_k \in \text{Mult}_{R, \square}$ ,  $\mathfrak{m}'_i \in \text{Mult}_R(\rho_i)$  is a second multisegment, we define

$$\mathfrak{m} * \mathfrak{m}' := \mathfrak{m}_1 * \mathfrak{m}'_1 + \cdots + \mathfrak{m}_k * \mathfrak{m}'_k,$$

extending the product  $*$  to  $*$ :  $\text{Mult}_{R, \square} \times \text{Mult}_{R, \square} \rightarrow \text{Mult}_{R, \square}$ .

**Lemma 3.4.2.** *Let  $\mathfrak{m} = \mathfrak{m}_1 + \cdots + \mathfrak{m}_k \in \text{Mult}_{R, \square}^{\text{ap}}$  as above. Then*

$$\mathcal{W}(\mathcal{S}_{\text{gen}, \psi}(\mathfrak{m}), \psi) \cong \mathcal{S}_{\text{gen}, \psi}(\mathfrak{m}_1) \times \cdots \times \mathcal{S}_{\text{gen}, \psi}(\mathfrak{m}_k)$$

*and hence for  $\mathfrak{m}' \in \text{Mult}_{R, \square}^{\text{ap}}$  a second multisegment we have*

$$\mathcal{W}(\mathcal{S}_{\text{gen}, \psi}(\mathfrak{m}) \times \mathcal{S}_{\text{gen}, \psi}(\mathfrak{m}'), \psi) = \mathcal{S}_{\text{gen}, \psi}(\mathfrak{m} * \mathfrak{m}').$$

*An analogous claim holds for  $\mathcal{S}_{\text{gen}, \psi}^{\cup}$  and if  $\mathfrak{n} \leq \mathfrak{m}$  then  $\mathcal{S}_{\text{gen}, \psi}^{\cup}(\mathfrak{n}) \subseteq \mathcal{S}_{\text{gen}, \psi}^{\cup}(\mathfrak{m})$ .*

*Proof.* There exists a nonzero map

$$\mathcal{S}_{\text{gen}, \psi}(\mathfrak{m}_1) \times \cdots \times \mathcal{S}_{\text{gen}, \psi}(\mathfrak{m}_k) \rightarrow \mathcal{W}(\mathcal{S}_{\text{gen}, \psi}(\mathfrak{m}), \psi).$$

Let  $\pi$  be the unique irreducible subrepresentation of the latter representation. It is then enough to show that  $\pi$  is the unique subrepresentation of the former representation. To see this note that by Theorem 2.2.1 and Lemma 2.2.1, the representation  $\pi_1 \times \cdots \times \pi_k$  is irreducible, where  $\pi_i$  is the unique irreducible subrepresentation of

$\mathcal{S}_{\text{gen},\psi}(\mathfrak{m}_i)$ . Since  $\pi_1 \times \cdots \times \pi_k$  is generic, it has to be isomorphic to  $\pi$ . Thus  $\pi$  is a subrepresentation of

$$\mathcal{S}_{\text{gen},\psi}(\mathfrak{m}_1) \times \cdots \times \mathcal{S}_{\text{gen},\psi}(\mathfrak{m}_k).$$

For the second point, note that if  $\mathfrak{m}$  and  $\mathfrak{m}'$  are in different cuspidal lines, then we just proved that  $\mathcal{W}(\mathcal{S}_{\text{gen},\psi}(\mathfrak{m}) \times \mathcal{S}_{\text{gen},\psi}(\mathfrak{m}'), \psi) = \mathcal{W}(\mathcal{S}_{\text{gen},\psi}(\mathfrak{m} + \mathfrak{m}'), \psi) = \mathcal{W}(\mathcal{S}_{\text{gen},\psi}(\mathfrak{m}' + \mathfrak{m}), \psi) = \mathcal{W}(\mathcal{S}_{\text{gen},\psi}(\mathfrak{m}') \times \mathcal{S}_{\text{gen},\psi}(\mathfrak{m}), \psi)$ . From this the second claim follows. Finally, if  $\mathfrak{n} \leq \mathfrak{m}$  with  $\mathfrak{m}_i, \mathfrak{n}_i \in \text{Mult}_R(\rho_i)$  as above, we have by definition  $\mathfrak{n} \leq \mathfrak{m}$  if and only if  $\mathfrak{n}_i \leq \mathfrak{m}_i$  for all  $i$ . From the first point and Proposition 3.3 it then follows that there exists an injection

$$\mathcal{S}_{\text{gen},\psi}^{\cup}(\mathfrak{n}) \hookrightarrow \mathcal{S}_{\text{gen},\psi}^{\cup}(\mathfrak{m}),$$

which by the uniqueness of the Whittaker model must be an inclusion.

The claim for  $\mathcal{S}_{\text{gen},\psi}^{\cup}$  follows the exact same pattern.  $\square$

#### 4. Standard modules

For the rest of this section, we fix an additive character  $\psi$ .

**Definition.** Two maps  $\mathcal{T}_{\psi}, \mathcal{T}_{\psi^{-1}} : \text{Mult}_R \rightarrow \mathfrak{Rep}$  are called of Whittaker type if they satisfy the following.

- (1) For each  $\mathfrak{m} \in \text{Mult}_R$ ,  $\mathcal{T}_{\psi}(\mathfrak{m})$  is a representation of  $G_{\deg(\mathfrak{m})}$  and is contained in  $\mathfrak{Rep}_{W,\psi}$ .
- (2)  $\mathcal{T}_{\psi}(\mathfrak{m})^{\epsilon} = \mathcal{T}_{\psi^{-1}}(\mathfrak{m}^{\vee})$ .
- (3) If  $\mathfrak{n} \leq \mathfrak{m}$  then  $\mathcal{T}_{\psi}(\mathfrak{n}) \subseteq \mathcal{T}_{\psi}(\mathfrak{m})$ .

Since by (2),  $\mathcal{T}_{\psi^{-1}}$  is determined by  $\mathcal{T}_{\psi}$ , we will usually omit  $\mathcal{T}_{\psi^{-1}}$ .

The map  $\mathcal{T}_{\psi}$  is called *L-standard* if it is of Whittaker type and for all  $\mathfrak{n}, \mathfrak{m} \in \text{Mult}_R^{\text{ap}}$

$$L(X, \mathcal{T}_{\psi}(\mathfrak{n}), \mathcal{T}_{\psi^{-1}}(\mathfrak{m})) = L(X, C'(\mathfrak{n}), C'(\mathfrak{m})).$$

Finally, an *L-standard* map  $\mathcal{T}$  is called *standard* if it moreover satisfies the following.

- (1) For each  $\mathfrak{m} \in \text{Mult}_R^{\text{ap}}$  there exists a nonzero map  $\mathcal{T}_{\psi}(\mathfrak{m}) \twoheadrightarrow \langle \mathfrak{m} \rangle$ .
- (2) For each  $\mathfrak{m} \in \text{Mult}_R^{\text{ap}}$ ,  $\dim_R \text{Hom}_{\Delta G_{\deg(\mathfrak{m})}}(\mathcal{T}_{\psi}(\mathfrak{m}) \otimes \mathcal{T}_{\psi^{-1}}(\mathfrak{m}^{\vee}), R) = 1$ .
- (3) The multiplicity of  $\langle \mathfrak{m} \rangle$  in  $\mathcal{T}_{\psi}(\mathfrak{m})$  is 1.

Replacing  $\text{Mult}_R$  by  $\text{Mult}_{R,\square}$  in the above definition, we obtain the notions of  $\square$ -Whittaker type,  $\square$ -L-standard and  $\square$ -standard.

It is easy to see that if  $\mathcal{T}_{\psi}$  is standard and  $\ell \neq 2$ , the nonzero map  $\mathcal{T}_{\psi}(\mathfrak{m}) \twoheadrightarrow \mathcal{T}_{\psi^{-1}}(\mathfrak{m}^{\vee})^{\vee}$ , which is unique up to a scalar, factors through  $\langle \mathfrak{m} \rangle$ . Indeed, since  $\ell \neq 2$ ,

we have that  $\langle \mathfrak{m} \rangle^c \cong \langle \mathfrak{m} \rangle^\vee$  is a quotient of  $\mathcal{T}_{\psi^{-1}}(\mathfrak{m}^\vee)$ , and since Aubert duality commutes with taking duals, we have a nonzero map

$$\mathcal{T}_\psi(\mathfrak{m}) \twoheadrightarrow \langle \mathfrak{m} \rangle \hookrightarrow \mathcal{T}_{\psi^{-1}}(\mathfrak{m}^\vee)^\vee.$$

For  $\mathcal{T}$  a standard map, the representations  $\mathcal{T}_{\psi^{-1}}(\mathfrak{m})$  with  $\mathfrak{m}$  aperiodic are called *standard modules*.

**4.1. Standard modules over  $\overline{\mathbb{Q}}_\ell$ .** For this section we set  $R = \overline{\mathbb{Q}}_\ell$ . For an arranged form  $(\Delta_1, \dots, \Delta_k)$  of a multisegment  $\mathfrak{m}$  we denote

$$T_\psi(\mathfrak{m}) := \mathcal{W}(\langle \Delta_1 \rangle) \times \cdots \times \mathcal{W}(\langle \Delta_k \rangle, \psi),$$

whose isomorphism type is independent of the arranged order; see, for example, [32, Theorem 6.1]. If  $\mathfrak{m}$  is moreover integral we write

$$T_\psi^{\text{en}}(\mathfrak{m}) := \mathcal{W}^{\text{en}}(\langle \Delta_1 \rangle, \psi) \times \cdots \times \mathcal{W}^{\text{en}}(\langle \Delta_k \rangle, \psi).$$

**Theorem 4.1.1.** *The map  $\mathcal{S}_\psi : \text{Mult}_{\overline{\mathbb{Q}}_\ell} \rightarrow \mathfrak{Rep}$  given by  $\mathfrak{m} \mapsto \mathcal{W}(T(\mathfrak{m}), \psi)$  is standard.*

*Proof.* For the computation of  $L$ -factors see [12, Theorem 8.2]. The other properties are proved in [32, §6, §7, §9]. □

We let  $\mathcal{S}_\psi(\mathfrak{m})^{\text{en}}$  be the subspace of  $\overline{\mathbb{Z}}_\ell$ -valued function in  $\mathcal{S}_\psi(\mathfrak{m})$ .

**Lemma 4.1.1.** *For all  $\mathfrak{m} \in \text{Mult}_{\overline{\mathbb{Q}}_\ell}$ ,  $\mathcal{S}_{\text{gen}, \psi}(\mathfrak{m}) = \mathcal{S}_\psi(\mathfrak{m})$ .*

*Proof.* We argue by induction on  $\text{deg}(\mathfrak{m})$ . We first treat the case where  $\mathfrak{m}$  is one segment  $[0, b]_\rho$ . Then by [32, Proposition 9.5] and Frobenius reciprocity we have  $\rho \times \langle [1, b]_\rho \rangle \twoheadrightarrow \langle [0, b]_\rho \rangle$ , which is a generic representation by the uniqueness of the Whittaker model. Thus  $\mathcal{W}(\rho \times \langle [1, b]_\rho \rangle, \psi) = \mathcal{W}(\langle [0, b]_\rho \rangle, \psi)$ . By induction on  $b$  the left-hand side equals  $\mathcal{S}_{\text{gen}, \psi}([0, b]_\rho)$  by Proposition 3.3.

We come to the general case. Write  $\mathfrak{m} = \rho * \mathfrak{m}'$  as follows. Let  $[0, b]_\rho$ ,  $\rho \in \mathfrak{C}$ , be a longest segment in  $\mathfrak{m}$  such all segments  $\Delta$  in  $\mathfrak{m}$  with  $a_\rho(\Delta) = 1$  satisfy  $l(\Delta) < b$ . Taking  $\mathfrak{m}' = \mathfrak{m} - [0, b] + [1, b]_\rho$  we obtain  $\rho * \mathfrak{m}' = \mathfrak{m}$ . One can choose an arranged form  $(\Delta_1, \dots, \Delta_k)$  of  $\mathfrak{m}'$  such that for  $j < i$ , where  $i$  is the minimal  $i$  such that  $\Delta_i = [1, b]_\rho$ , either  $\Delta_j$  has cuspidal support not intersecting  $\mathbb{Z}[\rho]$  or  $a_\rho(\Delta_j)$  is not equal to 1. In any case,  $[0, 0] + \Delta_j$  is unlinked for  $j < i$ . We recall that by the uniqueness of the Whittaker model, we have for three representations  $\pi_1, \pi_2, \pi_3$  of Whittaker type

$$\mathcal{W}(\pi_1 \times \pi_2 \times \pi_3, \psi) = \mathcal{W}(\pi_1 \times \mathcal{W}(\pi_2, \psi) \times \pi_3, \psi).$$

Thus by the case of a single segment

$$\begin{aligned} \mathcal{W}(\rho \times \langle \Delta_1 \rangle \times \cdots \times \langle \Delta_k \rangle, \psi) &= \mathcal{W}(\langle \Delta_1 \rangle \times \cdots \times \rho \times \langle \Delta_i \rangle \times \cdots \times \langle \Delta_k \rangle, \psi) \\ &= \mathcal{W}(\langle \Delta_1 \rangle \times \cdots \times \mathcal{W}(\rho \times \langle \Delta_i \rangle, \psi) \times \cdots \times \langle \Delta_k \rangle, \psi) \\ &= \mathcal{W}(\langle \Delta_1 \rangle \times \cdots \times \langle {}^+ \Delta_i \rangle \times \cdots \times \langle \Delta_k \rangle, \psi). \end{aligned}$$

It is easy to see that  $(\Delta_1, \dots, {}^+ \Delta_i, \dots, \Delta_k)$  is an arranged form of  $\mathfrak{m}$ ; hence the right-hand side is equal to  $\mathcal{S}_\psi(\mathfrak{m})$ . By induction and Proposition 3.3 the left-hand side is equal to  $\mathcal{S}_{\text{gen}, \psi}(\mathfrak{m})$ .  $\square$

**4.2. Standard modules over  $\overline{\mathbb{F}}_\ell$ .** We come to our definition of standard modules over  $\overline{\mathbb{F}}_\ell$ . We saw in the introduction that if  $\tilde{\mathfrak{m}}$  is a lift of  $\mathfrak{m} \in \text{Mult}_{\overline{\mathbb{F}}_\ell}$  (Section 2.3), Theorem 3.2.2 says that  $\mathcal{S}_\psi(\tilde{\mathfrak{m}})$  can be equipped with a natural integral structure, whose reduction mod  $\ell$  is again of Whittaker type and is denoted by  $\overline{\mathcal{S}_\psi(\tilde{\mathfrak{m}})}$ . For  $\mathfrak{m} \in \text{Mult}_{\overline{\mathbb{F}}_\ell}$  we then defined the intersection

$$\mathcal{S}_\psi(\mathfrak{m}) := \bigcap_{\tilde{\mathfrak{m}} \text{ lift of } \mathfrak{m}} \overline{\mathcal{S}_\psi(\tilde{\mathfrak{m}})}$$

in the space of Whittaker functions. The representation is nonzero if  $\mathfrak{m} \in \text{Mult}_{R, \square}$ : indeed, each  $\overline{\mathcal{S}_\psi(\tilde{\mathfrak{m}})}$  contains with multiplicity one and as a unique subrepresentation the degenerate representation  $Z(\mathfrak{s})$ , where  $\mathfrak{s} = \text{cusp}(\mathfrak{m})$ . Then  $Z(\mathfrak{s})$  is a subrepresentation of  $\mathcal{S}_\psi(\mathfrak{m})$  since  $\dim_R \text{Hom}_{G_n}(Z(\mathfrak{s}), \mathcal{W}(\psi)) = 1$ .

**Definition.** Let  $\mathcal{T}_\psi$  a map of Whittaker type. We call  $\mathcal{T}$  *extending* if for all  $\mathfrak{m} \in \text{Mult}_{R, \square}$  and  $\rho \in \mathfrak{C}^\square$

$$\mathcal{W}(\rho \times \mathcal{T}_\psi(\mathfrak{m}), \psi) \subseteq \mathcal{T}_\psi(\rho * \mathfrak{m}), \quad \mathcal{W}(\mathcal{T}_\psi(\mathfrak{m}) \times \rho, \psi) \subseteq \mathcal{T}_\psi(\mathfrak{m} * \rho)$$

**Lemma 4.2.1.** *The maps  $\mathcal{S}_\psi$  and  $\mathcal{S}_{\text{gen}, \psi}^\cup$  are extending.*

*Proof.* For  $\mathcal{S}_{\text{gen}, \psi}^\cup$  the claim is a consequence of Proposition 3.3. The claim for  $\mathcal{S}_\psi$  in the case  $R = \overline{\mathbb{F}}_\ell$  follows by noting that on the one hand, for every lift  $\tilde{\mathfrak{m}}'$  of  $\rho * \mathfrak{m}$ , one can find lifts  $\tilde{\rho}$  and  $\tilde{\mathfrak{m}}$  of  $\rho$  and  $\mathfrak{m}$  such that  $\tilde{\rho} * \tilde{\mathfrak{m}} = \tilde{\mathfrak{m}}'$  and vice versa. By Lemma 3.2.2 the surjective, and hence nonzero, map  $\mathcal{W}(\tilde{\rho}, \psi) \times \mathcal{S}_\psi(\tilde{\mathfrak{m}}) \rightarrow \mathcal{W}(\rho \times \mathcal{S}_\psi(\tilde{\mathfrak{m}}), \psi) = \mathcal{W}(\mathcal{S}_\psi(\tilde{\rho} * \tilde{\mathfrak{m}}), \psi)$  reduces to a nonzero map

$$\overline{\mathcal{W}(\tilde{\rho}, \psi)} \times \overline{\mathcal{S}_\psi(\tilde{\mathfrak{m}})} \rightarrow \overline{\mathcal{W}(\mathcal{S}_\psi(\tilde{\rho} * \tilde{\mathfrak{m}}), \psi)}.$$

Taking the intersection in the space of Whittaker models we obtain a nonzero map

$$\rho \times \mathcal{S}_\psi(\mathfrak{m}) \rightarrow \mathcal{S}_\psi(\rho * \mathfrak{m}).$$

To see that the map on the intersection is nonzero, it suffices to note that the map

$$\overline{\mathcal{W}(\tilde{\rho}, \psi)} \times \overline{\mathcal{S}_\psi(\tilde{\mathfrak{m}})} \rightarrow \overline{\mathcal{W}(\mathcal{S}_\psi(\tilde{\rho} * \tilde{\mathfrak{m}}), \psi)}$$

does not vanish on the subrepresentation  $\overline{\mathcal{W}(\tilde{\rho}, \psi)} \times Z(\text{cusp}(\mathfrak{m}))$ , where  $Z(\text{cusp}(\mathfrak{m}))$  is the unique irreducible subrepresentation of  $\mathcal{S}_\psi(\tilde{\mathfrak{m}})$  that is independent of the lift  $\tilde{\mathfrak{m}}$ .

By the uniqueness of the Whittaker model we thus have that  $\mathcal{W}(\rho \times \mathcal{S}_\psi(\tilde{\mathfrak{m}}), \psi) \subseteq \mathcal{S}_\psi(\rho \times \mathfrak{m})$ . The other inclusion follows by the same argument.  $\square$

**Corollary 4.2.1.** *The map  $\mathcal{S}_{\text{gen}, \psi}^{\cup} : \text{Mult}_{\mathbb{F}_\ell, \square} \rightarrow \mathfrak{Rep}$  is of  $\square$ -Whittaker type. For  $\mathfrak{m} \in \text{Mult}_{\mathbb{F}_\ell}$ ,  $\mathcal{S}_{\text{gen}, \psi}(\mathfrak{m}) \subseteq \mathcal{S}_\psi(\mathfrak{m})$  and  $\mathcal{S}_{\text{gen}, \psi}^{\cup}(\mathfrak{m}) \subseteq \mathcal{S}_\psi(\mathfrak{m})$ .*

*Proof.* We start with proving properties (1) and (2) of being a  $\square$ -Whittaker type for  $\mathcal{S}_{\text{gen}, \psi}$ , which proves them for  $\mathcal{S}_{\text{gen}, \psi}^{\cup}$ . The first claim is true by construction. The second claim follows from the following observation. Let  $\mathfrak{m} \in \text{Mult}_{\mathbb{F}_\ell, \square}$  and  $\rho \in \mathfrak{C}^{\square}$ . Then it follows from the construction of  $\mathfrak{m}_{\text{gen}}$  that  $(\rho * \mathfrak{m})^{\vee} = \mathfrak{m}^{\vee} * \rho^{\vee}$ . In particular, if  $\mathfrak{m}$  is of the form  $\rho_1 * \dots * \rho_k$  for suitable  $\rho_i \in \mathfrak{C}^{\square}$ , then  $\mathfrak{m}^{\vee} = \rho_k^{\vee} * \dots * \rho_1^{\vee}$ . Moreover,  $\text{Mult}_{\mathbb{F}_\ell, \square}$  is nonzero only if  $\ell \neq 2$  and hence  $\rho_i^{\vee} \cong \rho_i^{\vee}$ . We thus have that

$$\mathcal{S}_{\text{gen}, \psi}(\mathfrak{m})^{\vee} = \mathcal{W}(\rho_1 \times \dots \times \rho_k, \psi)^{\vee} = \mathcal{W}(\rho_k^{\vee} \times \dots \times \rho_1^{\vee}, \psi^{-1}) = \mathcal{S}_{\text{gen}, \psi^{-1}}(\mathfrak{m}^{\vee})$$

and the second claim follows. The third claim for  $\mathcal{S}_{\text{gen}, \psi}$  follows from Lemma 2.4.1.

Finally, to prove  $\mathcal{S}_{\text{gen}, \psi}^{\cup}(\mathfrak{m}) \subseteq \mathcal{S}_\psi(\mathfrak{m})$  it suffices to prove that for  $\mathfrak{n}$  aperiodic with  $\mathfrak{n} \leq \mathfrak{m}$  we have  $\mathcal{S}_{\text{gen}, \psi}^{\cup}(\mathfrak{n}) \subseteq \mathcal{S}_\psi(\mathfrak{m})$ . Moreover, by Lemma 3.4.2 it suffices to assume that  $\mathfrak{m} \in \text{Mult}_R(\rho)$ . Write  $\mathfrak{n} = \rho_1 * \dots * \rho_k$ . By the uniqueness of the Whittaker model it suffices to give a nonzero map  $\rho_1 \times \dots \times \rho_k \rightarrow \mathcal{S}_\psi(\mathfrak{m})$ . By Lemma 2.4.3 and the Geometric Lemma, we have that  $\rho_1 \otimes \dots \otimes \rho_k$  appears in  $r_{\overline{\rho}_\alpha}(\overline{\mathcal{S}_\psi(\tilde{\mathfrak{m}})})$ , for any lift  $\tilde{\mathfrak{m}}$  of  $\mathfrak{m}$ , as a subquotient. Since  $\mathfrak{m} \in \text{Mult}_R(\rho)$ , considering the central characters implies that it appears as a subrepresentation, which in turn implies that there exists a nonzero map  $\rho_1 \times \dots \times \rho_k \rightarrow \overline{\mathcal{S}_\psi(\tilde{\mathfrak{m}})}$ . Since  $\rho_1 \times \dots \times \rho_k$  is of Whittaker type,  $\mathcal{W}(\rho_1 \times \dots \times \rho_k, \psi)$  is contained in the intersection of the  $\overline{\mathcal{S}_\psi(\tilde{\mathfrak{m}})}$ , implying the claim.  $\square$

**Remark.** If  $\mathfrak{m}$  is a banal multisegment (see [22]), i.e., if for each cuspidal representation  $\rho$ , the cuspidal support of  $\mathfrak{m}$  does not contain  $\rho|-|^k$  for some  $k$ , the same argument as in Lemma 4.1.1 shows that  $\mathcal{S}_{\text{gen}, \psi}(\mathfrak{m})$  and  $\mathcal{S}_\psi(\mathfrak{m})$  agree.

**Lemma 4.2.2.** *Assume  $\mathcal{T}_\psi$  is an extending map of Whittaker type. Let  $\mathfrak{m} \in \text{Mult}_{R, \square}$  and let  $\Delta = [a, b]_\rho$ ,  $\rho \in \mathfrak{C}_m^{\square}$ , be a segment in  $\mathfrak{m}$ . Then  $\mathcal{T}_\psi(\mathfrak{m} - \Delta + {}^{-}\Delta) \subseteq \mathcal{T}_\psi(\mathfrak{m})^{(m)}$ .*

*Proof.* We show that  $\rho|-|^a * (\mathfrak{m} - \Delta + {}^{-}\Delta) \leq \mathfrak{m}$ . Let  $\Gamma = [a-1, b']_\rho$  be the longest segment in  $\mathfrak{m} - \Delta + {}^{-}\Delta$  with  $a_\rho(\Gamma) = a-1$ . If  $\Gamma = {}^{-}\Delta$  we have that the left side equals the right side. Otherwise, we have  $b' > b$  and that the left side equals

$$\mathfrak{m} - \Delta + {}^{-}\Delta - \Gamma + {}^{+}\Gamma,$$

which can be obtained from  $\mathfrak{m}$  via the elementary operation  $\Gamma + \Delta \mapsto {}^{+}\Gamma + {}^{-}\Delta$ . By the above observation and the properties of  $\mathcal{T}_\psi$ , we obtain that

$$\mathcal{W}(\rho|-|^a \times \mathcal{T}_\psi(\mathfrak{m} - \Delta + {}^{-}\Delta), \psi) \subseteq \mathcal{T}_\psi(\mathfrak{m}).$$

The claim then follows from Proposition 3.2.  $\square$

**4.3.  $L$ -factors of standard modules.** Fix an additive character  $\psi$ . The goal of this section is to show that  $\mathcal{S}_\psi$  and  $\mathcal{S}_{\text{gen},\psi}$  are  $L$ -standard.

**Theorem 4.3.1.** *Let  $\mathcal{T}_\psi$  be an extending map of Whittaker type such that  $\mathcal{T}_\psi(\mathfrak{m}) \subseteq \mathcal{S}_\psi(\mathfrak{m})$  for all  $\mathfrak{m} \in \text{Mult}_R$ . Then  $\mathcal{T}_\psi$  is  $L$ -standard, i.e., for  $\mathfrak{n}, \mathfrak{m} \in \text{Mult}_{R,\square}^{\text{ap}}$*

$$L(X, \mathcal{T}_\psi(\mathfrak{m}), \mathcal{T}_{\psi^{-1}}(\mathfrak{n})) = L(X, C'(\mathfrak{m}), C'(\mathfrak{n})).$$

This has been achieved in [16, Theorem 4.22] for generic representations if  $R$  is arbitrary. In the case  $R = \overline{\mathbb{Q}}_\ell$ , this follows from the work of [12] as mentioned in Section 4.1, yielding the following corollary of Lemma 3.3.2 and Lemma 3.3.3.

**Corollary 4.3.1.** *Let  $\mathfrak{n}, \mathfrak{m} \in \text{Mult}_R$ . Then*

$$L(X, \mathcal{T}_\psi(\mathfrak{m}), \mathcal{T}_{\psi^{-1}}(\mathfrak{n}))^{-1} | L(X, C'(\mathfrak{m}), C'(\mathfrak{n}))^{-1}.$$

Before we come to the proof of Theorem 4.3.1, we note the following useful lemmas, all of which follow easily from the definitions.

**Lemma 4.3.1.** *Let  $\mathfrak{m} \in \text{Mult}_R$  containing segments  $\Delta = [a, b]_\rho$ ,  $\Delta' = [a+1, b+1]_\rho$  for some  $\rho \in \mathfrak{C}^\square$ . Let  $\mathfrak{n} \in \text{Mult}_R$  and let  $\mathfrak{n}'$  be the submultisegment of  $\mathfrak{n}$  consisting of segments of the form  $[c, d]_{\rho'}$  with  $\rho' \cong \chi\rho^\vee$  for some unramified character  $\chi$  and  $c - d = b - a$ . Finally, set  $\mathfrak{m}' = \mathfrak{m} - \Delta - \Delta' + [a, b+1]_\rho + [a+1, b]_\rho$ . Then*

$$\frac{L(X, C'(\mathfrak{m}), C'(\mathfrak{n}))}{L(X, C'(\mathfrak{m}'), C'(\mathfrak{n}'))} = \prod_{[c,d]_{\chi\rho^\vee} \in \mathfrak{n}'} (1 - \chi(\varpi_F)q^{-a-d}X)^{-f(\rho)}.$$

**Lemma 4.3.2.** *Let  $\mathfrak{m}, \mathfrak{n} \in \text{Mult}_R$  and  $\Delta = [a, b]_\rho$  a segment of maximal length in  $\mathfrak{m}$ , where we assume  $\rho \in \mathfrak{C}^\square$ . Let  $\mathfrak{n}'$  be the submultisegment of  $\mathfrak{n}$  consisting of segments  $[c, d]_{\rho'}$  with  $\rho' \cong \chi\rho^\vee$  for some unramified character  $\chi$  and  $c - d \geq b - a$ . Finally set  $\mathfrak{m}' = \mathfrak{m} - \Delta + {}^-\Delta$ . Then*

$$\frac{L(X, C'(\mathfrak{m}), C'(\mathfrak{n}))}{L(X, C'(\mathfrak{m}'), C'(\mathfrak{n}'))} = \prod_{[c,d]_{\chi\rho^\vee} \in \mathfrak{n}'} (1 - \chi(\varpi_F)q^{-a-d}X)^{-f(\rho)}.$$

**Lemma 4.3.3.** *Let  $\mathfrak{m}, \mathfrak{n} \in \text{Mult}_R$  and  $\mathcal{T}_\psi$  a map of Whittaker type such that  $\mathcal{T}_\psi(\mathfrak{n}) \subseteq \mathcal{S}_\psi(\mathfrak{n})$ ,  $\mathcal{T}_\psi(\mathfrak{m}) \subseteq \mathcal{S}_\psi(\mathfrak{m})$ . Then*

$$\frac{L(X, \mathcal{T}_\psi(\mathfrak{m}), \mathcal{T}_{\psi^{-1}}(\mathfrak{n}))L(q^{-1}X^{-1}, C'(\mathfrak{m}^\vee), C'(\mathfrak{n}^\vee))}{L(q^{-1}X^{-1}, \mathcal{T}_\psi(\mathfrak{m}^\vee), \mathcal{T}_{\psi^{-1}}(\mathfrak{n}^\vee))L(X, C'(\mathfrak{m}), C'(\mathfrak{n}))}$$

is a unit in  $R[X, X^{-1}]$ .

*Proof.* Combining Lemmas 3.3.1 and 3.3.2 shows that the above fraction equals

$$r_\ell(\epsilon(X, \mathcal{S}_\psi(\tilde{\mathfrak{m}}), \mathcal{S}_{\psi^{-1}}(\tilde{\mathfrak{n}}), \psi)) \epsilon(X, \mathcal{T}_\psi(\mathfrak{m}), \mathcal{T}_{\psi^{-1}}(\mathfrak{n}), \psi)^{-1}$$

for suitable lifts  $\tilde{\mathfrak{m}}$  and  $\tilde{\mathfrak{n}}$  of  $\mathfrak{m}$  and  $\mathfrak{n}$ . Since  $\epsilon$ -factors are units in  $R[X, X^{-1}]$ , the claim follows.  $\square$

**Lemma 4.3.4.** *Let  $\alpha, \beta \in R$  such that  $\alpha^f \neq \beta^f$ . Then*

$$\gcd(1 - (\alpha X)^f, 1 - (\beta X)^f) = 1.$$

*Proof of Theorem 4.3.1.* In fact we will show the slightly stronger statement that if  $\mathfrak{m}, \mathfrak{n} \in \text{Mult}_R$  with at least one of them aperiodic, then

$$L(X, \mathcal{T}_\psi(\mathfrak{m}), \mathcal{T}_{\psi^{-1}}(\mathfrak{n})) = L(X, C'(\mathfrak{m}), C'(\mathfrak{n})).$$

We argue firstly by induction on  $\deg(\mathfrak{m})$  and  $\deg(\mathfrak{n})$  and for fixed  $\deg(\mathfrak{m})$  and  $\deg(\mathfrak{n})$  we argue moreover by induction on the order  $\leq$  on the set of multisegments. The base case is [16, Theorem 4.22].

By Corollary 4.3.1 we know that there exists  $P \in \overline{\mathbb{F}}_\ell[X]$  such that

$$L(X, \mathcal{T}_\psi(\mathfrak{m}), \mathcal{T}_{\psi^{-1}}(\mathfrak{n}))^{-1} P(X) = L(X, C'(\mathfrak{m}), C'(\mathfrak{n}))^{-1}.$$

Moreover, let  $\mathfrak{m}_b$  and  $\mathfrak{n}_b$  be the banal parts of  $\mathfrak{m}$  and  $\mathfrak{n}$ . By definition

$$L(X, C'(\mathfrak{m}), C'(\mathfrak{n})) = L(X, C'(\mathfrak{m}_b), C'(\mathfrak{n}_b))$$

and by Lemma 3.3.1 and Proposition 3.2 and the base case,

$$L(X, \mathcal{T}_\psi(\mathfrak{m}_b), \mathcal{T}_{\psi^{-1}}(\mathfrak{n}_b)) = L(X, \mathcal{T}_\psi(\mathfrak{m}), \mathcal{T}_{\psi^{-1}}(\mathfrak{n})),$$

hence it is enough to treat the case where  $\mathfrak{m} = \mathfrak{m}_b$ ,  $\mathfrak{n} = \mathfrak{n}_b$ .

We first assume without loss of generality that  $\mathfrak{m}$  is not aperiodic, i.e., it contains a multisegment

$$[a, b]_\rho + \cdots + [a + o(\rho) - 1, b + o(\rho) - 1]_\rho.$$

Choose  $i \in \{0, \dots, o(\rho) - 1\}$  and using the notation of Lemma 4.3.1, we set  $\Delta = [a + i, b + i]_\rho$ ,  $\Delta' = [a + i + 1, b + i + 1]_\rho$ . By assumption on,  $\mathcal{T}_\psi, \mathcal{T}_\psi(\mathfrak{m}') \subseteq \mathcal{T}_\psi(\mathfrak{m})$  and hence by induction on  $\leq$  we obtain by Lemma 4.3.1 that

$$P(X) \mid \prod_{\substack{[c, d]_{\chi\rho^\vee} \in \mathfrak{n} \\ d - c = b - a}} (1 - \chi(\varpi_{\mathbb{F}}) q^{-a-i-d} X)^{f(\rho)}.$$

Let  $I_i = \{\chi(\varpi_{\mathbb{F}}) q^{-a-i-d} : [c, d]_{\chi\rho^\vee} \in \mathfrak{n} : d - c = b - a\}$ . Assume that  $P(X)$  is nonzero and let  $(1 - \alpha X)$  be one of its nonzero factors with

$$(1 - \alpha X) \mid 1 - \chi(\varpi_{\mathbb{F}}) q^{-a-i-d} X)^{f(\rho)}$$

for some  $i$  and  $[c, d]_{\chi\rho^\vee} \in \mathfrak{n} : d - c = b - a$ . Let  $j \in \{1, \dots, o(\rho)\}$ . Since

$$(1 - \alpha X) \mid \prod_{\substack{[c, d]_{\chi\rho^\vee} \in \mathfrak{n} \\ d - c = b - a}} (1 - \chi(\varpi_{\mathbb{F}})q^{-a-j-d}X)^{f(\rho)},$$

it follows that  $(1 - \alpha X) \mid (1 - \chi'(\varpi_{\mathbb{F}})q^{-a-j-d'}X)^{f(\rho)}$  for some  $[c', d']_{\chi'\rho^\vee} \in \mathfrak{n}$ . Therefore, by Lemma 4.3.4,

$$\chi'(\varpi_{\mathbb{F}})^{f(\rho)}q^{f(\rho)(-a-j-d')} = \chi(\varpi_{\mathbb{F}})^{f(\rho)}q^{f(\rho)(-a-i-d)}.$$

It follows straightforwardly that this implies that for each  $j$ ,  $[c + j, d + j]_{\chi\rho^\vee}$  appears in  $\mathfrak{m}$ , contradicting the assumption on its being aperiodic.

Secondly, we assume that both  $\mathfrak{m}$  and  $\mathfrak{n}$  are aperiodic and, without loss of generality, that the length of the longest segment in  $\mathfrak{m}$  is greater than or equal than the length of the longest segment in  $\mathfrak{n}$ . We will now use the notation of Lemma 4.3.2, e.g.,  $\Delta = [a, b]_\rho$  is a longest segment in  $\mathfrak{m}$ . Now Lemma 4.2.2 and the induction hypothesis in combination with Proposition 3.2 and Lemma 3.3.3 give

$$P(X) \mid \prod_{[c, d]_{\chi\rho^\vee} \in \mathfrak{n}'} (1 - \chi(\varpi_{\mathbb{F}})q^{-a-d}X)^{f(\rho)}.$$

If  $\mathfrak{n}'$  is empty, we are done; hence we can assume that the longest segment in  $\mathfrak{m}$  and the longest segment in  $\mathfrak{n}$  have the same length.

Replacing  $\mathfrak{m}$  and  $\mathfrak{n}$  by  $\mathfrak{m}^\vee$  and  $\mathfrak{n}^\vee$  we obtain a polynomial  $P^\vee(X)$  such that by Lemma 4.3.3

$$\frac{P(X)}{P^\vee(q^{-1}X^{-1})} = rX^k, \quad r \in \overline{\mathbb{F}}_\ell, \quad k \in \mathbb{Z}.$$

Applying the same reasoning as above to  $P^\vee(X)$  we obtain

$$P^\vee(X) \mid \prod_{[c, d]_{\chi\rho^\vee} \in \mathfrak{n}'} (1 - \chi(\varpi_{\mathbb{F}})^{-1}q^{b+c}X)^{f(\rho)}.$$

We now assume that  $P(X)$  is not a constant, and has a zero at  $(\chi(\varpi_{\mathbb{F}})q^{-a-d})^{-1}$  with  $[c, d]_{\chi\rho^\vee} \in \mathfrak{n}'$ . Then  $P^\vee(q^{-1}X^{-1})$  has to have a zero also at  $\chi(\varpi_{\mathbb{F}})q^{-a-d}$ , implying that there exists  $[c', d']_{\chi'\rho^\vee} \in \mathfrak{n}'$  such that

$$\chi'(\varpi_{\mathbb{F}})q^{-b-c'+1} = \chi(\varpi_{\mathbb{F}})q^{-a-d}.$$

Thus  $\chi' = \chi|_{-}^{-c'+1+c}$  and hence  $[c + 1, d + 1]_{\chi\rho^\vee} \in \mathfrak{n}'$ .

But now we can switch the roles of  $\mathfrak{m}$  and  $\mathfrak{n}$  as

$$L(X, \mathcal{T}_\psi(\mathfrak{m}), \mathcal{T}_{\psi^{-1}}(\mathfrak{n})) = L(X, \mathcal{T}_\psi(\mathfrak{n}), \mathcal{T}_{\psi^{-1}}(\mathfrak{m}))$$

and apply Lemma 4.3.2 with the longest segment  $[c + 1, d + 1]_{\chi\rho^\vee}$ , yielding that  $[a - 1, b - 1]_\rho$  has to be a segment of  $\mathfrak{m}$ , since  $(\chi(\varpi_{\mathbb{F}})q^{-a-d})^{-1}$  is a zero of  $P(X)$ .

Repeating this process we obtain that for all  $i \in \mathbb{Z}_{\geq 0}$  the segment  $[a - i, b - i]_\rho$  is contained in  $\mathfrak{m}$ , a contradiction to the assumption that  $\mathfrak{m}$  is aperiodic.  $\square$

Recall now the  $C$ -parameters of [16], i.e., the image of the injective map constructed in [16]

$$C : \mathfrak{Irr}_n \rightarrow \{\text{semisimple Deligne } R\text{-representations of length } n\}.$$

For the precise definition we refer to [16]. The right hand side of the above map is equipped with a tensor product denoted by  $\otimes_{ss}$  and one can associate to two  $C$ -parameters  $C(\pi)$  and  $C(\pi')$  the three local factors

$$L(X, C(\pi) \otimes_{ss} C(\pi')), \epsilon(X, C(\pi) \otimes_{ss} C(\pi'), \psi), \gamma(X, C(\pi) \otimes_{ss} C(\pi'), \psi).$$

As a corollary to Theorem 4.3.1, one can prove exactly as in [16, §6.4] the following.

**Corollary 4.3.2.** *Let  $\pi = \langle \mathfrak{m} \rangle$ ,  $\pi' = \langle \mathfrak{m}' \rangle \in \mathfrak{Irr}$ . Then*

$$\begin{aligned} L(X, \mathcal{S}_\psi(\mathfrak{m}), \mathcal{S}_{\psi^{-1}}(\mathfrak{n})) &= L(X, C(\pi) \otimes_{ss} C(\pi')), \\ \epsilon(X, \mathcal{S}_\psi(\mathfrak{m}), \mathcal{S}_{\psi^{-1}}(\mathfrak{n}), \psi) &= \epsilon(X, C(\pi) \otimes_{ss} C(\pi'), \psi), \\ \gamma(X, \mathcal{S}_\psi(\mathfrak{m}), \mathcal{S}_{\psi^{-1}}(\mathfrak{n}), \psi) &= \gamma(X, C(\pi) \otimes_{ss} C(\pi'), \psi). \end{aligned}$$

If  $\mathfrak{m} \in \text{Mult}_{R, \square}$ , the same is true if one replaces  $\mathcal{S}$  by  $\mathcal{S}_{\text{gen}, \psi}^U$ .

**4.4. Quotients of standard modules.** Let  $\mathfrak{m} \in \text{Mult}_{R, \square}$ . We can define

$$J_{\mathfrak{m}} : \mathcal{S}_\psi(\mathfrak{m}) \otimes \mathcal{S}_{\psi^{-1}}(\mathfrak{m}^\vee) \otimes C_c^\infty(\mathbb{F}^n) \rightarrow R$$

by

$$(W, W', \phi) \mapsto L(X, \mathcal{S}_\psi(\mathfrak{m}), \mathcal{S}_{\psi^{-1}}(\mathfrak{m}^\vee))^{-1} I(X, W, W', \phi)|_{X=1}.$$

Let  $C_{c,0}^\infty(\mathbb{F}^n)$  be the subspace of  $C_c^\infty(\mathbb{F}^n)$  consisting of all function vanishing at 0.

**Proposition 4.1.** *The map  $J_{\mathfrak{m}}$  vanishes for all  $\phi \in C_{c,0}^\infty(\mathbb{F}^n)$ . In particular, we obtain a nonzero map*

$$J_{\mathfrak{m}} : \mathcal{S}_{\text{gen}, \psi}(\mathfrak{m}) \otimes \mathcal{S}_{\text{gen}, \psi^{-1}}(\mathfrak{m}^\vee) \hookrightarrow \mathcal{S}_\psi(\mathfrak{m}) \otimes \mathcal{S}_{\psi^{-1}}(\mathfrak{m}^\vee) \rightarrow R$$

given by

$$W \otimes W' \mapsto J_{\mathfrak{m}}(W, W', \phi),$$

where  $\phi$  is some fixed element in  $C_c^\infty(\mathbb{F}^n)$  such that  $\phi(0) \neq 0$ .

Moreover, if  $\tilde{\mathfrak{m}}$  is a lift of  $\mathfrak{m}$  then  $J_{\tilde{\mathfrak{m}}}$  restricts to a map

$$J_{\tilde{\mathfrak{m}}} : \mathcal{S}_\psi(\tilde{\mathfrak{m}})^{\text{en}} \otimes \mathcal{S}_{\psi^{-1}}(\tilde{\mathfrak{m}}^\vee)^{\text{en}} \rightarrow \overline{\mathbb{Z}}_\ell$$

whose reduction mod  $\ell$  equals  $J_{\mathfrak{m}}$ .

*Proof.* We start with the case  $R = \overline{\mathbb{Q}}_\ell$ . By [19, Proposition 4.6], if  $\tilde{\mathfrak{m}}$  is totally unlinked the map

$$J_{\tilde{\mathfrak{m}}} : \mathcal{S}_\psi(\tilde{\mathfrak{m}}) \otimes \mathcal{S}_{\psi^{-1}}(\tilde{\mathfrak{m}}^\vee) \otimes C_{c,0}^\infty(\mathbb{F}^n) \rightarrow \overline{\mathbb{Q}}_\ell$$

vanishes. More generally, let  $\tilde{\mathfrak{m}} = \Delta_1 + \dots + \Delta_k$  with the segments in an arranged order, choose  $s \in \mathbb{Z}^k$ , and set

$$\begin{aligned} \mathcal{S}_{\psi,s}(\tilde{\mathfrak{m}}) &= \mathcal{W}(\mathcal{W}(\langle \Delta_1 \rangle, \psi) | -|^s \times \dots \times \mathcal{W}(\langle \Delta_k \rangle, \psi) | -|^s, \psi), \\ \mathcal{S}_{\psi^{-1},s}(\tilde{\mathfrak{m}}^\vee) &= \mathcal{W}(\mathcal{W}(\langle \Delta_k^\vee \rangle, \psi^{-1}) | -|^s \times \dots \times \mathcal{W}(\langle \Delta_1^\vee \rangle, \psi^{-1}) | -|^s, \psi^{-1}). \end{aligned}$$

Fix flat sections

$$f_s \in \mathcal{S}_{\psi,s}(\tilde{\mathfrak{m}}), \quad f'_s \in \mathcal{S}_{\psi^{-1},s}(\tilde{\mathfrak{m}}^\vee)$$

in the sense of [5, §3]. We obtain for fixed  $\phi \in C_{c,0}^\infty(\mathbb{F}^n)$  by [12, Proposition 3.3] a rational function over  $\overline{\mathbb{Q}}_\ell$  such that  $P(X_1, \dots, X_k)$  such that  $J_{\tilde{\mathfrak{m}}}(f_s, f'_s, \phi) = P(q^{s_1}, \dots, q^{s_k})$ . As observed above, for all but finitely many  $s$ ,  $P(q^{s_1}, \dots, q^{s_k}) = 0$  and hence it vanishes everywhere.

The case  $R = \overline{\mathbb{F}}_\ell$  follows readily from the case  $R = \overline{\mathbb{Q}}_\ell$ . Let  $\tilde{\mathfrak{m}}$  be any lift of  $\mathfrak{m}$  to  $\overline{\mathbb{Q}}_\ell$ . Then the map

$$J_{\tilde{\mathfrak{m}}} : \mathcal{S}_\psi(\tilde{\mathfrak{m}}) \otimes \mathcal{S}_{\psi^{-1}}(\tilde{\mathfrak{m}}^\vee) \otimes C_{c,0}^\infty(\mathbb{F}^n) \rightarrow \overline{\mathbb{Q}}_\ell$$

vanishes. Since  $J_{\tilde{\mathfrak{m}}}$  obviously respects the integral structures, it reduces to the map

$$J_{\mathfrak{m}} : \mathcal{S}_\psi(\mathfrak{m}) \otimes \mathcal{S}_{\psi^{-1}}(\mathfrak{m}^\vee) \otimes C_{c,0}^\infty(\mathbb{F}^n) \rightarrow \overline{\mathbb{F}}_\ell,$$

which therefore also vanishes. Moreover, if  $\tilde{\mathfrak{m}}$  is any multisegment over  $\overline{\mathbb{Q}}_\ell$ , the map  $\mathcal{S}_\psi(\tilde{\mathfrak{m}}) \otimes \mathcal{S}_{\psi^{-1}}(\tilde{\mathfrak{m}}^\vee) \rightarrow \overline{\mathbb{Q}}_\ell$  factors through  $\langle \tilde{\mathfrak{m}} \rangle \otimes \langle \tilde{\mathfrak{m}}^\vee \rangle$  and in particular vanishes on  $\mathcal{S}_\psi(\tilde{\mathfrak{n}}) \otimes \mathcal{S}_{\psi^{-1}}(\tilde{\mathfrak{n}}^\vee)$  for any  $\tilde{\mathfrak{n}} < \tilde{\mathfrak{m}}$ . Hence  $J_{\tilde{\mathfrak{m}}}$  vanishes on

$$\mathcal{S}_\psi(\tilde{\mathfrak{n}}) \otimes \mathcal{S}_{\psi^{-1}}(\tilde{\mathfrak{n}}^\vee) \otimes C_c^\infty(\mathbb{F}^n).$$

We now argue that  $J_{\mathfrak{m}}$  does not vanish on the restriction to

$$\mathcal{S}_{\text{gen},\psi}(\mathfrak{m}) \otimes \mathcal{S}_{\text{gen},\psi^{-1}}(\mathfrak{m}^\vee) \otimes C_c^\infty(\mathbb{F}^n);$$

this will prove the claim. It suffices to show that for  $\mathfrak{n} < \mathfrak{m}$  the map  $J_{\mathfrak{m}}$  vanishes on  $\mathcal{S}_{\text{gen},\psi}^{\text{U}}(\mathfrak{n}) \otimes \mathcal{S}_{\text{gen},\psi^{-1}}^{\text{U}}(\mathfrak{m}^\vee) \otimes C_c^\infty(\mathbb{F}^n)$ . It suffices to show the claim for  $\mathfrak{n}$  maximal, i.e., that it is obtained from  $\mathfrak{m}$  via one elementary operation. But then we can choose a lift  $\tilde{\mathfrak{m}}$  of  $\mathfrak{m}$  and  $\tilde{\mathfrak{n}}$  of  $\mathfrak{n}$  with  $\tilde{\mathfrak{n}} < \tilde{\mathfrak{m}}$  and the claim follows from the observation above and the fact that  $\mathcal{S}_{\text{gen},\psi}^{\text{U}}(\mathfrak{n}) \subseteq \mathcal{S}_\psi(\mathfrak{n}) \subseteq \overline{\mathcal{S}_\psi(\tilde{\mathfrak{n}})}$ ; see Corollary 4.2.1. By an analogous argument we can also show that  $J_{\mathfrak{m}}$  vanishes on  $\mathcal{S}_{\text{gen},\psi}^{\text{U}}(\tilde{\mathfrak{m}}) \otimes \mathcal{S}_{\text{gen},\psi^{-1}}^{\text{U}}(\tilde{\mathfrak{n}}^\vee) \otimes C_c^\infty(\mathbb{F}^n)$  and hence it cannot vanish on  $\mathcal{S}_{\text{gen},\psi}(\mathfrak{m}) \otimes \mathcal{S}_{\text{gen},\psi^{-1}}(\mathfrak{m}^\vee) \otimes C_c^\infty(\mathbb{F}^n)$ . By the previous arguments we know that  $J_{\mathfrak{m}}$  vanishes on  $\mathcal{S}_{\text{gen},\psi}(\mathfrak{m}) \otimes \mathcal{S}_{\text{gen},\psi^{-1}}(\mathfrak{m}^\vee) \otimes C_{c,0}^\infty(\mathbb{F}^n)$ , which finishes the argument.  $\square$

**Theorem 4.4.1.** *Let  $\mathfrak{m} \in \text{Mult}_{R, \square}^{\text{ap}}$ . Then  $\langle \mathfrak{m} \rangle$  is the unique irreducible quotient of  $\mathcal{S}_{\text{gen}, \psi}(\mathfrak{m})$ .*

*Proof.* Let  $\mathfrak{m} \in \text{Mult}_{R, \square}^{\text{ap}}$ . We argue by induction on  $\text{deg}(\mathfrak{m})$  and  $\preceq$ . The base case is trivial. Let  $\pi = \langle \mathfrak{n} \rangle$  be a quotient of  $\mathcal{S}_{\text{gen}, \psi}(\mathfrak{m})$ . Note that  $\text{cusp}(\pi) = \text{cusp}(\mathfrak{m})$ . Since  $\pi$  has to be an irreducible subquotient of  $\mathcal{S}_{\psi}(\mathfrak{m})$ , we have by Lemma 2.2.2 that  $\mathfrak{n} \preceq \mathfrak{m}$ .

We write  $\mathfrak{m} = \mathfrak{m}' * \rho$ ,  $\rho \in \mathfrak{C}_m^{\square}$  and  $\mathfrak{m}' \in \text{Mult}_{R, \square}^{\text{ap}}$  and hence  $\mathcal{S}_{\text{gen}, \psi}(\mathfrak{m}') \times \rho \twoheadrightarrow \pi$  by Lemma 2.4.1. In particular  $\mathcal{S}_{\text{gen}, \psi}(\mathfrak{m}') \otimes \rho \hookrightarrow \overline{r(\text{deg}(\mathfrak{m}) - m, m)}(\pi)$ . By induction and Proposition 3.3 we obtain that  $\langle \mathfrak{m}' \rangle \otimes \rho$  appears in  $\overline{r(\text{deg}(\mathfrak{m}) - m, m)}(\mathcal{S}_{\psi}(\mathfrak{n}))$ . By the geometric lemma of Bernstein and Zelevinsky [2, Theorem 5.2; 32, Theorem 1.1], it follows that  $\mathfrak{m}' \preceq \mathfrak{n}'$ , where  $\mathfrak{n}'$  is of the form  $\mathfrak{n}' = \mathfrak{n} - [a, 0]_{\rho} + [a, -1]_{\rho}$  for a suitable segment  $[a, 0]_{\rho}$ . By the same argument as in Lemma 4.2.2 we have  $\mathfrak{n}' * \rho \preceq \mathfrak{n}$ . On the other hand by Lemma 2.4.1 we have  $\mathfrak{m} = \mathfrak{m}' * \rho \preceq \mathfrak{n}' * \rho \preceq \mathfrak{n}$  and hence  $\mathfrak{n} = \mathfrak{m}$ . Since  $\mathcal{S}_{\text{gen}, \psi}(\mathfrak{m}) \subseteq \mathcal{S}_{\psi}(\mathfrak{m})$ , and  $\langle \mathfrak{m} \rangle$  appears in the latter with multiplicity one by Lemma 2.3.1, this is also true for  $\mathcal{S}_{\text{gen}, \psi}$ .  $\square$

Let us state two corollaries to this result. By abuse of notation we will also write  $J_{\mathfrak{m}} : \mathcal{S}_{\psi}(\mathfrak{m}) \rightarrow \mathcal{S}_{\psi^{-1}(\mathfrak{m}^{\vee})}^{\vee}$  for the map obtained from Proposition 4.1. As a consequence of this proposition, it restricts to a map  $\mathcal{S}_{\text{gen}, \psi}(\mathfrak{m}) \rightarrow \mathcal{S}_{\text{gen}, \psi^{-1}(\mathfrak{m}^{\vee})}^{\vee}$ , which factors through  $\langle \mathfrak{m} \rangle$ .

**Corollary 4.4.1.** *Let  $\mathfrak{m} \in \text{Mult}_{R, \square}$ . Then  $\langle \mathfrak{m} \rangle$  appears in the image of  $J_{\mathfrak{m}}$  as a quotient.*

*Proof.* Let  $I(\mathfrak{m})$  denote the image of  $J_{\mathfrak{m}} : \mathcal{S}_{\psi}(\mathfrak{m}) \rightarrow \mathcal{S}_{\psi^{-1}(\mathfrak{m}^{\vee})}^{\vee}$ . Let  $\Sigma(\mathfrak{m})$  be the kernel of the map  $\mathcal{S}_{\text{gen}, \psi}(\mathfrak{m}) \rightarrow \langle \mathfrak{m} \rangle$ . Then  $\Sigma(\mathfrak{m}^{\vee}) = \Sigma(\mathfrak{m})^{\mathfrak{c}}$ . Since  $\mathcal{S}_{\text{gen}, \psi}$  is  $\square$ -standard, the kernel of the map  $J_{\mathfrak{m}}$  restricted to  $\mathcal{S}_{\text{gen}, \psi}(\mathfrak{m})$  is  $\Sigma(\mathfrak{m})$ ; thus  $I(\mathfrak{m})$  is a quotient of  $\Pi(\mathfrak{m}) := \Sigma(\mathfrak{m}) \setminus \mathcal{S}_{\psi}(\mathfrak{m})$ . We know that both  $I(\mathfrak{m})$  and  $\Pi(\mathfrak{m})$  contain  $\langle \mathfrak{m} \rangle$  with multiplicity 1 by Theorem 4.4.1 and  $\Pi(\mathfrak{m})$  contains it moreover as a subrepresentation. Since  $\Pi(\mathfrak{m})^{\mathfrak{c}} \cong \Pi(\mathfrak{m}^{\vee})$ , it also follows that  $\Pi(\mathfrak{m})$  admits  $\langle \mathfrak{m} \rangle$  as a quotient, and hence as a direct summand since it appears only with multiplicity 1. Thus  $\langle \mathfrak{m} \rangle$  is also a quotient of  $I(\mathfrak{m})$ .  $\square$

The second corollary follows readily from Theorem 4.4.1, Proposition 3.3, Lemma 2.2.2 and Corollary 4.2.1.

**Corollary 4.4.2.** *For  $\mathfrak{n}, \mathfrak{m} \in \text{Mult}_{R, \square}$  we have  $\mathcal{S}_{\text{gen}, \psi}^{\cup}(\mathfrak{n}) \subseteq \mathcal{S}_{\text{gen}, \psi}^{\cup}(\mathfrak{m})$  if and only if  $\mathfrak{n} \preceq \mathfrak{m}$ . We thus have an order-preserving injection*

$$\mathcal{S}_{\text{gen}, \psi}^{\cup} : \text{Mult}_{R, \square} \hookrightarrow \mathfrak{R}\text{ep}_{W, \psi},$$

*respecting the products.*

### Acknowledgments

I would like to express my sincere gratitude to Nadir Matringe for bringing the central problem of this paper to my attention and for kindly inviting me to NYU Shanghai in March 2024. I am also deeply grateful to Alberto Minguez for taking the time to read earlier versions of this paper and for providing valuable feedback. Their insights and support have been greatly appreciated. This work has been supported by the projects PAT4628923 and PAT4832423 of the Austrian Science Fund (FWF).

### References

- [1] A.-M. Aubert, “Dualité dans le groupe de Grothendieck de la catégorie des représentations lisses de longueur finie d’un groupe réductif  $p$ -adique”, *Trans. Amer. Math. Soc.* **347**:6 (1995), 2179–2189. MR
- [2] I. N. Bernstein and A. V. Zelevinsky, “Representations of the group  $GL(n, F)$ , where  $F$  is a local non-Archimedean field”, *Uspehi Mat. Nauk* **31**:3 (1976), 5–70. In Russian; translated in *Russ. Math. Surveys* **31**:3 (1976), 1–68. MR
- [3] I. N. Bernstein and A. V. Zelevinsky, “Induced representations of reductive  $p$ -adic groups, I”, *Ann. Sci. École Norm. Sup.* (4) **10**:4 (1977), 441–472. MR
- [4] K. Bongartz, “On degenerations and extensions of finite-dimensional modules”, *Adv. Math.* **121**:2 (1996), 245–287. MR
- [5] J. W. Cogdell and I. I. Piatetski-Shapiro, “Derivatives and L-functions for  $GL_n$ ”, pp. 115–173 in *Representation theory, number theory, and invariant theory*, Progr. Math. **323**, Birkhäuser, 2017. MR
- [6] J.-F. Dat, “ $v$ -tempered representations of  $p$ -adic groups, I:  $l$ -adic case”, *Duke Math. J.* **126**:3 (2005), 397–469. MR
- [7] B. Deng and J. Du, “Monomial bases for quantum affine  $\mathfrak{sl}_n$ ”, *Adv. Math.* **191**:2 (2005), 276–304. MR
- [8] J. Droschl, “On modular representations of inner forms of  $GL_n$  over a local non-archimedean field”, preprint, 2024. arXiv 2402.13969
- [9] M. Emerton and D. Helm, “The local Langlands correspondence for  $GL_n$  in families”, *Ann. Sci. Éc. Norm. Supér.* (4) **47**:4 (2014), 655–722. MR
- [10] M. Hanzer and G. Muić, “On an algebraic approach to the Zelevinsky classification for classical  $p$ -adic groups”, *J. Algebra* **320**:8 (2008), 3206–3231. MR
- [11] H. Jacquet and J. A. Shalika, “On Euler products and the classification of automorphic forms, II”, *Amer. J. Math.* **103**:4 (1981), 777–815. MR
- [12] H. Jacquet, I. I. Piatetski-Shapiro, and J. A. Shalika, “Rankin–Selberg convolutions”, *Amer. J. Math.* **105**:2 (1983), 367–464. MR
- [13] G. Kempken, *Eine Darstellung des Köchers  $\tilde{A}_k$* , Dissertation, Rheinische Friedrich-Wilhelms-Universität, Bonn, 1981. Bonner Mathematische Schriften, **137**. MR
- [14] R. Kurinczuk and N. Matringe, “Rankin–Selberg local factors modulo  $\ell$ ”, *Selecta Math. (N.S.)* **23**:1 (2017), 767–811. MR
- [15] R. Kurinczuk and N. Matringe, “Extension of Whittaker functions and test vectors”, *Res. Number Theory* **4**:3 (2018), art. id. 31, 18 pp. MR

- [16] R. Kurinczuk and N. Matringe, “The  $\ell$ -modular local Langlands correspondence and local constants”, *J. Inst. Math. Jussieu* **20**:5 (2021), 1585–1635. MR
- [17] B. Leclerc, J.-Y. Thibon, and E. Vasserot, “Zelevinsky’s involution at roots of unity”, *J. Reine Angew. Math.* **513** (1999), 33–51. MR
- [18] N. Matringe, “Derivatives and asymptotics of Whittaker functions”, *Represent. Theory* **15** (2011), 646–669. MR
- [19] N. Matringe, “On the local Bump–Friedberg  $L$ -function”, *J. Reine Angew. Math.* **709** (2015), 119–170. MR
- [20] N. Matringe and G. Moss, “The Kirillov model in families”, *Monatsh. Math.* **198**:2 (2022), 393–410. MR
- [21] A. Mínguez, “Sur l’irréductibilité d’une induite parabolique”, *J. Reine Angew. Math.* **629** (2009), 107–131. MR
- [22] A. Mínguez and V. Sécherre, “Représentations banales de  $GL_m(D)$ ”, *Compos. Math.* **149**:4 (2013), 679–704. MR
- [23] A. Mínguez and V. Sécherre, “Représentations lisses modulo  $\ell$  de  $GL_m(D)$ ”, *Duke Math. J.* **163**:4 (2014), 795–887. MR
- [24] A. Mínguez and V. Sécherre, “Types modulo  $\ell$  pour les formes intérieures de  $GL_n$  sur un corps local non archimédien”, *Proc. Lond. Math. Soc.* (3) **109**:4 (2014), 823–891. With an appendix by Sécherre, Vincent and Shaun Stevens. MR
- [25] A. Mínguez and V. Sécherre, “L’involution de Zelevinski modulo  $\ell$ ”, *Represent. Theory* **19** (2015), 236–262. MR
- [26] A. Pütz, “Degenerate affine flag varieties and quiver Grassmannians”, *Algebr. Represent. Theory* **25**:1 (2022), 91–119. MR
- [27] M. Reineke, “Generic extensions and multiplicative bases of quantum groups at  $q=0$ ”, *Represent. Theory* **5** (2001), 147–163. MR
- [28] C. M. Ringel, “The composition algebra of a cyclic quiver: towards an explicit description of the quantum group of type  $\tilde{A}_n$ ”, *Proc. London Math. Soc.* (3) **66**:3 (1993), 507–537. MR
- [29] M.-F. Vignéras, *Représentations  $l$ -modulaires d’un groupe réductif  $p$ -adique avec  $l \neq p$* , Progress in Mathematics **137**, Birkhäuser, 1996. MR
- [30] M.-F. Vignéras, “Induced  $R$ -representations of  $p$ -adic reductive groups”, *Selecta Math.* (N.S.) **4**:4 (1998), 549–623. MR
- [31] M.-F. Vignéras, “On highest Whittaker models and integral structures”, pp. 773–801 in *Contributions to automorphic forms, geometry, and number theory*, Johns Hopkins Univ. Press, Baltimore, MD, 2004. MR
- [32] A. V. Zelevinsky, “Induced representations of reductive  $p$ -adic groups, II: On irreducible representations of  $GL(n)$ ”, *Ann. Sci. École Norm. Sup.* (4) **13**:2 (1980), 165–210. MR
- [33] A. V. Zelevinsky, “The  $p$ -adic analogue of the Kazhdan–Lusztig conjecture”, *Funktsional. Anal. i Prilozhen.* **15**:2 (1981), 9–21. In Russian; translated in *Funct. Anal. Appl.* **15**:2 (1981), 83–92. MR

Received March 11, 2025. Revised April 10, 2026.

JOHANNES DROSCHL

VIENNA

AUSTRIA

johannes.carl.droschl@univie.ac.at



# TRANSVERSE MINIMAL FOLIATIONS ON UNIT TANGENT BUNDLES AND APPLICATIONS

SÉRGIO R. FENLEY AND RAFAEL POTRIE

**We show that given two transverse minimal foliations on the unit tangent bundle of a surface of genus  $\geq 2$ , their intersection either is an Anosov foliation or contains a Reeb surface. The existence of a Reeb surface is incompatible with partially hyperbolic foliations, so we deduce from this that certain partially hyperbolic diffeomorphisms in unit tangent bundles are collapsed Anosov flows. We also conclude that every volume preserving partially hyperbolic diffeomorphism of a unit tangent bundle is ergodic.**

## 1. Introduction

This article studies geometric and dynamical properties of one-dimensional subfoliations obtained as the intersection of two transverse minimal foliations on unit tangent bundles of higher genus surfaces. We prove some strong geometric properties that imply that under certain conditions, the foliation must be homeomorphic to the orbit foliation of the geodesic flow for a hyperbolic metric on the surface.

One big motivating example for us comes from partially hyperbolic dynamics in dimension 3: under very general orientability conditions there is a pair of two-dimensional branching foliations, which are approximated by regular foliations. The pair of foliations are transverse to each other, yielding a one-dimensional subfoliation of both. Suppose that one proves that the subfoliation is the flow foliation of a topological Anosov flow. Then the partially hyperbolic diffeomorphism is what is called a collapsed Anosov flow [5; 25]. This has some important consequences, such as accessibility and ergodicity in the volume preserving case [23; 24].

This naturally leads to the following very general question, which was the initial goal of this project:

**Question.** *Let  $\mathcal{F}_1$  and  $\mathcal{F}_2$  be transverse minimal foliations in a closed 3-manifold.*

---

Fenley was partially supported by Simons Foundation grant 280429 and 637554, National Science Foundation grant DMS-2054909, and the Institute for Advanced Study. Potrie was partially supported by CSIC I+D project “Estructuras topológicas de sistemas parcialmente hiperbólicos y aplicaciones”.  
*MSC2020:* primary 57R30, 37C86, 37D30; secondary 53C12, 57K30, 37C15.

*Keywords:* foliation, 3-manifold, partial hyperbolicity, Anosov flow, ergodicity.

*Are there simple conditions that guarantee that the intersection foliation is homeomorphic to an Anosov foliation?*

Here, by an *Anosov foliation* we mean the orbit foliation of a (topological) Anosov flow.<sup>1</sup> The scope of the question is quite general, and the conditions are somewhat more restrictive than studying vector fields tangent to a foliation (for instance, the horocycle flow is a subfoliation of the weak stable foliation of an Anosov flow but it is rarely obtained as the intersection of two foliations). Having a vector field tangent to a foliation already imposes some obstructions on the foliation and the manifold (see for instance [11; 30]), so we expect here to have even more. This very general question can be extended to the problem of understanding in general the one-dimensional foliations induced by intersecting two general transverse foliations in a closed 3-manifold; we have included minimality to simplify certain formulations<sup>2</sup> but it makes sense to ask this question in general. (In fact, a similar question for three transverse taut foliations is suggested in [39, § 7.1].)

We stress that the question of analyzing general transverse intersections of foliations in 3-manifolds is very natural, interesting in itself, and has appeared in other contexts, such as Anosov and pseudo-Anosov flows transverse to foliations [19; 20; 21; 39].


In this article we start the general study of geometric properties of one-dimensional subfoliations of a pair of transverse foliations in 3-manifolds. In this generality the problem is at this point complex (see Section 1.2 for recent progress). Here we restrict to a class of 3-manifolds: unit tangent bundles of surfaces of negative Euler characteristic. In this case we have strong rigidity for single foliations and this substantially helps study this problem. It is also relevant for us since many people working in partially hyperbolic dynamics are more familiar with this family of 3-manifolds. On the other hand many of the techniques introduced in this paper should be useful for the general problem. In fact, since this paper was released, much progress has been made in the problem, which we survey in Section 1.2, showing the impact it has had.

Although minimal foliations in unit tangent bundles are homeomorphic to the weak stable foliation of an Anosov flow (by a result of [34]; see also [29]), T. Barbot pointed out to us the paper [35], which gives a beautiful example showing that even in these manifolds, there may be obstructions to the intersection being an Anosov foliation. (In fact, [35] contains a triple of pairwise transverse minimal foliations.)

---

<sup>1</sup>A short definition of a topological Anosov flow is an expansive flow preserving a foliation; see [2] for a nice introduction. In this paper we work in unit tangent bundles, where every topological Anosov flow is orbit equivalent to the geodesic flow of some constant curvature metric (which is a smooth Anosov flow) so we will not differentiate between them.

<sup>2</sup>One can always blow up one of the foliations and the intersection will no longer be an Anosov foliation. Other phenomena can also arise; see for instance the examples constructed in [10].



**Figure 1.** A Reeb surface and its lift to the universal cover.

We show that, in unit tangent bundles, the conditions that obstruct the intersection to be homeomorphic to an Anosov foliation, are similar to those appearing in the example of [35]. In Section 7 we describe the example of [35] from the point of view of this paper as well as discuss some possible extensions, and in Section 9 we prove that the obstruction to intersect in a foliation homeomorphic to an Anosov foliation can be explained by behavior identical to the ones discussed in Section 7.

The two-dimensional foliations  $\mathcal{F}_1, \mathcal{F}_2$  we deal with have Gromov hyperbolic leaves. The strategy employed here to prove the Anosov behavior of the intersection foliation  $\mathcal{G} = \mathcal{F}_1 \cap \mathcal{F}_2$  in certain cases is the following: show geometric properties of the subfoliations inside the leaves of the two-dimensional foliations. More specifically we try to show that the subfoliations are by uniform quasigeodesics inside these two-dimensional leaves. We obtain a structure that is enough to identify the obstruction for this to happen: *Reeb surfaces*. These are surfaces in some leaf of  $\mathcal{F}_1$  or  $\mathcal{F}_2$  which are finitely covered by a two-dimensional annulus whose boundary circles are leaves of the foliation  $\mathcal{G}$  and the leaves of  $\mathcal{G}$  in the interior of the annulus spiral towards the boundary components in opposite directions (see Figure 1).

Our main result is the following:

**Theorem A.** *Let  $S$  be a closed orientable hyperbolic surface and let  $M = T^1S$ . Let  $\mathcal{F}_1, \mathcal{F}_2$  be minimal, two-dimensional foliations in  $M$  which are transverse to each other. Let  $\mathcal{G}$  be the intersection of  $\mathcal{F}_1$  with  $\mathcal{F}_2$ . Then either  $\mathcal{G}$  is homeomorphic to the orbit foliation of the geodesic flow of a hyperbolic metric on  $S$  or  $\mathcal{G}$  contains a Reeb surface.*

We remark that minimality is necessary as one can easily obtain counterexamples by blowing up weak stable and weak unstable foliations of Anosov flows. We note that we get a description of  $\mathcal{G}$  when there are Reeb surfaces: we refer the reader to Section 9 for precise formulations (see in particular, Corollary 9.20). The

description says that if there are Reeb surfaces, then some structure very similar to the Matsumoto–Tsuboi example has to occur. This is a rigidity result in the sense that counterexamples to Anosov behavior for  $\mathcal{G}$  have very few possibilities.

An important point is that in Theorem A there is no (a priori) dynamical hypothesis, but there is a strong dynamical consequence. On the two-dimensional level, there is an a priori result which is the starting point of our study here and which explains the choice to work with unit tangent bundles rather than general Seifert fibered manifolds: as mentioned earlier, Matsumoto [34] proved that each  $\mathcal{F}_i$  individually is topologically equivalent to the weak stable foliation<sup>3</sup> of the geodesic flow in  $M$ . While many arguments in our proof hold in more generality, we have chosen to avoid using too much terminology and background on foliations, so that someone willing to believe Theorem 2.1 could in principle follow the proof of the main theorem. It will also be helpful to convey the main strategy and ideas that we are trying to communicate. See Section 1.2 for updates on recent results.

As a consequence we prove the following result for partially hyperbolic diffeomorphisms (see Section 10 for precise definitions and more general results).

**Corollary B.** *Let  $f : M \rightarrow M$  be a volume preserving partially hyperbolic diffeomorphism in  $M = T^1S$  with  $S$  a closed orientable surface of genus  $g \geq 2$ . Then,  $f$  is a collapsed Anosov flow. Hence  $f$  is accessible and if  $f$  is  $C^2$ , then  $f$  is ergodic.*

**1.1. Outline of the strategy and organization of the paper.** A way to detect that a one-dimensional foliation is an Anosov foliation is to show that when seen as the orbits of a flow, this flow is expansive. This implies that the flow is pseudo-Anosov [31; 37], and since our one-dimensional foliation preserves a two-dimensional foliation, being pseudo-Anosov is enough to show it is an Anosov foliation [5, § 5].

In this article we will instead consider a more geometric point of view. As in [25; 5], the main strategy to show that the flow generated by the foliation is a topological Anosov flow, is to show that its flow foliation subfoliates the two-dimensional leaves by quasigeodesics. We say that the foliation is *leafwise quasigeodesic* when in each leaf of say  $\tilde{\mathcal{F}}_1$  the leaves of the intersected foliation  $\mathcal{G}$  are quasigeodesics (note that we do not ask the one-dimensional leaves to be quasigeodesics of  $\tilde{M}$ ).

The idea is to try extend arguments that work in a compact surface (see [32] or [30, Appendix A]). Note however that in closed surfaces, compactness gives many deck transformations that preserve the universal cover, while here, leaves of the foliation in the universal cover may typically have small stabilizer. Here,

---

<sup>3</sup>In contrast, for general Seifert manifolds, it was already known that minimal foliations come from representations of surface groups into  $\text{Homeo}(S^1)$  by [38; 13] (see also [16]), but what we use here is the rigidity result of Matsumoto that gives conjugacy to a Fuchsian representation. This is also based on previous work of Ghys [29].

the key point is to use minimality and compactness of  $M$  to bring leaves together. Transversality of the foliations allows to push some behavior to nearby leaves, and then, arguments like in [22] can help partially reproducing the arguments in the case of closed surfaces. However, there are some subtleties when trying to push behavior to nearby leaves associated with the transverse geometry of leaves in the universal cover. The fact that our two-dimensional foliations are *Anosov* (in the sense that they are homeomorphic to the weak stable foliation of an Anosov flow when  $M = T^1S$ ) is extremely useful. This allows for a very strong and precise form of “pushing” at most points in the leaf approaching the boundary at infinity (see Proposition 3.4). By this, we mean that if two leaves are close by in the leafspace, then, we can sort of copy the intersected foliation  $\tilde{\mathcal{G}}$  in one leaf to nearby leaves by following the intersection with leaves of the other foliation. Note that pushing requires having two transverse foliations and this is crucial in our arguments. In fact, for flows tangent to a foliation, more diverse behavior is possible as it is shown in [25] for the strong stable foliation of the examples constructed in [9].

The route taken for showing that the foliation  $\tilde{\mathcal{G}}$  is leafwise quasigeodesic follows the rough outline that was used in [25] for foliations that come from some special dynamical systems. Not all steps hold in full generality, as examples show, but we still describe here the main steps and explain under which assumptions they work.

**Landing.** In each leaf  $L \in \tilde{\mathcal{F}}_1$  we look at the restriction of  $\tilde{\mathcal{G}}$  to the leaf  $L$ . Then every ray of any given leaf of  $\tilde{\mathcal{G}}$  (which is always properly embedded in  $L$ ) has a well defined unique limit point in the compactification  $L \cup S^1(L)$  by the Gromov boundary of  $L$ . In Theorem 4.1 we show that in our setting, this holds in full generality.

**Small visual measure.** There are many ways a properly embedded ray in a hyperbolic plane can be extended to the boundary, in particular, we wish to rule out the possibility that it lands like horocycles do. For this, a technical property that we call small visual measure is relevant. An important consequence of small visual measure is that geodesic rays starting at a point of a ray  $r$  of a leaf  $c \in \tilde{\mathcal{G}}$  and landing at the same point as the ray  $r$  must be contained in a uniform neighborhood of  $r$ . This is also something that holds always in our setting as we prove in Section 5. We point out here that the fact that  $\mathcal{G}$  is obtained as the intersection of two transverse foliations is crucial, as the horocyclic flow of an Anosov flow subfoliates a minimal foliation of  $T^1S$  but its leaves do not satisfy the small visual measure property.

**No bubble leaves.** To get a quasigeodesic foliation we need to rule out the existence of leaves  $c$  of  $\tilde{\mathcal{G}}$  such that both rays of  $c$  land in the same point of  $S^1(L)$  (where  $L$  is the two-dimensional leaf containing  $c$ ). We call such a leaf  $c$  a *bubble leaf*. We note that the example in [35] contains bubble leaves, so we cannot expect to prove

the nonexistence of bubble leaves in general. But the existence of some nonbubble leaves is important in our analysis of the small visual measure and the construction of Reeb surfaces.

**Hausdorff leaf space.** Another consequence of a one-dimensional foliation subfoliating a two-dimensional foliation by Gromov hyperbolic leaves and being leafwise quasigeodesic is that the leaf space of the foliation  $\tilde{\mathcal{G}}$  in each leaf  $L \in \tilde{\mathcal{F}}_i$  must be Hausdorff. A priori, the nonexistence of bubble leaves is not enough to rule out non-Hausdorffness, so more analysis is needed. This is the content of Section 8 where we show that non-Hausdorffness of the leaf space leads to Reeb surfaces.

**Getting the quasigeodesic property.** Having Hausdorff leaf space is not enough to deduce the leafwise quasigeodesic property as the horocycle flow shows. In our context, we can show that it is enough and we do so in Section 6.

**Outline.** In Section 2 we study general properties of minimal foliations of  $T^1S$  and derive the consequences of [34] that we will use. In Section 3 we show some properties of pairs of transverse foliations, defining and describing the landing property and showing that in some settings it is possible to *push* behavior to nearby leaves. In Sections 4 and 5 we address landing and the small visual measure property. When the leaf space of  $\tilde{\mathcal{G}}$  is leafwise Hausdorff we show in Section 6 that the foliation  $\mathcal{G}$  must be an Anosov foliation. In Section 7 we revisit the examples from [35] and describe their properties from the point of view of this paper as well as some possible extensions. In Section 8 we complete the proof of Theorem A. In Section 9 we explore further properties that in some sense show that the examples of [35] are the only possible way to introduce Reeb surfaces. In Section 10 we study the applications of our result to the classification and ergodicity of partially hyperbolic dynamics. We note that the applications to partial hyperbolicity do not use Section 7 and Section 9, which can be skipped by the reader interested only in the applications to partial hyperbolicity.

**1.2. Recent developments.** Since this paper was released in early 2023, several new developments have been obtained that we explain here, trying to emphasize the influence of this particular paper. Let us comment on the papers [1; 27; 26].

In [1] we extended part of the results of this paper to a more general setting, in particular, we showed that if  $\mathcal{F}_1, \mathcal{F}_2$  are transverse  $\mathbb{R}$ -covered Anosov foliations in a 3-manifold which are uniformly equivalent, then, they either intersect in the orbit foliation of an Anosov flow, or they contain a Reeb surface. The goal of [1] was to present a completely different approach and also present results in higher dimensions (and some results in dimension 3 that are important in [26]), and while it reproves Theorem A of this paper, the proof is completely different and the

techniques do not allow to obtain the more precise results that we obtain here in Section 9, and which we consider to be one of the main contributions of this paper (and also, that up to now do not have any counterpart in any context).

The paper [27], which was released later than this paper, also has some overlap with it, though they are largely independent. In [27] we reversed the strategy presented in Section 1.1 by assuming from the start a very strong property on the intersected foliation: that the leaf space of the intersected foliation is Hausdorff. Under that assumption, we were able to work out the full outline of Section 1.1 by showing landing, small visual measure (assuming that the manifold has fundamental group which is not virtually solvable) and finally the quasigeodesic property of leaves. While [27] works in much wider generality than this paper (no assumptions on the topology of  $M$ ), its results would not materially shorten the current paper; they would only serve in reducing Section 6, by allowing us to apply directly the results in [27] (but that would be less natural, as our proof here is more direct).

Finally, in the recent [26], we proved a general statement needed for the classification of partially hyperbolic dynamics. Though it owes much to the ideas developed here, [26] does not use or depend on this paper. Its main result is about transverse foliations with Gromov hyperbolic leaves and says that the only obstruction to the intersected foliation being leafwise quasigeodesic is the presence of what we have called *generalized Reeb surfaces*, which extend the concept of Reeb surfaces used here. We note that [26] gives hope in progressing in the understanding of general transverse foliations, and in our opinion makes Section 9 of this paper even more relevant, since no analogue of this has been shown, and the existence of Reeb surfaces in some setting could be combined with the techniques in [26] to see if one can produce some incompressible torus in  $M$ . This could for instance be relevant in addressing the following question, which we believe may well have a positive answer (see also the question on page 39):

**Question.** *Let  $\mathcal{F}_1$  and  $\mathcal{F}_2$  be two transverse minimal foliations in a closed hyperbolic 3-manifold. Then, they intersect in the orbit foliation of a (topological) Anosov flow.*

## 2. Minimal foliations on unit tangent bundles

We consider  $M = T^1S$ , the unit tangent bundle of a closed orientable surface  $S$  of genus  $g \geq 2$ , together with a minimal foliation  $\mathcal{F}$  on  $M$ . Such foliations have been completely classified by Matsumoto [34]: each is homeomorphic to the weak stable foliation of the geodesic flow on  $S$  for a hyperbolic metric. We will expand on this as well as on previous results in [38; 13] to describe the foliations in a way that is useful for our purposes.

We will assume throughout the article that the foliations we consider are  $C^{0,1+}$ . This means that the leaves are  $C^1$  surfaces; see for instance [17]. This assumption

is mostly for convenience, as having smooth leaves simplifies the definition of distances and lengths inside leaves. In [15] it is shown that for each foliation there is a smooth structure in  $M$  which makes it  $C^{0,\infty+}$ , but it is unclear that this can be done simultaneously for both foliations. To avoid discretizations of distances and local problems, we will keep this assumption throughout.

**2.1. Unit tangent bundles.** Consider the universal covering map  $\pi : \tilde{S} \rightarrow S$  in which one can identify  $\tilde{S} \cong \mathbb{H}^2$  as follows: fix a discrete subgroup  $\Gamma$  of  $\text{Isom}_+(\mathbb{H}^2) \cong \text{PSL}(2, \mathbb{R})$  (acting on the right) where  $\Gamma \cong \pi_1(S)$ . We can identify  $\pi$  with the quotient map from  $\mathbb{H}^2 \rightarrow \mathbb{H}^2/\Gamma$ .

Since  $\text{PSL}(2, \mathbb{R})$  naturally identifies with  $T^1\mathbb{H}^2$  we obtain that  $T^1S$  is identified by this action with  $T^1\mathbb{H}^2/\Gamma$ . We parametrize  $T^1\mathbb{H}^2$  by  $\mathbb{H}^2 \times \partial\mathbb{H}^2$  by identifying a unit vector  $v \in T_x\mathbb{H}^2$  with the pair  $(x, v_+) \in \{x\} \times \partial\mathbb{H}^2$ , where  $v_+$  is the limit point of the geodesic in  $\mathbb{H}^2$  starting at  $x$  with speed  $v$ . The action of  $\text{PSL}(2, \mathbb{R})$  on  $\partial\mathbb{H}^2 \cong S^1$  is given by extending the action of isometries on geodesic rays (if one uses the upper half model of  $\mathbb{H}^2$  this action corresponds to the standard action by rational transformations on  $\mathbb{R} \cup \{\infty\}$ ).

We will denote by  $M = T^1S$ , and by  $\widehat{M} = T^1\tilde{S}$  its intermediate cover with deck transformations identified with the action of  $\Gamma$  in the coordinates given by the identification of  $T^1\tilde{S} \cong \mathbb{H}^2 \times \partial\mathbb{H}^2$ . We will denote by  $\tilde{M}$  the universal cover of  $M$  which covers  $\widehat{M}$  with deck transformation group associated with the center of  $\pi_1(M)$  (which corresponds to the deck transformation associated to the circle fiber of the circle bundle over  $S$ ).

**2.2. Horizontal foliations.** Consider the foliation  $\widehat{\mathcal{F}}_{ws}$  of  $T^1\tilde{S} \cong \mathbb{H}^2 \times \partial\mathbb{H}^2$  given by  $\widehat{\mathcal{F}}_{ws} = \{\mathbb{H}^2 \times \{\xi\}\}_{\xi \in \partial\mathbb{H}^2}$ . This foliation is  $\Gamma$ -invariant and since the  $\Gamma$  action on  $\partial\mathbb{H}^2$  is minimal it descends to a minimal foliation  $\mathcal{F}_{ws}$  of  $M$  which is exactly the weak stable foliation for the geodesic flow associated to the metric on  $S$  induced by the choice of  $\Gamma \subset \text{Isom}_+(\mathbb{H}^2)$ .

The following result from [34] will be very important in our study and says that  $\mathcal{F}_{ws}$  is the unique minimal foliation of  $M$  up to homeomorphisms isotopic to the identity on the base:

**Theorem 2.1** (Matsumoto [34]). *Let  $\mathcal{F}$  be a minimal foliation of  $M$ , then, there exists a homeomorphism  $h : M \rightarrow M$  inducing the identity on the base such that  $h(\mathcal{F}) = \mathcal{F}_{ws}$ .*

We note that the result of [34] is stated for  $C^2$  foliations without compact leaves, but all that is needed is that  $\mathcal{F}$  is *horizontal*, see also [29; 38].

To get the horizontal property for  $\mathcal{F}$  we use the fact that  $\mathcal{F}$  is minimal and apply Brittenham's theorem [13]. There is a slightly technical issue in Brittenham's result: in [13] a lamination is one that is carried by a branched surface, so technically a

foliation is not a lamination and must first be split along a finite set of leaves to produce an essential lamination  $\mathcal{L}$  which, due to [13], has a minimal sublamination which is either horizontal or vertical, which since  $\mathcal{F}$  is minimal this is  $\mathcal{L}$  itself. But  $\mathcal{F}$  cannot be vertical.<sup>4</sup> It follows that  $\mathcal{F}$  is horizontal and then one can apply Matsumoto's result. See also [16] for generalities on foliations on circle bundles.

That  $h$  induces the identity on the base means that if  $p : T^1S \rightarrow S$  is the projection, then the induced actions on the fundamental group satisfy  $p_* \circ h_* = p_*$ . Note however, that  $h$  may not be homotopic to identity as a map of  $M$  and therefore two minimal foliations  $\mathcal{F}_1$  and  $\mathcal{F}_2$  may not be *uniformly equivalent* in the sense that leaves in the universal cover may not be a bounded distance away from a leaf of the other foliation. (See [39] for discussion on this notion, which is different from the notion, also used sometimes, of having homotopic plane fields.)

We will use some other properties of minimal foliations on  $M$ . Some of these hold more generally for *Reebless foliations* (due to Novikov's theorem; see [16]). We state the properties we need in the setting we will use where the proofs are a direct consequence of the corresponding properties for  $\mathcal{F}_{ws}$  and Theorem 2.1 (the last point also uses [13] for smoothness):

**Corollary 2.2.** *Let  $\mathcal{F}$  be a minimal foliation on  $M = T^1S$  and denote by  $\widehat{\mathcal{F}}$  and  $\widetilde{\mathcal{F}}$  the lifts to the covers  $\widehat{M}$  and  $\widetilde{M}$ . Then:*

- (i) *If  $\tau$  is a curve transverse to  $\widetilde{\mathcal{F}}$  then it intersects each leaf at most once.*
- (ii) *The leaf space  $\widehat{\mathcal{L}} = \widehat{M}/\widehat{\mathcal{F}}$  of  $\widehat{\mathcal{F}}$  is homeomorphic to  $S^1$  and the leaf space  $\widetilde{\mathcal{L}}$  of  $\widetilde{\mathcal{F}}$  is homeomorphic to  $\mathbb{R}$ .*
- (iii) *There is a (smooth) isotopy of  $\mathcal{F}$  that makes every leaf transverse to the circle fibers of  $T^1S$ .*

Note that a transversal to  $\widetilde{\mathcal{F}}$  or  $\mathcal{F}$  is a continuous curve  $\tau : I \rightarrow M$  where  $I$  is some interval such that for every  $t \in I$ , there is  $\varepsilon > 0$  such that the curve  $\tau|_{(t-\varepsilon, t+\varepsilon)}$  is monotone in the leaf space of a foliation chart of  $\widetilde{\mathcal{F}}$  or  $\mathcal{F}$  (which is an interval) around  $\tau(t)$ . We will many times abuse notation and denote by  $\tau$  the image of a transversal.

**2.3. Universal circle.** One consequence of Matsumoto's result (Theorem 2.1) is that every minimal foliation in  $M$  is homeomorphic to our model and so we can compare the geometry of leaves with that of hyperbolic disks simultaneously. This allows to make natural projections into  $\mathbb{H}^2$  of lifts of leaves of a minimal foliation  $\mathcal{F}$  in  $M$  to the intermediate cover  $\widehat{M}$  or universal cover  $\widetilde{M}$ .

We consider  $p : M \rightarrow S$  to be the projection of the fiber bundle  $S^1 \rightarrow M = T^1S \rightarrow S$ . By our coordinate choices, the map  $p$  lifts to a projection  $\widehat{p} : \widehat{M} \rightarrow \mathbb{H}^2$ ,

---

<sup>4</sup>If it were, it would induce a nonsingular foliation in the base surface which has nonzero Euler characteristic.

that, given the identification  $\widehat{M} \cong \mathbb{H}^2 \times \partial\mathbb{H}^2$  corresponds to the map  $\hat{p}(x, \xi) = x$ . We also denote by  $\tilde{p} : \tilde{M} \rightarrow \mathbb{H}^2$  the lift to the universal cover.

For a minimal foliation  $\mathcal{F}$  on  $M$  we will denote by  $\widehat{\mathcal{F}}$  and  $\widetilde{\mathcal{F}}$  the lifts to  $\widehat{M}$  and  $\tilde{M}$ . For concreteness we will consider the following metric on  $\widehat{M} \cong \mathbb{H}^2 \times \partial\mathbb{H}^2$ , given by the fact that the metric on  $\mathbb{H}^2 \times \{\xi\}$  is the one that makes projection in the first coordinate an isometry, makes the sets  $\mathbb{H}^2 \times \{\xi\}$  and  $\{x\} \times \partial\mathbb{H}^2$  orthogonal and measures distances in  $\{x\} \times \partial\mathbb{H}^2$  via the visual metric; namely, we define the length of an interval  $[\xi_1, \xi_2] \subset \partial\mathbb{H}^2$  as the angle between the geodesic rays starting from  $x$  and landing on  $\xi_1$  and  $\xi_2$  respectively, and measured in the direction on which rays land in the interior of the interval. This metric is invariant under the action of  $\Gamma$  and thus produces a Riemannian metric on the quotient  $M$  which makes the projection  $p : M \rightarrow S$  to be a Riemannian submersion on  $S$  when given the metric induced by the action of  $\Gamma$  on  $\mathbb{H}^2$ . In coordinates  $\widehat{M} \cong \mathbb{H}^2 \times \partial\mathbb{H}^2$  the Riemannian metric is

$$(2-1) \quad \langle v, w \rangle_{(x, \xi)} = \left( \langle v_{\mathbb{H}^2}, w_{\mathbb{H}^2} \rangle_{T_x \mathbb{H}^2}^2 + \langle v_{\partial\mathbb{H}^2}, w_{\partial\mathbb{H}^2} \rangle_{(x, \xi)}^2 \right)^{1/2}.$$

where  $v_{\mathbb{H}^2}, w_{\mathbb{H}^2}, v_{\partial\mathbb{H}^2}, w_{\partial\mathbb{H}^2}$  are the projections of the vectors on the first and second coordinates respectively, the inner product  $\langle \cdot, \cdot \rangle_{T_x \mathbb{H}^2}$  is the standard inner product in the hyperbolic plane and  $\langle \cdot, \cdot \rangle_{(x, \xi)}$  is an inner product on  $T_{(x, \xi)}(\{x\} \times \partial\mathbb{H}^2)$  given by the identification of the landing of geodesic rays of  $T_x^1 \mathbb{H}^2$  with  $\partial\mathbb{H}^2$  at the point  $\xi \in \partial\mathbb{H}^2$ .

The metric on  $\tilde{M}$  will be the path metric associated with the pullback of the Riemannian metric above by the universal cover projection. This metric in  $\tilde{M}$  will be denoted by  $d : \tilde{M} \times \tilde{M} \rightarrow \mathbb{R}_{\geq 0}$ .

Given a leaf  $L$  of a foliation  $\mathcal{F}$  we consider the path distance in  $L$  induced by the restriction of the ambient Riemannian metric on  $L$ . The choices we have made are not important if we consider objects up to *quasi-isometry*: if one chooses another metric, one obtains a distance that is quasi-isometric to the first one. Recall that a (not necessarily continuous) map  $q : (X_1, d_1) \rightarrow (X_2, d_2)$  is a  $Q$ -quasi-isometry if for every  $x, y \in X_1$  one has

$$(2-2) \quad \frac{1}{Q} d_1(x, y) - Q \leq d_2(q(x), q(y)) \leq Q d_1(x, y) + Q$$

and the image of  $q$  is  $Q$ -dense in  $X_2$ . Being quasi-isometric (meaning there exists a quasi-isometry between the spaces) is an equivalence relation between metric spaces which is particularly relevant for Gromov hyperbolic spaces such as  $\mathbb{H}^2$ . We will use several basic properties of Gromov hyperbolic spaces and refer to [12] for proofs of those facts.

Theorem 2.1 implies the following which in particular shows that leaves of a minimal foliation are Gromov hyperbolic:

**Proposition 2.3.** *There is a uniform constant  $Q_0 := Q_0(\mathcal{F})$  such that for every  $L \in \widehat{\mathcal{F}}$  (or  $\widetilde{\mathcal{F}}$ ) it follows that the restriction  $\hat{p}|_L : L \rightarrow \mathbb{H}^2$  (resp.  $\tilde{p}|_L : L \rightarrow \mathbb{H}^2$ ) is a  $Q_0$ -quasi-isometry.*

*Proof.* After an isotopy one can assume the foliation is horizontal (Corollary 2.2(iii)), and thus, by choosing an appropriate metric, we can ensure that the projection is an isometry. Since  $M$  is compact it follows that the lift of either  $\hat{p}$  or  $\tilde{p}$  restricted to any leaf is a uniform quasi-isometry. Since the statement (up to changing the constants) is invariant under change of metric on  $M$ , we conclude.  $\square$

**Remark 2.4.** A far-reaching generalization is Candel’s uniformization theorem (see [16, Chapter 7]). It can be used to show that every minimal foliation on a 3-manifold with fundamental group of exponential growth admits a metric which makes every leaf of negative curvature (see [23, § 5] and [5, Appendix A]).

We can identify the Gromov boundary of  $L$  or, equivalently, the *circle at infinity*  $S^1(L)$  of each leaf  $L \in \widehat{\mathcal{F}}$  (or  $\widetilde{\mathcal{F}}$ ) with  $\partial\mathbb{H}^2$  in a canonical way. Notice that the *universal circle* of  $\mathcal{F}$  as defined in [39] is also canonically identified with  $\partial\mathbb{H}^2$  in this case.


**2.4. Nonmarker points.** Here we introduce the notion of marker and nonmarker points for the foliations we are interested in. More general definitions can be found in [16]. We start by analyzing  $\mathcal{F}_{ws}$  using the metric given by (2-1) and then in the next subsection we show similar properties for every minimal foliation (because by Theorem 2.1, they are all homeomorphic to  $\mathcal{F}_{ws}$ ).

An orientation will be fixed on  $\mathbb{H}^2$ . Then given an oriented geodesic  $\ell \in \mathbb{H}^2$  we denote  $H_+(\ell)$  and  $H_-(\ell)$  the half spaces determined by  $\ell$  (that is, the closure in  $\mathbb{H}^2$  of the connected components of  $\mathbb{H}^2 \setminus \ell$ ) with respect to the chosen orientations. Specifically  $H_+(\ell)$  is the half space to the left of  $\ell$  and  $H_-(\ell)$  is the half space to the right of  $\ell$ , with respect to the orientation in  $\mathbb{H}^2$ . For  $X \subset \mathbb{H}^2$  and  $C > 0$  denote  $B_C(X)$  to be the set of points  $x \in \mathbb{H}^2$  whose distance to  $X$  is less than or equal to  $C$ .

**Proposition 2.5.** *Given  $\varepsilon \in (0, \pi)$  there exists  $C := C(\varepsilon)$  such that for every interval  $[\eta, \xi] \in \partial\mathbb{H}^2$  the set of points  $x \in \mathbb{H}^2$  such that the visual length from  $x$  of the interval  $[\eta, \xi]$  is less than  $\varepsilon$  is  $H_-(\ell) \setminus B_C(\ell)$  where  $\ell$  is a geodesic joining  $\eta, \xi$  oriented so that the interval  $[\eta, \xi]$  is contained in the closure of  $H_+(\ell)$  in the compactification  $\mathbb{H}^2 \cup \partial\mathbb{H}^2$ .*

*Proof.* For every  $x \in H_+(\ell)$  the visual length from  $x$  of  $[\eta, \xi]$  is greater than or equal to  $\pi > \varepsilon$ , so the set we wish to describe does not intersect  $H_+(\ell)$ .

Now, fix a geodesic ray  $r : [0, \infty) \rightarrow \mathbb{H}^2$ , parametrized by unit speed, starting at some point  $r(0)$  in  $\ell$ , orthogonal to it and contained in  $H_-(\ell)$ . For  $t > t'$  the ideal geodesic triangle joining  $r(t), \xi, \eta$  contains the triangle joining  $r(t'), \xi, \eta$  so



**Figure 2.** The set  $D_\varepsilon(I)$  for  $I = [\xi, \eta]$  is the complement of the union of  $H_+(\ell)$  and  $B_C(\ell)$ . If  $\varepsilon$  is very small, then  $C$  is very large.

the inner angle at the point  $r(t)$  is smaller than the one at the point  $r(t')$  by Gauss–Bonnet formula. As it varies continuously, and since the angle is  $\pi$  at  $r(0)$  and tends to 0 at  $r(\infty)$ , it follows that every such geodesic ray has a unique point  $r(t_0)$  on which the angle is exactly  $\varepsilon$  and it follows also that  $d(r(t_0), \ell) = t_0 := C(\varepsilon)$ . To show that this does not depend on  $r(0)$  consider the one parameter family of isometries of  $\mathbb{H}^2$  fixing  $\xi, \eta$ ; these isometries map  $r(t_0)$  transitively along the boundary of  $B_{t_0}(\ell)$  preserving angles. Hence  $t_0$  does not depend on  $r(0)$ , and depends only on  $\varepsilon$ . This proves the proposition.  $\square$

We can parametrize leaves of  $\widehat{\mathcal{F}}_{ws}$  by  $\partial\mathbb{H}^2$  as these are of the form  $\mathbb{H}^2 \times \{\xi\}$  with  $\xi \in \partial\mathbb{H}^2$ . We denote by  $L_\xi = \mathbb{H}^2 \times \{\xi\}$  and call  $\xi$  the *nonmarker* point of  $L_\xi$ . This will be denoted as  $\alpha(L_\xi) = \xi$ . The key point is that the choices of coordinates make this point in  $\partial\mathbb{H}^2$  special with respect to the leaf  $L_\xi$  as it will now be explained.

Given  $\varepsilon > 0$  and an interval  $I \subset \partial\mathbb{H}^2$  we define the set (see Figure 2)

$$(2-3) \quad D_\varepsilon(I) = \{x \in \mathbb{H}^2 : d_{\widehat{M}}((x, \xi), L_\eta) < \varepsilon \text{ for all } \xi, \eta \in I\},$$

where  $d_{\widehat{M}}$  is the distance in  $\widehat{M}$ . The point  $(x, \xi)$  belongs to  $L_\xi$  by definition. We will always assume that  $I$  is not  $\partial\mathbb{H}^2$  nor a single point. For any interval  $I = [\xi_-, \xi_+]$ , the points in  $D_\varepsilon(I)$  form a subset of  $\mathbb{H}^2$  and the points in  $\partial\mathbb{H}^2$  for which the biggest distance between the corresponding leaves is achieved is when  $\{\eta, \xi\} = \{\xi_-, \xi_+\}$ . So it is enough to look at the distance of points of the form  $(x, \xi_-)$  to  $L_{\xi_+}$  (or  $(x, \xi_+) \in L_{\xi_-}$ ). From the previous proposition we deduce:

**Corollary 2.6.** *Given  $I \subset \partial\mathbb{H}^2$  an interval and a constant  $\varepsilon \in (0, \pi)$ , there exists  $C := C(I, \varepsilon)$  such that  $D_\varepsilon(I)$  is equal to the set  $H_-(\ell) \setminus B_C(\ell)$ . Here  $\ell$  is the geodesic joining the endpoints of  $I$  oriented so that  $I$  is contained in the closure of  $H_+(\ell)$  in  $\mathbb{H}^2 \cup \partial\mathbb{H}^2$ .*

We will denote the lift of the points of  $D_\varepsilon(I)$  in  $L_\xi$  by

$$(2-4) \quad D_\varepsilon(\xi, I) = \{(x, \xi) \in \mathbb{H}^2 \times \{\xi\} : x \in D_\varepsilon(I)\} \subset L_\xi.$$

**2.5. Minimal foliations in  $M = T^1S$ .** Now, let  $\mathcal{F}$  be an arbitrary minimal foliation on  $M$ . By Theorem 2.1 there is a homeomorphism  $h : M \rightarrow M$  mapping  $\mathcal{F}$  to  $\mathcal{F}_{ws}$ .

We will produce maps  $\Phi_L$  for  $L$  in  $\tilde{\mathcal{F}}$  which are uniform quasi-isometries (Proposition 2.3). First fix an isotopy of  $M$ , which restricted to each individual leaf of  $\mathcal{F}$  is as regular as the leaves of  $\mathcal{F}$  are, to get to a foliation  $\mathcal{F}'$  which is transverse to the standard Seifert fibration of  $M = T^1S$  (Corollary 2.2(iii)). This is the Seifert fibration whose fibers are the unit vectors over a given point in  $S$ . The lift of this isotopy to  $\tilde{M}$  is denoted by  $\nu$ . For any leaf of  $\mathcal{F}'$  project to  $\mathbb{H}^2$  using the lift of this Seifert fibration, this projection is  $\tilde{p}$ . Since the angle between  $\mathcal{F}'$  and the Seifert fibration is bounded below, this projection is a uniform quasi-isometry. The composition of the initial isotopy (lifted to  $\tilde{M}$ ) and the projection is well defined up to a bounded distortion (depending on  $\mathcal{F}$ ). We denote this composition by  $\Phi_L$ . The conclusion is that we obtain an equivariant collection of uniform quasi-isometries  $\Phi_L$  from leaves  $L \in \tilde{\mathcal{F}}$  to  $\mathbb{H}^2$ . Also, if  $\tilde{p} : \tilde{M} \rightarrow \mathbb{H}^2$  is the standard projection, there is a uniform constant  $C_0 > 0$  such that  $d_{\mathbb{H}^2}(\Phi_L(x), \tilde{p}(x)) < C_0$ . Note that  $\Phi_L = \tilde{p} \circ \nu$  restricted to  $L$ .

This allows us to:

- Associate to each  $L \in \tilde{\mathcal{F}}$  a point, which we will denote throughout the article by  $\alpha(L) \in \partial\mathbb{H}^2$ , and that we will call the *nonmarker point* of  $L$ . It also identifies the leaf space of  $\tilde{\mathcal{F}}$  with  $\partial\mathbb{H}^2 \cong S^1$ . This uses both  $h$  and the quasi-isometries above. First the map  $h$  shows that for any leaf  $F$  of  $\mathcal{F}$  there is one ideal direction which is transversely noncontracting (the nonmarker direction), and all other directions are contracting. In particular, by lifting to appropriate covers, this is also true for leaves in  $\tilde{\mathcal{F}}$  or  $\hat{\mathcal{F}}$ . The uniform quasi-isometries then allow for  $L \in \tilde{\mathcal{F}}$  to obtain the unique nonmarker point  $\alpha(L)$  in  $\partial\mathbb{H}^2$ . Finally using  $h$  again one can show that the map  $\alpha$  is a homeomorphism when considered as a map from the leaf space of  $\hat{\mathcal{F}}$  and  $\partial\mathbb{H}^2 \cong S^1$ .
- For each  $L$  in  $\tilde{\mathcal{F}}$  (or  $\hat{\mathcal{F}}$ ) Identify the (Gromov) boundary at infinity  $S^1(L)$  of  $L$  with  $\partial\mathbb{H}^2$  using the fact that  $\Phi_L$  is a quasi-isometry. Importantly, the induced map  $\Phi_L : S^1(L) \rightarrow \partial\mathbb{H}^2$  is independent on the choice of the isotopy of  $\mathcal{F}$  to a horizontal foliation.
- We have maps

$$(2-5) \quad \Phi_L : L \cup S^1(L) \rightarrow \mathbb{H}^2 \cup \partial\mathbb{H}^2,$$

As explained before, each such map, when restricted to  $L$  is a diffeomorphism,

for each  $L$ ; and the diffeomorphisms have uniformly bounded derivatives (thus, it is a uniform quasi-isometry, independent on  $L$ ).

Note that  $\alpha(L)$  belongs to  $\partial\mathbb{H}^2$  but we can identify it canonically with a point in  $S^1(L)$  via  $\Phi_L$  and sometimes we will go back and forth with these identifications. Also, these facts hold for leaves  $L \in \widehat{\mathcal{F}}$  and we will abuse notation and denote by  $\Phi_L : L \cup S^1(L) \rightarrow \mathbb{H}^2 \cup \partial\mathbb{H}^2$  all such maps.

**2.6. Nearby sets in distinct leaves.** For  $\varepsilon > 0$ , a leaf  $L \in \widetilde{\mathcal{F}}$  (or  $\widehat{\mathcal{F}}$  using the metric  $d_{\widehat{M}}$ ) and an interval  $I$  of the leaf space of  $\widetilde{\mathcal{F}}$  (or  $\widehat{\mathcal{F}}$ ) we define

$$(2-6) \quad \widehat{D}_\varepsilon(L, I) = \{y \in L : \forall E \in I, d(y, E) < \varepsilon\}$$

Corollary 2.6 leads to the next result, thanks to the map  $\Phi_L$  and the fact that  $\mathcal{F}$  is homeomorphic to  $\mathcal{F}_{ws}$ :

**Proposition 2.7.** *There is a constant  $C > 0$  independent of  $L$  such that the set  $\widehat{D}_\varepsilon(L, I)$  it is at Hausdorff distance less than  $C$  from  $\Phi_L^{-1}(D_\varepsilon(I))$ . In particular, since  $\Phi_L$  is a quasi-isometry, there is another constant  $C' > 0$  such that the Hausdorff distance between  $\Phi_L(\widehat{D}_\varepsilon(L, I))$  and  $D_\varepsilon(I)$  is less than  $C'$ . Moreover, if  $L_1 \in I$  and  $x_n \in \widehat{D}_\varepsilon(L, I)$  is a sequence of points converging to some point  $\xi \in S^1(L)$  which is not  $\alpha(L')$  for some  $L' \in I$  then we have that  $d(x_n, L_1) \rightarrow 0$ .*

*Proof.* Recall that we start with an isotopy from  $\mathcal{F}$  to a horizontal foliation, then project using the Seifert fibration. These are the maps  $\Phi_L$ . Choose  $\varepsilon'$  depending on  $\varepsilon$  and such that if points are within  $\varepsilon'$  then after undoing the isotopy the points are at most  $\varepsilon$  from each other. This shows that

$$\Phi_L^{-1}(D_{\varepsilon'}(I)) \subset \widehat{D}_\varepsilon(L, I).$$

Since  $D_{\varepsilon'}(I)$  and  $D_\varepsilon(I)$  are a bounded Hausdorff distance from each other, there is  $C_1$  with  $B_{C_1}(\Phi_L^{-1}(D_\varepsilon(I))) \subset \widehat{D}_\varepsilon(L, I)$ .

Conversely, given  $\varepsilon$ , since the isotopy is the lift of a compact isotopy there is  $\varepsilon_1 = \varepsilon_1(\varepsilon)$  such that  $\Phi_L(\widehat{D}_\varepsilon(L, I)) \subset D_{\varepsilon_1}(I)$ . But  $D_{\varepsilon_1}(I)$  is a bounded Hausdorff distance from  $D_\varepsilon(I)$ , so there is  $C_2 > 0$  such that

$$\Phi_L(\widehat{D}_\varepsilon(L, I)) \subset B_{C_2}(D_\varepsilon(I)).$$

Taking  $\Phi_L^{-1}$ , and noticing that it is a quasi-isometry, produces  $C_3 > 0$  such that  $\widehat{D}_\varepsilon(L, I) \subset B_{C_3}(\Phi_L^{-1}(D_\varepsilon(I)))$ . This finishes the proof of the proposition.  $\square$

This proposition will combine well with Corollary 2.6 to control the geometry of the sets  $D_\varepsilon(L, I)$ .

**2.7. Minimality of the action in the universal circle.** In the next proposition we collect some facts about the action of the fundamental group of  $S$  in the boundary  $\partial\mathbb{H}^2$ . Recall that we have chosen a fixed hyperbolic metric on  $S$  which is induced by a subgroup  $\Gamma \cong \pi_1(S)$  of isometries of  $\mathbb{H}^2$  that induces an action on  $\partial\mathbb{H}^2$  (recall that for such groups every non identity element acts as a hyperbolic isometry, so it has exactly two fixed points in  $\partial\mathbb{H}^2$ , one attracting and one repelling).

The fundamental group  $\pi_1(M)$  of  $M$  is a central extension of  $\pi_1(S)$  (that is, there is a surjective morphism  $\pi_1(M) \rightarrow \pi_1(S)$  such that the preimage of the identity is the center of the group  $\pi_1(M)$  and is generated by the homotopy class of the fibers) and its action on  $\partial\mathbb{H}^2$  is induced by the action of its projection on  $\pi_1(S)$ .

**Proposition 2.8.** *The action of  $\Gamma \cong \pi_1(S)$  in  $\partial\mathbb{H}^2$  is minimal (i.e., every orbit is dense). Given open sets  $U, V$  in  $\partial\mathbb{H}^2$  there exists an element  $\gamma \in \Gamma$  such that*

- $\gamma(U) \cap V \neq \emptyset$ , and
- the fixed points of  $\gamma$  are one contained in  $U$  and one contained in  $V$ .

This result is classical; a proof can be found for instance in [22] where we prove an extension to general uniform  $\mathbb{R}$ -covered foliations ([22, Proposition 5.3] applied to the foliation by compact surfaces in  $S \times S^1$  gives the previous result). Note that since we can identify the leaf space of  $\widehat{\mathcal{F}}$  with  $\partial\mathbb{H}^2$  this also gives information about the action of  $\Gamma$  acting on the leaf space of  $\widehat{\mathcal{F}}$  on  $T^1\mathbb{H}^2$  which is a circle identified with  $\partial\mathbb{H}^2$  via the map  $\alpha$  sending a leaf into its nonmarker point. If we go to the universal cover, then, the central extension of  $\Gamma$  that is  $\pi_1(M)$  provides all lifts of the action of  $\Gamma$  in  $\partial\mathbb{H}^2$  to the universal cover (in particular, there are always lifts with fixed points and the action is minimal).

**2.8. Some plane topology.** We will use the following standard consequence of the classical Schoenflies theorem (see [36, § 9], for instance). We will always be using piecewise smooth curves, so the proof is simpler.

**Proposition 2.9.** *Let  $c$  be a properly embedded curve in the plane. Then, the complement of  $c$  is the union of two topological open disks whose boundary in the plane is exactly  $c$ .*

Since we will always be working on  $\mathbb{H}^2$ , where we have a natural compactification  $\mathbb{H}^2 \cup \partial\mathbb{H}^2$  homeomorphic to a disk, we want to understand the complements of properly embedded curves in this compactification. For a set  $K \subset \mathbb{H}^2$  the *limit set* of  $K$  is the closure of  $K$  in  $\mathbb{H}^2 \cup \partial\mathbb{H}^2$  intersected with  $\partial\mathbb{H}^2$ . We consider a circular order in  $\partial\mathbb{H}^2$ . Let  $c$  be a properly embedded curve in  $\mathbb{H}^2$  and denote by  $I_1$  and  $I_2$  the limit sets of the two rays of  $c$  (i.e., consider  $x \in c$  and denote by  $c_1, c_2$  the connected components of  $c \setminus \{x\}$ , then  $I_i$  is limit set of  $c_i$ ). Notice that both  $I_1, I_2$  are connected subsets of  $\partial\mathbb{H}^2$  and are either  $\partial\mathbb{H}^2$  or an interval. In the case that

they (or one of them) are not  $\partial\mathbb{H}^2$  we denote them by:  $I_1 := [a, b]$ ,  $I_2 := [c, d]$  in the circular order of  $\partial\mathbb{H}^2$ .<sup>5</sup>

**Corollary 2.10.** *Let  $c$  be a properly embedded curve in  $\mathbb{H}^2$  as above. Let  $D^+$  and  $D^-$  denote the connected components of  $\mathbb{H}^2 \setminus c$ , and let  $J^+$ ,  $J^-$  be their respective limit sets. Then*

- *If  $I_1 \cup I_2 = \partial\mathbb{H}^2$  then both  $J^+$  and  $J^-$  coincide with  $\partial\mathbb{H}^2$  (this includes the case where one of the intervals  $I_1$  or  $I_2$  coincides with  $\partial\mathbb{H}^2$ ).*
- *Assume that  $[a, b] \cap [c, d] = \emptyset$ , and suppose that  $a, b, c, d$  are circularly ordered in  $\partial\mathbb{H}^2$ . Then one of  $J^+$ ,  $J^-$  is  $[a, d]$  and the other is  $[c, b]$ .*
- *Finally suppose  $[a, b]$ ,  $[c, d]$  intersect (and their union is not  $\partial\mathbb{H}^2$ ). Then one of  $J^+$ ,  $J^-$  is  $\partial\mathbb{H}^2$  and the other is  $[a, b] \cup [c, d]$ .*

*Proof.* The limit set  $J^\pm$  of  $D^\pm$  is a compact connected subset of the boundary  $\partial\mathbb{H}^2$ . Since  $c$  is the boundary in  $\mathbb{H}^2$  of both  $D^+$  and  $D^-$  it follows that  $I_1 \cup I_2$  is contained in both  $J^+$  and  $J^-$ . This already proves the first point.

For the second and third items, it is implicitly assumed that none of  $I_1, I_2$  are  $\partial\mathbb{H}^2$ . Since  $D^+ \cup D^-$  must accumulate in all of  $\partial\mathbb{H}^2$ , it follows that if one considers a point  $\xi \in \partial\mathbb{H}^2 \setminus (I_1 \cup I_2)$  then it has a neighborhood  $N$  in the compactification  $\mathbb{H}^2 \cup \partial\mathbb{H}^2$ , so that  $N \cap \mathbb{H}^2$  which is contained in either  $D^+$  or  $D^-$ . To finish we must show that  $(b, c)$  is contained in one of  $J^+$ ,  $J^-$  and  $(d, a)$  is contained in the other. To do that consider a geodesic  $\mu$  in  $L$  with one ideal point in  $(b, c)$  and the other in  $(d, a)$ . The ideal points are disjoint from  $I_1 \cup I_2$  so  $\mu$  has rays contained in  $D_1 \cup D_2$ . If both rays are contained in say  $D_1$  it follows that both rays of  $c$  are also contained in the same complementary component of  $\mu$ . This contradicts that  $c$  limits on both  $[a, b]$  and  $[c, d]$ . Hence only one ideal point is in  $J^+$  and the other is in  $J^-$ , and consequently one of  $(b, c)$ ,  $(d, a)$  is contained in  $J^+$  and the other in  $J^-$ . This proves the second statement.


For the third statement: since  $I_1$  intersects  $I_2$ , then the above fact implies that  $\partial\mathbb{H}^2 \setminus (I_1 \cup I_2)$  is contained in one and only one of  $J^+$  or  $J^-$ . Since  $I_1 \cup I_2 \subset J^+ \cap J^-$ , the third statement follows.  $\square$

We will also need the following consequence of Proposition 2.9

**Proposition 2.11.** *Let  $r_1, r_2$  be two disjoint properly embedded rays in  $\mathbb{H}^2$  which limit in intervals  $I_1$  and  $I_2$  respectively. Assume that  $I_1 \neq \partial\mathbb{H}^2$ , then  $I_2$  cannot be contained in the interior of  $I_1$ .*

*Proof.* Consider a point  $o \in \mathbb{H}^2$  and two geodesic rays  $g_1, g_2$  from  $o$  landing at points  $\xi_1, \xi_2$  in the interior of  $I_1$  and a ray  $g_3$  landing at a point  $\xi_3 \notin I_1$ . We claim that there are arcs  $v_n$  of  $r_1$  joining  $g_1, g_2$  which converge in the topology of  $\mathbb{H}^2 \cup \partial\mathbb{H}^2$  to the

<sup>5</sup>We allow  $a = b$  or  $c = d$ , in which case the interval is a point.



**Figure 3.** The limit set of  $r_1$  is the interval  $I_1$  with endpoints  $a, b$ .  $g_1, g_2$  are geodesic rays with ideal points in the interior of  $I_1$ . The segments  $v_n$  are segments in  $r_1$  which limit to  $[a, b]$ . The points  $z_k$  are in  $r_2$  and have to connect outside the  $v_n$  to one of  $g_1$  or  $g_2$  in  $D_{n_k}$ . So the intersections with (say)  $g_1$  limit to an arbitrary point in the interior of  $I_1$ .

interval  $[\xi_1, \xi_2]$  contained in  $I_1$ . This is because given  $R > 0$  we have that outside the disk  $B_R(o)$  of radius  $R$  centered in  $o$  there are sequences of points  $x_n, y_n$  of  $r_1$  converging to points in  $I_1 \setminus [\xi_1, \xi_2]$  and such that the arc  $J_n$  joining  $x_n$  to  $y_n$  is completely outside  $B_R(o)$  and cannot intersect  $g_3$ . Thus, there must be an arc inside  $J_n$  that we call  $v_n$  joining  $g_1$  and  $g_2$  and which is outside  $B_R(o)$  and contained in the region bounded by  $\partial B_R(o) \cup g_1 \cup g_2$  which accumulates in  $[\xi_1, \xi_2]$ . This proves our claim.

We refer to Figure 3 for the situation in this Lemma.

Let  $g'_i$  be the subray of  $g_i$  starting in  $v_n \cap g_i$ . If we consider the regions  $D_n$  obtained as the closure in  $\mathbb{H}^2 \cup \partial\mathbb{H}^2$  bounded by  $v_n \cup g_1 \cup g_2$  and accumulating in  $[\xi_1, \xi_2]$  we get that after subsequence we can assume it is a nested sequence  $D_{n+1} \subset D_n$  of disks such that  $\bigcap D_n = [\xi_1, \xi_2]$ . If  $r_2$  is a properly embedded ray disjoint from  $r_1$  which accumulates in a point  $\xi \in (\xi_1, \xi_2)$  we can see that there is a sequence of points  $z_k \in r_2$  converging to  $\xi$  and thus must be contained in some sets of the form  $D_{n_k} \setminus D_{n_{k+1}}$ . Since  $r_2$  is connected, to join  $z_k$  with  $z_{k+1}$  it must exit  $D_{n_k}$  without intersecting  $D_{n_{k+1}}$  thus there must be an arc  $t_k$  of  $r_2$  that joins  $z_k$  to some point  $w_k$  in either  $g_1$  or  $g_2$ . Up to subsequence, we can assume without loss of generality that  $w_k$  is in  $g_1$  and we deduce that  $r_2$  must limit in  $[\xi_1, \xi]$ . Since  $\xi_1, \xi_2$  were arbitrary, we deduce that  $r_2$  cannot limit in an interval completely contained in the interior of  $I_1$  as desired.  $\square$

**Remark 2.12.** In Proposition 2.11, if  $I_1 = \partial\mathbb{H}^2$ , we cannot consider the ray  $g_3$  which forces the curves  $v_n$  to join  $g_1, g_2$  on the “correct” side. It is possible however to restrict the possible landing regions of disjoint properly embedded rays in this setting. This will be used later, but we will not give a precise general statement.

### 3. Pairs of transverse foliations

In this section we will consider two transverse minimal foliations  $\mathcal{F}_1$  and  $\mathcal{F}_2$  in  $M = T^1S$ . We will call  $\mathcal{G} := \mathcal{F}_1 \cap \mathcal{F}_2$  the one-dimensional foliation by the connected components of intersections of leaves of  $\mathcal{F}_1$  and  $\mathcal{F}_2$ . Note that since  $S$  is orientable it induces orientations on  $M$ ,  $\mathcal{F}_1$  and  $\mathcal{F}_2$  and thus also on  $\mathcal{G}$ , we fix one such orientation and lift it to  $\tilde{M}$ .

We consider the maps  $\Phi_L^1$  and  $\Phi_E^2$  from the compactification of leaves  $L \in \widehat{\mathcal{F}}_1$  (or  $\tilde{\mathcal{F}}_1$ ) and  $E \in \widehat{\mathcal{F}}_2$  (or  $\tilde{\mathcal{F}}_2$ ) as defined in (2-5). Since the foliation will be implicit from the leaf, we will usually omit the superscript, writing  $\Phi_L$  for  $\Phi_L^1$  when it is clear that  $L$  lies in  $\tilde{\mathcal{F}}_1$  or  $\widehat{\mathcal{F}}_1$ .

**3.1. Some general properties.** Given a leaf  $c \in \mathcal{G}$  and  $x \in c$  we will denote by

$$(3-1) \quad c_x^+ \cup c_x^- = c \setminus \{x\}$$

the two rays of  $c$ , where  $c_x^+$  is the one in the positive orientation starting at  $x$ . We will denote by  $c^+$  and  $c^-$  the *ray class*, meaning the equivalence class of rays of  $c$  such that any two equivalent rays coincide in a subray.

Given a curve  $c \in \widehat{\mathcal{G}}$  (or  $\tilde{\mathcal{G}}$ ) we know that  $c$  is a connected component of  $L \cap E$  where  $L \in \widehat{\mathcal{F}}_1$  and  $E \in \widehat{\mathcal{F}}_2$  (or in the corresponding lifts to the universal cover).<sup>6</sup> The following is a direct consequence of the quasi-isometric properties of  $\Phi_L^1$  and  $\Phi_E^2$ :

**Lemma 3.1.** *The curves  $\Phi_L^1(c)$  and  $\Phi_E^2(c)$  are proper and a bounded distance away from each other in  $\mathbb{H}^2$ .*

In particular, if we denote by  $\bar{c}_L$  the closure of  $c$  in  $L \cup S^1(L)$  and by  $\bar{c}_E$  the respective closure in  $E \cup S^1(E)$  we get the following important property:


**Corollary 3.2.** *The sets  $\Phi_L(\bar{c}_L \setminus c)$  and  $\Phi_E(\bar{c}_E \setminus c)$  are contained in  $\partial\mathbb{H}^2$  and coincide.*

We will then write, for  $c \in \widehat{\mathcal{G}}$ ,

$$(3-2) \quad \partial\Phi(c) := \Phi_L(\bar{c}_L \setminus c) = \Phi_E(\bar{c}_E \setminus c) \subset \partial\mathbb{H}^2.$$

Since  $c$  is properly embedded we can write  $\partial\Phi(c) = \partial^+\Phi(c) \cup \partial^-\Phi(c)$  where each denotes the accumulation points of the positive and negative rays of  $c$  once a point is removed (it is easy to see that this is independent on the removed point and so it is a property of the ray class). Note that each of  $\partial^+\Phi(c)$  and  $\partial^-\Phi(c)$  are compact connected and nonempty subsets of  $\partial\mathbb{H}^2$  (see Figure 4). Now we define a fundamental property to be analyzed in this article:

<sup>6</sup>For the remainder of this section we will omit saying that things hold both in the cover  $\widehat{M}$  and the universal cover  $\tilde{M}$ .



**Figure 4.** An example where the negative ray of  $c$  lands and the positive ray of  $c$  accumulates in an interval which is not a point.

**Definition 3.3.** Given  $c \in \widehat{\mathcal{G}}$  we say that the *positive ray lands* (resp. the *negative ray lands*) if  $\partial^+\Phi(c)$  is a point (resp.  $\partial^-\Phi(c)$  is a point).

For notational simplicity, we also denote by  $\partial\Phi(r)$  the accumulating set of a ray when  $r$  is a ray of a leaf  $c \in \mathcal{G}_L$ . In addition if  $r$  lands, we say that  $\partial\Phi(r)$  is the landing point of  $r$ . If  $r$  in  $L$  lands, then  $r$  also limits in a single point  $b$  in  $S^1(L)$  (in particular  $\Phi_L(b) = \partial\Phi(r)$ ). We also say that  $r$  lands in  $b$ .

**3.2. Pushing to nearby leaves.** If  $\varepsilon$  is sufficiently small it is smaller than charts on which  $\mathcal{F}_1$  is horizontal and  $\mathcal{F}_2$  vertical and this means that the whole structure of  $\mathcal{G}_L$  that one sees in a leaf  $L$  of (say)  $\widetilde{\mathcal{F}}_1$  can be pushed isotopically to nearby leaves  $L'$  of  $\widetilde{\mathcal{F}}_1$  in the set  $\widehat{D}_\varepsilon(L, I)$ . The roles of the foliations can be reversed.

**Proposition 3.4.** *There exists  $\varepsilon > 0$  and  $C > 0$  such that if  $L, E$  are leaves of  $\widetilde{\mathcal{F}}_1$  which are contained in an interval  $I$  of the leaf space of  $\widetilde{\mathcal{F}}_1$  and  $r$  is a segment of a leaf of  $\widetilde{\mathcal{G}}$  which is contained in  $\widehat{D}_\varepsilon(L, I)$ , as in equation (2-6), such that  $r \subset F \cap L$  where  $F \in \widetilde{\mathcal{F}}_2$ , then, there exists a segment  $r'$  of a leaf in  $\widetilde{\mathcal{G}}$  contained in  $F \cap E$  such that  $\Phi_L^1(r)$  and  $\Phi_E^1(r')$  are at distance less than  $C$ . In particular, if  $r$  is a ray, then the landing sets  $\partial\Phi(r)$ ,  $\partial\Phi(r')$  of  $r$  and  $r'$  coincide.*

*Proof.* This is a direct consequence of transversality of the foliations. If  $\varepsilon$  is small enough, then as long as points remain at distance less than  $\varepsilon$  the intersection of  $F$  with  $L$  and  $E$  will happen in a closeby set. Since the maps  $\Phi_L^1$  and  $\Phi_E^1$  are quasi-isometries and a bounded distance away from the projection to  $\mathbb{H}^2$ , the uniform bound  $C$  is obtained.  $\square$

**3.3. Nonseparated leaves.** Given a leaf  $L \in \tilde{\mathcal{F}}_i$  ( $i = 1, 2$ ) we defined the foliation  $\mathcal{G}_L = \tilde{\mathcal{G}}|_L$ . Denote by  $\mathcal{L}_L = L/\mathcal{G}_L$  its leaf space which is a one-dimensional (possibly) non-Hausdorff manifold.

When  $\mathcal{L}_L$  is non-Hausdorff one has *nonseparated leaves*, that is, distinct leaves  $c_1, c_2 \in \mathcal{G}_L$  which are accumulated by a sequence  $d_n$  of leaves of  $\mathcal{G}_L$  (since these are foliations, this is equivalent to having sequences  $x_n, y_n \in d_n$  such that  $x_n \rightarrow x_\infty \in c_1$  and  $y_n \rightarrow y_\infty \in c_2$ ).

**Remark 3.5.** Two leaves  $c_1, c_2 \in \mathcal{G}_L$  may be separated while there is no transversal to  $\mathcal{G}_L$  which intersects both, in that case, there must be leaves  $e_1, e_2$  (one of which could coincide with  $c_1$  or  $c_2$ ) which are nonseparated and separate  $c_1$  from  $c_2$  in the sense that they lie in different connected components of  $L \setminus \{e_i\}$ .


In principle, there is no a priori relation between  $\mathcal{L}_L$  and  $\mathcal{L}_E$  for different leaves  $L, E$ . However:

**Proposition 3.6.** *Assume that there is some  $L \in \tilde{\mathcal{F}}_i$  ( $i = 1, 2$ ) such that the leaf space  $\mathcal{L}_L$  of  $\mathcal{G}_L$  is Hausdorff. Then, for every  $E \in \tilde{\mathcal{F}}_1$  and  $F \in \tilde{\mathcal{F}}_2$  the leaf spaces  $\mathcal{L}_E$  and  $\mathcal{L}_F$  are Hausdorff.*

To show this, we first need the following useful property (see Figure 5):

**Lemma 3.7.** *Fix  $L \in \tilde{\mathcal{F}}_1$ . The leaf space  $\mathcal{L}_L$  of  $\mathcal{G}_L$  is non-Hausdorff if and only if there is  $E \in \tilde{\mathcal{F}}_2$  such that  $L \cap E$  is not connected. Moreover, if two distinct leaves  $c_1, c_2 \in \mathcal{G}_L$  are nonseparated then, there is  $E \in \tilde{\mathcal{F}}_2$  such that  $c_1 \cup c_2 \subset L \cap E$ .*

*Proof.* Assume first that  $L \cap E$  is not connected, therefore, there are leaves  $c_1, c_2 \in \mathcal{G}_L$  which belong to  $E$ . Assume that the leaf space  $\mathcal{L}_L$  of  $\mathcal{G}_L$  is Hausdorff, this means



**Figure 5.** Intersection in more than one connected component forces non-Hausdorff leaf space of the induced one-dimensional foliation.

that there exists a transversal to  $\mathcal{G}_L$  joining  $c_1$  and  $c_2$ . This transversal is a transversal to  $\tilde{\mathcal{F}}_2$  intersecting  $E$  twice, in contradiction with Corollary 2.2.

Now, assume that  $c_1$  and  $c_2$  are nonseparated leaves in  $\mathcal{G}_L$ . Let  $d_n \in \mathcal{G}_L$  be a sequence of leaves converging to both  $c_1$  and  $c_2$ . It follows that  $d_n \in E_n \cap L$  where  $E_n$  is a sequence of leaves of  $\tilde{\mathcal{F}}_2$ . Consider  $x_n, y_n \in d_n$  such that  $x_n \rightarrow x_\infty \in c_1$  and  $y_n \rightarrow y_\infty \in c_2$ . It follows that  $E_n$  converges to both the leaf through  $x_\infty$  and the leaf through  $y_\infty$ . Since the leaf space of  $\tilde{\mathcal{F}}_2$  is Hausdorff (see Corollary 2.2), these leaves must coincide, and thus  $c_1 \cup c_2 \subset E \cap L$  where  $E \in \tilde{\mathcal{F}}_2$  is the leaf containing  $x_\infty$  (and  $y_\infty$ ).  $\square$

*Proof of Proposition 3.6.* By the previous lemma, if there is a leaf  $L \in \tilde{\mathcal{F}}_1$  for which  $\mathcal{L}_L$  is non-Hausdorff then the same holds for some leaf  $E \in \tilde{\mathcal{F}}_2$  and vice versa.

This implies that it is enough to show that if there is a leaf  $L \in \tilde{\mathcal{F}}_1$  for which  $\mathcal{L}_L$  is non-Hausdorff then the same holds for every  $L' \in \tilde{\mathcal{F}}_1$ .

Since  $\mathcal{F}_1$  is minimal, given an open interval  $I$  in the leaf space  $\mathcal{L}_1 = \tilde{M}/\tilde{\mathcal{F}}_1$  and a leaf  $L' \in \tilde{\mathcal{F}}_1$  there is some deck transformation  $\gamma \in \pi_1(M)$  such that  $\gamma L' \in I$ . Thus, since having non-Hausdorff leaf space is invariant under deck transformations, it is enough to show that there is an interval in the leaf space where every leaf there has non-Hausdorff leaf space.

Take  $L \in \tilde{\mathcal{F}}_1$  and  $E \in \tilde{\mathcal{F}}_2$  such that  $L \cap E$  is not connected and let  $c_1, c_2 \in \mathcal{G}_L$  be two different connected components. Note that  $c_1, c_2$  also belong to  $\mathcal{G}_E$ . Consider a transversal  $\tau : (-\varepsilon, \varepsilon) \rightarrow E$  to  $\mathcal{G}_E$  through some point  $x \in c_1$  (i.e.,  $\tau(0) = x$ ) and denote by  $c_t$  the leaf of  $\mathcal{G}_E$  through the point  $\tau(t)$ . Up to changing orientation in  $\tau$ , we can assume that for  $t > 0$  the leaves  $c_t$  and  $c_2$  belong to different connected components of  $E \setminus c_1$ . This is because  $c_2$  is in one component of  $E \setminus c_1$ . Denote  $L_t$  the leaf of  $\tilde{\mathcal{F}}_1$  through  $\tau(t)$ . Given a point  $y \in c_2$  it follows that for small enough  $t \in (-\varepsilon, \varepsilon)$  the leaf  $L_t$  intersects  $E$  close to  $y$ , thus, in the same connected component of  $E \setminus c_1$  as  $c_2$ . It follows that for small  $t > 0$  the leaf  $L_t$  intersects  $E$  in two connected components and thus  $\mathcal{G}_{L_t}$  has non-Hausdorff leaf space as we wanted to show.  $\square$

**3.4. Nonseparated rays and landing.** If  $c \in \mathcal{G}_L$ , recall that a ray of  $c$  is a connected component of  $c \setminus \{x\}$  for some  $x \in c$  and a ray class is one equivalence class of rays by the relation of one of them being contained in the other; there are exactly two ray classes for each  $c \in \mathcal{G}_L$ .

**Lemma 3.8.** *Let  $c_1, c_2 \in \mathcal{G}_L$  be two nonseparated leaves, then, there is exactly one ray class  $r$  of  $c_1$  and transversals  $\tau_1, \tau_2 : [0, \varepsilon) \rightarrow L$  with  $\tau_i(0) \in c_i$  and an increasing homeomorphism  $\varphi : (0, t_1] \rightarrow (0, t_2]$  for some small  $t_1, t_2$  with the property that if  $0 < t \leq t_1$  and  $e_t$  is the ray class of the leaf of  $\mathcal{G}_L$  through  $\tau_1(t)$  oriented as  $r$  then  $e_t$  intersects  $\tau_2$  at the point  $\tau_2(\varphi(t))$ .*

*Proof.* The existence of such segments is because being nonseparated implies that there are leaves  $e_n \in \mathcal{G}_L$  accumulating in both, so if we fix two transversals  $\tau$  and  $\eta$  to  $c_1$  and  $c_2$  we have segments of  $d_n$  of  $e_n$  intersecting  $\tau$  and  $\eta$  in points  $x_n$  and  $y_n$  and such that  $x_n \rightarrow c_1$  and  $y_n \rightarrow c_2$ . The segments  $d_n$  have length going to infinity.

Assume first that there are segments  $d_n$  and  $d_m$  with opposite orientation or which cut  $\tau$  or  $\eta$  in different connected components of  $\tau \setminus c_1$  or  $\eta \setminus c_2$ . We get a Jordan curve  $\mathcal{J}$  made by  $d_n \cup d_m$  and two segments of  $\tau$  and  $\eta$  joining the points  $x_n$  and  $x_m$  and  $y_n$  and  $y_m$ . Assume without loss of generality that  $x_n$  is closer to  $c_1$  than  $x_m$ . Consider the ray  $r_m$  of  $e_m$  starting from  $x_m$  which does not contain  $d_m$ . By construction,  $r_m$  enters the interior of the curve  $\mathcal{J}$  and since it cannot intersect  $d_n$  or  $d_m$  it must exit  $\mathcal{J}$  intersecting either  $\tau$  or  $\eta$  in some point different from  $x_m$  or  $y_m$  which is a contradiction with Corollary 2.2 (in this case, it is just the Poincaré–Bendixson theorem that is being used since we are arguing inside  $L$ ).

Now, the same argument implies that between two segments  $d_n$  and  $d_m$  every leaf intersecting the transversal  $\tau$  must exit through  $\eta$  thus completing the proof of the lemma by choosing convenient parametrizations  $\tau_1$  and  $\tau_2$  of  $\tau$  and  $\eta$  respectively.  $\square$

Define a *nonseparated ray class* of a leaf  $c \in \mathcal{G}_L$  to be one which is nonseparated from some other leaf in  $\mathcal{G}_L$  as in the previous Lemma. Now we can prove:

**Proposition 3.9.** *Let  $c_1, c_2 \in \mathcal{G}_L$  be nonseparated leaves of  $\mathcal{G}_L$  such that  $c_1 \cup c_2 \subset L \cap E$  with  $L \in \tilde{\mathcal{F}}_1$  and  $E \in \tilde{\mathcal{F}}_2$ . Then, if  $c_1^+$  is the nonseparated ray class of  $c_1$  with  $c_2$  then one has that the positive ray lands and  $\partial^+ \Phi(c_1) = \alpha(E)$ .*

For the following proof, a crucial fact<sup>7</sup> will be that if  $\xi \neq \alpha(L)$  and  $\varepsilon > 0$  is small, then there is a neighborhood  $I$  of  $L$  in the leaf space and a neighborhood  $J$  of  $\xi$  in  $S^1(L)$  such that in  $L \cup S^1(L)$  the set  $\widehat{D}_\varepsilon(L, I)$  contains  $N \cap L$ , where  $N$  is a neighborhood of  $J$  in  $L \cup S^1(L)$  (see Proposition 2.7).

If we assume that  $\alpha(E) \notin \partial^+ \Phi(c_1)$  then it is easier to obtain a contradiction, since we can fix a transversal  $\tau$  to  $c_1^+$  sufficiently close to some  $\xi \in \partial^+ \Phi(c_1)$  so that the full ray from this point is contained in  $\widehat{D}_\varepsilon(E, I)$  and thus, we can show that if  $E'$  is a leaf nearby to  $E$  intersecting  $\tau$  in the direction of  $c_2$ , then the  $E' \cap L$  must follow closely the curve  $c_1$  all along the ray, but on the other hand, since  $c_1$  and  $c_2$  are nonseparated the curve must joint a point close to  $\tau$  to some point in a transversal to  $c_2$ , and since leaves of  $\tilde{\mathcal{F}}_2$  cannot intersect a transversal twice, one gets a contradiction. The main difficulty in what follows is that  $c_1^+$  could approach  $\alpha(E)$  and then we need to argue more carefully to get control on the nearby intersection.

*Proof of Proposition 3.9.* Assume that  $\partial^+ \Phi(c_1) \neq \{\alpha(E)\}$ ; this means that it contains some point  $\xi \neq \alpha(E)$  in  $\partial \mathbb{H}^2$ .

<sup>7</sup>This property is true for general foliations homeomorphic to weak stable foliations of Anosov flows, but not true for general minimal foliations.

Fix a sequence  $x_k \in c_1^+$  such that  $\Phi_L(x_k) \rightarrow \xi$  which also implies that  $\Phi_E(x_k) \rightarrow \xi$  by Lemma 3.1. We can assume by taking some subsequence that for some small  $\varepsilon$  smaller than the one given by Proposition 3.4 and some small intervals of leaves  $I$  of  $\tilde{\mathcal{F}}_2$  containing  $E$  in the boundary we have that  $x_k \in \widehat{D}_\varepsilon(E, I)$ .

We fix a sequence of transversals  $\tau_k : [0, \varepsilon) \rightarrow L$  to  $\mathcal{G}_L$  in  $L$ , through  $c_1$  with  $\tau_k(0) = x_k$  of length  $\varepsilon$  for  $\varepsilon$  as above. We assume that the transversals intersect the component of  $L \setminus c_1$  which contains  $c_2$ . We also fix a transversal  $\eta : [0, \varepsilon) \rightarrow L$  such that  $\eta(0) \in c_2$  as in Lemma 3.8.

Since  $\xi$  is a marker point for  $\tilde{\mathcal{F}}_2$  (i.e.,  $\xi \neq \alpha(E)$ ) by our choice of  $I$  we know that if a leaf  $E' \in I$ , it intersects  $\tau_1([0, \delta))$  for some small  $\delta$ , and by Proposition 3.4 it contains a curve which remains at distance less than  $\varepsilon/2$  from  $E$  and which limits in  $\xi$  (in other words, whose image under  $\Phi_{E'}$  limits in  $\xi$ ). Note that this curve has nothing to do with the ray  $c_1^+$ .

In particular, we know that  $E'$  will intersect  $\tau_k$  for all sufficiently large  $k$  (in fact, Proposition 3.4 also implies that  $E'$  will intersect  $\tau_k$  in points arbitrarily close to  $x_k$ ). Note however that it could be that the intersection of  $E'$  with  $\tau_k$  happens in a different connected component of  $E' \cap L$  since we do not know that  $c_1^+$  stays far from  $\alpha(E)$ . Our goal in what follows is to show that this “splitting” does not take place.

Since  $c_1^+$  is nonseparated from  $c_2$  it follows from Lemma 3.8 that, up to reducing  $\delta$ , for every  $t \in (0, \delta]$  the leaf  $e_t$  of  $\mathcal{G}_L$  through  $\tau_1(t)$  intersects  $\eta$  in some point defining a segment  $d_t$  which joins both transversals.

We claim that for every large  $k$ , every point in  $\tau_k((0, \varepsilon))$  must belong to some  $d_t$ . To see this, first take the segment  $d_{t_1}$  and consider  $k$  so large that  $\tau_k((0, \varepsilon))$  is completely contained in the region of  $L$  bounded by  $c_1^+$ , the arc of the transversals  $\tau_1$  joining  $\tau_1(0)$  with  $\tau_1(t_1)$ ,  $d_{t_1}$ , the arc of  $\eta$  joining  $\eta(0)$  with the intersection point with  $d_{t_1}$  with  $\eta$  and the corresponding ray of  $c_2$  (this bounds a region by Proposition 2.9 and by the choice of the transversals  $\tau_k$  their image is contained in this region). Now, fix some point  $\tau_k(s)$  with  $s \in (0, \varepsilon)$ . By continuity of  $\mathcal{G}_L$  there is  $t \in (0, \delta)$  such that  $e_t$ , the curve of  $\mathcal{G}_L$  through  $\tau_1(t)$  intersects  $\tau_k$  in some  $\tau_k(s')$  with  $s' < s$ . Now consider the region  $\mathcal{J}$  bounded by the Jordan curve formed by  $d_t$ ,  $d_{t_1}$  and the arcs of  $\eta$  and  $\tau_1$  joining their endpoints. Then we get that the curve  $e \in \mathcal{G}_L$  passing through  $\tau_m(s)$  must escape  $\mathcal{J}$  intersecting once the transversal  $\tau_1$  and once the transversal  $\eta$ , and the intersection point with  $\tau_1$  is in a point  $\tau_1(t')$  with  $t < t' < t_1$ , which shows the claim.

For different values of  $t$  the segments  $d_t$  belong to distinct leaves of  $\tilde{\mathcal{F}}_2$ , since each leaf intersects a transversal at most once (see Corollary 2.2). On the other hand, given some  $t$ , since  $d_t$  is compact and the points  $x_k$  escape to infinity, it follows that  $d_t$  cannot intersect  $\tau_k$  for very large  $m$  because  $c_1$  is properly embedded. This contradicts the fact that the leaf  $E' \in \tilde{\mathcal{F}}_2$  through some point  $\tau_1(t)$  described before must intersect  $\tau_k$  for all  $k$ . This contradiction then implies that  $c_1$  cannot limit on  $\xi \neq \alpha(E)$ , or in other words  $\partial^+ \Phi(c_1) = \{\alpha(E)\}$ .  $\square$

**Remark 3.10.** If  $L \in \tilde{\mathcal{F}}_1$  and two curves  $c_1, c_2 \in \mathcal{G}_L$  are nonseparated, then, as proved in Lemma 3.7, they belong to  $L \cap E$  with some  $E \in \tilde{\mathcal{F}}_2$ . However, it should be noted that it does not follow from Lemma 3.7 that  $c_1$  and  $c_2$  must be nonseparated in the leaf space of  $\mathcal{G}_E$  (only that there is no transversal in  $E$  from  $c_1$  to  $c_2$ , recall Remark 3.5) and so it does not follow that  $\alpha(L) = \alpha(E)$ . See Section 7 for concrete examples.

#### 4. Landing of rays

In this section we will show:

**Theorem 4.1.** *Let  $\mathcal{F}_1, \mathcal{F}_2$  be two transverse foliations on  $M = T^1S$  and let  $\mathcal{G}$  denote the foliation obtained by intersection. Then, every ray of a leaf of  $\tilde{\mathcal{G}}$  lands (cf. Definition 3.3).*

The strategy of the proof is as follows. We show that if enough rays land, then every ray should land. In Proposition 3.9 we have shown that if rays are nonseparated then they should land in the nonmarker point of one of the foliations. We will use this to reduce to the case where leaves have Hausdorff leaf space, where landing can also be shown rather directly by “pushing” to close-by leaves.

**4.1. Some rays land then every ray lands.** We will first prove the following:

**Proposition 4.2.** *Assume that there is a leaf  $c \in \mathcal{G}_L$  for some  $L \in \tilde{\mathcal{F}}_i$  ( $i = 1$  or  $2$ ) having one ray that lands in a point which is not  $\alpha(L)$  (i.e.,  $\partial^+ \Phi(c) = \{\xi\} \neq \{\alpha(L)\}$  or  $\partial^- \Phi(c) = \{\xi\} \neq \{\alpha(L)\}$ ). Then, every ray of any leaf of  $\tilde{\mathcal{G}}$  lands in both foliations.*

We will use minimality of the foliation to show that there is a dense set of points in  $S^1(L)$  which are landing points of leaves of  $\mathcal{G}_L$ .

**Lemma 4.3.** *Under the assumptions of Proposition 4.2, for every leaf  $E \in \tilde{\mathcal{F}}_i$  and every  $I \subset \partial \mathbb{H}^2$  nontrivial interval, there is some leaf  $e \in \mathcal{G}_E$  such that either  $\partial^+ \Phi(e) = \{\eta\}$  or  $\partial^- \Phi(e) = \{\eta\}$  for some  $\eta \in I$ .*

*Proof.* Let us work in  $\widehat{M}$ , else, we can apply some deck transformations in the center of  $\pi_1(M)$  so that everything makes sense. Let  $\Gamma = \pi_1(M)/\mathbb{Z}$  where  $\mathbb{Z}$  is the center, notice that  $\Gamma \cong \pi_1(S)$  (cf. Section 2.1).

Up to considering a subinterval of  $I$  we can assume that the closure of  $I$  is disjoint from  $\alpha(E)$ . Consider a deck transformation  $\gamma \in \Gamma$  of  $\widehat{M} \rightarrow M$  such that  $\gamma\xi \in I$  (see Proposition 2.8). Again, up to reducing  $I$  we can assume that  $\gamma\alpha(L) \notin I$ .

Fix some open interval  $J$ , such that the closures of  $J, I$  are disjoint, and in addition such that  $\alpha(E), \gamma\alpha(L)$  are in the interior of  $J$ .

Fix  $\varepsilon > 0$  given by Proposition 3.4. There is a subray  $r \subset c$  landing in  $\xi$  such that the projection of  $\gamma r$  is completely contained in  $\widehat{D}_\varepsilon(E, J)$ . This is because  $\gamma r$  lands in  $\gamma\xi \in I \subset J^c$  (see (2-6)) and  $\alpha(E), \gamma\alpha(L)$  are in  $J$ . Here  $\gamma r \subset \gamma L$  and the pushed through ray is in  $E$  and has ideal point in  $\gamma\xi \in I$ . This finishes the proof.  $\square$

*Proof of Proposition 4.2.* Let  $E \in \tilde{\mathcal{F}}_i$ . Suppose that some ray  $c$  in  $\mathcal{G}_E$  does not land, and let  $I$  be its limit set, which is then not a point.

Let  $J$  be a closed nondegenerate interval whose closure is strictly contained in the interior of  $I$  (possibly  $I = \partial\mathbb{H}^2$ ).

By Lemma 4.3 one can find in  $E$  at least two rays  $e_1, e_2$  of  $\mathcal{G}_E$  landing in different points  $\xi_1 \neq \xi_2$  in the interior of  $J$ . We can further assume that  $e_1, e_2$  are disjoint. Fix a curve joining the starting points of  $e_1$  and  $e_2$  and this defines two regions  $D^+$  and  $D^-$  whose limit sets in  $\partial\mathbb{H}^2$  are the two closed intervals  $[\xi_1, \xi_2]$  and  $[\xi_2, \xi_1]$  in  $\partial\mathbb{H}^2$  with respect to the circular order (cf. Corollary 2.10). Since every ray of  $\mathcal{G}_E$  is properly embedded and rays are disjoint or contained in one another, it follows that up to removing a compact piece, then the ray  $c$  must belong to either  $D^+$  or  $D^-$ , and so can only accumulate on  $[\xi_1, \xi_2]$  or  $[\xi_2, \xi_1]$ . This contradicts the fact that  $c$  accumulates on all of  $I$ , and  $I$  is an interval intersecting the interior of both  $[\xi_1, \xi_2]$  and  $[\xi_2, \xi_1]$ . This proves the proposition.  $\square$

The following stronger statement will be used later:

**Lemma 4.4.** *Let  $e \subset L \cap E$  be such that  $\partial^+\Phi(e) = I$  where  $I$  is not a point (in other words, the positive ray of  $e$  does not land). Then  $\alpha(L) \in I$ . In addition if  $I \neq \partial\mathbb{H}^2$ , then for every  $\xi$  in the interior of  $I$  there is no ray of a leaf of  $\mathcal{G}_E$  which lands in  $\xi$ . The same holds for  $\partial^-\Phi(e)$ .*

*Proof.* In this lemma we also work in  $\widehat{M}$ , but the arguments can be easily adapted to do the same in  $\widetilde{M}$ . All leaves in this lemma are in the same foliation. First notice that if  $I = \partial\mathbb{H}^2$  then there is nothing to prove.

First assume that  $\alpha(L) \notin I$ . Consider  $\gamma \in \pi_1(M)$  such that  $\gamma$  acting on  $\partial\mathbb{H}^2$  has a repelling fixed point  $\gamma^- \notin I$  (and close to  $\alpha(L)$ ) and an attracting fixed point  $\gamma^+ \in I$  (Proposition 2.8).

Now, consider the leaf  $E'$  associated to the repelling fixed point, that is, such that  $\alpha(E') = \gamma^-$ . It follows that this leaf is invariant under  $\gamma$ . Consider a ray  $r$  of  $e$ , such that  $\partial\Phi(r) = I$  which is contained in  $\widehat{D}_\varepsilon(E, J)$  where  $J$  is some interval of the leaf space containing  $E'$  and  $E$  and  $\varepsilon > 0$  given by Proposition 3.4. Using Proposition 3.4 we deduce that there is a ray  $r'$  of a leaf of  $\mathcal{G}_{E'}$  such that  $\partial\Phi(r) = I$ . Moreover,  $\gamma r'$  is a ray in  $\mathcal{G}_{E'}$  with  $\partial\Phi(\gamma(r)) = \gamma(I)$ , which is strictly contained in  $I$ . This contradicts Proposition 2.11.

The second statement follows immediately from Proposition 2.11.  $\square$

**4.2. Finding landing rays.** In this section we will show the following:

**Proposition 4.5.** *Let  $L \in \tilde{\mathcal{F}}_1$  and  $\gamma \in \pi_1(M) \setminus \{\text{id}\}$  such that  $\gamma L = L$ . Then, there is some  $c \in \mathcal{G}_L$  which has at least one ray which lands.*

We consider a closed curve in the annulus  $A = L/\langle\gamma\rangle$  not homotopically trivial (for instance take a closed geodesic in  $A$  with some metric in  $L$  induced by  $M$ ) and

denote by  $\ell$  a lift to  $L$  of this closed curve, which is thus  $\gamma$  invariant, that is  $\gamma\ell = \ell$ . For  $K > 0$  we denote as  $B_K(\ell)$  the closed tubular neighborhood of radius  $K$  around  $\ell$  for some  $\gamma$ -invariant metric on  $L$ ; note that the quotient  $B_K(\ell)/\langle\gamma\rangle$  is compact.

**Lemma 4.6.** *Assume that  $\gamma$  does not fix any leaf of  $\mathcal{G}_L$ . For every  $K > 0$  there is  $K' > 0$  such that if  $c \in \mathcal{G}_L$  then any connected component of  $c \cap B_K(\ell)$  has length less than  $K'$ .*

*Proof.* Assume, toward a contradiction, that there are sequences of arcs  $d_n$  of leaves  $c_n \in \mathcal{G}_L$  of length going to infinity such that  $d_n \subset B_K(\ell)$ . There is a sequence  $\gamma^{k_n}$  of iterates of  $\gamma$  with  $k_n \in \mathbb{Z}$  such that  $\gamma^{k_n}d_n$  has its midpoint in a compact fundamental domain of  $B_K(\ell)$  which is  $\gamma$ -invariant by choice. It follows that in the limit,  $\gamma^{k_n}d_n$  converges to at least one leaf  $c_\infty \in \mathcal{G}_L$  which is completely contained in  $B_K(\ell)$ .

Let  $\pi_\gamma : L \rightarrow L/\langle\gamma\rangle$  be the projection. Then  $\pi_\gamma(c_\infty)$  is contained in a compact annulus. It follows that each ray of  $c_\infty$  either projects to a closed curve or to a curve asymptotic to a closed curve. This is a contradiction to hypothesis.  $\square$

Note that if there is a leaf  $c \in \mathcal{G}_L$  which is fixed by  $\gamma$  then immediately we have that the landing points of  $c$  are  $\partial^\pm\Phi_L(c) = \{\gamma^+, \gamma^-\}$  the attractor and repeller of  $\gamma$  acting at infinity.

Now we show that under the assumptions of the previous lemma, rays must keep intersecting  $B_K(\ell)$  indefinitely.

**Lemma 4.7.** *Under the assumptions of Lemma 4.6 we have that there is  $K > 0$  such that if  $c \in \mathcal{G}_L$  and  $r$  is a ray of  $c$  that does not land, then  $r$  must intersect  $B_K(\ell)$ .*

We stress that  $K$  is independent of  $c$  and  $r$ . Note that since  $r$  cannot be contained in  $B_K(\ell)$  by the previous lemma, we deduce that every ray needs to enter and leave  $B_K(\ell)$  infinitely many times.

*Proof.* Fix  $\varepsilon > 0$  given by Proposition 3.4. Now fix some large  $K > 0$  so that if  $I$  is a closed interval of the leaf space of  $\tilde{\mathcal{F}}_1$  containing  $L$  as an endpoint, then  $\widehat{D}_\varepsilon(L, I)$  contains the connected component of the complement of  $B_K(\ell)$  which does not accumulate in  $I$  (see Proposition 2.7). Note that this value of  $K$  is independent of the interval  $I$  as long as  $L$  is an endpoint and  $I$  is sufficiently small (so that it is contained in the closure of one of the components of the complement of  $\ell$ ).

Assume for a contradiction that  $r$  is completely contained in the complement of  $B_K(\ell)$  which then must be contained in a unique connected component because  $r$  is connected. Therefore, the limit set  $I = \partial\Phi_L(r)$  is a proper interval contained in  $\partial\mathbb{H}^2$  which does not intersect the interior of the limit set of the image by  $\Phi_L$  of the connected component of  $L \setminus B_K(\ell)$  which does not contain  $r$ . Now pick an interval  $J$  of the leaf space with  $L$  as an endpoint so that  $\alpha(E) \in J$  only if  $E = L$ . For any  $E \neq L$  in  $I$  we can then push the ray  $r$  to  $E$  applying Proposition 3.4 and

we deduce that there is a ray  $r'$  in  $E$  which lands in an interval which is disjoint from  $\alpha(E)$  contradicting Lemma 4.4.  $\square$

*Proof of Proposition 4.5.* As noted, if  $\gamma$  fixes some leaf of  $\mathcal{G}_L$  then this leaf must land in the attractor and repeller of  $\gamma$  so the proposition holds. Therefore, we can assume we are under the assumptions of Lemma 4.6.

Let  $c$  be a leaf of  $\mathcal{G}_L$  and  $r$  a ray of  $c$ . If  $r$  lands, the Proposition is proved, so assume that  $r$  does not land. Then we know that  $r$  must keep intersecting  $B_K(\ell)$  for  $K$  as in Lemma 4.7. Consider a sequence  $d_n$  of segments of  $r$  completely contained in the complement of  $B_K(\ell)$  except for the endpoints which belong to  $\partial B_K(\ell)$ . Using Lemma 4.6 we can choose the segments  $d_n$  with arbitrarily large length, since otherwise we would get arbitrarily large segments of  $r$  contained in  $B_{K'}(\ell)$  for some larger  $K'$ .

Let  $x_n, y_n$  be the endpoints of  $d_n$ . We claim first that  $d(x_n, y_n)$  must be bounded: if not, then up to considering an inverse we can assume that  $\gamma x_n$  is contained in the segment  $J$  of  $\partial B_K(\ell)$  joining  $x_n$  and  $y_n$ . Since  $d_n \cup J$  is a Jordan curve and  $\gamma y_n$  does not belong to  $J$  we deduce that  $\gamma d_n$  intersects  $d_n$  which is not possible since they belong to different leaves of  $\mathcal{G}_L$ .

There is a sequence  $k_n \in \mathbb{Z}$  such that  $\gamma^{k_n} x_n$  belongs to a compact set. Thus, up to taking subsequences both  $\gamma^{k_n} x_n$  and  $\gamma^{k_n} y_n$  converge to points  $x_\infty$  and  $y_\infty$  in  $\partial B_K(\ell)$ . Note that  $x_\infty$  and  $y_\infty$  cannot be in the same leaf of  $\mathcal{G}_L$  and being accumulated by  $\gamma^{k_n} d_n$  their leaves must be nonseparated. Therefore the proposition follows from Proposition 3.9.  $\square$


**4.3. Proof of Theorem 4.1.** To prove Theorem 4.1 we want to apply Proposition 4.2. For this, it will be useful to use both foliations. We first show that if there is a leaf of  $\mathcal{G}_L$  which has both rays landing in the same point, then after going to a closeby leaf of the other foliation, we will be able to apply Proposition 4.2 and we will get landing for all leaves in both foliations:

**Lemma 4.8.** *Let  $L \in \tilde{\mathcal{F}}_1$  be such that there is  $c \in \mathcal{G}_L$  both of whose rays land in the same point, i.e.,  $\partial^+ \Phi_L^1(c) = \partial^- \Phi_L^1(c) = \{\xi\}$ . Then, there is some leaf  $E \in \tilde{\mathcal{F}}_2$  with  $\alpha(E) \neq \xi$  such that it contains a leaf  $c' \in \mathcal{G}_E$  with  $\partial^+ \Phi_E^1(c') = \partial^- \Phi_E^1(c') = \{\xi\}$ .*

*Proof.* Consider the Jordan curve obtained by  $c \cup \{\xi\}$  in  $L \cup S^1(L)$  and denote by  $D$  the disk bounded by it.

Every curve  $\hat{c} \in \mathcal{G}_L$  intersecting  $D$  must be contained in  $D$  and thus has both rays landing on  $\xi$ . Given a transversal  $\tau : [0, t_0) \rightarrow L$  to  $\mathcal{G}_L$  with  $\tau(0) \in c$  and  $\tau(t) \in D$  for  $t > 0$  we denote  $E_t \in \tilde{\mathcal{F}}_2$  to the leaf through  $\tau(t)$ .

If we denote by  $c_t$  the curve of  $\mathcal{G}_L$  containing  $\tau(t)$  we get  $c_t \subset L \cap E_t$ . Since in  $L$  both rays of  $c_t$  land and the landing point is  $\xi$ , we get by Corollary 3.2 that for every small  $t$ , then  $c_t$  lands in  $E_t$  and the landing point is  $\xi$ . Since the nonmarker point  $\alpha(E_t)$  varies monotonically we can choose  $t$  so that  $\alpha(E_t) \neq \xi$ .  $\square$



**Figure 6.** A depiction of the objects in the proof of Lemma 4.9.

The argument can be slightly extended to get:

**Lemma 4.9.** *Let  $c \in \mathcal{G}_L$  satisfy  $\partial^+ \Phi_L(c) = \alpha(L)$  and  $\partial^- \Phi_L(c) \neq \partial \mathbb{H}^2$ . Then all rays of  $\tilde{\mathcal{G}}$  land.*

*Proof.* Let  $I = \partial^- \Phi_L(c)$ . Suppose first that  $I = \alpha(L)$ , then both rays of  $c$  land in the same point  $\alpha(L)$ . The previous lemma implies that there is some ray of  $\tilde{\mathcal{G}}$  in some leaf  $E$  of  $\tilde{\mathcal{F}}_2$  landing in a point different from  $\alpha(E)$ . Then Proposition 4.2 implies that all rays of  $\tilde{\mathcal{G}}$  land and the lemma is proved.

If  $I$  is a single point  $\xi \neq \alpha(L)$ , then again Proposition 4.2 also implies that all rays land, and the lemma is also proved.

Finally from now on assume that  $I$  is not a single point. Since  $I \neq \partial \mathbb{H}^2$ , Lemma 4.4 implies that  $\alpha(L) \in I$ .

The curve  $c$  separates  $L$  in two connected components, each diffeomorphic to a disk (see Proposition 2.9). Denote these components by  $D^+$  and  $D^-$ . It follows that up to relabeling we can assume that the closure of  $\Phi_L(D^+)$  in  $\mathbb{H}^2 \cup \partial \mathbb{H}^2$  is equal to  $\Phi_L(c \cup D^+) \cup I$  while the closure of  $\Phi_L(D^-)$  in  $\mathbb{H}^2 \cup \partial \mathbb{H}^2$  is equal to  $\Phi_L(c \cup D^-) \cup \partial \mathbb{H}^2$ . See Figure 6.

Consider a transversal  $\tau : (-\varepsilon, \varepsilon) \rightarrow L$  to  $\mathcal{G}_L$  with  $\tau(0) \in c$ . Denote by  $c_t$  the leaf of  $\mathcal{G}_L$  through the point  $\tau(t)$ . We can assume that if  $t > 0$  then  $\tau(t) \in D^+$ , thus, the limit set  $\partial^\pm \Phi_L(c_t) = I_t$  of  $c_t$  must be contained in  $I$ . Again as in the beginning of the proof we either finish the proof of the lemma or we are in the case that, the interval  $I_t$  is nondegenerate for any  $t > 0$ . Hence  $I_t$  contains  $\alpha(L)$  by Lemma 4.4. In addition  $I_t$  is weakly monotonically decreasing, meaning that if  $t > t'$  then  $I_t \subset I_{t'}$  (they could coincide).

Denote by  $E_t$  the leaf of  $\tilde{\mathcal{F}}_2$  such that  $c_t \subset L \cap E_t$ . Then  $\alpha(E_0) = \alpha(L)$ ; otherwise

again Proposition 4.2 would imply that all rays of  $\tilde{\mathcal{G}}$  land in their appropriate leaves, but at this point we are assuming that the negative ray of  $c$  does not land. This contradiction shows that  $\alpha(E_0) = \alpha(L)$ .

In addition Lemma 4.4, applied to  $\mathcal{F}_2$ , implies that  $\alpha(E_t)$  is contained in  $I_t$ . For small  $t > 0$ ,  $\alpha(E_t)$  is contained in the interior of  $I$ . Notice that  $\alpha(E_t)$  varies continuously with  $t$ . We fix some small  $t_0 > 0$ . Now fix a  $t_1$  with  $0 < t_1 < t_0$  and such that  $E_{t_1}$  is invariant under some deck transformation  $\gamma$  which has a fixed point outside  $I$  (such leaves are dense, so there must be one). The other fixed point of  $\gamma$  is necessarily  $\alpha(E_{t_1})$ . We can arrange that this point is contained in the interior of  $I_{t_0} \subset I_{t_1} \subset I$ . This is because by choosing  $t_1$  small we can ensure that  $\alpha(E_{t_1})$  is in the interior of  $I_{t_0}$ , and since  $I_{t_0} \subset I_{t_1}$ , the claim follows.


We get that  $c_{t_1}$  has a ray converging to an interval  $I_{t_1}$  which contains only one of the fixed points of  $\gamma$ . We proved before that  $\alpha(L) = \alpha(E_0) \in I_t$  for any  $t$ . In addition since  $I_{t_0} \subset I_{t_1}$  we get that  $\alpha(E_{t_0})$  is also in  $I_{t_1}$ , so the interval  $[\alpha(E_0), \alpha(E_{t_0})]$  contained in  $I$  is contained in  $I_{t_1}$ . We conclude that  $\alpha(E_{t_1})$  is an interior point of  $I_{t_1}$ , and  $I_{t_1}$  does not contain the other fixed point of  $\gamma$ . Applying  $\gamma$  to  $E_{t_1}$  we obtain two rays whose landing sets are proper intervals one contained in the interior of the other ( $I_{t_1}$  and  $\gamma(I_{t_1})$ ), contradicting Proposition 2.11 and completing the proof.  $\square$

*Proof of Theorem 4.1.* Using Proposition 4.5 we know that there is at least one  $L \in \tilde{\mathcal{F}}_1$  and one  $c \in \mathcal{G}_L$  having one landing ray. Up to orientation we can assume that  $\partial^+ \Phi_L(c) = \xi$ . Using Proposition 4.2, either the result holds or  $\xi = \alpha(L)$ . Using Lemma 4.9 we can further assume that  $\partial^- \Phi_L(c) = \partial \mathbb{H}^2$  (it is the only possibility of that lemma that does not yield that all rays land), in particular this assumption means that there is a ray of  $\tilde{\mathcal{G}}$  that does not land.

Note that all these results apply to both  $\tilde{\mathcal{F}}_1$  and  $\tilde{\mathcal{F}}_2$  so we will now work with  $\tilde{\mathcal{F}}_2$ . Showing landing in both foliations is equivalent because of Lemma 3.1.

Let  $\tau : (-\varepsilon, \varepsilon) \rightarrow L$  be a transversal to  $\mathcal{G}_L$  with  $\tau(0) \in c$ . Call  $c_t$  the leaf of  $\mathcal{G}_L$  passing through  $\tau(t)$  and  $E_t \in \tilde{\mathcal{F}}_2$  such that  $c_t \subset L \cap E_t$ . Notice that the  $\{E_t\}$  are pairwise distinct leaves.

We know that  $\alpha(E_0) = \alpha(L)$  because  $\partial^+ \Phi_{E_0}(c) = \partial^+ \Phi_L(c) = \alpha(L)$  (else we can apply Proposition 4.2 to  $\tilde{\mathcal{F}}_2$ ). We choose the transversal  $\tau$  so small that  $\alpha(E_t)$  is injective in  $(-\varepsilon, \varepsilon)$  and  $\alpha(E_t)$  is very close to  $\alpha(L)$ . Let  $I_t$  be the short closed interval in  $\partial \mathbb{H}^2$  from  $\alpha(L)$  to  $\alpha(E_t)$  and  $J_t$  the closure of the complementary interval in  $\partial \mathbb{H}^2$ . Note that for  $t > 0$  since  $c_t \in L \cap E_t$ , if it has a landing ray, then Proposition 4.2 applied to both foliations implies it must land  $\alpha(L)$  and in  $\alpha(E_t)$  which are different, so the limit set of both rays is a nontrivial connected set. Lemma 4.4 applied to both foliations implies that the limit set of each ray of  $c_t$  must contain both  $\alpha(L)$  and  $\alpha(E_t)$  thus it is some nontrivial connected set containing  $I_t$  or  $J_t$ .



**Figure 7.** A depiction of some parts of the argument of the proof of Theorem 4.1: inclusion of sets  $D_s \subset D_t$  and the “pairing” configuration for rays.

For any  $E_t$  which is a deck translate of  $E_0$  there is a ray of  $\mathcal{G}_{E_t}$  which lands (i.e., the image of the landing ray of  $c_0$ ). By Proposition 4.2 again, it follows that this ray must land in  $\alpha(E_t)$ . Let  $r_t$  be one such ray. Consider

$$\mathcal{B} = \{t \in (-\varepsilon, \varepsilon) : E_t = \gamma E_0 \text{ for some } \gamma \in \pi_1(M)\}.$$

By minimality  $\mathcal{B}$  is dense in  $(-\varepsilon, \varepsilon)$ . We will consider  $\mathcal{B}^+ = \mathcal{B} \cap (0, \varepsilon)$ .

Consider  $t \in \mathcal{B}$  and let  $g_t$  be a ray of  $c_t$ . Denote by  $I_g$  its limit set (that is  $\partial\Phi_{E_t}(g_t)$ ) in  $\partial\mathbb{H}^2$ . Recall that  $I_g$  must contain  $I_t$  or  $J_t$  (in particular, it not a point). The ray  $g_t$  is disjoint from  $r_t$ , because  $r_t$  lands and no ray of  $c_t$  lands. If  $I_g \neq \partial\mathbb{H}^2$ , then applying Lemma 4.4 (first to  $L$  and  $c_0^+$  and then to  $E_t$  and  $r_t$ ) we deduce that  $I_g$  coincides with either  $I_t$  or  $J_t$  because there cannot be a ray landing in its interior. If  $I_g = \partial\mathbb{H}^2$  let us show a contradiction with a similar argument: in  $L$  the ray  $g_t$  has to be disjoint from the ray of  $c_0$  which lands in  $\alpha(L)$ , in particular there cannot be a sequence of segments in  $g_t$  which limit to an interval in  $S^1(L)$  with  $\alpha(L)$  in its interior. When seen in  $E_t$  then  $g_t$  has to be disjoint from  $r_t$ , so again it cannot have a sequence of segments with a limit which is an interval in  $S^1(E_t)$  with  $\alpha(E_t)$  in its interior. It follows that  $I_g$  could not be all of  $\partial\mathbb{H}^2$  and thus  $I_g$  is either  $I_t$  or  $J_t$ .

Recall that  $c_0^+$  is a ray which lands in  $\alpha(L)$ . Let  $D_t$  be the component of  $L \setminus c_t$  which does not contain  $c_0 = c$ , in particular  $D_t$  is an open set in  $L$  which has boundary (in  $L$ ) equal to  $c_t$ . Notice that if  $s > t > 0$ , then  $D_s \subset D_t$ , and in particular that  $c_s$  is contained in  $D_t$ . For each  $t > 0$  let  $c_t^+, c_t^-$  be the subrays of  $c_t$  determined by  $\tau(t)$  and oriented coherently with  $c_0$ .

Consider  $t \in \mathcal{B}^+$ . We know that  $\partial\Phi_L(c_t^-)$  can only be  $I_t$  or  $J_t$  for each such  $t$ . There are also two possibilities depending on where  $\alpha(E_t)$  is (so in total, four possibilities). We say that  $\alpha(E_t)$  and  $\tau(t)$  cross if we consider a ray  $r$  from  $\tau(0)$  to


$\xi \in J_t$  avoiding  $c_0^+ \cup \tau([0, t])$  and there for every curve in  $L \cup S^1(L)$  joining  $\tau(t)$  with  $\alpha(E_t)$  the curve intersects  $c_0^+ \cup r$ . Note that this is independent on the choice of  $r$ , and it is also independent on  $t > 0$ . We say they *pair* otherwise (i.e., there is a curve joining  $\tau(t)$  with  $\alpha(E_t)$  and avoiding  $c_0^+ \cup r$ ).

We suppose first that  $\tau(t), \alpha(E_t)$  pair, see this in Figure 7. Suppose first that for some  $t \in \mathcal{B}^+$  we have that  $\partial\Phi_L(c_t^-) = I_t$ . In this case, note that the limit of  $\Phi_L(D_t)$  is also  $I_t$ . For any  $s > t$  we have  $D_s \subset D_t$ , so we now obtain that  $\partial\Phi_L(c_s^-) \subset I_t$ . But for such  $s$  the limit set of  $c_s^-$  has to contain  $\alpha(E_s)$  which is not contained in  $I_t$  a contradiction. See Figure 7.


Now assume that  $\tau(t), \alpha(E_t)$  pair but  $\partial\Phi_L(c_t^-) = J_t$ . Then  $c_t^-$  has points which limit to  $\alpha(E_t)$  but  $c_t^-$  does not limit in any point in the interior of  $I_t$ . For any  $s > t$  in  $\mathcal{B}^+$  we cannot have that  $\partial\Phi_L(c_s^-)$  is  $J_s$ . This is because the ray  $c_t^-$  limiting to  $\alpha(E_t)$  forces the ray  $c_s^-$  to limit to some point in the interior of  $I_s$ . Thus, we can apply the previous analysis to  $s$  to get a contradiction. This finishes the analysis in the case where the transversal  $\tau$  pairs.

We now consider the case where  $\tau(t), \alpha(E_t)$  cross. Suppose first that for some  $t \in \mathcal{B}^+$  the ray  $c_t^-$  limits to  $I_t$ . We depict this in Figure 8, left. Then for  $s < 0$ , it follows that the limit sets of both  $c_s^+, c_s^-$  have to be contained in  $I_t$ . This property is impossible, because for  $s < 0$  contained in  $\mathcal{B}$  the limit sets would have to be either  $I_s$  or  $J_s$  and neither is contained in  $I_t$ .

The remaining case is that for any  $t \in \mathcal{B}^+$  we have that  $\tau(t), \alpha(E_t)$  cross and  $\partial\Phi_L(c_t^-) = J_t$ . Fix some  $t \in \mathcal{B}^+$ . We will need to consider a different disk  $Z_t$  in  $L$  whose boundary is made up of  $c_t^-$ , the arc  $\tau([0, t])$  and  $c_0^+$  and chosen so that it contains  $c_t^+$ . Notice that this disk limits only on  $J_t$  (see Figure 8, right). Now consider  $s$  with  $0 < s < t$ , and  $s \in \mathcal{B}^+$ . Then  $c_s^+$  is contained in  $Z_t$ , so its limit set



**Figure 8.** A depiction of the cases that  $\tau(t), \alpha(t)$  cross. Left: the case that the limit set of  $c_t^-$  is  $I_t$ . Right: the case that the limit set is  $J_t$ . For simplicity of viewing we depict  $\partial\mathbb{H}^2$  in a line segment, concentrating on what is happening near  $\alpha(L)$ .



**Figure 9.** The shadow of a set from a point  $x \in L$  is the set of points in  $S^1(L)$  blocked by this set (when seen in  $\mathbb{H}^2$ ). To have small visual measure means that if an arc of  $\mathcal{G}_L$  is very far from  $x$  then its shadow has to be very small (independently of its length).

has to be also contained in  $J_t$ . On the other hand it has to be either  $J_s$  or  $I_s$  but neither is contained in  $J_t$ , which is a contradiction.

This finishes the proof of Theorem 4.1.  $\square$

## 5. Small visual measure

Let  $\mathcal{F}_1, \mathcal{F}_2$  be two transverse minimal foliations on  $M = T^1S$  and  $\mathcal{G} = \mathcal{F}_1 \cap \mathcal{F}_2$  the intersection foliation.

We define the *shadow* of a subset  $X \subset L$  seen from  $x \in L$  as the set of points  $\xi \in \partial\mathbb{H}^2$  for which there is a geodesic ray in  $\mathbb{H}^2$  starting at  $\Phi_L(x)$  and passing through some point  $y \in \Phi_L(X)$  with ideal point  $\xi$ .

As in [25, § 4.3] we will say that  $\mathcal{G}$  has the *small visual measure* property if for every  $\varepsilon > 0$  there exists  $R > 0$  such that if  $L \in \tilde{\mathcal{F}}_i$ ,  $x \in L$  and  $\ell$  a segment of some leaf of  $\mathcal{G}_L$  such that  $\ell \cap B_R(x) = \emptyset$ , then the shadow of  $\ell$  seen from  $x$  has length smaller than  $\varepsilon$  (meaning that the angle of the interval of vectors such that geodesic rays in  $\mathbb{H}^2$  from  $\Phi_L(x)$  to the shadow of  $\ell$  is less than  $\varepsilon$ ).

In this section we will show:

**Proposition 5.1.** *Let  $\mathcal{F}_1, \mathcal{F}_2$  be two transverse minimal foliations on  $M = T^1S$  and let  $\mathcal{G}$  denote the foliation obtained by intersection. Then,  $\mathcal{G}$  has the small visual measure property.*

Before we proceed with the proof let us explore some of its consequences. In [25, Lemma 5.9] the following is proved.

**Proposition 5.2.** *Assume that  $\mathcal{G}$  has the small visual measure property, then, there exists  $a_0 > 0$  such that for every segment  $\ell$  of leaf of  $\mathcal{G}_L$  we have that the geodesic segment joining the endpoints of  $\ell$  is contained in  $B_{a_0}(\ell)$ .*

We stress that  $B_{a_0}(\ell)$  is the neighborhood of  $\ell$  of size  $a_0$  in  $L$ , rather than in  $\tilde{M}$ .

Note that from this we get that small visual measure allows to control endpoints of rays of leaves of  $\mathcal{G}_L$ . That is, if one considers a point  $x \in L$  which belongs to some leaf  $c \in \mathcal{G}_L$  and considers points  $y_n$  in  $c_x^+$  going to infinity. It follows that the geodesic segments joining  $x$  and  $y_n$  are all contained in the  $a_0$  neighborhood of  $c_x^+$ , thus, it follows that the geodesic ray joining  $x$  with  $\partial^+\Phi(c)$  is contained in the  $a_0$ -neighborhood of  $c_x^+$ .

### 5.1. Bubble leaves.

**Definition 5.3.** (bubble leaves) We say that  $c \in \tilde{\mathcal{G}}$  is a *bubble leaf* if  $\partial^+\Phi(c) = \partial^-\Phi(c) = \{\xi\}$  where  $\xi$  is some point in  $\partial\mathbb{H}^2$ .

We have dealt with such leaves in Lemma 4.8 showing that nearby leaves of the other foliation must also have bubble leaves with the same endpoint. We will perform a similar argument now, but trying to control the size of the interval on the leaf space where this holds.

The goal of this subsection is to show the following:


**Proposition 5.4.** *There exists a leaf  $c \in \tilde{\mathcal{G}}$  such that  $\partial^+\Phi(c) \neq \partial^-\Phi(c)$  (see (3-2)).*

We will need to control the place where the bubble leaves land. For this, we will separate the leaf  $L$  in *bubble regions*. To introduce this, let us first make some definitions. Notice first that by Corollary 2.10 we know that if  $c \in \mathcal{G}_L$  is a bubble leaf with  $\partial^\pm\Phi(c) = \{\xi\}$  we can define  $D_c$  to be the disk in the complement of  $c$  such that the accumulation of  $\Phi_L(D_c)$  in  $\partial\mathbb{H}^2$  is exactly  $\xi$ .

**Definition 5.5.** (bubble region of  $c$ ) Given  $L \in \tilde{\mathcal{F}}_i$  and  $c \in \mathcal{G}_L$  a bubble leaf with  $\partial^\pm\Phi(c) = \{\xi\}$  we denote by  $\mathcal{B}(c, L)$  the *bubble region of  $c$*  in  $L$  as the union of all leaves  $c' \in \mathcal{G}_L$  such that there is some  $c'' \in \mathcal{G}_L$  which is a bubble leaf and is such that  $D_c \cup D_{c'} \subset D_{c''}$ . We call  $\xi$  the *landing point* of the bubble region.

We also remark that in general the bubble region of  $c$  is not the union of all bubble leaves  $c' \in \mathcal{G}_{L_i}$  such that  $\partial\Phi^\pm(c') = \{\xi\}$ . For example it could be that  $c, c'$  bound maximal disjoint disks  $D_c, D_{c'}$ . By maximal for  $c$  we mean there is no leaf  $c''$  of  $\mathcal{G}_{L_i}$  distinct from  $c$ , and such that  $D_{c''} \supset D_c$ . In this case  $D_c, D_{c'}$  are two disjoint bubble regions with the same ideal point  $\xi$ .

**Lemma 5.6.** *Every bubble region is either open or closed. Each leaf contains at most countably many distinct bubble regions. If every leaf in  $\mathcal{G}_L$  is a bubble leaf, then there is a unique open bubble region  $\mathcal{B}$  such that  $\Phi_L(\mathcal{B})$  accumulates in all of  $\partial\mathbb{H}^2$ .*



**Figure 10.** Depiction of a leaf  $L$  such that every leaf of  $\mathcal{G}_L$  is a bubble leaf. The painted regions are the closed bubble regions and their complement is the open bubble region.

In Figure 10 a depiction of the conclusion of this lemma is presented.

*Proof.* Each bubble region is a set saturated by  $\mathcal{G}_L$ . There is a natural partial order between the leaves in a bubble region given by  $c_1 < c_2$  if  $D_{c_1} \subset D_{c_2}$ . It follows that if there is a maximal element in the bubble region, then it is unique (by definition of bubble region, every pair of elements is smaller or equal than some element in the region) and the bubble region is closed, while if there is no maximal element then the bubble region is open. As bubble regions have nonempty interior we know that there are at most countably many bubble regions.

Assume now that every leaf of  $\mathcal{G}_L$  belongs to some bubble region. If there is a unique bubble region, then it is an open bubble region since there cannot be a maximal element. We next prove that not every bubble is closed: Let  $B$  be a closed bubble with boundary leaf  $c$  and let  $\tau$  be a small transversal to  $\mathcal{G}$  in  $L$  with an endpoint in  $c$  and not intersecting the interior of  $B$ . Parametrize  $\tau$  as  $x_t$ ,  $0 \leq t \leq 1$  with  $x_0$  in  $c$ . Suppose a bubble region  $B'$  intersects  $\tau$  not in  $x_0$ . Then its boundary intersects  $\tau$  (as  $x_0$  is not in  $B'$ ). In addition its boundary intersects  $\tau$  in a single point, because if it intersected  $\tau$  in two points, one would produce a bubble leaf intersecting  $\tau$  in two points, contradiction. It follows that there is a single bubble region  $B'$  containing  $\tau \setminus \{x_0\}$  and  $B'$  is an open bubble.

Finally, given an open bubble region  $\mathcal{B}$  we can consider  $\widehat{\mathcal{B}}$  to be  $\mathcal{B}$  together with all the closed bubbles whose boundary intersects the boundary of  $\mathcal{B}$ . It follows that

$\widehat{\mathcal{B}}$  is an open set and that if  $\mathcal{B}'$  is another open bubble region, then  $\widehat{\mathcal{B}} \cap \widehat{\mathcal{B}}' = \emptyset$ , thus, we get there is a unique open bubble region.

To conclude, using Corollary 2.10 we know that the complement of a closed bubble region accumulates in all of  $S^1(L)$ , so the same holds for the complement of countably many closed bubble regions, thus the open bubble region accumulates everywhere, also using the properties derived above.  $\square$

We remark that the uniqueness of open bubble regions in a leaf  $L$  of  $\widetilde{\mathcal{F}}_i$  does not necessarily work if not every leaf in  $\mathcal{G}_L$  is a bubble leaf.

Since open bubble regions are special, it is natural to expect that their landing point will also be special:

**Lemma 5.7.** *If every leaf in  $\widetilde{\mathcal{G}}$  is a bubble leaf then in each  $L \in \widetilde{\mathcal{F}}_i$  the unique open bubble region of  $\mathcal{G}_L$  has  $\alpha(L)$  as its landing point.*

*Proof.* The proof is by contradiction. Assume that there is a leaf  $L \in \widetilde{\mathcal{F}}_i$  where the unique open bubble region  $\mathcal{O}$  has landing point  $\xi \neq \alpha(L)$ . We fix a closed interval  $I$  in the leaf space of  $\widetilde{\mathcal{F}}_i$  containing  $L$  in its interior and such that for every  $L' \in I$  we have that  $\alpha(L') \neq \xi$ . Denote by  $J \subset \partial\mathbb{H}^2$  the interval  $J = \{\alpha(L') : L' \in I\}$ . We consider  $\varepsilon > 0$  from Proposition 3.4 and fix the region  $\widehat{D}_\varepsilon(L, I)$ .

We first notice that Lemma 4.3 shows that the existence of a leaf whose landing point is  $\xi \neq \alpha(L)$  implies that in each leaf  $L$  the set of landing points of rays in the leaf is dense in  $\partial\mathbb{H}^2$ . Thus, for a given leaf  $L$  there must be rays converging to this dense set of points and thus also closed bubble regions in  $L$  the union of whose landing points is a dense in  $\partial\mathbb{H}^2$ . Pick one closed bubble region  $\mathcal{B}$  whose ideal point is neither  $\alpha(L)$  nor  $\xi$ .

Up to shrinking  $I$  to a smaller interval we assume that  $\mathcal{B}$  completely contained in  $\widehat{D}_\varepsilon(L, I)$ . Moreover, if we call  $c$  the maximal element of  $\mathcal{B}$ , it is a leaf of  $\mathcal{G}_L$  whose endpoints are  $\partial^\pm\Phi(c) = \{\eta\}$  with  $\eta \notin J$ . We can assume that every closed bubbles leaf whose landing point is in the interval  $K$  joining  $\xi$  and  $\eta$  and not intersecting  $J$  is contained in  $\widehat{D}_\varepsilon(L, I)$ . Indeed any such bubble region  $\mathcal{B}'$  not contained in  $\widehat{D}_\varepsilon(L, I)$  has points limiting to points in the interval  $K$  as well as points a bounded distance from the geodesic  $g$  in  $L$  with ideal points the endpoints of  $I$ . Suppose there are infinitely many of these. They cannot accumulate to some point in  $L$ , hence segments of the boundary of these limit on a nondegenerate open interval of  $S^1(L)$ . This contradicts that the set of ideal points of leaves of  $\mathcal{G}_L$  is dense in  $S^1(L)$ . It follows that there are only finitely many bubbles in  $L$  satisfying this property and we take the last one satisfying the property.

Since  $\mathcal{O}$  is an open bubble, we can find a sequence of nested curves  $e_n$  in  $\mathcal{O}$  which accumulates on  $c$ . More precisely, if  $D_{e_n}$  are the disks defined by  $e_n$  and contained in  $\mathcal{O}$  it follows that  $D_{e_n} \subset D_{e_{n+1}}$  and  $\mathcal{O} = \bigcup_n D_{e_n}$ . It follows that there are rays  $r_n$  of  $e_n$  contained in  $\widehat{D}_\varepsilon(L, I)$  and that accumulate on  $c$  (note that the full

curve  $e_n$  must enter  $\widehat{D}_\varepsilon(L, I)$ , but it cannot happen that for large  $n$  both rays are contained in  $\widehat{D}_\varepsilon(L, I)$ , since the union of the disks  $D_{e_n}$  is all of  $\mathcal{O}$ ).

Now we fix a leaf  $E \in I$  that is invariant under some  $\gamma \in \pi_1(M) \setminus \{\text{id}\}$  (cf. Section 2.7) and such that the fixed points of  $\gamma$  are different from  $\xi$  (note that one must be contained in  $J$ ). If we apply Proposition 3.4 we deduce that the leaf  $c$  pushes to a leaf  $c'$  of  $\mathcal{G}_E$  whose endpoints are  $\eta$  and the rays  $r_n$  push to rays  $r'_n$  which have one endpoint in  $\xi$  and accumulate on a ray of  $c'$ . If  $e'_n$  are the leaves of  $\mathcal{G}_E$  containing the rays  $r'_n$  we deduce that they must belong to an open bubble region since every leaf of  $\mathcal{G}_E$  is a bubble leaf and the leaves  $e'_n$  are all landing in  $\xi$ . It follows that  $E$  has an open bubble region with limit point  $\xi$ . Applying  $\gamma$  we deduce that there is more than one open bubble region in  $E$  contradicting the previous lemma.  $\square$

To prove Proposition 5.4 we will proceed by contradiction and therefore assume that all leaves of  $\widetilde{\mathcal{G}}$  are bubble leaves. This will allow us to use the previous results. To be able to get a contradiction we will need to construct uncountably many bubble leaves with different landing points in a leaf of the other foliation.

*Proof of Proposition 5.4.* Assume for a contradiction that every leaf of  $\widetilde{\mathcal{G}}$  is a bubble leaf.

We claim that for each  $L \in \widetilde{\mathcal{F}}_1$  there is a nontrivial interval of the leaf space of  $\widetilde{\mathcal{F}}_2$  which contains bubble leaves whose limit points are  $\alpha(L)$ . This is just because Lemma 5.7 states that the open bubble region  $\mathcal{O}_L$  of  $L$ , which exists due to Lemma 5.6, must land in  $\alpha(L)$ . Considering a transversal  $\tau$  to  $\mathcal{G}_L$  inside  $\mathcal{O}_L$  gives a non trivial interval  $I_L$  of the leaf space of  $\widetilde{\mathcal{F}}_2$  (i.e., those leaves which intersect  $\tau$ ) containing bubble leaves whose landing point is  $\alpha(L)$ . The interval  $I_L$  also depends on  $\tau$ , but we are omitting this dependence. Notice that we are not claiming that the elements of  $\{I_L, L \in \widetilde{\mathcal{F}}_1\}$  are pairwise distinct.

Pick  $x_n$  dense and countable in the leaf space of  $\widetilde{\mathcal{F}}_2$  (which is homeomorphic to  $\mathbb{R}$ ). If for every  $x_n$  there are only countably many  $L \in \widetilde{\mathcal{F}}_1$  such that  $x_n \in I_L$ , then the set of those  $L \in \widetilde{\mathcal{F}}_1$  for which  $I_L$  does not contain any  $x_n$  is still uncountable. Hence this set is nonempty, contradicting the density of  $\{x_n\}$ , since any  $I_L$  is an open set in the leaf space of  $\widetilde{\mathcal{F}}_2$ .

We deduce that there is a leaf  $E \in \widetilde{\mathcal{F}}_2$  for which there are uncountably many  $L \in \widetilde{\mathcal{F}}_1$  such that  $E \in I_L$  with pairwise distinct  $\alpha(L)$ . It follows that  $E$  contains uncountably many bubble leaves which land in pairwise different points. This contradicts the fact that a leaf can contain at most countably many distinct bubble regions (see Lemma 5.6). This contradiction completes the proof.  $\square$

**5.2. Proof of Proposition 5.1.** The proof will be by contradiction. Using what we have proved we will have a ray landing at a marker point. Then, assuming that the small visual property fails, we can use the following lemma and minimality to get a contradiction.

**Lemma 5.8.** *For every  $\delta > 0$  there exists  $K := K(\delta) > 0$  such that if*

- *$I$  is a segment in  $\partial\mathbb{H}^2$ ,*
- *$x \in \mathbb{H}^2$ , and*
- *$r$  is a geodesic ray starting from  $x$  and landing in some point  $\xi$  of  $I$  separating  $I$  in two intervals of visual measure  $> \delta/2$  seen from  $x$ ,*

*then if  $y \in r$  satisfies  $d_{\mathbb{H}^2}(x, y) > K$ , it follows that the visual measure of  $I$  seen from  $y$  is larger than  $2\pi - \delta$ . Here  $d_{\mathbb{H}^2}$  is the hyperbolic metric in  $\mathbb{H}^2$ .*

*Proof.* Since isometries are transitive in  $T^1\mathbb{H}^2$  we can assume in the upper half space model of  $\mathbb{H}^2$ , that  $x = i$ , the geodesic ray  $r$  is the ray  $\{ai : a \geq 1\}$  and the interval  $I$  contains, for some large  $t$  depending on  $\delta$  a set of the form  $A_t = [t, +\infty) \cup \{\infty\} \cup (-\infty, -t]$ . The lemma then reduces to computing the visual measure of  $A_t$  from a point of the form  $ai$  which goes to  $2\pi$  with  $a \rightarrow \infty$ .  $\square$

*Proof of Proposition 5.1.* Consider  $c \in \tilde{\mathcal{G}}$  given by Proposition 5.4 and let  $L \in \tilde{\mathcal{F}}_1$ ,  $E \in \tilde{\mathcal{F}}_2$  be such that  $c \subset L \cap E$ . We will argue for  $L$  (the case for  $E$  is symmetric).

Let  $r$  be a ray of  $c$  with landing point  $\partial\Phi(r) = \xi \neq \alpha(L)$ . Let  $I$  be a closed interval of the leaf space of  $\tilde{\mathcal{F}}_1$  containing  $L$  in its interior and such that if  $L' \in I$  then  $\alpha(L') \neq \xi$ . Up to reducing the ray  $r$  we can assume that  $r \subset \widehat{D}_\varepsilon(L, I)$  for  $\varepsilon$  as given in Proposition 3.4.

Now assume for a contradiction that there is  $\delta > 0$  and a sequence  $\ell_n$  of segments of  $\mathcal{G}_{L_n}$  with  $L_n \in \tilde{\mathcal{F}}_1$  and points  $x_n \in L_n$  such that  $d_{L_n}(x_n, g_n) > 5n$  and  $\ell_n$  has shadow in  $S^1(L_n)$  of length  $> 10\delta$  when seen from  $x_n$ . Up to cutting the segments  $\ell_n$  we can assume that the shadow has length  $> 5\delta$  and is disjoint from  $\alpha(L_n)$ .

Consider for each  $n$  a geodesic ray from  $x_n$  to the middle point of the shadow of  $\ell_n$  and a point  $y_n$  in the geodesic ray and at distance  $= 2n$  from  $x_n$ . By minimality of  $\mathcal{F}_1$  and up to changing slightly the geodesic ray, we can assume that there is  $\gamma_n$  sending  $y_n$  to a point very close to  $L$  and in a compact set of  $\tilde{M}$ . Using Lemma 5.8 we get that, up to a subsequence,

- $d_{L_n}(y_n, g_n) > 2n$ ,
- the visual measure of  $\ell_n$  seen from  $y_n$  is larger than  $2\pi - a_n$  with  $a_n \rightarrow 0$  (this is after identifying  $S^1(L)$  with  $\partial\mathbb{H}^2$ ), and
- $\gamma_n y_n \rightarrow y_\infty \in L$ .

We chose  $\ell_n$  so that  $\alpha(L_n)$  does not belong to the shadow of  $\ell_n$  from  $x_n$  and thus we get that the shadow from  $\gamma_n y_n$  of  $\gamma_n g_n$  contains  $\xi$  in its interior if  $n$  is big enough. This is because  $\gamma_n L_n \rightarrow L$  and  $\alpha(\gamma_n L_n) \rightarrow \alpha(L)$ . Since for large  $n$  the leaf  $\gamma_n L_n$  is close to  $L$  we can assume that  $\gamma_n L_n$  is in  $I$  and we can push arcs  $\ell'_n$  of  $\gamma_n g_n$  to  $L$  so that  $\Phi_L(\ell'_n)$  accumulates in an interval containing  $\xi$  in its interior. This forces  $\ell'_n$  to eventually intersect  $r$  which is a contradiction.  $\square$

## 6. Leafwise Hausdorff leaf space implies quasigeodesic

Here we will show the following:

**Theorem 6.1.** *Let  $\mathcal{F}_1$  and  $\mathcal{F}_2$  be transverse minimal foliations of  $M = T^1S$  and let  $\mathcal{G}$  denote the foliation obtained by intersection. Assume that in some leaf  $L \in \tilde{\mathcal{F}}_1$  the leaf space of  $\mathcal{G}_L$  is Hausdorff. Then, the foliation  $\mathcal{G}$  is an Anosov foliation.*

This also follows from [27] which gives an alternative proof. Here we present a proof in our restricted setting, since it is vastly simpler and can show some of the ideas in a more transparent way.

Recall that an *Anosov foliation* is a one-dimensional foliation in  $M$  which is homeomorphic to the orbit foliation of a topological Anosov flow. As follows from [5, § 6], due to minimality, it is enough to show that the one-dimensional foliation is *quasigeodesic*, that is, for every leaf  $L \in \tilde{\mathcal{F}}_i$  the leaves of  $\mathcal{G}_L$  are quasigeodesics in  $L$ .

Here we are working with  $M = T^1S$  so it makes sense to compare our one-dimensional foliation with the one of the geodesic flow in negative curvature: this foliation, when restricted to a leaf  $L$  of the weak-stable or weak-unstable foliation has the following properties that we will try to produce to show Theorem 6.1:

- One of the landing points of every leaf of  $\mathcal{G}_L$  is  $\alpha(L)$ .
- Given any point  $\xi \in \partial\mathbb{H}^2 \setminus \{\alpha(L)\}$  there is a (unique) leaf whose landing points are  $\xi$  and  $\alpha(L)$ .

Thanks to some classical results, showing these properties will be enough to establish that the foliation is an Anosov foliation (see [28]). This implies that it is quasigeodesic (see [18]). We refer the reader to [5, § 5] for details.

**6.1. Bubble leaves and landing.** First we need to show that bubble leaves (recall Section 5.1) produce non-Hausdorff behavior. For this, the small visual measure property is crucial, in particular Proposition 5.2.

We recall here that the assumptions of Theorem 6.1 imply that in *every* leaf of  $\tilde{\mathcal{F}}_1$  and  $\tilde{\mathcal{F}}_2$  the restriction of  $\tilde{\mathcal{G}}$  has Hausdorff leaf space because of Proposition 3.6.

**Lemma 6.2.** *If  $\mathcal{G}_L$  has a bubble leaf for some  $L \in \tilde{\mathcal{F}}_i$ , then there is a leaf  $L' \in \tilde{\mathcal{F}}_i$  such that  $\mathcal{G}_{L'}$  has non-Hausdorff leaf space.*

*Proof.* Consider a bubble leaf  $c \in \mathcal{G}_L$  (assume  $L \in \tilde{\mathcal{F}}_1$ ) and fix a geodesic ray  $r$  starting at some point in  $c$  which has the same landing point  $\xi \in \mathbb{H}^2$  as  $c$  (i.e.,  $\partial\Phi(r) = \partial^\pm\Phi(c) = \{\xi\}$ ). Let  $x_0$  be the starting point of  $r$  in  $c$  and consider the rays  $c_{x_0}^+$  and  $c_{x_0}^-$ . Passing to the limit in Proposition 5.2 we know that there is some  $a_0 > 0$  such that  $r$  is contained in the  $a_0$ -neighborhood of  $c_{x_0}^+$  and  $c_{x_0}^-$ .

Fix a sequence  $y_n \in r$  such that  $\Phi_L(y_n) \rightarrow \xi$  and consider points  $p_n \in c_{x_0}^+$  and  $q_n \in c_{x_0}^-$  at distance less than  $a_0$  from  $y_n$  in  $L$ . It follows that

- $d_L(p_n, q_n) \leq 2a_0$ ,
- $\Phi_L(p_n) \rightarrow \xi$  and  $\Phi_L(q_n) \rightarrow \xi$ , and
- the length of the segment of  $c$  joining  $p_n$  and  $q_n$  goes to infinity.

Now consider deck transformations  $\gamma_n \in \pi_1(M)$  which map  $p_n$  to a compact fundamental domain  $K \subset \tilde{M}$ . Since  $\gamma_n$  are isometries and  $d_L(p_n, q_n)$  is bounded, the points  $\gamma_n q_n$  also belong to a compact set. Up to subsequence we can assume that both  $\gamma_n p_n$  and  $\gamma_n q_n$  converge to points  $p_\infty$  and  $q_\infty$ . Note that since the distance in  $L$  from  $p_n$  and  $q_n$  is bounded, both points  $p_\infty$  and  $q_\infty$  belong to the same leaf  $L' \in \tilde{\mathcal{F}}_i$ .

Since the length from  $p_n$  to  $q_n$  along  $c$  goes to infinity it follows that  $p_\infty$  and  $q_\infty$  cannot belong to the same leaf of  $\mathcal{G}_{L'}$ . On the other hand  $p_n, q_n$  belong to the same leaf of  $\tilde{\mathcal{G}}$  and thus the same leaf  $E_n$  of  $\tilde{\mathcal{F}}_2$  which since  $p_n, q_n$  are close to  $L'$  it follows that either  $E_n \cap L'$  is not connected (in which case, the leaf space of  $\mathcal{G}_{L'}$  is not Hausdorff) or  $E_n \cap L'$  is a sequence of leaves of  $\mathcal{G}_{L'}$  converging to both the leaf through  $p_\infty$  and  $q_\infty$  and again we get that  $\mathcal{G}_{L'}$  does not have Hausdorff leaf space.

Since  $p_\infty, q_\infty$  do not belong to the same leaf of  $\mathcal{G}_{L'}$  but  $p_n, q_n$  belong to the same leaf of  $\tilde{\mathcal{F}}_2$  thus,  $\mathcal{G}_{L'}$  does not have Hausdorff leaf space as we wanted to show.  $\square$


Now we show that at least one of the endpoints of each leaf in  $\mathcal{G}_L$  must be the nonmarker point. Fix an orientation in  $L$  and consider, for a given  $c \in \mathcal{G}_L$  which we know separates  $L$  in two open disks  $D_c^\pm$ , each one such that the limit set of  $\Phi_L(D_c^\pm)$  in  $\partial\mathbb{H}^2$  is one of the intervals joining the landing points of  $c$  (see Corollary 2.10).

**Lemma 6.3.** *Let  $L \in \tilde{\mathcal{F}}_i$  and  $c \in \mathcal{G}_L$  then either  $\partial^+\Phi(c) = \{\alpha(L)\}$  or  $\partial^-\Phi(c) = \{\alpha(L)\}$ .*

*Proof.* Let  $L \in \tilde{\mathcal{F}}_i$  and  $c \in \mathcal{G}_L$  and assume that  $\alpha(L) \notin \partial^+\Phi(c) \cup \partial^-\Phi(c)$ .

Proposition 3.4 lets us assume that the leaf  $L$  is fixed by some  $\gamma \in \pi_1(M) \setminus \{\text{id}\}$ , and such that both points in  $\partial^+\Phi(c) \cup \partial^-\Phi(c)$  belong to the same connected component of  $\partial\mathbb{H}^2$  minus the fixed points of  $\gamma$ . It is enough to consider an interval  $I$  of the leaf space of  $\tilde{\mathcal{F}}_i$  containing  $L$  in the interior so that the nonmarker points of leaves in  $I$  are never in  $\partial^+\Phi(c) \cup \partial^-\Phi(c)$ : hence given an even smaller interval we can push the entire leaf  $c$  to nearby leaves of  $\tilde{\mathcal{F}}_i$  along the leaf of the other foliation with Proposition 3.4. Then since leaves with nontrivial stabilizer are dense (see Section 2.7), we can push to one of them in  $I$  to get the same property in such a leaf. To get that the fixed points of  $\gamma$  do not link with  $\partial^+\Phi(c) \cup \partial^-\Phi(c)$  we can use for instance that the set of closed geodesics in  $S$  is dense in  $S$  (Section 2.7).

We now choose orientations so that  $D_c^+$  is the complementary component of  $c$  which does not limit on  $\gamma^+, \gamma^-$ . If there is a transversal to  $\mathcal{G}_L$  in  $L$  from  $c$  to  $\gamma c$ , then the image of this under  $\gamma$  is a transversal from  $\gamma c$  to  $\gamma^2 c$ . Notice



**Figure 11.** If no ray lands in  $\alpha(L)$  the leaf space cannot be Hausdorff.

that  $D_c^+$ ,  $\gamma(D_c^+)$ ,  $\gamma^2(D_c^+)$  are pairwise disjoint. It now follows that  $c$ ,  $\gamma^2 c$  cannot intersect a common transversal and the leaf space cannot be Hausdorff. (See Figure 11.)  $\square$

**6.2. Construction of the Anosov foliation.** To complete the proof that the foliation  $\mathcal{G}$  is an Anosov foliation we need to show that for every  $L \in \tilde{\mathcal{F}}_i$  and  $\xi \in \partial\mathbb{H}^2 \setminus \{\alpha(L)\}$  there is a leaf  $c \in \mathcal{G}_L$  such that  $\partial\Phi(c) = \{\xi, \alpha(L)\}$  and that such a leaf is unique; this will produce an equivariant equivalence with the leaf space of the orbit flow of the geodesic flow in negative curvature [28] which is known to be enough to show that the foliation has the desired properties.

We divide this into some steps:

**Lemma 6.4.** *Let  $L \in \tilde{\mathcal{F}}_i$  and  $\xi \in \partial\mathbb{H}^2$ , then, there exists  $c \in \mathcal{G}_L$  with a ray landing in  $\xi$  (i.e., such that  $\xi \in \partial^\pm\Phi(c)$ ).*

*Proof.* Recall from Lemma 4.3 that the set of points in  $\partial\mathbb{H}^2$  for which there is a ray in  $L$  which lands in that point is dense.

Consider a sequence  $c_n$  of leaves of  $\mathcal{G}_L$  converging to some leaf  $c \in \mathcal{G}_L$  (which is unique because the leaf space of  $\mathcal{G}_L$  is Hausdorff). Let  $\xi_n$  be the landing point of  $c_n$  different from  $\alpha(L)$  and let  $\xi$  and  $\alpha(L)$  be the landing points of  $c$ . Assume that (up to taking a subsequence)  $\xi_n \rightarrow \eta \neq \xi$ . One can then consider a ray which lands in a point which belongs to the interval between  $\xi$  and  $\eta$  not containing  $\alpha(L)$ . This ray belongs to some leaf  $\ell \in \mathcal{G}_L$ . By the previous lemma, the leaf  $\ell$  has one ideal point in  $\alpha(L)$ . It follows that  $\ell$  must then separate the curves  $c_n$  with large  $n$  from  $c$ , contradicting the fact that  $c_n \rightarrow c$ .

This continuity plus the fact that the landing points are dense implies that every point different from  $\alpha(L)$  is a landing point as we wanted to show.  $\square$

Now we show uniqueness as a consequence of minimality of the foliations, and derive from Theorem 2.1 that the nonmarker points of leaves vary monotonically (Section 2.4).

**Lemma 6.5.** *Let  $L \in \tilde{\mathcal{F}}_i$  and  $\xi \in \partial\mathbb{H}^2 \setminus \{\alpha(L)\}$ , then, the leaf  $c \in \mathcal{G}_L$  such that  $\partial^\pm\Phi(c) = \{\xi, \alpha(L)\}$  is unique.*

*Proof.* Assume that in a leaf  $L \in \tilde{\mathcal{F}}_1$  (the case where  $L \in \tilde{\mathcal{F}}_2$  is symmetric) there are two leaves of  $\mathcal{G}_L$  landing in  $\xi$  and  $\alpha(L)$ . Since the leaf space of  $\mathcal{G}_L$  is Hausdorff and one of the landing points of every  $c \in \mathcal{G}_L$  is  $\alpha(L)$  we deduce that there is an interval  $I$  in the leaf space of  $\tilde{\mathcal{F}}_2$  consisting on leaves  $E$  such that  $E \cap L$  is a leaf  $c \in \mathcal{G}_L$  such that  $\partial^\pm\Phi_L(c) = \{\xi, \alpha(L)\}$ . From the previous lemma applied to  $\tilde{\mathcal{F}}_2$  we know that if  $E \in I$  and if  $c' \in \mathcal{G}_E$  then  $\alpha(E) \in \partial^\pm\Phi_E(c')$ . Considering  $c \in E \cap L$  we deduce that for every  $E \in I$  we have that  $\alpha(E) \in \{\xi, \alpha(L)\}$ . Since the point  $\alpha(E)$  varies continuously with  $E$  we deduce that there is an interval of  $\tilde{\mathcal{F}}_2$  which has the same nonmarker point while the nonmarker point varies monotonically.  $\square$

Now we are ready to prove the main result of this section.


*Proof of Theorem 6.1.* From our previous results we have a bijection between the space of leaves of  $\widehat{\mathcal{G}}$  (the lift of the foliation  $\mathcal{G}$  to  $\widehat{M}$  the intermediate cover of  $M$ ) and the set of pairs of distinct points of  $\partial\mathbb{H}^2$ . This bijection is continuous and equivariant under deck transformations which is enough to show that the foliation is homeomorphic to the orbit foliation of the geodesic flow of a hyperbolic metric on  $S$ . (See [28] or [5] for details.)  $\square$

### 7. The Matsumoto–Tsuboi example revisited

Here we revisit the example from [35] from the point of view of our results. In [35] an example of a pair of transverse foliations<sup>8</sup> of  $\mathbb{T}^2 \times [-1, 1]$  is constructed in such a way that the boundaries match with the weak stable and weak unstable foliations seen in the lift to  $T^1S$  of a simple closed geodesic. That is, if  $\ell$  is a closed geodesic in  $S$ , it lifts to a torus  $T \subset T^1S$  which contains two periodic orbits of the geodesic flow (associated to the orbits associated to  $\gamma$  with both orientations). Note that the weak stable and weak unstable foliations are horizontal, so they are transverse to  $T$ . Figure 12 depicts the intersection of the foliations with the torus in its universal cover (which become tangent at the two periodic orbits that are common to both weak foliations).

This way one can cut  $T^1S$  along such a torus and glue this new foliation of  $\mathbb{T}^2 \times [-1, 1]$  to obtain a new pair of transverse foliations of  $T^1S$ . From the way this pair of foliations is constructed, it is clear that the new foliations, called  $\mathcal{F}_1, \mathcal{F}_2$ ,

<sup>8</sup>In [35] triples of transverse foliations are considered, but for the interests of our work we will ignore the third foliation.



**Figure 12.** The intersection of the transverse tori with the weak stable and weak unstable of the geodesic flow.

will continue to be minimal and their intersection foliation  $\mathcal{G}$  will be such that for every leaf  $L \in \tilde{\mathcal{F}}_i$  the foliation  $\mathcal{G}_L$  will not have Hausdorff leaf space. In this paper we will not use this construction, but we thought it could be relevant to spend some time explaining the example from our point of view which was relevant for us to formulate the right statements in the next two sections. We plan to study this example as well as other examples of transverse foliations with Reeb surfaces in a future work.

**7.1. The construction and its possible variants.** Let  $S$  be a closed surface with a hyperbolic metric and let  $\varphi_t : T^1S \rightarrow T^1S$  be the geodesic flow. The metric induces an action of  $\Gamma = \pi_1(S)$  on  $\mathbb{H}^2$  and on  $T^1\mathbb{H}^2 = \widehat{M}$  such that the weak stable foliation of  $\varphi_t$  is the foliation  $\mathcal{F}_{ws}$  defined in Section 2 by identifying  $T^1\mathbb{H}^2$  with  $\mathbb{H}^2 \times \partial\mathbb{H}^2$ . The weak unstable foliation will be called  $\mathcal{F}_{wu}$  and is obtained in a similar process, only that the identification of  $T^1\mathbb{H}^2$  with  $\mathbb{H}^2 \times \partial\mathbb{H}^2$  is made by identifying the point at infinity as the limit of the geodesic in the backwards direction. Both  $\mathcal{F}_{ws}$  and  $\mathcal{F}_{wu}$  are transverse to the fibers.


Fix any simple closed geodesic  $\ell$  in  $M = T^1S$  for the metric  $g$  and let  $T_\ell$  be the torus obtained by looking at the unit vectors tangent at points of  $\ell$ . This torus intersects both  $\mathcal{F}_{ws}$  and  $\mathcal{F}_{wu}$  transversally and the flow is transverse to  $T_\ell$  except at the two orbits of  $\varphi_t$  contained in  $T_\ell$ . See Figure 12.

In [35] a construction of two transverse foliations  $\mathcal{S}, \mathcal{U}$  of  $T_\ell \times [0, 1]$  are given intersecting the boundary tori with the following properties:


- Each leaf  $S \in \mathcal{S}$  is either a cylinder (i.e., homeomorphic to  $S^1 \times [0, 1]$ ) or a band (i.e., homeomorphic to  $\mathbb{R} \times [0, 1]$ ), and the intersection of  $S$  with each boundary torus  $T_\ell \times \{0\}$  and  $T_\ell \times \{1\}$  are the same (in the trivialization  $T_\ell \times [0, 1]$ ) and coincides with the intersection of some leaf of  $\mathcal{F}_{ws}$  with  $T_\ell$ .

- Each leaf  $U \in \mathcal{U}$  is either a cylinder or a band, and the intersection with each boundary torus  $T_\ell \times \{0\}$  and  $T_\ell \times \{1\}$  is the same and coincides with the intersection of some leaf of  $\mathcal{F}_{wu}$  with  $T_\ell$ .
- There are exactly two cylinder leaves  $S_1, S_2$  of  $\mathcal{S}$  and  $U_1, U_2$  of  $\mathcal{U}$  corresponding to the periodic orbits of the flow in  $T_\ell$ .
- Each leaf  $S \in \mathcal{S} \setminus \{S_1, S_2\}$  intersects either  $U_1$  or  $U_2$  in two connected components that are infinite lines and bound a Reeb band.
- Each leaf  $U \in \mathcal{U} \setminus \{U_1, U_2\}$  intersects either  $S_1$  or  $S_2$  in two connected components that are infinite lines and bound a Reeb band.
- $S_1$  intersects  $U_1$  in at least three circles,
- $S_2$  intersects  $U_2$  in at least three circles and  $U_1$  in at least four circles.

We depict one possibility in Figures 13 and 14.




**Figure 13.** The horizontal lines represent the cylinders  $U_1$  and  $U_2$  (and their translations up by deck transformations) and the red curves represent the cylinders  $S_1$  and  $S_2$  (and their translations up by deck transformations). The figure depicts how the leaves intersect and each intersection corresponds to a circle in the leaf. The rest of the leaves of the  $\mathcal{U}$  foliation are also horizontal leaves, but when going around the holonomy of the compact leaves, they intersect in a different height (in particular, the corresponding leaf is an infinite band whose intersection with the torus is an infinite family of horizontal segments which accumulate in  $U_1$  and  $U_2$  when going forward and backward respectively). The other leaves of the  $\mathcal{S}$  foliation interpolate the traces of  $S_1$  and  $S_2$  and also change as going around the torus in the flow direction.




**Figure 14.** Left: the lift of one of the cylinders ( $U_2$ ) and the intersected foliation lifted to this cover. Right: the same for the other cylinder ( $U_1$ ). These intersections correspond to intersecting the leaves as one moves in the direction of the flow (which makes leaves of the  $\mathcal{S}$  foliation approach in the forward or backward direction to the leaves  $S_1$  and  $S_2$ ). The straight lines corresponds to the intersection with the leaves  $S_1$  and  $S_2$  which are invariant under moving one step up.

We now consider the foliations  $\mathcal{F}_1, \mathcal{F}_2$  in  $M_0$  obtained by first cutting  $T^1S$  foliated with  $\mathcal{F}_{ws}$  and  $\mathcal{F}_{wu}$  along  $T_\gamma$  and gluing a copy of  $T_\ell \times [0, 1]$  foliated by  $\mathcal{S}$  and  $\mathcal{U}$  and gluing the foliation  $\mathcal{F}_{ws}$  with  $\mathcal{S}$  and  $\mathcal{F}_{wu}$  with  $\mathcal{U}$ . The manifold  $M_0$  is still diffeomorphic to  $T^1S$  (we have changed  $T_\gamma$  for  $T_\gamma \times [0, 1]$  with the trivial identification of  $T_\gamma \times \{0\}$  and  $T_\gamma \times \{1\}$  with the two copies of  $T_\gamma$  one gets after “cutting”  $T^1S$ ) and the foliations  $\mathcal{F}_1$  and  $\mathcal{F}_2$  are everywhere transverse. We note that it is possible to choose the foliations  $\mathcal{S}$  and  $\mathcal{U}$  in order that the resulting foliations  $\mathcal{F}_1$  and  $\mathcal{F}_2$  are *not* uniformly equivalent, that is, the homeomorphisms  $h_1$  and  $h_2$  given by Theorem 2.1 sending  $\mathcal{F}_1$  and  $\mathcal{F}_2$  to  $\mathcal{F}_{ws}$  cannot be homotopic to each other (see Figures 15 and 16). The example in [35] provides examples for which both foliations are uniformly equivalent.

**Remark 7.1.** One can glue more than one block  $T_\ell \times [0, 1]$ , concatenating consecutive blocks. It is also possible to do the cut and paste process in several disjoint simple closed geodesic to obtain variants of these examples. More generally, take any Anosov flow in a closed 3-manifold with an embedded Birkhoff torus with only two Birkhoff annuli (that is, when the intersection of the weak stable and weak unstable foliations of the Anosov flow intersect the tori exactly as in this



**Figure 15.** How the cylinders of each foliation can intersect so as to yield a pair of non-uniformly equivalent foliations intersecting transversally.



**Figure 16.** Left: A diagram of the foliations in the cylinder leaves when crossing the tori. Right: the foliation interpolating both foliations of a leaf that accumulates in both cylinders in the different directions.

case). Then, one can glue some of the foliations of  $\mathbb{T}^2 \times [0, 1]$  and produce similar examples. We leave to a forthcoming work the (similar) analysis of this case. One can also do this with embedded Birkhoff tori with more Birkhoff annuli, but then the building blocks have to be accordingly adjusted: one needs at least the same

number of Birkhoff annuli, which can be achieved by finite covers of  $\mathbb{T}^2 \times [0, 1]$ . This will also be pursued in the future.

**7.2. Properties of the intersected foliation.** From the point of view of this paper, it is relevant for us to understand the properties of the foliation  $\mathcal{G}$  obtained by intersecting  $\mathcal{F}_1$  and  $\mathcal{F}_2$ . The main observations are the following:

- Given  $L \in \tilde{\mathcal{F}}_1$  we have that the foliation  $\mathcal{G}_L$  coincides with the foliation by geodesics as long as  $L$  does not intersect the lift to the universal cover of  $T_\ell \times [0, 1]$ . This lift has the property that it intersects every leaf in some regions which are bounded by two curves at bounded distance and landing at given points of  $\partial\mathbb{H}^2$  independent of the leaf  $L$ .
- Inside each of the regions of intersection of the lift of  $T_\ell \times [0, 1]$  to  $\tilde{M}$  the foliation  $\mathcal{G}_L$  has some design that depends on the specific foliation (see for instance Figure 14 for a depiction of the leaves in some specific examples). However, it is always the case that in these regions there is a pair of nonseparated leaves at bounded distance of each other and landing at the two points at infinity of the region. Every leaf that enters the region must land in one of the endpoints of the region. In particular each ray of a leaf that enters this region does not leave this region.
- Given two nonseparated leaves  $c_1, c_2 \in \mathcal{G}_L$  they must accumulate in one of the regions mentioned in the previous point. Note that if the nonseparated leaves are not contained in the region modified between the two tori, the nonseparated leaves may be at unbounded distance inside their leaf.

These properties will become more apparent in Section 9 which actually shows that behavior like this is the only possible.


## 8. A dichotomy Hausdorff vs Reeb surfaces

Let  $\mathcal{F}_1, \mathcal{F}_2$  be two transverse minimal foliations on  $M = T^1S$  and let  $\mathcal{G}$  be the foliation obtained by intersecting them.

We define a *Reeb surface* to be a compact surface with boundary  $R$  satisfying the following properties:

- $R$  is contained in a leaf  $S$  of  $\mathcal{F}_i$  (with  $i = 1, 2$ ),
- the boundary components of  $R$  are closed leaves of  $\mathcal{G}$ ,
- if  $L$  is a lift of  $S$  to the universal cover, then, the lift of  $R$  to  $L$  is a surface  $B$  whose boundary consists of exactly two leaves  $c_1, c_2$  of  $\tilde{\mathcal{G}}$  at bounded distance which are nonseparated in  $\mathcal{G}_L$ .

In particular, it is easy to see that the surface must be either an annulus or a Möbius band.



**Figure 17.** The interior of a non-Hausdorff bigon may not be trivially foliated if it is not  $\gamma$ -invariant for some  $\gamma \in \pi_1(M)$ .

We will define for a leaf  $L \in \tilde{\mathcal{F}}_i$  a *non-Hausdorff bigon* to be a pair of nonseparated leaves  $c_1, c_2 \in \mathcal{G}_L$  such that  $\partial^+ \Phi(c_1) \neq \partial^- \Phi(c_1)$  and  $\partial^\pm \Phi(c_1) = \partial^\pm \Phi(c_2)$ . It does not follow that the leaf space of  $\mathcal{G}_L$  inside the bigon is the reals (see Figure 17). For example one can have bubble leaves  $c, c'$  in the bigon so that the disks  $D_c, D_{c'}$  are disjoint. On the other hand it is easy to see that if the boundaries of the bigon are invariant by a nontrivial deck transformation, then the leaf space of  $\mathcal{G}_L$  inside the bigon is the reals. In particular, the existence of a Reeb surface is equivalent to having a non-Hausdorff bigon invariant under some deck transformation  $\gamma \neq \text{id}$ .

Here we show:

**Theorem 8.1.** *Either there is a leaf  $L \in \tilde{\mathcal{F}}_1$  such that the leaf space of  $\mathcal{G}_L$  is Hausdorff, or there exists a Reeb surface.*

Theorem 8.1 together with Theorem 6.1 is enough to conclude the proof of Theorem A. The purpose of this section is to prove Theorem 8.1.

**8.1. Some analysis on nonseparated leaves.** We first find a useful criterion to obtain Reeb surfaces:

**Proposition 8.2.** *Let  $L \in \tilde{\mathcal{F}}_i$  containing two nonseparated leaves  $c_1, c_2 \in \mathcal{G}_L$  such that there is  $\gamma \in \pi_1(M) \setminus \{\text{id}\}$  which fixes  $L$  (i.e.,  $\gamma L = L$ ). Assume that the nonseparated rays  $r_1, r_2$  of  $c_1, c_2$  (cf. Section 3.4) satisfy  $\partial \Phi(r_1) = \partial \Phi(r_2) = \xi$  with  $\gamma \xi = \xi$ . Then,  $\mathcal{F}_i$  has a Reeb surface in the projection of  $L$  to  $M$ .*

Note in particular that if  $\xi = \alpha(L)$  then the hypothesis that  $\gamma \xi = \xi$  holds automatically.

*Proof.* Let  $\ell \subset L$  be a  $\gamma$ -invariant curve joining  $\xi$  with the other fixed point of  $\gamma$  in  $S^1(L)$ , without loss of generality we can assume that  $\ell$  intersects  $r_1$  and  $r_2$ . The small visual measure property (Proposition 5.2) implies that there is  $a_0$  such that every ray  $r$  of  $\mathcal{G}_L$  which lands in  $\xi$  and intersects  $\ell$  satisfies the condition that the  $a_0$ -neighborhood of  $r$  contains a ray of  $\ell$ . This applies to  $r_1, r_2$  but also their iterates  $\gamma^k r_i$  for  $i = 1, 2$  and  $k \in \mathbb{Z}$ . Since  $\gamma^j r_1$  is nonseparated with  $\gamma^j r_2$  it follows that in a given fundamental domain, one can have at most finitely many such rays. This implies that there is  $k \in \mathbb{Z} \setminus \{0\}$  such that  $\gamma^k r_i \cap r_i \neq \emptyset$  and thus  $\gamma^k c_i = c_i$  for  $i = 1, 2$ .

This implies that  $c_1$  and  $c_2$  join  $\xi$  with the other fixed point of  $\gamma$  and project to a (simple) closed curve in  $M$ . If  $\gamma$  preserves orientation, it follows that both  $c_1$  and  $c_2$  are  $\gamma$ -invariant, else, it can be that  $\gamma$  permutes them and  $\gamma^2$  preserves them. In both cases we deduce that they bound a Reeb surface.  $\square$

**8.2. Proof of Theorem 8.1.** The following proposition will be proved in the next subsection.

**Proposition 8.3.** *Let  $L \in \tilde{\mathcal{F}}_i$  and  $c_1, c_2 \in \mathcal{G}_L$  be two nonseparated leaves such that their nonseparated rays  $r_1, r_2$  satisfy  $\partial\Phi(r_1) = \partial\Phi(r_2) = \xi \neq \alpha(L)$ . Then, there is  $L' \in \tilde{\mathcal{F}}_i$  and a deck transformation  $\gamma \in \pi_1(M) \setminus \{\text{id}\}$  such that*

- $\gamma\xi = \xi$  and  $\gamma L' = L'$ , and
- *there is a  $\gamma$ -invariant non-Hausdorff bigon in  $L'$  (which has  $\xi$  as one of its endpoints).*

We explain how Theorem 8.1, and thus Theorem A, follow from this proposition. As shown in Proposition 3.6, if some leaf  $L \in \tilde{\mathcal{F}}_i$  has the property that the leaf space of  $\mathcal{G}_L$  is non-Hausdorff then the same holds for every leaf of both foliations. By Proposition 8.2, if there is a leaf  $L \in \tilde{\mathcal{F}}_i$  with nontrivial stabilizer for which the nonseparated rays of a pair of non separated leaves land in  $\alpha(L)$  then we can conclude the existence of a Reeb surface. So, we can assume that there is a leaf  $L \in \mathcal{F}_1$  which has nonseparated leaves  $c_1, c_2$  such that their nonseparated rays  $r_1, r_2$  satisfy  $\partial\Phi(r_1) = \partial\Phi(r_2) \neq \alpha(L)$ . Thus, we can apply Proposition 8.3 to deduce the existence of a  $\gamma$ -invariant non-Hausdorff bigon in some leaf  $L' \in \mathcal{F}_1$  which implies the existence of a Reeb surface by Proposition 8.2. This completes the proof of Theorem 8.1.

For the purposes of the study we will make in Section 9, it is useful to obtain the following additional information:

**Addendum 8.4.** *Under the assumptions of Proposition 8.3 one can choose  $L'$  so that the rays  $r_1, r_2$  are asymptotic to  $L'$  in  $\tilde{M}$ .*

Here, by “asymptotic” to  $L'$  in  $M$  we mean that if we parametrize  $r_i$  in  $M$  by arc length as  $r_i(t)$ , then, as  $t$  goes to infinity,  $d(r_i(t), L')$  converges to zero. In other words the projection  $\pi(r_i)$  of  $r_i$  to  $M$  is asymptotic to a closed curve of  $\mathcal{G}$  in  $\pi(L')$ .

**8.3. Proof of Proposition 8.3.** The proof of this proposition has two main steps: we first analyze some impossible configurations of returns of the curves  $r_i$  when projected to  $M$  and then we use the small visual measure to determine that the rays  $\pi(r_1)$  and  $\pi(r_2)$  need to accumulate as desired.

Let  $r_1, r_2$  be nonseparated rays of  $\mathcal{G}_L$  converging to  $\xi \in S^1(L)$ . Let  $c_i$  be the leaves of  $\mathcal{G}_L$  containing them. Let  $\sigma : [0, 1] \rightarrow L$  be a properly embedded segment satisfying the following conditions:

- (1)  $\sigma(0) \in r_1, \sigma(1) \in r_2$ .
- (2)  $\sigma$  intersects  $c_1 \cup c_2$  only in the endpoints and  $\sigma$  transverse to  $\mathcal{G}_L$  except at one interior point.
- (3) There is a small transverse arc  $\beta$  to  $\mathcal{G}_L$  starting in  $\alpha(0)$  and going in the direction of  $c_2$  so that if a leaf of  $\mathcal{G}_L$  intersects the interior of  $\alpha$  then it intersects  $\beta$ .

Let  $\mathcal{C} = r_1 \cup r_2 \cup \sigma([0, 1])$  which is a Jordan curve and let  $\mathcal{B} = \mathcal{B}(r_1, r_2, \sigma)$  be the closure of the connected component of the complement of  $\mathcal{C}$  which accumulates only on  $\xi$  in  $S^1(L)$ . We say that  $\mathcal{B}$  is a *good half-band* if  $\mathcal{B}$  is the closure of the union of the arcs of leaves of  $\mathcal{G}_L$  contained in  $\mathcal{B}$  and joining points of  $\sigma = \sigma([0, 1])$ . Equivalently, we say that  $\mathcal{B}$  is a good half-band if the boundary contains two nonseparated rays and it does not contain any complete leaf in its interior.

**Lemma 8.5.** *Let  $c'_1, c'_2 \in \mathcal{G}_L$  be two nonseparated leaves in  $L \in \tilde{\mathcal{F}}_i$  such that their nonseparated rays  $r'_1, r'_2$  satisfy  $\partial\Phi(r'_1) = \partial\Phi(r'_2) = \xi$ . Then, there exist nonseparated leaves  $c_1, c_2 \in \mathcal{G}_L$  with nonseparated rays  $r_1, r_2$  satisfying  $\partial\Phi(r_1) = \partial\Phi(r_2) = \xi$  and moreover it bounds a good half-band.*

*Proof.* Applying the consequence of the small visual measure property given in Proposition 5.2 we get that for any geodesic ray  $\ell_0$  landing in  $\partial\Phi(r'_1) = \partial\Phi(r'_2) = \xi$  one has that  $\ell_0$  has a subray contained in the  $a_0 + a_1$ -neighborhood of every ray  $\beta$  of  $\mathcal{G}_L$  that lands in  $\xi$ : this is because there is a geodesic ray with starting point the starting point of  $\beta$ , ideal point  $\xi$  and contained in the  $a_0$  neighborhood of  $\beta$ . Then by Gromov hyperbolicity, there is a global constant  $a_1 > 0$  such that any two geodesic rays with ideal point  $\xi$ , have subrays which are  $< a_1$  Hausdorff distant from each other in  $L$ .

A priori there could be some leaves of  $\mathcal{G}_L$  nonseparated from  $c'_1$  and  $c'_2$  and between  $c'_1, c'_2$  (i.e., if we pick a curve  $\sigma'$  as above and define  $\mathcal{B}' = \mathcal{B}(c'_1, c'_2, \sigma')$ , there may be curves nonseparated from  $c'_1$  and  $c'_2$  which have both endpoints in  $\xi$  and intersect  $\mathcal{B}'$  – and hence contained in  $\mathcal{B}'$ ). But since all of these keep intersecting the  $a_0 + a_1$  neighborhood of  $\ell_0$ , it follows that there are only finitely many of them.

Therefore up to replacing  $c'_2$  if necessary by the leaf nonseparated from  $c'_1$ ,  $c'_2$  and closest to  $c'_1$  we assume there are no nonseparated leaves from  $c'_1$ ,  $c'_2$  and between them. This way, up to replacing the curves and choosing the rays conveniently one can produce  $\sigma$  so that it produces a good half-band as above.  $\square$

We can then start the analysis with a good half-band  $\mathcal{B}_L = \mathcal{B}(r_1, r_2, \sigma)$ . Denote by  $E = E(r_1, r_2)$  the leaf of  $\tilde{\mathcal{F}}_2$  such that  $r_1 \cup r_2 \subset L \cap E$ . Choosing  $\alpha$  in a similar way as  $\sigma$  but contained in  $E$  and such that the curve  $\beta$  in the last condition is small, this implies that every curve of  $\mathcal{G}$  intersecting the interior of the band intersects  $\mathcal{B}_L$  in a compact set. It also implies that every leaf of  $\tilde{\mathcal{F}}_2$  that intersects the interior of  $\mathcal{B}_L$  must intersect a transversal to  $E$ , in particular,  $E$  cannot intersect the interior of  $\mathcal{B}_L$ , unless  $\gamma E = E$ , in which case we get the conclusion of Proposition 8.3.

We consider an arc  $\tau$  in  $E$  joining the endpoints of  $\sigma$ . Note that there is a uniform  $a_1$  such that we can choose  $\tau$  to have length less than  $a_1$  because leaves of  $\tilde{\mathcal{F}}_2$  are uniformly properly embedded.<sup>9</sup> The  $r_i$  are contained in leaves  $c_i$  of  $\tilde{\mathcal{G}}$ . We also choose  $\tau$  to that it does not intersect  $c_1 \cup c_2$  in its interior. We denote by  $\mathcal{B}_E = \widehat{\mathcal{B}}(r_1, r_2, \tau)$  the half-band in  $E$  whose boundary in the compactification of  $E$  is  $r_1 \cup r_2 \cup \tau \cup \xi$ . Note that this may not be a “good half-band” as we defined before, because there is no reason for  $r_1$  and  $r_2$  to be nonseparated in  $\mathcal{G}_E$ .

**Lemma 8.6.** *There is an embedded disk  $D$  in  $\tilde{M}$  whose boundary is  $\sigma \cup \tau$  intersecting  $\mathcal{B}_L \cup \mathcal{B}_E$  only in the boundary.*

*Proof.* The bands  $\mathcal{B}_L$  and  $\mathcal{B}_E$  only intersect at their boundaries  $r_1 \cup r_2$ , and thus  $\mathcal{B}_L \cup \mathcal{B}_E$  is a properly (and tamely) embedded copy of  $S^1 \times [0, \infty)$  (in fact, it is piecewise  $C^1$ ). Taking the one point compactification of  $\tilde{M}$  to  $S^3$  we get that  $\hat{D} = \mathcal{B}_L \cup \mathcal{B}_E \cup \{\infty\}$  is a tamely embedded disk whose boundary is  $\tau \cup \sigma$  and therefore, there is a homeomorphism of  $S^3$  sending  $\hat{D}$  to the standard disk in a hemisphere (see [36, § 17]). Thus one can homotope the disk  $\hat{D}$  rel  $\partial \hat{D}$  away from  $\infty$  and so that it does not intersects  $\hat{D}$  in the interior and get an embedded disk  $D$  in  $\tilde{M}$  with the desired properties.  $\square$

Let  $\mathcal{V} := \mathcal{V}(r_1, r_2, \sigma, \tau)$  be the region defined in  $\tilde{M}$  bounded by  $D \cup \mathcal{B}_L \cup \mathcal{B}_E$  such that in each leaf inside, it limits only on  $\xi$ . This is a topological ball.<sup>10</sup> We call such a region a *good half-region*.

We analyze a good half-region  $\mathcal{V} = \mathcal{V}(r_1, r_2, \sigma, \tau)$  with boundary  $D \cup \mathcal{B}_L \cup \mathcal{B}_E$  where  $\mathcal{B}_L$  is a good half-band in  $L \in \tilde{\mathcal{F}}_1$  and  $\mathcal{B}_E$  is a half-band in  $E$ . Let  $p_i \in \sigma \cap \tau \cap r_i$  the corner points of  $D$  and small foliation boxes  $B_1^i$  and  $B_2^i$  of  $\tilde{\mathcal{F}}_1$  and  $\tilde{\mathcal{F}}_2$  respectively

<sup>9</sup>This follows from Theorem 2.1 in our case, but is also a general fact for  $\mathbb{R}$ -covered foliations; see, e.g., [16, Lemma 4.48].

<sup>10</sup>Again, this is a consequence of Schoenflies theorem in dimension 3, see [36, § 17]. Note that the surfaces we are constructing are all tamely embedded by construction (they are piecewise leaves of foliations, and  $D$  can be chosen smooth).

around  $p_i$  (for  $i = 1, 2$ ). We consider  $I_1^i, I_2^i$  small nondegenerate closed intervals in the leaf spaces of  $\tilde{\mathcal{F}}_1$  and  $\tilde{\mathcal{F}}_2$  consisting on the leaves intersecting  $B_1^i$  and  $B_2^i$  respectively. We can assume that the boxes are small enough that (for  $i = 1, 2$  and  $j = 1, 2$ ),

- every leaf of  $\tilde{\mathcal{F}}_1$  through  $B_1^i$  intersects  $E$ ,
- every leaf of  $\tilde{\mathcal{F}}_2$  through  $B_2^i$  intersects  $L$ , and
- the set  $\widehat{D}_\varepsilon(L, I_1^i)$  contains<sup>11</sup> $\mathcal{B}_L$  (see Proposition 2.7).


By slightly changing  $B_1^i$  and  $B_2^i$  we can and will assume that  $I_1^1 = I_1^2 = I_1$  and  $I_2^1 = I_2^2 = I_2$ . These conditions on the sets  $B_i^j$  and the intervals  $I_1, I_2$  will be assumed in what follows. We can also consider  $I_1, I_2$  small enough that they are disjoint from their images by the deck transformation associated to the center of  $\pi_1(M)$  (i.e., they correspond to an interval in the leaf space  $\widehat{\mathcal{L}}$  in  $\widehat{M}$ ; see Corollary 2.2). This implies that every  $\gamma \in \pi_1(M) \setminus \{\text{id}\}$  can have at most two fixed points in  $I_i$  and if it has two, one must be attracting and one repelling.

We are going to show that there are some restrictions on how  $\mathcal{V}$  can intersect with its translates under deck transformations. See Figure 18.

**Lemma 8.7.** *Assume that  $\gamma \in \pi_1(M)$  is such that*

- *there is a point  $z_1 \in r_1$  such that  $\gamma z_1 \in B_1^1 \cap B_2^1$ ,*
- *there is a point  $z_2 \in r_2$  such that the distance in  $L$  from  $z_1$  to  $z_2$  is less than  $2a_0 + 1$  and such that  $\gamma z_2 \in B_1^2 \cap B_2^2$ ,*
- *$\gamma(I_1) \subset \text{Int}(I_1)$ , and*
- *$I_2 \subset \text{Int}(\gamma(I_2))$ .*

*Then,  $\gamma L$  cannot intersect the interior of  $\mathcal{V}$ , unless  $\gamma E = E$ , in which case we achieve the conclusion of Proposition 8.3.*



**Figure 18.** Forbidden returns of a good half-region according to Lemma 8.7.

<sup>11</sup>This uses the fact that  $\xi \neq \alpha(L)$ .

*Proof.* We assume for a contradiction that  $\gamma L$  intersects the interior of  $\mathcal{V}$ . From how we chose the intervals it follows that there is a (unique) fixed point of  $\gamma$  in  $I_1$  (which is attracting) and one in  $I_2$  (which is repelling). Let  $L'$  be the fixed point by  $\gamma$  in  $I_1$  and similarly  $E'$  the fixed point of  $\gamma$  in  $I_2$ . Note that since  $\gamma$  is attracting on  $I_1$  then either  $L' = L$  or  $L'$  belongs to the same connected component of  $I_1 \setminus \{L\}$  as  $\gamma L$  and since  $\gamma$  is expanding on  $I_2$ , either  $E' = E$  or  $E'$  belongs to the connected component of  $I_2 \setminus \{E\}$  not containing  $\gamma E$  (recall that the intervals were chosen small enough that the action of  $\gamma$  has a unique fixed point in  $I_1$  and  $I_2$ ). In particular, both  $E'$  and  $L'$  intersect  $B_i^j$  for  $i = 1, 2$ ;  $j = 1, 2$  and using Corollary 2.2 we know that they intersect each such box in a unique plaque.

Since we have assumed that  $\gamma L$  intersects the interior of  $\mathcal{V}$  then  $\gamma L \neq L$ , and so by our choice of  $I_1$ , it follows that  $L'$  also intersects the interior of  $\mathcal{V}$ , since it belongs to the connected component of  $I_1 \setminus L$  containing  $\gamma L$  (the set of leaves of  $\tilde{\mathcal{F}}_1$  which intersect the interior of  $\mathcal{V}$ ). Since  $L' \in I_1$  and  $\mathcal{B}_L \subset \widehat{D}_\varepsilon(L, I_1)$  we have that  $L$  and  $L'$  are asymptotic in  $\mathcal{B}_L$  (i.e.,  $d(L \cap \mathcal{B}_L, L') < \varepsilon$ , meaning every point in  $\mathcal{B}_L$  is  $< \varepsilon$  from  $L'$ ) and we can apply Proposition 3.4. Denote by  $\mathcal{B}_{L'}$  the band in  $L'$  obtained by pushing from  $\mathcal{B}_L$  to  $L'$  in the sense of Proposition 3.4 (i.e., the band  $\mathcal{B}_{L'}$  is the one bounded by the intersections of  $E$  with  $L'$  which are close to  $r_1$  and  $r_2$ ). Note that this implies that  $\mathcal{B}_{L'}$  is a good half-band, and its boundary rays that we denote as  $r'_1, r'_2$  (and are contained in  $L' \cap E$ ) land in  $\xi$  and are nonseparated.

Let us first assume that  $\gamma E$  intersects  $\mathcal{V}$ . In this case, since  $\gamma z_1 \in B_1^1 \cap B_2^1$  and  $\gamma z_2 \in B_1^2 \cap B_2^2$  it follows that  $\gamma \mathcal{B}_L$  and  $\gamma \mathcal{B}_E$  must intersect  $\mathcal{V}$ . We obtain that  $\mathcal{V}$  projects to a solid torus in  $M_\gamma = \tilde{M}/\langle \gamma \rangle$ . Then that  $\gamma \mathcal{B}_L$ , whose boundaries are  $\gamma r_1$  and  $\gamma r_2$ , has a definite width in  $\gamma L$ , meaning that the distance from one curve to the other is bounded below by some uniform constant due to the fact that they belong to the same leaf of  $\tilde{\mathcal{F}}_2$ . This implies that the band cannot disappear and is completely contained in  $\mathcal{V}$ . It must thus be asymptotic to an infinite strip strictly inside  $\mathcal{B}_{L'}$  which must be invariant under  $\gamma$  and such that the boundaries are curves which limit on  $\xi$  and separate  $r'_1$  from  $r'_2$  contradicting the fact that  $\mathcal{B}_{L'}$  is a good half-band.

Suppose now that  $\gamma E = E$ . The argument in the previous case still applies so the band  $\mathcal{B}_L$  is asymptotic to a band in  $L'$  which is invariant under  $\gamma$ . This produces a bigon in  $L'$  and  $\mathcal{B}_L$  is asymptotic to this  $\gamma$ -invariant bigon. This achieves the conclusion of Proposition 8.3 in this case.

Finally we treat the case that  $\gamma E$  does not intersect  $\mathcal{V}$ . This implies that  $E'$ , the fixed point of  $\gamma$  in  $I_2$  belongs to the connected component of  $I_2 \setminus \{E\}$  not containing  $\gamma E$ . Thus,  $E'$  must intersect  $\mathcal{V}$ . Since  $\mathcal{B}_{L'}$  is a good half-band, each connected component of  $E' \cap \mathcal{B}_{L'}$  is a piece of a leaf of  $\mathcal{G}_{L'}$  and must be a compact interval by definition of good half-band. The same holds for  $E' \cap \mathcal{B}_L$  due to Proposition 3.4. However, since  $\gamma$  fixes both  $L'$  and  $E'$  and because  $E'$  intersects  $B_1^2$  in a unique

plaque we get that  $\gamma$  must fix the connected component of the intersection  $L' \cap E'$  intersecting  $B_1^2$  and thus projects to a closed curve which lifts to a curve separating  $r'_1$  from  $r'_2$  we have shown had to be nonseparated. This is a contradiction and completes the proof.  $\square$

Now, we will show that it is possible to find deck transformations with the properties required by Lemma 8.7.

Consider a geodesic ray  $\ell_0$  in  $L$  that limits in  $\xi$ . The small visual measure property implies that there is some  $a_0 > 0$  so that for each  $x$  in  $\ell_0$  there is  $z$  in  $r_i$  with  $d_L(x, z) < a_0$  (see Proposition 5.2).

**Lemma 8.8.** *Let  $\mathcal{V} = \mathcal{V}(r_1, r_2, \sigma, \tau)$  be a good half-region and  $p_1 = \sigma \cap r_1 \cap \tau$ . Consider  $B_1^1, B_2^1$  and  $I_1, I_2$  as above and assume that there exists a sequence of points  $z_n$  in  $r_1$ , with  $d_L(z_n, x_n) < a_0$  where the  $x_n$  are in a geodesic ray  $\ell_0$  as above and  $x_n \rightarrow \xi$ . Assume moreover that there are deck transformations  $\gamma_n$  such that  $\gamma_n z_n \in B_1^1 \cap B_2^1$ . Then, for large enough  $n$  we have that  $\gamma_n I_1 \subset \text{Int}(I_1)$  and  $\gamma_n^{-1} I_2 \subset \text{Int}(I_2)$ .*

*Proof.* This will use that  $\xi \neq \alpha(L)$ , but  $\xi = \alpha(E)$  by Proposition 3.9. Up to subsequence assume that  $\gamma_n z_n$  and  $\gamma_n x_n$  converge.

Fix a transversal  $\zeta$  to  $\mathcal{F}_1$  through  $p_1$  intersecting exactly the leaves of  $\tilde{\mathcal{F}}_1$  in  $I_1$ . Let  $L_1, L_2$  be the leaves of  $\tilde{\mathcal{F}}_1$  through the endpoints of  $\zeta$ . Recall that the ideal point of  $r_1, r_2$  and hence that of  $\ell_0$  is  $\xi \neq \alpha(L)$  so it is a contracting direction for  $\tilde{\mathcal{F}}_1$ . So for big enough  $n$  there is a transversal to  $\tilde{\mathcal{F}}_1$  of arbitrarily short length, through  $x_n$  and connecting the leaves  $L_1, L_2$ . By assumption  $d_L(x_n, z_n)$  is bounded by  $a_0$  independently of  $n$ , so using the local product structure of the foliation  $\mathcal{F}_1$  we obtain that there is also a transversal  $\beta_n$  to  $\tilde{\mathcal{F}}_1$  through  $z_n$  of very small length and connecting  $L_1$  to  $L_2$ . Then  $\gamma_n \beta_n$  is a transversal of very small length passing through  $\gamma_n z_n$  which is very close to  $p$ . In particular for  $n$  sufficiently  $\gamma_n \beta_n$  intersects a set of leaves of  $\tilde{\mathcal{F}}_1$  which is strictly contained in  $I_1$ . This implies that  $\gamma_n I_1 \subset \text{Int}(I_1)$  for  $n$  big enough.

Next consider  $I_2$  and  $\tilde{\mathcal{F}}_2$ . We first have to verify that in  $E$  the points  $z_n$  are a bounded distance from a geodesic ray in  $E$ . First recall that Proposition 2.3 proves that there is a constant  $Q_0 > 0$  such that for any  $F$  a leaf of  $\tilde{\mathcal{F}}_i$  then  $\Phi_F^i : F \rightarrow \mathbb{H}^2$  is a  $Q_0$  quasi-isometry.

The  $z_n$  are a bounded distance from the geodesic ray  $\ell_0$  in  $L$ . By Proposition 2.3 the image  $\Phi_L^1(\ell_0)$  is a quasigeodesic and hence the points  $\Phi_L^1(z_n)$  are a bounded distance from a geodesic ray in  $\mathbb{H}^2$  which we denote by  $\ell_1$ . Lemma 3.1 implies that there is  $a_1 > 0$  such that for any point  $x$  in  $\tilde{M}$ , if  $x \in L \in \tilde{\mathcal{F}}_1$  and  $x \in E \in \tilde{\mathcal{F}}_2$ , then  $d_{\mathbb{H}^2}(\Phi_L^1(x), \Phi_E^2(x)) < a_1$ . Hence  $\Phi_E^2(z_n)$  are a bounded distance from  $\ell_1$ . Applying  $(\Phi_E^2)^{-1}$  shows that  $z_n$  are a bounded distance from  $(\Phi_E^2)^{-1}(\ell_1)$ . The last

curve is a quasigeodesic in  $E$  (again Proposition 2.3) and we conclude that  $z_n$  are a bounded distance from a geodesic ray in  $E$  as claimed.

We know that  $\ell_1$  converges to  $\alpha(E)$ , so holonomy of  $\tilde{\mathcal{F}}_2$  along  $\ell_1$  is expanding. We have chosen  $B_2^1$  sufficiently small that the transversal through  $z_n$  intersecting the leaves corresponding to  $B_2^1$  will have length much bigger than length of a transversal through  $p_1$ . Then one gets that  $\text{Int}(\gamma_n I_2) \supset I_2$ , or equivalently  $\gamma_n^{-1} I_2 \subset \text{Int}(I_2)$  as claimed.  $\square$

Before we show that an accumulation point must be fixed by some deck transformation we will show some general property about geodesic rays in leaves of  $\tilde{\mathcal{F}}_i$ .

**Lemma 8.9.** *Let  $\ell$  be a geodesic ray in some leaf  $L \in \tilde{\mathcal{F}}_i$  with  $\partial\Phi(\ell) = \xi \in \partial\mathbb{H}^2$  then one of the following holds:*

- (1) *there is  $\gamma \in \pi_1(M) \setminus \{\text{id}\}$  and a leaf  $L' \in \tilde{\mathcal{F}}_i$  such that  $\gamma L' = L'$  and  $\gamma\xi = \xi$  and the projection of  $\ell$  spirals towards the projection of  $L'$ , or,*
- (2) *for every  $\varepsilon_1 > 0$  there is a foliated box  $U$  in  $\tilde{M}$  of diameter less than  $\varepsilon_1$ , a sequence of points  $y_n \in \ell$  going to infinity and deck transformations  $\eta_n \in \pi_1(M)$  for which  $\eta_n y_n \in U$  are in different leaves of  $\tilde{\mathcal{F}}_i$  and such that  $\eta_n L$  accumulate in infinitely many distinct leaves of  $\tilde{\mathcal{F}}_i$ .*

*Proof.* Up to changing  $\ell$  by a uniformly bounded amount which does not affect the result, one can assume that  $\mathcal{F}_i$  is the weak stable foliation of the geodesic flow on  $M = T^1 S$  for a hyperbolic metric on  $S$  (cf. Theorem 2.1). Note that since this is the case, the accumulation set of  $\ell$  when projected to  $M$  is the same as the accumulation of the orbit of the geodesic flow from  $\alpha(L)$  to  $\xi$  inside  $L$  traversed in the direction opposite to the flow. In particular it consists of a compact connected set saturated by orbits that we shall denote by  $\Lambda$ .

First, note that unless  $L = \gamma L$  and  $\gamma\xi = \xi$  for some  $\gamma \in \pi_1(M) \setminus \{\text{id}\}$  (in which case  $\Lambda$  is the unique closed geodesic in  $L$  and we are in the first option) we have that all returns of the projection of  $\ell$  to  $M$  to a foliation box must happen in distinct plaques of  $\mathcal{F}_1$ . This is the same for any backward orbit of the geodesic flow, if it limits in a point  $x$  and it is not periodic, then it must intersect a transversal to  $x$  in infinitely many distinct plaques of  $\mathcal{F}_1$  (because  $\mathcal{F}_1$  is the weak stable foliation of the geodesic flow by construction).

Next, we assume that there is some  $\varepsilon_1 > 0$  such that for every  $x \in \Lambda$  the two-dimensional transversal of size  $\varepsilon_1$  to the geodesic flow through the point  $x$  does not intersect  $\Lambda$  except perhaps at  $x$  itself. In this case, we get that  $\Lambda$  is a closed geodesic which belongs to some leaf  $\mathcal{F}_1$  and considering some lift  $L'$  of this leaf to  $\tilde{M}$  gives us the first option of the lemma.

Finally, assume that  $\Lambda$  is not a periodic orbit. Then, there is  $x \in \Lambda$  such that every transversal to the flow through  $x$  intersects  $\Lambda$  in infinitely many distinct

leaves. For every foliated neighborhood  $U$  of  $x$  we get that the returns need to have accumulation points every point of  $\Lambda \cap U$  which has infinitely many different leaves obtaining the second option of the lemma.  $\square$

Using the small visual measure property, one can extend this to rays of  $\mathcal{G}_L$  for every  $L$ .

**Corollary 8.10.** *Let  $r$  be a ray in some leaf of  $\mathcal{G}_L$  for  $L \in \tilde{\mathcal{F}}_i$  with ideal point  $\xi$  and let  $\ell_0$  be a geodesic ray in  $L$  with ideal point  $\xi$ . Suppose that  $\ell_0$  does not spiral towards a leaf  $L' \in \tilde{\mathcal{F}}_i$  invariant under some deck transformation. Then, for every  $\varepsilon_2 > 0$  there is a sequence of points  $x_n \in r$  going to infinity and deck transformations  $\eta_n \in \pi_1(M)$  such that for every  $n_0 > 0$  there are  $n_1 < n_2 < n_3$  such that  $\eta_{n_1}x_{n_1}, \eta_{n_2}x_{n_2}, \eta_{n_3}x_{n_3}$  are  $\varepsilon_2$  distance apart and  $\eta_{n_2}L$  is between  $\eta_{n_1}L$  and  $\eta_{n_3}L$ .*

*Proof.* Fix  $a_0$  given by the small visual measure so that for every  $y \in \ell_0$  there is a point  $x \in r$  such that  $d(x, y) < a_0$ . For  $y \in \tilde{M}$  denote by  $D_y$  to the disk of  $\tilde{\mathcal{F}}_i$  of radius  $a_0$  centered in  $y$ .

We fix  $\varepsilon_2$  and choose  $\varepsilon_1$  sufficiently small that if two points  $z, w$  are  $\varepsilon_1$ -close, then, the disks  $D_z$  and  $D_w$  are at Hausdorff distance less than  $\varepsilon_2$ . Now, choose  $k$  sufficiently large that if one chooses  $k$ -points  $z_1, \dots, z_k$  in a transversal to  $\tilde{\mathcal{F}}_i$  of length  $\varepsilon_1$  and picks one point  $w_i$  in each  $D_{z_i}$  then it holds that at least 3 of them are  $\varepsilon_2$  apart.

Note that the assumption on  $r$  implies that  $\ell_0$  satisfies the second alternative in the previous lemma, implying that for every  $\varepsilon_1 > 0$  there is a sequence of points  $y_n \in \ell_0$  going to infinity in  $\ell_0$  and deck transformations  $\eta_n$  such that all points  $\eta_n y_n$  belong to an open set  $U$  of diameter less than  $\varepsilon_1$  and that the leaves  $\eta_n L$  have infinitely many accumulation points. In particular, we can assume that  $\eta_n y_n$  accumulate in  $k$ -points  $z_1, \dots, z_k \in U$  belonging to different leaves of  $\tilde{\mathcal{F}}_i$ .

Now, given  $n_0$ , we can choose  $n_1 < n_2 < n_3$  and points  $y_{n_1}, y_{n_2}$  and  $y_{n_3}$  such that the  $\eta_{n_i} y_{n_i}$  are in  $U$ , the leaf  $\eta_{n_2} L$  is between the leaves  $\eta_{n_1} L$  and  $\eta_{n_3} L$ , and the corresponding points  $x_{n_i}$  in  $r$  are  $\varepsilon_2$  close. This completes the proof.  $\square$

The next lemma completes the proof of Proposition 8.3 and Addendum 8.4.

**Lemma 8.11.** *Let  $c_1, c_2 \in \mathcal{G}_L$  be two nonseparated leaves in  $L \in \tilde{\mathcal{F}}_i$  whose nonseparated rays  $r_1, r_2$  satisfy  $\partial\Phi(r_1) = \partial\Phi(r_2) = \xi \neq \alpha(L)$ . Then there exists some  $\gamma \in \pi_1(M) \setminus \{\text{id}\}$  and  $L' \in \tilde{\mathcal{F}}_i$  such that  $\gamma L' = L'$  and  $\gamma\xi = \xi$ . The rays  $r_i$ , when projected to  $M$ , spiral towards the projection of  $L'$ , and there exists a  $\gamma$ -invariant bigon in  $L'$ .*

*Proof.* Assume without loss of generality (cf. Lemma 8.5) that  $r_1$  and  $r_2$  are boundaries of a good half-band  $\mathcal{B}_L$  in  $L$ .

Let  $\mathcal{V} := \mathcal{V}(r_1, r_2, \sigma, \tau)$  a good half-region with boundary  $D \cup \mathcal{B}_L \cup \mathcal{B}_E$  constructed as in Lemma 8.6. We can choose an ordering of the leaf space of  $\tilde{\mathcal{F}}_1$  so that leaves that intersect  $\mathcal{V}$  are above  $L$ .

By Proposition 8.2 we can assume that  $r_1$  satisfies the assumptions of Corollary 8.10. Orient  $r_1$  and given  $x \in r_1$  consider  $r_1(x)$  to be the ray starting at  $x$ . We can also define  $\mathcal{V}_x := \mathcal{V}(r_1(x), r_2, \sigma_x, \tau_x)$  by changing the arcs to intersect in  $x$ . Now, using Corollary 8.10 given  $\varepsilon_2$  we can find points  $x, y$  in  $r_1$  such that  $y$  is in  $r_1(x)$  and a deck transformation  $\eta$  such that  $x$  and  $\eta y$  are very close and that  $\eta L$  is above  $L$ . This contradicts Lemma 8.7.

It follows that the point  $\xi$  is invariant under some  $\gamma$  and that the ray  $r_1$  converges to a closed leaf  $L'$  of  $\tilde{\mathcal{F}}_1$  invariant under  $\gamma$ .

In addition, the region between  $r_1$  and  $r_2$  in  $L$  is asymptotic to a region in  $L'$ . This region in  $L'$  has ideal point  $\xi \neq \alpha(L')$ . The rays  $r_1$  and  $r_2$  are thus converging to some limit rays  $e_1$  and  $e_2$  in  $L'$ . These rays are nonseparated in  $L'$  and they have ideal point  $\xi$  such that  $\gamma\xi = \xi$  and  $\gamma L' = L'$ . Proposition 8.2 implies that  $e_1, e_2$  are the boundaries of a  $\gamma$ -invariant bigon. This completes the proof of Lemma 8.11.  $\square$

**Remark 8.12.** Note that in the proof we also obtain that the endpoint  $\xi$  is not equal to  $\alpha(L')$  for the leaf  $L'$  we have found.


## 9. Consequences of the existence of Reeb surfaces

The next result gives more structure and improves Theorem A. In a certain sense, it says that if the intersected foliation  $\mathcal{G}$  is not an Anosov foliation, then the foliations should look very much like the ones studied in Section 7.

**Theorem 9.1.** *Assume that  $\mathcal{F}_1$  and  $\mathcal{F}_2$  have a Reeb surface and let  $B$  be its lift to  $L \in \tilde{\mathcal{F}}_1$  (a non-Hausdorff bigon). Denote by  $\xi, \eta$  its limit points in  $\partial\mathbb{H}^2$ . Then, for every  $L' \in \tilde{\mathcal{F}}_i$  there is a pair of nonseparated leaves  $c_1, c_2 \in \mathcal{G}_{L'}$  such that  $\partial^\pm\Phi(c_i) = \{\xi, \eta\}$  for  $i = 1, 2$ .*

To prove this, we first provide some stability properties of non-Hausdorff bigons in nearby leaves. Let  $B$  be a non-Hausdorff bigon in a leaf  $L \in \tilde{\mathcal{F}}_i$  with boundaries  $c_1, c_2 \in \mathcal{G}_L$ , we denote by  $\xi_B^+$  and  $\xi_B^-$  the points in  $\partial\mathbb{H}^2$  given by  $\partial^\pm\Phi(c_i) = \{\xi_B^+, \xi_B^-\}$  for  $i = 1, 2$  and such that  $\{\xi_B^\pm\}$  is the ideal point of the rays of  $c_1, c_2$  which are nonseparated from each other. See Figure 19.

**9.1. Persistence and extension to the closure.** We next consider a Reeb surface lifting to a non-Hausdorff bigon  $B$  in some leaf  $L \in \tilde{\mathcal{F}}_i$  which is  $\gamma$ -invariant, and let  $I_\gamma^\pm$  be the connected components of  $\partial\mathbb{H}^2 \setminus \{\gamma^\pm\}$  where  $\gamma^\pm$  are the fixed points of  $\gamma$ . We will show that for one of the intervals  $I_\gamma^\pm$ , say  $I_\gamma^+$ , it holds that for every  $L' \in \tilde{\mathcal{F}}_i$  such that  $\alpha(L') \in I_\gamma^+$  the leaf  $L'$  has a non-Hausdorff bigon  $B'$  sharing the same endpoints as  $B$ . To do this, we will need to prove several stability properties,



**Figure 19.** The ideal points of a non-Hausdorff bigon  $B$ .

and along the way we will obtain some useful information that will help us to prove Theorem 9.1.

We first show the easy consequence of Proposition 3.4 that bigons persist in nearby leaves when both endpoints are marker points.

**Lemma 9.2.** *Let  $B$  be a non-Hausdorff bigon in a leaf  $L \in \tilde{\mathcal{F}}_i$  with boundaries  $c_1, c_2 \in \mathcal{G}_L$  such that  $\alpha(L) \notin \{\xi_B^+, \xi_B^-\}$ . Then, there is a neighborhood  $I$  of  $L$  in the leaf space of  $\tilde{\mathcal{F}}_i$  such that for every  $L' \in I$  there is a non-Hausdorff bigon  $B'$  in  $L'$  which shares both endpoints with  $B$ .*

*Proof.* This is just an application of Proposition 3.4 since the non-Hausdorff bigon is contained in  $\widehat{D}_\varepsilon(L, I)$  (see Proposition 2.7) for some  $\varepsilon$  for which Proposition 3.4 applies and some small neighborhood  $I$  of  $L$  in the leaf space of  $\tilde{\mathcal{F}}_i$ .  $\square$

Such bigons extend to the boundary  $\gamma$ -invariant leaves:

**Lemma 9.3.** *Let  $B$  be a non-Hausdorff bigon in a leaf  $L \in \tilde{\mathcal{F}}_i$  with boundaries  $c_1, c_2 \in \mathcal{G}_L$  such that  $\alpha(L) \notin \{\xi_B^+, \xi_B^-\}$ . Assume that there is  $\gamma \in \pi_1(M)$  whose fixed point at  $\partial\mathbb{H}^2$  are exactly  $\xi_B^+$  and  $\xi_B^-$ . Then, for every  $L' \in \tilde{\mathcal{F}}_i$  such that  $\alpha(L')$  is in the same connected component as  $\alpha(L)$  in  $\partial\mathbb{H}^2 \setminus \{\xi_B^+, \xi_B^-\}$  the leaf  $L'$  has a non-Hausdorff bigon with the same endpoints as  $B$ . Moreover, this property passes to the closure of that set of leaves and thus extends to the boundary leaves  $L_0, L_1$  such that  $\alpha(L_0) = \xi_B^+$  and  $\alpha(L_1) = \xi_B^-$ .*

We note that up to composing  $\gamma$  with a power of the deck transformation associated to fibers we can assume that the leaves  $L_0$  and  $L_1$  in the conclusion are  $\gamma$ -invariant.

*Proof.* We assume that  $L \in \tilde{\mathcal{F}}_1$ . The set of leaves for which  $\alpha(L')$  belongs to the same connected component of  $\alpha(L)$  in  $\partial\mathbb{H}^2 \setminus \{\xi_B^+, \xi_B^-\}$  is a countable union of open intervals. For any pair of such intervals, there is power of the deck transformation associated to the fiber (which induces the identity in  $\partial\mathbb{H}^2$ ) which maps one interval to the other, so it is enough to show that one such open interval  $I_0$  (the one containing  $L$ ) satisfies the desired property. We can also assume that  $\gamma$  fixes  $I_0$ .

Let  $E \in \tilde{\mathcal{F}}_2$  be the leaf such that  $E \cap L$  contains the boundaries of the non-Hausdorff bigon and consider the set of leaves  $I_1 \subset I_0$  consisting of leaves  $L'$  such that  $L' \cap E$  contains curves that bound a non-Hausdorff bigon joining  $\{\xi_B^+, \xi_B^-\}$ . Note that since  $\xi_B^+$  is the point where the nonseparation happens, it follows from Proposition 3.9 that  $\alpha(E) = \xi_B^+$  and is thus  $\gamma$ -invariant. We claim that this implies that  $\gamma E = E$ : note that since  $\gamma\alpha(E) = \alpha(E)$  then either  $E$  is fixed, or it is mapped to a leaf  $E'$  obtained by acting on  $E$  with a power of the deck transformation associated to the fiber and therefore  $\gamma$  acts freely on the leaf space of  $\tilde{\mathcal{F}}_2$  (recall that we are assuming that  $\gamma$  is the element of  $\pi_1(M) \setminus \{\text{id}\}$  which also fixes  $I_0$ , which is an interval in the leaf space of  $\tilde{\mathcal{F}}_1$ ). To show that  $\gamma E = E$  we therefore note that  $\gamma^n(c_1), \gamma^n(c_2)$  are boundaries in bigons in  $\gamma^n(L)$  and since  $\gamma$  fixes  $I_0$  which is a bounded interval in the leaf space of  $\tilde{\mathcal{F}}_1$  the iterates  $\gamma^n(L)$  converges to some leaf  $L' \in \tilde{\mathcal{F}}_1$ . By the small visual measure property it follows that  $\gamma^n(c_1)$  must remain intersecting a compact set in  $\tilde{M}$  and hence  $\gamma^n(E)$  which contains  $\gamma^n(c_1)$  must also intersect a compact set in  $\tilde{M}$  for all  $n$ . This shows that  $\gamma$  cannot act freely on the leaf space of  $\tilde{\mathcal{F}}_2$  and since  $\alpha(E)$  is  $\gamma$ -invariant we deduce  $\gamma E = E$ .

Applying the argument in the previous lemma and using the fact that as long as  $\alpha(L') \notin \{\xi_B^+, \xi_B^-\}$  we can push the non-Hausdorff bigon to nearby leaves we deduce that  $I_1$  is open in  $I_0$ .

To complete the proof we will show that  $I_1$  is also closed in  $I_0$ .

Consider  $L_n \rightarrow L' \in I_0$  with  $L_n$  in an open interval  $I$  contained in  $I_1$ . The leaf  $L_n$  contains a non-Hausdorff bigon  $B_n$  bounded by leaves  $r_1^n, r_2^n$  of  $\mathcal{G}_{L_n}$  contained in  $L_n$  but also in  $E$ , because pushing preserves the  $\tilde{\mathcal{F}}_2$  leaves they are in. Denote by  $D_n$  the region in  $E$  bounded by  $r_1^n$  and  $r_2^n$ . We can assume that the sequence  $D_n$  is monotonic. If the region  $D_n$  decreases with  $n$  (i.e., if  $D_{n+1} \subset D_n$ ) then since  $D_n$  must contain a non-Hausdorff bigon of  $E$  we get that in the limit the curves  $r_1^n$  and  $r_2^n$  converge to curves joining  $\xi_B^+$  and  $\xi_B^-$ , as we want to show.

We assume then that the sets  $D_n$  increase with  $n$  and consider the set  $D_\infty = \overline{\bigcup_n D_n}$  whose closure in  $E \cup S^1(L)$  is compact, connected, and cannot be the whole  $E \cup S^1(E)$  (because of the small visual measure property). Let  $R'_1 = \{\ell_1, \dots, g_k, \dots\}$  be the (possibly finite, but countable) collection of all limits of the curves  $r_1^n$  in  $\mathcal{G}_E$ . Similarly, denote by  $R'_2 = \{\ell_1^2, \dots, \ell_m^2, \dots\}$  the limits of  $r_2^n$ . Since the  $r_1^n, r_2^n$  are in  $L_n \cap E$ , the limits are in  $E \cap L'$ . Hence they belong to  $\mathcal{G}_E$  and  $\mathcal{G}_{L'}$ . By Proposition 3.9 the limit points of the leaves  $\ell_i^j$  can be either the ones of  $r_j^n$  (that is,

$\xi_B^+ = \alpha(E)$  or  $\xi_B^-$ ), or  $\alpha(L')$  if there is more than one limit curve.

The set  $R'_1 \cup R'_2$  in  $E \cup S^1(E)$  bounds the set  $D_\infty$  which is  $\mathcal{G}_E$ -saturated and has nonempty interior since the region between  $r_1^n$  and  $r_2^n$  must contain a non-Hausdorff bigon which belongs to all the  $D_n$ . Since the possible limit points of any leaf in the boundary are only  $\xi_B^+$ ,  $\xi_B^-$ ,  $\alpha(L')$ , It follows that  $R'_1$  or  $R'_2$  must contain at least one leaf with an ideal point in  $\xi_B^+$  and one with an ideal point in  $\xi_B^-$  (they could be the same leaf).

Therefore, if  $L'$  does not contain a non-Hausdorff bigon joining  $\xi_B^+$  and  $\xi_B^-$  bounded by leaves in  $R'_1 \cup R'_2 \subset L' \cap E$ , there must be a region whose limit in  $\partial\mathbb{H}^2$  is either exactly  $\{\xi_B^+, \xi_B^-, \alpha(L')\}$ ; or two regions, one with ideal points  $\{\xi_B^+, \alpha(L')\}$ , and one with ideal points  $\{\xi_B^-, \alpha(L')\}$ . We analyze the first possibility, the second one is similar. Since  $E$  is  $\gamma$ -invariant and since  $\gamma$  does not fix  $\alpha(L')$ , then applying  $\gamma$  or  $\gamma^{-1}$  to a curve with ideal points  $\xi_B^-, \alpha(L')$  we obtain a curve with one ideal point  $\xi_B^-$  and another in the open interval  $(\xi_B^+, \alpha(L'))$  which does not contain  $\xi_B^-$ . This gives a contradiction since the curves in  $\mathcal{G}_E$  cannot cross.

Note that the boundary leaves bound a non-Hausdorff bigon in  $L'$ . Indeed these boundary leaves have ideal points  $\xi_B^-, \xi_B^+$ , and Proposition 3.4 implies that the same local picture has to be seen in  $L'$  as for  $L_n$  for large  $n$ . In other words the bigons in  $L_n$  push through to  $L'$ .

Note that in the case of  $L_n \rightarrow L_i$  with  $i = 0, 1$  we get the same conclusion (even simpler, since we do not have the possibility to have three limit points), only that when  $\alpha(L_i) = \alpha(E)$  we cannot ensure that the region bounded by the leaves is a non-Hausdorff bigon. However, since there is no transversal intersecting both boundary leaves, it is easy to see that there must be one non-Hausdorff bigon in between.  $\square$

When the endpoints of the bigon contain the nonmarker point of  $L$ , stability is harder to establish. There are two cases depending on whether  $\xi_B^+ = \alpha(L)$  or  $\xi_B^- = \alpha(L)$ .

**9.2. Half interval stability: the case where  $\alpha(L) = \xi_B^+$ .** The goal of this subsection is to give a proof of the following

**Proposition 9.4.** *Let  $B$  be a non-Hausdorff bigon in a leaf  $L \in \tilde{\mathcal{F}}_i$  with boundaries  $c_1, c_2 \in \mathcal{G}_L$  such that  $\alpha(L) = \xi_B^+$ . Assume moreover that for some  $\gamma \in \pi_1(M) \setminus \{\text{id}\}$  we have that  $\gamma B = B$ . Then, there is a half neighborhood  $I$  of  $L$  in the leaf space of  $\tilde{\mathcal{F}}_i$  (i.e.,  $I$  is a connected component of  $J \setminus \{L\}$  where  $J$  is a neighborhood of  $L$  in the leaf space) such that for every  $L' \in I$  there is a non-Hausdorff bigon  $B'$  in  $L'$  which shares both endpoints with  $B$ .*

Let us assume that  $L \in \tilde{\mathcal{F}}_1$ . Since  $c_1$  and  $c_2$  are nonseparated in  $\mathcal{G}_L$ , we know that there is  $E \in \tilde{\mathcal{F}}_2$  such that  $c_1 \cup c_2 \subset E \cap L$ .

Let  $B_E$  denote the region in  $E$  bounded by  $c_1 \cup c_2$  which is an infinite band with bounded width, limiting on  $\xi_B^\pm$  (cf. Lemma 3.1). This set  $B_E$  is not necessarily a non-Hausdorff bigon since  $c_1$  and  $c_2$  could be separated in  $\mathcal{G}_E$ , but there must exist some non-Hausdorff bigon contained in  $B_E$  and limiting in the same points. By assumption  $\alpha(L) = \xi_B^+$  and using Proposition 3.9 we have that  $\alpha(E) = \xi_B^+ = \alpha(L)$ . Up to considering the inverse, we can assume that  $\gamma$  acts as an expansion on  $\xi_B^+$ . (Since  $\alpha(E) = \alpha(L)$  we have  $E = \gamma E$ .)

Fix small transversals  $\tau_1$  and  $\tau_2$  to  $\mathcal{G}_E$  in  $E$  parametrized in such a way that the leaf  $L_t \in \tilde{\mathcal{F}}_1$  through  $\tau_1(t)$  also passes through  $\tau_2(t)$  and that  $\tau_i(0)$  belongs to  $c_i$ . We also assume these are parametrized so that for  $t > 0$  we have that  $\tau_1(t)$  belongs to  $B_E$ .

For  $t < 0$  denote by  $c_1^t$  and  $c_2^t$  the leaves of  $\mathcal{G}_E$  through  $\tau_1(t)$  and  $\tau_2(t)$  respectively, which belong to  $L_t \cap E$ . Note that these cannot coincide because they are separated by  $c_1$  (and  $c_2$ ). Note also that the rays of  $c_1^t$  and  $c_2^t$  in the direction of  $\xi_B^-$  must approximate  $c_1$  and  $c_2$  for  $t$  small because a neighborhood of  $\xi_B^-$  in  $L$  is contained in the set where Proposition 3.4 applies.

The leaves  $c_1^t$  and  $c_2^t$  cannot be connected by a transversal in  $L_t$  (since they are both in  $E$ ). Between them (maybe coinciding with one of them), there is a pair of leaves  $e_1^t, e_2^t$  which is nonseparated and such that  $e_1^t, e_2^t$  both separate  $c_1^t$  from  $c_2^t$  (unless they coincide with them). It follows that both have  $\xi_B^-$  as an ideal point.

We call  $E_t \in \tilde{\mathcal{F}}_2$  the leaf such that  $e_1^t \cup e_2^t \subset L_t \cap E_t$ . We will show that the other landing point of  $e_1^t, e_2^t$  is in  $\xi_B^+$  and thus there is a non-Hausdorff bigon in  $L_t$  with ideal points  $\xi_B^-, \xi_B^+$  as desired. Notice that a priori  $E_t$  does not vary continuously with  $t$ , even though we will show in the next lemma that  $E_t$  is continuous with  $t$  when  $t = 0$ .

The other rays of  $e_1^t$  and  $e_2^t$  must land in  $\alpha(E_t)$  (because the rays we showed limit in  $\xi_B^-$  are separated by Proposition 3.4 and thus the other rays are nonseparated; therefore Proposition 3.9 applies). In particular, the curves  $e_1^t$  and  $e_2^t$  bound a non-Hausdorff bigon  $B_t$  in  $L_t$  which limits in  $\xi_t = \alpha(E_t)$  and in  $\xi_B^-$ . Let  $L_0 = L$  and  $\xi_0 = \alpha(L_0) = \alpha(E)$ .

**Lemma 9.5.** *The function  $\xi_t$  of  $t$  is continuous at zero. More precisely, for every  $J$  neighborhood of  $\xi_B^+$  there is  $\delta > 0$  such that for every  $t \in (\delta, 0]$  we have that  $\xi_t \in J$ .*

*Proof.* To see this fix some small neighborhood  $U$  of  $\xi_0$  in  $\partial\mathbb{H}^2$ . Now choose a transversal  $\eta$  to  $\mathcal{G}_L$  starting at a point in  $c_1$  and entering  $B$ .

We choose  $\eta$  small enough such that if a leaf  $\ell$  of  $\mathcal{G}_{L_0}$  intersects  $\eta$ , then the leaf  $\ell$  is contained in a leaf  $E' \in \tilde{\mathcal{F}}_2$  such that  $\alpha(E') \in U$ . Denote by  $\ell_0$  the leaf of  $\mathcal{G}_L$  through the endpoint of  $\eta$  which is not in  $c_1$ . In particular  $\ell_0$  is contained in the interior of  $B$ . Now choose a neighborhood  $I$  of  $\alpha(L)$  such that  $\ell_0 \cup \eta$  is contained in  $\widehat{D}_\varepsilon(L, I)$  (this set is defined as in (2-6), where  $\varepsilon$  is chosen so that Proposition 3.4

holds). Recall that  $L = L_0$ . Let  $\ell_t$  be the push through to  $L_t$  of the leaf  $\ell_0 \subset L$ . In other words  $\ell_t$  is the component of  $\tilde{\mathcal{F}}_2(\ell_0) \cap L_t$  which is  $\varepsilon$ -close to  $\ell_0$ . In the same way  $\eta$  can be pushed to a transversal  $\eta_t$  to  $\mathcal{G}_{L_t}$  in  $L_t$  starting in  $c_1^t$  and ending in  $\ell_t$ . For each  $t$ ,  $\ell_t$  is contained in between  $c_1^t, c_2^t$  in  $L_t$ . In addition there is a transversal  $\eta_t$  to  $\mathcal{G}_{L_t}$  from  $c_1^t$  to  $\ell_t$ . Therefore  $\ell_t$  is in between  $e_1^t, e_2^t$  in  $L_t$ . In particular this implies that  $e_1^t$  is in between  $c_1^t$  and  $\ell_t$  in  $L_t$ . We have that  $c_1^t \subset E$ ,  $e_1^t \subset E_t$  and  $\ell_t \subset \tilde{\mathcal{F}}_2(\ell_0)$ , with  $\alpha(\tilde{\mathcal{F}}_2(\ell_0))$  in  $U$ . Since  $\alpha(E)$  is in  $U$ , it now follows that  $\alpha(E_t)$  is in  $U$  all well, as long as  $\alpha(L_t)$  is in  $I$ . Since  $\xi_t = \alpha(E_t)$  this completes the proof.  $\square$

Note that we have also shown that there is a transversal to  $\mathcal{G}_{L_t}$  from  $e_1^t$  to  $c_1^t$  which  $\varepsilon$  pushes to  $L = L_0$  and similarly there is a transversal from  $e_2^t$  to  $c_2^t$  which also pushes to  $L$ . It now makes sense to talk about monotonicity of  $\xi_t$ , we can indeed show:

**Lemma 9.6.** *The point  $\xi_t$  varies in a weakly monotonic way, that is, for small  $t, t' \in (-\delta, 0]$  we have that if  $t' < t$  then  $\xi_{t'} \leq \xi_t$  for the orientation of  $J$  making  $\xi_B^+$  the maximal point.*

*Proof.* For small  $\delta' > 0$ , consider  $\eta : (-\delta', 0] \rightarrow L$  such that  $\eta$  is transverse to  $\mathcal{G}_L$  and such that  $\eta(0) \in c_1$  and  $\eta(s) \in B$  for all  $s \in (-\delta', 0)$ . Note that as we have shown in Proposition 3.9, that if  $\delta$  is small enough, we know that for every  $t \in (-\delta, 0)$  we have that  $e_1^t$  belongs to the same leaf of  $\tilde{\mathcal{F}}_2$  as  $\eta(s)$  for some  $s \in (-\delta', 0]$ . This identification will be recorded by a function  $\rho : (-\delta, 0] \rightarrow (-\delta', 0]$  such that  $\rho(t) = s$ .

If  $\delta$  is small enough, then the image of  $\eta$  is contained in  $\widehat{D}_\varepsilon(L, I_\delta)$  where  $I_\delta$  is the interval of  $\partial\mathbb{H}^2$  made by  $\alpha(L_t)$  with  $t \in (-\delta, 0]$ , so, applying Proposition 3.4 we find transversals  $\eta' : (-\delta', 0] \rightarrow L_t$  intersecting the same  $\tilde{\mathcal{F}}_2$  leaves. Denote by  $\ell_s \in \mathcal{G}_L$  the leaf of  $\mathcal{G}_L$  through the point  $\eta(s)$  and by  $E'_s \in \tilde{\mathcal{F}}_2$  the leaf that contains  $\ell_s$ . We note that all  $\ell_s$  with  $s \in (-\delta', 0)$  are bubble leaves with endpoint in  $\xi_B^-$ .

For the function  $\rho$  defined above, we have  $E_t = E'_{\rho(t)}$ . Consider  $e_t^t$  to be the leaf of  $\mathcal{G}_{L_t}$  containing  $\eta'(\rho(t'))$ . We get that  $e_t^t = e_1^t$  by definition.

We can thus restate the claim stating that whenever  $t' < t < 0$  one has that  $\rho(t') \leq \rho(t)$ . We assume by contradiction that this does not hold, that is, for a pair  $t' < t < 0$  we have  $\rho(t') > \rho(t)$ . Then we get that the leaf  $e_t^{t'}$  is a bubble leaf with endpoint  $\xi_B^-$ . On the other hand  $e_t^t = e_1^t$  is a leaf in  $L_t$  with one ideal point  $\xi_B^-$ . It follows that the leaf  $E_t = E'_{\rho(t)}$  must intersect  $L_{t'}$  and  $L_0$  in bubble leaves while it intersects  $L_t$  in at least  $e_1^t \cup e_2^t$ . We will show that this is impossible: Denote by  $c$  the corresponding bubble intersection of  $E_t$  with  $L$ . By the remark after the proof of Lemma 9.5, we know that there is a small transversal  $\beta$  to  $\mathcal{G}_{L_{t'}}$  in  $L_{t'}$  from  $e_t^{t'}$  to  $c_2^{t'}$  and this pushes to  $L_0$ , through  $\tilde{\mathcal{F}}_2(e_t^{t'}) = E_t$ . Then  $E_t$  intersects  $L_t$  near the push through of  $\beta$ . The same happens for  $e_2^t$ . Hence the local leaf of  $E_t$  passes through  $e_2^t$  and also  $c$ . In other words there is a small transversal  $v_2$  to  $\mathcal{G}_{E_t}$  in  $E_t$  intersecting

$e'_1$ ,  $e'_2$  and  $c$  in turn. In the same way there is a small transversal  $v_1$  to  $\mathcal{G}_{E_t}$  in  $E_t$  from  $e'_1$  to  $e'_2$  to  $c$ . Consider the closed curve in  $E_t$  which is the concatenation of  $v_1$ , a segment in  $c$ ,  $v_2$  and a segment in  $e'_1$ . This closed curve does not bound a disk in  $E_t$  because  $e'_1$ ,  $e'_2$  intersect this curve in a single point. This would show that  $E_t$  is not a plane. This contradiction completes the proof.  $\square$

Note that Lemma 9.2 applies directly as soon as  $\xi_t \neq \alpha(L_t)$ , so, we get:

**Lemma 9.7.** *For every  $t \in (-\delta, 0]$  either  $\xi_s$  is locally constant near  $t$  or  $\xi_t = \alpha(L_t)$ .*

Finally, we show:

**Lemma 9.8.** *If  $\xi_t = \alpha(L_t)$  for  $t \neq 0$  then, one cannot have that  $L_t$  is invariant under some  $\gamma_t \in \pi_1(M) \setminus \{\text{id}\}$ .*

*Proof.* First notice that if this is the case then also one has that  $\gamma_t \xi_t = \xi_t$  since  $\gamma_t L_t = L_t$  implies that  $\alpha(L_t)$  is  $\gamma_t$ -invariant. On the other hand, by definition,  $L_t$  contains a non-Hausdorff bigon  $B_t$  whose endpoints are  $\xi_t$  and  $\xi_B^-$ . Since  $t \neq 0$  and  $\xi_t = \alpha(L_t)$  then  $\xi_t \neq \alpha(L_0)$ . But as  $\xi_B^+ = \alpha(L_0)$ , then  $\xi_t \neq \xi_B^+$ . Also we know that  $\xi_B^+$  and  $\xi_B^-$  are the fixed points of  $\gamma = \gamma_0$  (the deck transformation leaving  $L = L_0$  invariant) then  $\gamma_t \xi_B^- \neq \xi_B^-$ . Let  $B_t$  be a bigon in  $L_t$  with ideal points  $\xi_t$  and  $\xi_B^-$ .

Now we argue as in Proposition 8.2: Due to the small visual measure property there is a uniform bound on the number of distinct non-Hausdorff bigons that share an endpoint in a given leaf of  $\tilde{\mathcal{F}}_t$ . On the other hand, applying  $\gamma_t^n$  to  $B_t$  we obtain infinitely many disjoint non-Hausdorff bigons in  $L_t$  sharing one of the endpoints, namely  $\xi_t$ . This gives a contradiction and proves the lemma.  $\square$

*Proof of Proposition 9.4.* We show that  $\xi_t$  must be constant and equal to  $\xi_B^+$ , as desired. Assume first that we have that  $\xi_t = \alpha(L_t)$  in an open interval  $I \subset (-\delta, 0)$ . Since leaves with nontrivial stabilizer are dense, it follows that for some  $t$  close to 0 we have that  $L_t$  is  $\gamma_t$  invariant for some  $\gamma_t \in \pi_1(M) \setminus \{\text{id}\}$  contradicting Lemma 9.8.

Now, assume that for some  $t \in (-\delta, 0)$  we have that  $\xi_t \neq \alpha(L_t)$ . Consider  $A = \{s \in (-\delta, 0) : \xi_s = \xi_t\}$ . Note that Lemma 9.5 implies  $\xi_t$  is continuous at  $t = 0$  we know that  $A$  avoids a neighborhood of 0. Therefore, if  $s_0 < 0$  is the supremum of  $A$ , it follows from Lemma 9.7 that  $\xi_{s_0} = \alpha(L_{s_0})$ . In  $L_{s_0}$  we have a bigon  $B_{s_0}$  with  $\xi_{B_{s_0}}^+ = \xi_{s_0}$ . Applying Proposition 8.3 to  $L_s$  with  $s$  in the interior of the interval (so  $s \neq s_0$ ) we deduce that  $\xi_{s_0}$  is fixed by some deck transformation  $\hat{\gamma}_s \in \pi_1(M)$  and not acting trivially on  $\partial\mathbb{H}^2$ . Since  $\alpha(L_{s_0}) = \xi_{s_0}$  then up to changing  $\hat{\gamma}_s \in \pi_1(M)$  by some power of the deck transformation generated by the fiber we get some  $\gamma_s \in \pi_1(M) \setminus \{\text{id}\}$  which fixes  $L_{s_0}$  again contradicting Lemma 9.8, unless  $s_0 = 0$ . In other words the interval where  $\xi_s$  is constant has an endpoint in 0.

This completes the proof that  $\xi_t$  must be constant equal to  $\xi_B^+$  for  $t \in (-\delta, 0)$  and thus completes the proof of Proposition 9.4.  $\square$

**Remark 9.9.** We have in fact showed the following: If  $B$  is a  $\gamma$ -invariant bigon in a leaf  $L \in \tilde{\mathcal{F}}_1$ , and  $E \in \tilde{\mathcal{F}}_2$  is the leaf which intersects  $B$  in the boundary  $c_1, c_2 \in \mathcal{G}_L$ , is such that the nonseparated rays of  $c_1, c_2$  in  $L$  land in  $\alpha(L)$ , then for every leaf  $L' \in \tilde{\mathcal{F}}_1$  close to  $L$  intersecting  $E$  outside the region  $B_E$  bounded by  $c_1 \cup c_2$ , the leaf  $L'$  contains a bigon  $B'$  whose endpoints coincide with those of  $B$  and its boundaries correspond to the intersection with  $E$ .

**9.3. Half interval stability: the nonseparated side is a marker point.** This means that  $\xi_B^+ \neq \alpha(L)$ , but  $\xi_B^- = \alpha(L)$ . When the endpoint of the bigon is a marker point for  $L$ , we can also push it to one side, but the argument is different:

**Proposition 9.10.** *Let  $B$  be a non-Hausdorff bigon in a leaf  $L \in \tilde{\mathcal{F}}_i$  with boundaries  $c_1, c_2 \in \mathcal{G}_L$  such that  $\alpha(L) = \xi_B^-$ . Assume moreover that for some  $\gamma \in \pi_1(M) \setminus \{\text{id}\}$  we have that  $\gamma B = B$ . Then, there is a half neighborhood  $I$  of  $L$  in the leaf space of  $\tilde{\mathcal{F}}_i$  such that for every  $L' \in I$  there is a non-Hausdorff bigon  $B'$  in  $L'$  which shares both endpoints with  $B$ .*

*Proof.* Assume for concreteness that  $L \in \tilde{\mathcal{F}}_1$ . Let  $E$  be the leaf in  $\tilde{\mathcal{F}}_2$  such that  $c_1 \cup c_2 \subset L \cap E$ . Then  $\alpha(L) = \xi_B^-$  by assumption and  $\alpha(E) = \xi_B^+$  by Proposition 3.9.

Denote by  $B_E$  the region in  $E$  bounded between  $c_1$  and  $c_2$ . Suppose that  $B'$  is a non-Hausdorff bigon contained in  $B_E$ . We claim that  $B'$  must have its nonseparated rays land in  $\xi_B^+$  (and in particular, this shows that  $c_1$  and  $c_2$  cannot be nonseparated in  $E$  because they belong to  $L$  and  $\alpha(L) = \xi_B^-$  and Proposition 3.9 would imply that the nonseparated rays land there). To prove the claim, consider first some  $E'$  very close to  $E$  in such a way that it intersects  $B$  in a curve which is very close to  $c_1 \cup c_2$ . If  $C$  is a non-Hausdorff bigon in  $B_E$  limiting in  $\xi_B^-$ , first let  $L''$  be the  $\tilde{\mathcal{F}}_1$  leaf containing  $\partial C$ , and then can take a leaf  $L'$  of  $\tilde{\mathcal{F}}_1$  very close to  $L''$  and in such a way that the intersection of  $L'$  with  $E$  is a bubble leaf with points very close to  $E'$ . It follows that the intersection of  $L'$  and  $E'$  in the region between  $B$  and  $B_E$  must be compact, which is a contradiction because it would give a circle leaf in  $\mathcal{G}_L$  (and  $\mathcal{G}_{E'}$ ). This proves the claim.

The same argument shows that any bigon in  $B_E$  cannot have a boundary leaf in  $c_1$  or  $c_2$ . This implies that inside  $B_E$  both  $c_1, c_2$  have a neighborhood that does not intersect a bigon. Starting from  $c_1$ , consider leaves  $e$  of  $\mathcal{G}_E$  inside  $B_E$  and near  $c_1$ . There is a small interval of leaves where  $\gamma$  acts as a contraction or expansion in this interval. Similarly for  $c_2$ . Hence there is a maximal interval  $[c_1, e_1]$  where  $\gamma$  fixes the endpoints and no other leaf of  $\mathcal{G}_E$  in between. Similarly there is a maximal interval  $[c_2, e_2]$  with the same properties. The leaves  $e_1, e_2$  are in the same leaf  $L'$  of  $\tilde{\mathcal{F}}_1$  and it is invariant under  $\gamma$ . By construction  $\alpha(L') = \xi_B^+$ . Finally any leaf  $L''$  of  $\tilde{\mathcal{F}}_1$  between  $L$  and  $L'$  has a bigon  $B_{L''}$  which is asymptotic to  $B$  in one direction and to a bigon  $B_{L'}$  in the other direction.

This provides the interval required by the proposition. □

**Remark 9.11.** As a consequence of the proof we have the following property: Let  $B$  a  $\gamma$ -invariant non-Hausdorff bigon in a leaf  $L \in \tilde{\mathcal{F}}_1$  bounded by curves  $c_1, c_2$  such that  $c_1 \cup c_2 \subset L \cap E$  for some  $E \in \tilde{\mathcal{F}}_2$ . Let  $\{\xi_B^+, \xi_B^-\}$  the endpoints of the bigon, where as above,  $\xi_B^+$  denotes the nonseparated point of the curves  $c_1, c_2$ . Then, we have that all non-Hausdorff bigons in  $E$  (note that there is at least one since there is no transversal in  $E$  from  $c_1$  to  $c_2$ ) that are contained between  $c_1$  and  $c_2$  have its nonseparated point in  $\xi_B^+$ .

In contrast with Remark 9.9 we notice the following:

**Remark 9.12.** If  $B$  is a  $\gamma$ -invariant bigon in a leaf  $L \in \tilde{\mathcal{F}}_1$  and  $E \in \tilde{\mathcal{F}}_2$  is the leaf which intersects  $B$  in the boundary  $c_1, c_2 \in \mathcal{G}_L$  is such that the nonseparated rays of  $c_1, c_2$  in  $L$  land in a point *different from*  $\alpha(L)$ , then, for every leaf  $L' \in \tilde{\mathcal{F}}_1$  close to  $L$  intersecting  $E$  *inside* the region  $B_E$  bounded by  $c_1 \cup c_2$  contains a bigon  $B'$  whose endpoints coincide with those of  $B$  and its boundaries correspond to the intersection with  $E$ .

**9.4. Putting all stability together.** Let  $B$  be a non-Hausdorff bigon in  $L \in \tilde{\mathcal{F}}_1$  with boundaries  $c_1, c_2 \in \mathcal{G}_L$  and endpoints  $\xi_B^+$  and  $\xi_B^-$  (recall that  $\xi_B^+$  denotes the endpoint which is the limit of the nonseparated rays of  $c_1$  and  $c_2$ ). Consider a transversal  $\tau : [0, \delta) \rightarrow L$  to  $\mathcal{G}_L$  such that  $\tau(0) \in c_1$ , so that  $\tau((0, \delta))$  is contained in  $B$ . Let then  $E_t \in \tilde{\mathcal{F}}_2$  be the leaf through the point  $\tau(t)$ . We will denote by  $\mathcal{J}_B$  the closed interval in  $\partial\mathbb{H}^2$  which is the closure of the connected component of  $\partial\mathbb{H}^2 \setminus \{\xi_B^+, \xi_B^-\}$  containing  $\alpha(E_t)$ ,  $t > 0$ ; that is,

$$(9-1) \quad \mathcal{J}_B = \overline{\text{cc}_{\alpha(E_t)}(\partial\mathbb{H}^2 \setminus \{\xi_B^+, \xi_B^-\})}.$$

We note that the definition of  $\mathcal{J}_B$  is independent on the choice of  $\tau$  and  $t > 0$  (and also works if  $\tau(0) \in c_2$  instead of  $c_1$ ).

**Lemma 9.13.** *Let  $B$  be a  $\gamma$ -invariant non-Hausdorff bigon in a leaf  $L \in \tilde{\mathcal{F}}_1$  with boundaries  $c_1, c_2 \in \mathcal{G}_L$  such that  $\alpha(L) = \xi_B^-$  and  $\gamma \in \pi_1(M) \setminus \{\text{id}\}$ . Then, for every  $L' \in \tilde{\mathcal{F}}_1$  such that  $\alpha(L') \in \mathcal{J}_B$  (see (9-1)) we have that  $L'$  contains a non-Hausdorff bigon  $B'$  with endpoints  $\xi_B^-, \xi_B^+$ . In the same way if  $\alpha(L) = \xi_B^+$  we also get that for every leaf  $L' \in \tilde{\mathcal{F}}_1$  such that  $\alpha(L') \in \mathcal{J}_B$  we have that  $L'$  contains a non-Hausdorff bigon  $B'$  with endpoints  $\xi_B^-, \xi_B^+$ .*

*Proof.* Both  $\mathcal{F}_1$  and  $\mathcal{F}_2$  are transversally orientable. We choose an orientation such that if  $\mu : (-\varepsilon, \varepsilon) \rightarrow \tilde{M}$  is a positively oriented path transverse to *both* foliations  $\tilde{\mathcal{F}}_1, \tilde{\mathcal{F}}_2$ . Then, if  $L_t \in \tilde{\mathcal{F}}_1$  and  $E_t \in \tilde{\mathcal{F}}_2$  are the leaves through  $\mu(t)$  then  $\alpha(L_t)$  and  $\alpha(E_t)$  move both clockwise. To see that this is possible, recall that when lifted to  $\tilde{M}$  it follows that the transverse orientation induce a direction on which the point  $\alpha(L_t)$  varies as  $L_t$  varies in the leaf space of  $\tilde{\mathcal{F}}_1$ . Note that since both foliations inherit the orientation of the base (because they are horizontal) the transverse orientations

match (recall that from Theorem 2.1 we know that there are homeomorphisms inducing the identity in the base which maps the foliations  $\mathcal{F}_i$  to  $\mathcal{F}_{ws}$ ), and so we can speak of moving in the clockwise or counterclockwise direction in  $\partial\mathbb{H}^2$ .

Let  $E$  be the  $\tilde{\mathcal{F}}_2$  leaf containing  $c_1 \cup c_2$ , and let  $B_E$  be the region in  $E$  bounded by  $c_1 \cup c_2$ . Let  $\tau$  be a transversal to  $\mathcal{G}_L$  starting at  $c_1$  and entering  $B$ .

The following happens: moving  $E_t$  (the leaf through  $\tau(t)$ , so  $E_0 = E$ ) moves  $\alpha(E_t)$  in one direction of  $\partial\mathbb{H}^2$ , starting at  $\xi_B^+$ , by definition  $\alpha(E_t)$  moves into  $\mathcal{J}_B$ . Without loss of generality, we can assume that  $\alpha(E_t)$  moves clockwise, equivalently the leaves of  $\tilde{\mathcal{F}}_2$  intersecting  $B$  are *above*  $E_0$ . In other words  $B$  is above  $E_0$ . Next consider a curve  $\eta : (-\delta, \delta) \rightarrow E_0$  transverse to  $\mathcal{G}_{E_0}$ , such that  $\eta(0) \in c_1$ , and parametrized clockwise; i.e., if  $L_s$  is the leaf of  $\tilde{\mathcal{F}}_1$  through the point  $\eta(s)$  then  $\alpha(L_s)$  moves clockwise as  $s$  increases.

Now if  $\alpha(L_0) = \xi_B^-$ , the previous lemma tells us that the leaves  $L_s$  which have bigons bounded by the curves in  $E \cap L_s$  are the leaves intersecting the region  $B_E$  (see Remark 9.12). We saw above that the region  $B$  is above  $E_0$  and therefore  $B_E$  is *below*  $L_0$ .

Therefore the leaves  $L_t$  which have these bigons, satisfy that the point  $\alpha(L_t)$  is locally counterclockwise to  $\xi_B^-$  and thus in the same connected component as  $\alpha(E_s)$  (for small  $s$ ) in  $\partial\mathbb{H}^2 \setminus \{\xi_B^+, \xi_B^-\}$ .

Applying Lemma 9.3 we get that the full closed interval between  $\xi_B^+$  and  $\xi_B^-$ , has a bigon with ideal points  $\xi_B^-, \xi_B^+$ . If on the other hand we assume that  $\alpha(E_t)$  moves counterclockwise when  $t$  increases, we get that  $B$  is *below*  $E_0$ , and as above it will follow that  $B_E$  is above  $L_0$ . One obtains the same result as above.

Finally we consider the case that  $\alpha(L) = \xi_B^+$ . In this case the important fact to note is that in the proof of Proposition 9.4 we obtain the half neighborhood by moving in the opposite direction (see Remark 9.9), where we explain that the  $L_s$  near  $L_0$  which intersect  $E_0$  forming a bigon in  $L_s$  intersect  $E_0$  *outside*  $B_E$  (as opposed to inside  $B_E$  in the previous case). Therefore with the conventions as in the previous case we have the following: if  $\alpha(E_t) > \alpha(E_0)$  (for  $t > 0$ ), we produce bigons in leaves  $L_s$  above  $L_0$ , so  $\alpha(L_s) > \alpha(L_0)$ . Therefore  $\alpha(E_t)$  moves clockwise and  $\alpha(L_t)$  moves clockwise. But  $\alpha(E_t)$  moving clockwise when  $t$  increases means that  $\alpha(E_t)$  moves inside  $\mathcal{J}_B$  for  $t > 0$ , and henceforth  $\alpha(L_s)$  moves inside  $\mathcal{J}_B$  for  $t > 0$ . Thus we obtain the second statement of the lemma.  $\square$

Putting together what we have shown, we can deduce:

**Proposition 9.14.** *Let  $\mathcal{F}_1, \mathcal{F}_2$  be two transverse minimal foliations of  $M = T^1S$  and let  $\mathcal{G}$  be their intersection. Then if  $\mathcal{G}$  is not homeomorphic to the foliation given by the geodesic flow of a hyperbolic metric, it follows that there are finitely many disjoint simple closed curves  $s_1, \dots, s_k$  in  $S$  such that for every periodic non-Hausdorff bigon  $B$  in a leaf  $L \in \tilde{\mathcal{F}}_i$  we have that  $\partial\mathcal{J}_B$  (equation (9-1)) corresponds*

to the endpoints of a lift of one of the curves  $s_j$  to  $\mathbb{H}^2$ . In particular:

- If two periodic non-Hausdorff bigons share one endpoint, then they must share both endpoints.
- The endpoints of two different periodic non-Hausdorff bigons cannot be linked.
- Up to deck transformations, there are finitely many periodic non-Hausdorff bigons. Equivalently there are finitely many Reeb surfaces of  $\mathcal{G}$  in  $\mathcal{F}_i$ .

*Proof.* Assume that a leaf  $L \in \tilde{\mathcal{F}}_1$  has a  $\gamma$ -periodic non-Hausdorff bigon  $B$ . Due to Propositions 9.4 or 9.10 we know that we can push such a bigon to nearby leaves. Then, thanks to Lemma 9.3 we know that every leaf  $L' \in \tilde{\mathcal{F}}_1$  with  $\alpha(L') \in \mathcal{J}_B$  has a bigon joining the endpoints  $\xi_B^+$  and  $\xi_B^-$  of  $B$ .

First, assume that  $\mathcal{J}_B$  and  $\mathcal{J}_{B'}$  share an endpoint, then, since  $\gamma$  and  $\gamma'$  must fix those endpoints, we deduce that  $\gamma'$  and  $\gamma$  belong to the same cyclic group of  $\pi_1(M)$ . Thus, we deduce that both endpoints must coincide.


Now assume that there are two distinct non-Hausdorff bigons  $B, B'$  such that  $B$  is  $\gamma$ -periodic and  $B'$  is  $\gamma'$ -periodic and that the ideal points of  $B, B'$  are linked.

Again using Propositions 9.4 and 9.10, we can find bigons with same ideal points as  $B$  in all leaves  $L''$  in  $\tilde{\mathcal{F}}_1$  in the interval defined by  $\alpha(L'')$  in the closure of one complementary component of  $\xi_B^+, \xi_B^-$ . The set of such  $\alpha(L'')$  produces an interval  $I_B$  of  $\partial\mathbb{H}$ . The same holds for  $B'$ , with corresponding interval  $I_{B'}$ . So if the ideal points link, it follows that the interiors of  $I_B, I_{B'}$  intersect, and we can find a leaf  $L'$  which has both a bigon with same endpoints as  $B$  and one which has both endpoints as  $B'$ . This is a contradiction since bigons cannot cross (since they are bounded by leaves of  $\mathcal{G}_{L'}$  which is a foliation).

Finally, note that since a  $\gamma$ -invariant non-Hausdorff bigon cannot cross with its translates by other deck transformations, each such non-Hausdorff bigon corresponds to a simple closed curve in  $S$ . Similarly, distinct periodic non-Hausdorff bigons correspond to disjoint curves, and at most finitely many such disjoint curves can exist in  $S$ .

To prove the final property: we may assume periodic bigons  $B_1, B_2$  are associated with same simple closed curve  $s_j$  of  $S$ , and are both bigons in say  $\tilde{\mathcal{F}}_1$ . There is a unique element  $\gamma$  of  $\pi_1(M)$  which projects to  $s_j$  in  $S$  and acts with fixed points in  $\tilde{\mathcal{F}}_i$ . This  $\gamma$  has a discrete set of fixed points in the leaf space of  $\tilde{\mathcal{F}}_1$ . So we may assume up to fiber translates, that  $B_1, B_2$  are in the same leaf of  $\tilde{\mathcal{F}}_1$ . But then there are finitely many such. This proves finiteness of Reeb surfaces in  $M$ .  $\square$

**9.5. Creating new bigons.** As in the examples of [35], it is possible that some bigons do not have continuations beyond one half-interval of the leaf space. Thus, to show that there are bigons in every leaf, we need to construct new bigons (i.e., which do not come from varying continuously from the original one) in order to



**Figure 20.** How the nonmarker points move on the other foliation as one considers the leaf associated to a transversal of  $\mathcal{G}_L$ .

produce bigons in the other half interval. This requires a careful analysis of the orientation of the foliations.

We first show that the existence of some periodic bigon forces landing points of rays to be rather restricted. We state this in somewhat more generality.

**Lemma 9.15.** *Assume that there is a leaf  $L \in \tilde{\mathcal{F}}_i$  that contains a leaf  $e \in \mathcal{G}_L$  such that  $\alpha(L) \notin \{\partial^+ \Phi(e) \cup \partial^- \Phi(e)\}$ . Assume also that  $\partial^+ \Phi(e) \neq \partial^- \Phi(e)$ . Let  $J$  be a nontrivial interval in the leaf space of  $\tilde{\mathcal{F}}_i$ . Then there is a leaf  $L'$  in  $J$  fixed by a nontrivial deck transformation  $\beta$  such that no leaf of  $\mathcal{G}_{L'}$  has one ideal point fixed by  $\beta$ .*

*Proof.* Decrease  $J$  if necessary so that  $\alpha(L')$  never attains either of  $\partial^+ \Phi(e)$  or  $\partial^- \Phi(e)$  for  $L' \in J$ . In particular,  $\alpha$  is injective in  $J$ . Let  $a_1 = \partial^+ \Phi(e)$ ,  $a_2 = \partial^- \Phi(e)$ .

Let  $I$  be an open interval in  $\partial \mathbb{H}^2$  contained in the connected component of  $\partial \mathbb{H}^2 \setminus \{a_1, a_2\}$  which does not contain  $\alpha(L)$ .

Choose a nontrivial deck transformation  $\beta$  with one fixed point in  $I$  and one in  $\alpha(J)$ . Let  $L'$  be a leaf of  $\tilde{\mathcal{F}}_i$  fixed by  $\beta$ , and we can assume that  $L'$  is in  $J$ . This leaf satisfies the conclusion. Suppose that  $\mathcal{G}_{L'}$  has a leaf  $c$  which is not a bubble leaf and has one ideal point fixed by  $\beta$ . Without loss of generality assume that  $c$  has an ideal point in  $I$  which is fixed by  $\beta$ . Then iterating by  $\beta$  or  $\beta^{-1}$  we eventually obtain a leaf  $c'$  of  $\mathcal{G}_{L'}$  with one ideal point in  $I$  and one ideal point in  $\alpha(J)$ . But  $a_1, a_2$  link with  $I, \alpha(J)$ . This contradicts that  $c'$  is a leaf of  $\mathcal{G}_{L'}$  with ideal points  $a_1, a_2$ , which would cause crossing of different leaves of  $\mathcal{G}_{L'}$ .

If  $c$  is a bubble leaf landing in a point fixed by  $\beta$ , one can iterate and produce a leaf which is not a bubble leaf and has the same characteristics.  $\square$

**Remark 9.16.** If the leaf space of  $\tilde{\mathcal{G}}$  is not Hausdorff, it follows from Theorem 8.1 that there are some non-Hausdorff bigons in some leaf, and by Proposition 9.14 that an open set of leaves contains non-Hausdorff bigons. Using minimality, we deduce that every leaf  $L \in \tilde{\mathcal{F}}_i$  has infinitely many non-Hausdorff bigons with distinct endpoints in  $\partial\mathbb{H}^2$  and therefore we are in the hypothesis of the previous lemma for every  $L \in \tilde{\mathcal{F}}_i$ .

To produce new non-Hausdorff bigons we need to push to the other side of the bigons which is more delicate. We first show the following:

**Lemma 9.17.** *Let  $L \in \tilde{\mathcal{F}}_1$  and let  $E \in \tilde{\mathcal{F}}_2$  with  $\alpha(L) \neq \alpha(E)$  be such that there is  $\gamma \in \pi_1(M) \setminus \{\text{id}\}$  fixing  $L$  and  $E$ . Let  $\ell_0 \in E \cap L$  be a connected component of the intersection such that  $\{\partial^+\Phi(\ell_0), \partial^-\Phi(\ell_0)\} = \{\alpha(L), \xi_0\}$  for some  $\xi_0 \neq \alpha(L)$ . Let  $\tau : [0, \delta) \rightarrow L$  be a transversal to  $\mathcal{G}_L$ , with  $\tau(0) \in \ell_0$ , and such that if  $E_t$  denotes the leaf through  $\tau(t)$  for  $t > 0$  we have that  $\alpha(E_t)$  and  $\tau(t)$  are in different connected components of  $L \setminus \ell_0$ . Then, for small  $t > 0$ , if  $\ell_t$  denotes the curve of  $\mathcal{G}_L$  through  $\tau(t)$  it follows that  $\{\partial^+\Phi(\ell_t), \partial^-\Phi(\ell_t)\} \subset \{\alpha(L), \xi_0\}$  and at least one of the points is  $\alpha(L)$ .*

*Proof.* Consider  $I_t$  the interval of the leaf space of  $\tilde{\mathcal{F}}_2$  made of the leaves  $E_s$  with  $s \in (0, t)$  and fix a small  $\varepsilon$  such that Proposition 3.4 applies for  $\tilde{\mathcal{F}}_2$ . For  $t$  small we identify  $I_t$  with  $\bigcup\{\alpha(E_s), s \in (0, t)\}$ . Consider then the set  $\widehat{D}_\varepsilon(E, I_t)$  for this foliation. It follows (see Proposition 2.7) that the set  $\widehat{D}_t$  which is the projection of  $\widehat{D}_\varepsilon(E, I_t)$  to  $L$  contains a complementary component of a neighborhood of uniform size around the geodesic joining the points  $\alpha(E) = \alpha(E_0)$  and  $\alpha(E_t)$ . This complementary component has  $\alpha(L)$  in its closure.

In particular, one can choose  $t$  so small that  $\widehat{D}_t$  is disjoint from the image of  $\tau$ . In particular, one gets that one of the rays of  $\ell_t$  must converge to  $\alpha(L)$ . We now want to understand the other ray. If  $\alpha(E) \neq \xi_0$  then, the same argument shows that the other ray of  $\ell_t$  converges to  $\xi_0$ , so we will assume in what follows that  $\alpha(E) = \xi_0$ .

We assume by contradiction that the other ray of  $\ell_t$  starting at  $\tau(t)$ , that we call  $r_t$ , lands in some point  $\xi \notin \{\alpha(L), \alpha(E)\}$ . It cannot land in any point of the interior of  $I_t$  since it would need to intersect  $\ell_0$ .

We can thus apply Proposition 3.4 to the leaves  $E_s$  for  $s \in (t, 0)$  to obtain a family of rays  $r_s$  in  $\widehat{L}_t$  that start at  $\tau(s)$  and always land in  $\xi$ . To see this, note that while  $r_s$  intersects  $\widehat{D}_t$  then the ray varies continuously with  $s$ , but if stops intersecting  $\widehat{D}_t$  it could in principle split into more than one ray. However, due to Proposition 3.9 the landing should occur in  $\alpha(E_s)$  which is impossible since it would force the ray to intersect  $\ell_0$ . Finally, note that when  $s \rightarrow 0$  there could be splitting, but in this

case we obtain that there is a curve in  $E_0 \cap L$  (note that  $E_0 = E$ ) which goes from  $\alpha(E)$  to  $\xi$ .

In conclusion, we have shown that if the result does not hold, then  $E \cap L$  must have a curve  $\ell'$  from one of the fixed points of  $\gamma$  to  $\xi$ . Since  $E$  and  $L$  are  $\gamma$ -invariant we can iterate this intersected curve  $\ell'$  to obtain a sequence of distinct intersections of  $E \cap L$  landing in the same point (and since they do not admit a common transversal, they must have some distance in between). This contradicts the small visual measure property and completes the proof of the lemma.  $\square$

There is a similar phenomenon when the splitting goes in the opposite direction:

**Lemma 9.18.** *Assume that  $\tilde{\mathcal{G}}$  is not Hausdorff.<sup>12</sup> Let  $L \in \tilde{\mathcal{F}}_1$  and  $E \in \tilde{\mathcal{F}}_2$  be such that  $\alpha(E) = \alpha(L)$  so that there is  $\gamma \in \pi_1(M) \setminus \{\text{id}\}$  fixing both  $L$  and  $E$ . Let  $\ell_0 \in E \cap L$  be a connected component of the intersection such that  $\{\partial^+ \Phi(\ell_0), \partial^- \Phi(\ell_0)\} = \{\alpha(L), \xi_0\}$  for some  $\xi_0 \neq \alpha(E) = \alpha(L)$ . Let  $\tau : [0, \delta) \rightarrow L$  be a transversal to  $\mathcal{G}_L$ , with  $\tau(0) \in \ell_0$ . Then, for small  $t > 0$ , if  $\ell_t$  denotes the leaf of  $\mathcal{G}_L$  through  $\tau(t)$  it follows that  $\{\partial^+ \Phi(\ell_0), \partial^- \Phi(\ell_0)\} \subset \{\alpha(L), \xi_0\}$  and at least one of the points is  $\xi_0$ .*

*Proof.* Since  $E$  and  $L$  are  $\gamma$ -invariant then, one must have that  $\ell_0$  also is. Otherwise  $\ell_0$  is not  $\gamma$  periodic, so we would get infinitely many distinct connected components of  $E \cap L$ , all of them sharing the endpoint  $\alpha(L) = \alpha(E)$ . But this contradicts small visual measure since no two of such components can intersect a common transversal. It follows that  $\xi_0, \alpha(L)$  are the two fixed points of  $\gamma$ .

Consider  $I_\delta$  the interval of the leaf space of  $\tilde{\mathcal{F}}_2$  which consists of the leaves  $E_t$  through  $\tau(t)$  with  $t \in (0, \delta)$ . The curve  $\ell_t \subset E_t \cap L$  is the curve of  $\tilde{\mathcal{G}}$  passing through  $\tau(t)$ . Since  $\xi_0 \neq \alpha(E)$  we can apply Proposition 3.4 to deduce that one of the rays of  $\ell_t$  converges to  $\xi_0$ . We now want to understand the other ray.

Consider for  $t \in [0, \delta)$  the other ray of  $\ell_t$  starting at  $\tau(t)$ , which we call  $r_t$ , and let  $\xi_t \in \partial\mathbb{H}^2$  be the landing point of  $\xi_t$ . As in the proof of Proposition 9.4 we claim that the point  $\xi_t$  is weakly monotonic as we move  $t$ , that it is continuous at  $t = 0$  and that either it is  $\xi_t = \alpha(E_t)$  or  $\xi_t$  is locally constant. Weak monotonicity is simpler in this case, since if  $t' > t$  we have that  $\ell_t$  separates  $\ell_{t'}$  from  $\ell_0$ . Using the pushing argument of Proposition 3.4 we get that if  $\xi_t$  is not  $\alpha(E_t)$  then  $\xi_t$  must be locally constant. Finally for continuity at 0: we assume that  $\xi_t \neq \xi_0$  for some small  $t$  (else we get the result if  $\xi_t = \xi_0$  for all  $t$  sufficiently small). Assume that  $\alpha(L)$  is the attracting fixed point of  $\gamma$ . Then  $\gamma^n(\ell_t)$  converges to  $\alpha(L)$  when  $n \rightarrow \infty$ . Since  $\gamma^n(\ell_t) = \ell_{t_n}$  with  $t_n \rightarrow 0$ , continuity of  $\xi_t$  at  $t = 0$  follows.

Now suppose that  $\xi_t = \alpha(E_t)$  for a nontrivial interval  $J$  in  $(0, t)$ . We identify  $J$  with an interval in the leaf space of  $\tilde{\mathcal{F}}_2$ , which we can identify with an interval in  $\partial\mathbb{H}^2$  as well. As noted in Remark 9.16 we satisfy the hypothesis of Lemma 9.15. Hence

<sup>12</sup>This is a standing assumption in this section, but we emphasize it here because the proof makes crucial use of Lemma 9.15.

there is a leaf  $E_t$  in  $J$  such that  $E_t$  is fixed by a nontrivial deck transformation  $\beta$  and no leaf of  $\mathcal{G}_{E_t}$  has one ideal point fixed by  $\beta$ . However we know that  $\xi_t = \alpha(E_t)$  and  $\alpha(E_t)$  is fixed by  $\beta$ . This is a contradiction since  $\ell_t$  has one ideal point  $\xi_t$ .

Exactly as in Proposition 9.4, we deduce that if  $\xi_s$  is not constant and equal to  $\alpha(E) = \alpha(L)$  then either it is constant and equal to  $\xi_0$  or the function  $\xi_s$  is locally constant with jumps in a discrete set of  $(0, \delta)$  in points  $a_n \rightarrow 0$  such that for each  $n$  we have  $\ell_{a_n}$  lands in  $\xi_n := \alpha(E_{a_n})$ . This implies that the intersection of  $E_{a_n} \cap L$  contains, besides  $\ell_{a_n}$  a curve joining  $\xi_{n-1}$  and  $\xi_n$ . These curves are accumulated by the curves  $\ell_t$  with  $t \in (a_{n-1}, a_n)$  and therefore, due to Proposition 8.3 we know that there is some  $\gamma_n \in \pi_1(M) \setminus \{\text{id}\}$  fixing  $E_{a_n}$ . This implies that  $E_{a_n}$  has a non-Hausdorff bigon joining  $\xi_n$  with some point in the interval from  $\xi_0$  to  $\xi_{n-1}$  not containing  $\xi_n$ . This contradicts Proposition 9.14 because it produces bigons whose endpoints are linked or have a unique common endpoint. This completes the proof.  $\square$

We can now prove the main result of this section:

**Proposition 9.19.** *Let  $B$  be a  $\gamma$ -periodic non-Hausdorff bigon in a leaf  $L \in \tilde{\mathcal{F}}_i$  for some  $\gamma \in \pi_1(M) \setminus \{\text{id}\}$ . Then, one of the following options hold:*

- (1) *there are  $\gamma$ -invariant bigons  $B_1$  and  $B_2$  in  $L$  such that  $\xi_{B_1}^+ = \xi_{B_2}^+$  and such that  $\mathcal{J}_{B_1} \cup \mathcal{J}_{B_2} = \partial \mathbb{H}^2$ , or*
- (2) *there is an even number of ordered<sup>13</sup>  $\gamma$ -invariant bigons  $B_1, B_2, \dots, B_{2k}$  in  $L$  such that consecutive ones have different nonseparated points. Moreover, if the order is chosen so that  $B_1$  is the bigon such that  $\xi_{B_1}^- = \alpha(L)$  then one has that  $\mathcal{J}_{B_1}$  is the interval which is separated from  $B_1$  by  $B_2$ .*

*Proof.* Due to small visual measure, there are finitely many  $\gamma$ -invariant non-Hausdorff bigons in  $L$ , which we denote by  $B_1, \dots, B_k$  in order (i.e.,  $B_j$  separates  $B_{j-1}$  from  $B_{j+1}$  in  $L$ ). Each such  $\gamma$ -periodic non-Hausdorff bigon  $B_j \in L$  corresponds to the intersection of  $L$  with some  $E_j \in \tilde{\mathcal{F}}_2$  such that  $\alpha(E_j) \in \{\xi, \eta\}$ , which are the fixed points of  $\gamma$  (and also the separated and nonseparated points of  $B$ ). Since  $L$  is  $\gamma$ -invariant we know that  $\alpha(L) \in \{\xi, \eta\}$ . Moreover, if  $\alpha(E_j) = \alpha(L)$  then it follows that  $\xi_{B_j}^+ = \alpha(L)$  while if  $\alpha(E_j) \neq \alpha(L)$  then  $\xi_{B_j}^+ \neq \alpha(L)$ .

There are also a finite number of  $\gamma$ -invariant leaves of  $\mathcal{G}_L$ , say  $\ell_0, \dots, \ell_m$ , corresponding to intersections with leaves  $F_i \in \tilde{\mathcal{F}}_2$  such that  $\alpha(F_i) \in \{\xi, \eta\}$ . (Indeed, these leaves are at some minimal distance away from each other and each one must be uniformly close to the geodesic joining  $\xi$  and  $\eta$ ). We also consider these leaves ordered in the same direction. Some of them are contained in the boundaries of the bigons  $B_j$ . We know that  $m \geq 1$  since there is at least one  $\gamma$ -periodic non-Hausdorff bigon.

<sup>13</sup>This means that in  $L$  the bigon  $B_j$  separates  $B_{j-1}$  from  $B_{j+1}$  for all  $j = 2 \dots k - 1$ .

Let  $D$  be the closure of the region in  $L$  bounded by  $\ell_0, g_m$ . Since  $m \geq 1$  then  $D$  is nonempty, with nonempty interior. Fix a transversal orientation to  $\mathcal{F}_2$ .

Consider, for every  $i \in \{0, \dots, m - 1\}$ , a transversal  $\tau_i : [0, \delta) \rightarrow L$  such that  $\tau_i(0) \in \ell_i$  and that  $\tau_i(t)$  is between  $\ell_i$  and  $\ell_{i+1}$ . Denote by  $\mathcal{C}_i$  the (closed) interval from  $\xi$  to  $\eta$  and containing  $\alpha(H_t)$  where  $H_t \in \tilde{\mathcal{F}}_2$  is the leaf such that  $\tau_i(t) \in H_t$ . This interval is well defined and is independent of the choice of  $\tau_i$  and  $t$ .

**Claim 1.**  $\mathcal{C}_i \cup \mathcal{C}_{i+1} = \partial\mathbb{H}^2$ . *The region between  $\ell_i$  and  $\ell_{i+1}$  is a bigon if and only if  $\alpha(F_i) = \alpha(F_{i+1})$ .*

*Proof.* The interior of  $\mathcal{C}_i$  is exactly the set of points  $v$  in  $\partial\mathbb{H}^2$  such that there is a leaf  $E \in \tilde{\mathcal{F}}_2$  that intersects the region between  $\ell_i$  and  $\ell_{i+1}$  in  $L$  and such that  $\alpha(E) = v$ . This is because both sets are  $\gamma$  invariant and contain small neighborhoods of the endpoints. Hence if one considers a transversal to  $\mathcal{G}_L$  intersecting in the interior the leaf  $\ell_{i+1}$  (which is a connected component of  $L \cap F_{i+1}$ ), one sees that  $\mathcal{C}_i$  and  $\mathcal{C}_{i+1}$  correspond to distinct intervals and thus  $\mathcal{C}_i \cup \mathcal{C}_{i+1} = \partial\mathbb{H}^2$  as desired.  $\square$

To see the last property, notice that if the region is a bigon then  $F_i = F_{i+1}$  so the  $\alpha$ 's coincide. Otherwise  $F_i, F_{i+1}$  are distinct, fixed by  $\gamma$ , and there is no other  $\gamma$  invariant leaf between  $F_i$  and  $F_{i+1}$ . Hence when acting on the leaf space of  $\tilde{\mathcal{F}}_2$ , it follows that up to inverse,  $\gamma$  is attracting in  $F_i$  and repelling in  $F_{i+1}$ . This implies that  $\alpha(F_i) \neq \alpha(F_{i+1})$ .

**Claim 2.** *Consider  $B_j, B_{j+1}$  consecutive non-Hausdorff bigons in  $L$ . Then if  $\xi_{B_j}^+ = \xi_{B_{j+1}}^+$  we have  $\mathcal{J}_{B_j} \cup \mathcal{J}_{B_{j+1}} = \partial\mathbb{H}^2$ , and if  $\xi_{B_j}^+ = \xi_{B_{j+1}}^-$  then  $\mathcal{J}_{B_j} = \mathcal{J}_{B_{j+1}}$ .*

*Proof.* In the first case one has between the boundaries of  $B_j, B_{j+1}$  an even (possibly zero, if  $B_j \cap B_{j+1}$  intersect in some leaf  $\ell_j$ ) number of regions between consecutive curves  $\ell_i$ . Since each intersection makes a half turn in  $\partial\mathbb{H}^2$  without changing orientation (because this only happens when one crosses a non-Hausdorff bigon) we obtain that the leaves of  $\tilde{\mathcal{F}}_2$  intersecting the interior of  $B_j$  and  $B_{j+1}$  correspond to different intervals in  $\partial\mathbb{H}^2$  whose boundaries are  $\{\xi, \eta\}$  thus  $\mathcal{J}_{B_j} \cup \mathcal{J}_{B_{j+1}} = \partial\mathbb{H}^2$ . The other case is similar, since one has an odd number of such regions in between.  $\square$

**Conclusion.** Claim 2 implies that the only way that all non-Hausdorff bigons with the same nonseparated point have the same associated interval is that adjacent non-Hausdorff bigons must have distinct nonseparated points. Note that we still could have that there is a unique such non-Hausdorff bigon.

To obtain that there are an even number of such bigons, we will use Lemmas 9.17 and 9.18. Using Lemma 9.18 we will deduce that  $\ell_0$  and  $\ell_m$  satisfy  $\alpha(F_0) = \alpha(F_m) \neq \alpha(L)$ . Indeed, suppose to the contrary that  $\alpha(L) = \alpha(F_0)$ . Then we can apply Lemma 9.18 to the curve  $\ell_0$  contained in  $L \cap F_0$ . It has ideal points  $\xi, \eta$ , which are distinct; one of them is  $\alpha(L) = \alpha(F_0)$ , the other we call  $\xi_0$ . Now consider a transversal  $\tau$  starting in  $\ell_0$  and exiting  $D$  and  $\ell'$  a curve of  $\mathcal{G}_L$  intersecting  $\tau$

outside  $D$  and near  $\tau(0)$ . Apply Lemma 9.18 to  $\ell_0$ : it implies that the ideal points of  $\ell'$  are contained in  $\alpha(L)$ ,  $\xi_0$ . But that is impossible given the choice of  $\ell_0$ : no curve outside  $D$  has ideal points contained in  $\{\xi, \eta\}$ . This shows that  $\alpha(F_0) \neq \alpha(L)$ . This also applies to  $\ell_m$ , showing that  $\alpha(L) \neq \alpha(F_m)$ . Hence  $\alpha(F_m) = \alpha(F_0) \neq \alpha(L)$ .

Moreover Lemma 9.17 implies that  $\mathcal{C}_0$  is the interval of  $\partial\mathbb{H}^2 \setminus \{\xi, \eta\}$  which is in the opposite connected component of  $\partial\mathbb{H}^2 \setminus \{\eta, \xi\}$  of  $\ell_m$  in  $L \setminus \ell_0$ . Similarly,  $\mathcal{C}_{m-1}$  is the opposite interval, that is, the interval of  $\partial\mathbb{H}^2 \setminus \{\xi, \eta\}$  which is in the opposite connected component to  $\ell_0$  in  $L \setminus \ell_m$ . In particular, the two intervals  $\mathcal{C}_0, \mathcal{C}_{m-1}$  are different.

We will now check the progress along leaves  $\tilde{\mathcal{F}}_2$  when we cross the region  $D$  starting from  $\ell_0$  all the way through  $\ell_m$ . Notice that crossing each periodic non-Hausdorff bigon changes the orientation in which one makes progress in the leaves of  $\tilde{\mathcal{F}}_2$ . Lemma 9.17 implies that in both  $\ell_0$  and  $\ell_m$  when one crosses into  $D$ , then the  $\alpha$ 's of the  $\tilde{\mathcal{F}}_2$  leaves move in the opposite direction (e.g., when crossing  $\ell_0$  it is in the complementary component of  $\ell_0$  which does not contain  $D$ ). In particular, crossing through  $\ell_0$  into  $D$  and through  $\ell_m$  outside  $D$ , one needs to cross  $\tilde{\mathcal{F}}_2$  leaves with the same orientation. Thus there must be an even number of non-Hausdorff bigons in between.

Finally, if we choose the order so that  $B_1$  is the one such that  $\xi_{B_1}^- = \alpha(L)$  we have that one needs to intersect an even number of curves  $\ell_i$  after  $\ell_0$  to get to  $B_1$  (possibly  $\ell_0$  is the boundary of  $B_1$ , in which case we do not intersect any). We get that the boundary of  $B_1$  is of the form  $\{\ell_{2i} \cup \ell_{2i+1}\}$  for some  $i \geq 0$ . This implies that  $\mathcal{C}_{2i} = \mathcal{C}_0$ , because if  $G$  is the leaf of  $\tilde{\mathcal{F}}_2$  containing  $\partial B_1$ , then  $\alpha(G) = \xi_{B_1}^+$ . This completes the proof.  $\square$

## 9.6. Proof of Theorem 9.1 and applications.

*Proof of Theorem 9.1.* By assumption we know that there is at least one leaf  $L$  with a non-Hausdorff bigon  $B$  and some  $\gamma \in \pi_1(M) \setminus \{\text{id}\}$  for which we have  $\gamma B = B$  and if we denote the endpoints as  $\{\xi, \eta\} \subset \partial\mathbb{H}^2$  these are also  $\gamma$ -invariant. Without loss of generality we assume that  $L \in \tilde{\mathcal{F}}_1$ . In particular,  $\alpha(L) \in \{\xi, \eta\} = \{\xi_B^+, \xi_B^-\}$ . Let  $E \in \tilde{\mathcal{F}}_2$  be such that the boundary of  $B$  is contained in  $E \cap L$ .

Denote by  $\hat{L}$  a leaf in  $\tilde{\mathcal{F}}_1$  such that  $\alpha(\hat{L}) \in \{\xi, \eta\}$  and that  $\alpha(\hat{L}) \neq \alpha(L)$ . The leaf  $\hat{L}$  must also be  $\gamma$ -invariant. It follows from applying either Proposition 9.4 or Proposition 9.10 that  $\hat{L}$  contains at least one non-Hausdorff bigon which up to changing  $\hat{L}$  by a deck transformation associated to the fiber we can assume is bounded by leaves of  $E \cap \hat{L}$ .

We now apply Proposition 9.19. If the first condition of the proposition happens for either  $L$  or  $\hat{L}$ , we get that there are  $\gamma$ -periodic non-Hausdorff bigons  $B_i, B_j$  in  $L$  (or  $\hat{L}$ ) with the same nonseparated point and such that  $\mathcal{J}_{B_i} \cup \mathcal{J}_{B_j} = \partial\mathbb{H}^2$ . Applying

Lemma 9.13 we deduce that for every  $L' \in \tilde{\mathcal{F}}_1$  there must be a non-Hausdorff bigon joining  $\xi$  and  $\eta$  in  $L'$ . This completes this case.

Thus, we can assume that both for  $L$  and  $\hat{L}$  we have that the second option of Proposition 9.19 holds. We need to show:

**Claim 3.** *If we consider  $B_1, \dots, B_{2k}$  the  $\gamma$ -invariant non-Hausdorff bigons in  $L$  with a chosen order and use the same order in  $\hat{L}$  and get  $\gamma$ -invariant non-Hausdorff bigons  $\hat{B}_1, \dots, \hat{B}_{2m}$  we have that  $k = m$  and that the nonseparated point of  $B_1$  and  $\hat{B}_1$  coincide.*

*Proof.* Using Lemma 9.13 we know that there is a closed interval  $\mathcal{J} \subset \partial\mathbb{H}^2$  (with boundary =  $\{\xi, \eta\}$ ) such that each bigon  $B_i$  in  $L$  pushes to a bigon in every  $L'$  with  $\alpha(L') \in \mathcal{J}$ . Moreover, if  $E_i$  is the leaf of  $\tilde{\mathcal{F}}_2$  such that  $E_i$  intersects  $L$  in the boundary of  $B_i$ , then we have that every leaf  $L'$  as above has the corresponding bigon contained in between the intersection of  $L'$  and  $E_i$ . Note that Remark 9.11 implies that the nonseparated point of the corresponding bigons are in the same direction.

Since the intersection of  $E_i$  with the leaves  $L'$  from  $L$  to  $\hat{L}$  separates the region between  $L$  and  $\hat{L}$ , the order of the bigons cannot be reversed. We deduce that in  $\hat{L}$  we have bigons in the same order and the same directions. It remains to show that there cannot be bigons in  $\hat{L}$  that do not come from pushing those in  $L$ , but this just follows by a symmetric argument.  $\square$

**Conclusion.** Assume that  $B_1$  satisfies  $\xi_{B_1}^- = \alpha(L)$ , the other case is symmetric. In this case we deduce from Claim 1 that  $\xi_{\hat{B}_1}^+ = \alpha(\hat{L})$ , and hence  $\xi_{\hat{B}_{2k}}^- = \alpha(\hat{L})$ . By Proposition 9.19 it follows that  $\mathcal{J}_{B_1} \cup \mathcal{J}_{\hat{B}_{2k}} = \partial\mathbb{H}^2$ ; to see this, first apply it to  $L$  so  $\mathcal{J}_{B_1}$  is the interval opposite to  $B_i, i > 1$ . When seen in  $\hat{L} \cup S^1(\hat{L})$  we have that  $\mathcal{J}_{B_1}$  is the interval in the same complementary component of  $\hat{L} \setminus \hat{B}_{2k}$  that contains  $\hat{B}_1$ . Now apply it to  $\hat{L}$ : then  $\mathcal{J}_{\hat{B}_{2k}}$  is the component opposite from  $\hat{B}_1$  when seen from  $\hat{B}_{2k}$ . Hence  $\mathcal{J}_{B_1} \cup \mathcal{J}_{\hat{B}_{2k}} = \partial\mathbb{H}^2$ .

Finally, applying Lemma 9.13, we deduce that for every  $L' \in \tilde{\mathcal{F}}_1$  there must be a non-Hausdorff bigon joining  $\xi$  and  $\eta$  in  $L'$ .  $\square$

**Corollary 9.20.** *Assume that  $\mathcal{F}_1$  and  $\mathcal{F}_2$  are two transverse minimal foliations of  $M = T^1S$  such that the foliation  $\mathcal{G}$  obtained as their intersection is not homeomorphic to the orbit foliation of the geodesic flow for a hyperbolic metric on  $S$ . Then, there are finitely many disjoint simple closed curves  $s_1, \dots, s_k$  in  $S$  such that for every lift  $\tilde{s}_j$  of some of these curves, if  $\xi, \eta$  denote the endpoints of  $\tilde{s}_j$  in  $\partial\mathbb{H}^2$ , they satisfy the following: for every leaf  $L \in \tilde{\mathcal{F}}_i$  we have that  $L$  has a non-Hausdorff bigon joining  $\xi$  and  $\eta$ .*

Even if we fix the homotopy classes of the curves  $s_i$  which contain Reeb surfaces, it could be that the intersected foliations are not equivalent since there are possible

variants to the Matsumoto–Tsuboi construction (see Remark 7.1). Thus we can add several copies of  $\mathbb{T}^2 \times I$  associated with the same  $s_i$  and they will not produce equivalent foliations. This statement gives a combinatorial way to describe the possible intersected foliations.

*Proof.* It follows from Theorem 9.1 that once a leaf  $L$  contains a non-Hausdorff bigon invariant by some  $\gamma \in \pi_1(M) \setminus \{\text{id}\}$  then every leaf contains a non-Hausdorff bigon joining the same endpoints. Note that since everything is equivariant under the action of  $\pi_1(M)$  we get that every leaf must contain a non-Hausdorff bigon joining the fixed points of all elements in  $\pi_1(M)$  conjugated to  $\gamma$ . Since non-Hausdorff bigons cannot intersect, these form a lamination in  $\partial\mathbb{H}^2$  (by this we mean a collection of pairs of distinct points in  $\partial\mathbb{H}^2$  which are pairwise not linked) and thus corresponds to a simple closed curve in  $S$  (see also Proposition 9.14). Moreover, if there is another  $\gamma'$ -invariant non-Hausdorff bigon, it must be also disjoint from the first one, so we get that it is associated to a disjoint simple closed curve in  $S$ . Since there are only finitely many homotopy classes of disjoint simple closed curves in a surface  $S$  of genus  $g \geq 2$  we obtain the result.  $\square$

This section motivates the following:

**Question.** *Let  $\mathcal{F}_1, \mathcal{F}_2$  be two transverse minimal foliations in a closed 3-manifold  $M$ . Assume that the lift  $\tilde{\mathcal{G}}$  of the intersection foliation  $\mathcal{G} = \mathcal{F}_1 \cap \mathcal{F}_2$  to the universal cover does not have Hausdorff leaf space. Is it true that  $M$  contains a  $\pi_1$ -injective torus  $T$  such that every leaf  $L \in \tilde{\mathcal{F}}_i$  contains non-Hausdorff bigons joining the endpoints of the intersections of lifts of  $T$  with  $L$ ? In particular, is it true that if  $M$  is atoroidal then two transverse foliations intersect with Hausdorff leaf space in the universal cover?*

## 10. Application to partially hyperbolic diffeomorphisms

A diffeomorphism  $f : M \rightarrow M$  will be said *partially hyperbolic* if it admits a  $Df$ -invariant splitting  $TM = E^s \oplus E^c \oplus E^u$  such that there is some  $n > 0$  such that for  $x \in M$  and unit vectors  $v^s \in E^s(x)$ ,  $v^c \in E^c(x)$  and  $v^u \in E^u(x)$  we have

$$(10-1) \quad \|Df^n v^s\| < \frac{1}{2} \min\{1, \|Df^n v^c\|\}, \quad \|Df^n v^u\| > 2 \max\{1, \|Df^n v^c\|\}.$$

The problem of classification of partially hyperbolic diffeomorphisms in 3-manifolds introduced in [7; 14] has seen a lot of activity in the last few years. In particular, we point to [6; 4; 25] for the analysis when the 3-manifold is hyperbolic, and more generally for diffeomorphisms homotopic to the identity in general 3-manifolds. In [5] we have introduced a class of systems called *collapsed Anosov flows* which would provide a natural and useful notion of classification of such systems. In [25] we have shown that this class contains all partially hyperbolic

diffeomorphisms in hyperbolic 3-manifolds. The proof involves a careful study of pairwise transverse foliations, but also dynamics is introduced at several points in a crucial way. As mentioned in the introduction, the goal of this paper is to see to which extent we can extract dynamical information by using only the geometric properties of transverse foliations.

We note that besides the classical examples of time one maps of geodesic flows (or more generally, *discretized Anosov flows* ([6; 33]), unit tangent bundles admit many other classes of examples (see [8; 9]), some of which have been studied in [3; 25].

The goal of this section is to prove Corollary B which states that every conservative partially hyperbolic diffeomorphism in  $M = T^1S$  is a collapsed Anosov flow up to finite cover and it is thus accessible (see [24]). This section will assume some familiarity with standard results and notions of partial hyperbolicity; all of them can be found in [5].

**10.1. Preliminary results and precise statement.** Recall from [14] that when a partially hyperbolic diffeomorphism has orientable bundles whose orientation is preserved by  $f$ , then it preserves transverse *branching foliations*  $\mathcal{W}^{cs}$  and  $\mathcal{W}^{cu}$  tangent respectively to  $E^s \oplus E^c$  and  $E^c \oplus E^u$ . In [30, Theorem 3.1] some conditions are obtained which imply that these foliations do not have vertical leaves (for instance, being volume preserving is one of such conditions). Vertical means the leaf can be homotoped to be a union of Seifert fibers in  $M = T^1S$ . If no vertical leaves exist, then, arguments like in [6; 4] (see in particular [25, Proposition 8.3]) will allow us to prove the following:

**Theorem 10.1.** *Let  $f : M \rightarrow M$  be a partially hyperbolic diffeomorphism on  $M = T^1S$  where  $S$  is a surface of genus  $g \geq 2$ . Assume moreover that  $f$  preserves branching foliations which do not have vertical leaves. Then,  $f$  is a collapsed Anosov flow.*

Here, being a collapsed Anosov flow means that there is a semiconjugacy between  $f$  and a self-orbit equivalence of an Anosov flow of  $M$  (which by a classical result of Ghys must be orbit equivalent to a geodesic flow, see [28]). The semiconjugacy is required to have some technical properties relating the flow and the center direction. We obtain the strongest such condition, called *strong collapsed Anosov flow* in [5]. Since we will deduce a property of the leaves that implies such condition by [5, Theorem D] we will refer the reader to [5] for the actual definition of a collapsed Anosov flow.

These are the two results we shall use to be able to apply Theorem A to partially hyperbolic diffeomorphisms:

**Theorem 10.2** (Burago and Ivanov). *Let  $f : M \rightarrow M$  be a partially hyperbolic diffeomorphism preserving an orientation of its invariant bundles. Then,  $f$  admits*

branching foliations  $\mathcal{W}^{cs}$  and  $\mathcal{W}^{cu}$  which are approximated by foliations  $\mathcal{F}^{cs}$  and  $\mathcal{F}^{cu}$ . Moreover, if the branching foliations  $\mathcal{W}^{cs}$  and  $\mathcal{W}^{cu}$  are minimal, then one can choose  $\mathcal{F}^{cs}$  and  $\mathcal{F}^{cu}$  to also be minimal.

We refer the reader to [5, § 3] for discussion of this result, in particular the final property. The approximation is such that there is a *collapsing* map  $h : M \rightarrow M$  which sends leaves of  $\mathcal{F}^{cs}$  to leaves of  $\mathcal{W}^{cs}$  (same for  $cu$ ), is homotopic to the identity, is  $C^1$  along leaves, is  $\varepsilon$ -close to the identity, and has derivatives along the leaves which is  $C^1$  close to the identity along the leaves, and has further properties. Many properties transfer between the two foliations, in particular the topological types of leaves of both  $\mathcal{W}^{cs}$  and  $\mathcal{F}^{cs}$  is the same.

To obtain Theorem 10.1 we will then apply the following criterion given by [5, Theorem D]:

**Theorem 10.3.** *Let  $f : M \rightarrow M$  be a partially hyperbolic diffeomorphism preserving an orientation of its invariant bundles. If leaves of the foliation obtained by intersecting  $\mathcal{F}^{cs}$  and  $\mathcal{F}^{cu}$  are quasigeodesics in the universal cover of their respective leaves, then  $f$  is a collapsed Anosov flow.*

Accessibility and ergodicity then follow from [24, Theorem A]. We note that Corollary B follows since the branching foliations of volume preserving partially hyperbolic diffeomorphisms are what we call  $f$ -minimal, so they cannot have vertical leaves, and therefore satisfy the assumptions of Theorem 10.1. We note in fact that  $f$ -minimality is studied in [30] in many situations, including the case where  $f$  is *chain-recurrent* (something weaker than volume preserving or transitive) or belongs to certain isotopy classes. It is also shown in [5, Proposition 4.8] that being  $f$ -minimal is an open and closed condition on partially hyperbolic diffeomorphisms, so that if some  $f$  is known to be isotopic to a chain recurrent one along partially hyperbolic diffeomorphisms, then it will be in the hypothesis of Theorem 10.1.

**10.2. Partially hyperbolic foliations do not admit Reeb surfaces.** We now explain why existence of a Reeb surface in the approximating foliations is not possible if the branching foliations come from a partially hyperbolic diffeomorphism. This will reduce the proof of Theorem 10.1 to showing that under its assumptions the branching foliations exist and are minimal.

**Proposition 10.4.** *Let  $f : M \rightarrow M$  be a partially hyperbolic diffeomorphism preserving branching foliations  $\mathcal{W}^{cs}$  and  $\mathcal{W}^{cu}$  which are approximated in the sense of Theorem 10.2 by true transverse foliations  $\mathcal{F}^{cs}$  and  $\mathcal{F}^{cu}$ . Then, the approximating foliations do not have Reeb surfaces.*

*Proof.* A Reeb surface of  $\mathcal{F}^{cs}$  is finitely covered by an annulus whose boundary components are leaves of  $\mathcal{F}^{cs} \cap \mathcal{F}^{cu}$  and whose interior is made of infinitely many curves spiraling to the boundary. When collapsing to  $\mathcal{W}^{cs}$  and  $\mathcal{W}^{cu}$ , these annuli

collapse to “branched” annuli  $S$  contained in some leaf of  $\mathcal{W}^{cs}$  or  $\mathcal{W}^{cu}$  (say  $\mathcal{W}^{cs}$ ) which is an annulus, the boundaries are leaves of the one-dimensional branching foliation induced by the intersection and no transversal from one side intersects the other. Using the Poincaré–Bendixson theorem we deduce that every flow transverse to the boundaries of  $S$  must have a periodic orbit. Applying this to the flow generated by a unit vector field tangent to  $E^s$  we deduce the existence of a closed curve tangent to  $E^s$  which is a contradiction with partial hyperbolicity (because  $f$  would contract the curve until you find a circle tangent to  $E^s$  in an arbitrarily small ball).  $\square$

**10.3. Proof of Theorem 10.1.** Using Theorem 8.1, Theorem 6.1, and Theorem 10.3 it is enough to show that the approximating foliations  $\mathcal{F}^{cs}$  and  $\mathcal{F}^{cu}$  given by Theorem 10.2 are minimal. For this, it is enough to show that this is true for the branching foliations under the assumption that there are no vertical leaves for such foliations.

Theorem 10.2 shows that the collapsing map can be chosen to be a bijection between the sets of leaves of  $\mathcal{F}^{cs}$  and  $\mathcal{F}_{ws}$  (say), and the collapsing map preserves their homotopic properties. Therefore  $\mathcal{F}^{cs}$ ,  $\mathcal{F}^{cu}$  do not have vertical leaves by assumption.

Since  $\mathcal{F}^{cs}$  is horizontal, the leaf space of  $\tilde{\mathcal{F}}^{cs}$  is homeomorphic to the reals  $\mathbb{R}$ , and  $\mathcal{F}^{cs}$  blows down to a minimal foliation (see [20]). By Theorem 2.1 this foliation has only planes and annuli leaves. Hence  $\mathcal{W}^{cs}$  has only planes and annuli leaves (here, the topology of the leaf is, by definition, the topology of the quotient of the leaf in the universal cover by the deck transformations that fix the given leaf).

Since every leaf of  $\mathcal{W}^{cs}$  is a cylinder or a plane and the foliation is  $\mathbb{R}$ -covered we can argue exactly as in [25, Proposition 8.3] to get a contradiction with partial hyperbolicity (the quasigeodesic property is used in the proof of [25, Proposition 8.3] only to show that leaves are cylinders or planes). This completes the proof of minimality.

**10.4. Proof of Corollary B.** Take a regular finite cover  $M_1$  and iterate in order to have orientability of the bundles as well as their preservation. As explained in [30, § 7], once we take a finite cover  $M_1$ , since the foliations will not have vertical leaves they need to be horizontal. To show that there are no vertical leaves and that the branching foliations are minimal in the cover  $M_1$ , we use the volume preservation assumption as in [30]. In [30, Lemma 7.1] and [30, Subsection 6.4] it is proved that deck translations associated with the cover  $M_1 \rightarrow M$  preserve the orientations of all the bundles. This implies that the original bundles were horizontal and the orientability conditions were satisfied in  $M$ .

Hence the partially hyperbolic diffeomorphism  $f$  is a collapsed Anosov flow. It follows that  $f$  is accessible, and if  $f$  is  $C^2$  (and volume preserving) then  $f$  is ergodic (see [24]). This proves Corollary B.

### Acknowledgements

We thank Thierry Barbot for pointing out [35], which was crucial to this work, and particularly Section 9. We also thank the several referees, who gave important feedback on the paper.

### References

- [1] T. Barbot, S. Fenley, and R. Potrie, “On transverse R-covered minimal foliations”, preprint, 2025. arXiv 2501.14489
- [2] T. Barthelmé and K. Mann, “Pseudo-Anosov flows, a plane approach”, preprint, 2025. arXiv 2509.15375
- [3] T. Barthelmé, S. R. Fenley, S. Frankel, and R. Potrie, “Dynamical incoherence for a large class of partially hyperbolic diffeomorphisms”, *Ergodic Theory Dynam. Systems* **41**:11 (2021), 3227–3243. MR
- [4] T. Barthelmé, S. R. Fenley, S. Frankel, and R. Potrie, “Partially hyperbolic diffeomorphisms homotopic to the identity in dimension 3, II: Branching foliations”, *Geom. Topol.* **27**:8 (2023), 3095–3181. MR
- [5] T. Barthelmé, S. R. Fenley, and R. Potrie, “Collapsed Anosov flows and self orbit equivalences”, *Comment. Math. Helv.* **98**:4 (2023), 771–875. MR
- [6] T. Barthelmé, S. R. Fenley, S. Frankel, and R. Potrie, “Partially hyperbolic diffeomorphisms homotopic to the identity in dimension 3, I: The dynamically coherent case”, *Ann. Sci. Éc. Norm. Supér. (4)* **57**:2 (2024), 293–349. MR
- [7] C. Bonatti and A. Wilkinson, “Transitive partially hyperbolic diffeomorphisms on 3-manifolds”, *Topology* **44**:3 (2005), 475–508. MR
- [8] C. Bonatti, A. Gogolev, and R. Potrie, “Anomalous partially hyperbolic diffeomorphisms, II: Stably ergodic examples”, *Invent. Math.* **206**:3 (2016), 801–836. MR
- [9] C. Bonatti, A. Gogolev, A. Hammerlindl, and R. Potrie, “Anomalous partially hyperbolic diffeomorphisms, III: Abundance and incoherence”, *Geom. Topol.* **24**:4 (2020), 1751–1790. MR
- [10] C. Bonatti, J. Bowden, and R. Potrie, “Some remarks on projective Anosov flows in hyperbolic 3-manifolds”, pp. 359–369 in 2018 *MATRIX annals*, MATRIX Book Ser. **3**, Springer, [2020] ©2020. MR
- [11] S. Boyer and Y. Hu, “Taut foliations in branched cyclic covers and left-orderable groups”, *Trans. Amer. Math. Soc.* **372**:11 (2019), 7921–7957. MR
- [12] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, Grundle Math. Wissen. **319**, Springer, 1999. MR
- [13] M. Brittenham, “Essential laminations in Seifert-fibered spaces”, *Topology* **32**:1 (1993), 61–85. MR
- [14] D. Burago and S. Ivanov, “Partially hyperbolic diffeomorphisms of 3-manifolds with abelian fundamental groups”, *J. Mod. Dyn.* **2**:4 (2008), 541–580. MR
- [15] D. Calegari, “Leafwise smoothing laminations”, *Algebr. Geom. Topol.* **1** (2001), 579–585. MR
- [16] D. Calegari, *Foliations and the geometry of 3-manifolds*, Oxford University Press, 2007. MR
- [17] A. Candel and L. Conlon, *Foliations*, Graduate Studies in Mathematics **23**, Amer. Math. Soc., 2000. MR
- [18] S. R. Fenley, “Anosov flows in 3-manifolds”, *Ann. of Math. (2)* **139**:1 (1994), 79–115. MR

- [19] S. R. Fenley, “Foliations with good geometry”, *J. Amer. Math. Soc.* **12**:3 (1999), 619–676. MR
- [20] S. R. Fenley, “Foliations, topology and geometry of 3-manifolds:  $\mathbb{R}$ -covered foliations and transverse pseudo-Anosov flows”, *Comment. Math. Helv.* **77**:3 (2002), 415–490. MR
- [21] S. R. Fenley, “Geometry of foliations and flows, I: Almost transverse pseudo-Anosov flows and asymptotic behavior of foliations”, *J. Differential Geom.* **81**:1 (2009), 1–89. MR
- [22] S. R. Fenley and R. Potrie, “Minimality of the action on the universal circle of uniform foliations”, *Groups Geom. Dyn.* **15**:4 (2021), 1489–1521. MR
- [23] S. R. Fenley and R. Potrie, “Ergodicity of partially hyperbolic diffeomorphisms in hyperbolic 3-manifolds”, *Adv. Math.* **401** (2022), art. id. 108315, 45 pp. MR
- [24] S. R. Fenley and R. Potrie, “Accessibility and ergodicity for collapsed Anosov flows”, *Amer. J. Math.* **146**:5 (2024), 1339–1359. MR
- [25] S. R. Fenley and R. Potrie, “Partial hyperbolicity and pseudo-Anosov dynamics”, *Geom. Funct. Anal.* **34**:2 (2024), 409–485. MR
- [26] S. Fenley and R. Potrie, “Partial hyperbolicity, ergodicity, and transverse foliations in dimension 3”, preprint, 2025. arXiv 2510.15176
- [27] S. R. Fenley and R. Potrie, “Intersection of transverse foliations in 3-manifolds: Hausdorff leaf space implies leafwise quasigeodesic”, *J. Reine Angew. Math.* **822** (2025), 1–48. MR
- [28] E. Ghys, “Flots d’Anosov sur les 3-variétés fibrées en cercles”, *Ergodic Theory Dynam. Systems* **4**:1 (1984), 67–80. MR
- [29] E. Ghys, “Groupes d’homéomorphismes du cercle et cohomologie bornée”, pp. 81–106 in *The Lefschetz centennial conference* (Mexico City, 1984), Contemp. Math. **58**-III, Amer. Math. Soc., 1987. MR
- [30] A. Hammerlindl, R. Potrie, and M. Shannon, “Seifert manifolds admitting partially hyperbolic diffeomorphisms”, *J. Mod. Dyn.* **12** (2018), 193–222. MR
- [31] T. Inaba and S. Matsumoto, “Nonsingular expansive flows on 3-manifolds and foliations with circle prong singularities”, *Japan. J. Math. (N.S.)* **16**:2 (1990), 329–340. MR
- [32] G. Levitt, “Foliations and laminations on hyperbolic surfaces”, *Topology* **22**:2 (1983), 119–135. MR
- [33] S. Martinchich, “Global stability of discretized Anosov flows”, *J. Mod. Dyn.* **19** (2023), 561–623. MR
- [34] S. Matsumoto, “Some remarks on foliated  $S^1$  bundles”, *Invent. Math.* **90**:2 (1987), 343–358. MR
- [35] S. Matsumoto and T. Tsuboi, “Transverse intersections of foliations in three-manifolds”, pp. 503–525 in *Essays on geometry and related topics*, Monogr. Enseign. Math. **38**, Enseignement Math., Geneva, 2001. MR
- [36] E. E. Moise, *Geometric topology in dimensions 2 and 3*, Graduate Texts in Mathematics **47**, Springer, 1977. MR
- [37] M. Paternain, “Expansive flows and the fundamental group”, *Bol. Soc. Brasil. Mat. (N.S.)* **24**:2 (1993), 179–199. MR
- [38] W. P. Thurston, *Foliations of 3-manifolds which are circle bundles*, Ph.D. thesis, University of California, Berkeley, 1972, available at <https://www.proquest.com/docview/302592092>. MR
- [39] W. P. Thurston, “Three-manifolds, foliations and circles, I”, preprint, 1997. arXiv 9712268

Received June 11, 2024. Revised October 30, 2025.

SÉRGIO R. FENLEY  
DEPARTMENT OF MATHEMATICS  
FLORIDA STATE UNIVERSITY  
TALLAHASSEE, FL  
UNITED STATES  
fenley@math.fsu.edu

RAFAEL POTRIE  
CENTRO DE MATEMÁTICA  
UNIVERSIDAD DE LA REPÚBLICA  
MONTEVIDEO  
URUGUAY

and

IRL-IFUMI  
LABORATORIO DEL PLATA (CNRS)  
MONTEVIDEO  
URUGUAY  
rpotrie@cmat.edu.uy

## REFINED BOUNDS FOR THE EIGENVALUES OF THE STOKES OPERATOR

ZHENGCHAO JI AND TÜRKAY YOLCU

**We analyze bounds for the sums of eigenvalues of the Stokes operator restricted to a bounded domain  $\Omega \subset \mathbb{R}^d$  with  $d \geq 2$ . We improve upon existing lower bound estimates due to Ilyin as well as those by the second author and S. Yıldırım Yolcu, while preserving sharpness in the sense of Weyl asymptotics.**

### 1. Introduction

Let  $\Omega$  be an open bounded set in  $\mathbb{R}^d$ ,  $d \geq 2$ . We derive sharper estimates for the eigenvalues  $\{\lambda_k\}_{k=1}^\infty$  of Stokes problem defined by

$$(1) \quad \begin{cases} -\Delta \mathbf{u}_k + \nabla p_k = \lambda_k \mathbf{u}_k & \text{in } \Omega, \\ \operatorname{div} \mathbf{u}_k = 0 & \text{in } \Omega, \\ \mathbf{u}_k = 0 & \text{on } \partial\Omega. \end{cases}$$

The eigenvalues of the Stokes problem in (1) are important due to their numerous applications in fluid mechanics. They can be interpreted as the eigenvalues associated with linear (small or infinitesimal) self-oscillations of the fluid within the domain  $\Omega$  [4]. When the eigenfunctions are required to meet specific conditions that are crucial from a physical perspective in fluid dynamics, exact eigenvalues for the Stokes problem are not obtainable. Consequently, both theoretical and practical aspects necessitate the close identification of the eigenvalues. The literature concerning the eigenvalues of the Stokes problem is vast, and recent studies aimed at deriving estimates for these eigenvalues are documented in [6; 9; 10; 12; 24].

Before presenting the results, we first review some basic facts about the theory of the Stokes operator. Let  $\mathcal{U}$  denote the set of smooth, divergence-free vector functions with compact supports. Specifically,

$$\mathcal{U} = \{\mathbf{u} : \Omega \rightarrow \mathbb{R}^d, \mathbf{u} \in C_0^\infty(\Omega), \operatorname{div} \mathbf{u} = 0\}$$

---

*MSC2020:* primary 35P15; secondary 35P30.

*Keywords:* Stokes operator, Berezin–Li–Yau inequality, eigenvalue, inequality, Weyl asymptotics.

and let the closure of  $\mathcal{U}$  in  $L^2(\Omega)$  and  $\mathbf{H}_0^1(\Omega)$  be denoted by  $L$  and  $U$ , respectively. In particular,

$$U \subseteq \{\mathbf{u} \in \mathbf{H}_0^1(\Omega), \operatorname{div} \mathbf{u} = 0\}$$

and if  $\Omega$  is open, bounded and locally Lipschitz, we have the equality [3]:

$$U = \{\mathbf{u} \in \mathbf{H}_0^1(\Omega), \operatorname{div} \mathbf{u} = 0\}.$$

We note that  $L^2(\Omega)$  can be written as  $L^2(\Omega) = L \oplus L^\perp$ , where

$$L^\perp = \{\mathbf{u} \in L^2(\Omega), \mathbf{u} = \nabla p, p \in L_{\text{loc}}^2(\Omega)\}$$

(see [3; 4; 19]). The Stokes operator is formally expressed as

$$A\mathbf{u} = -P_L(\Delta\mathbf{u}),$$

where the linear operator  $P_L : L^2(\Omega) \rightarrow L$ , defined by

$$P_L(\mathbf{v}) = \mathbf{v} - \nabla\Delta^{-1}(\operatorname{div} \mathbf{v}),$$

is called the Leray projection [4], i.e.,  $P_L(P_L(\mathbf{v})) = P_L(\mathbf{v})$ . Notice that  $P_L$  becomes the identity operator for the divergence-free vector fields because  $P_L(\mathbf{v}) = \mathbf{v}$  for  $\operatorname{div} \mathbf{v} = 0$ . Furthermore,  $P_L$  can be understood as the projection onto divergence-free vector fields. This projection is particularly employed to remove some terms and components in the Stokes and Navier–Stokes equations. Going back to the Stokes problem in (1), one can also see that

$$A\mathbf{u} = -\Delta\mathbf{u} + \nabla p, \quad p = \Delta^{-1}(\operatorname{div} \Delta\mathbf{u}),$$

and that for all  $\mathbf{u}, \mathbf{w}$  in  $U$ ,  $A$  is defined by

$$(A\mathbf{u}, \mathbf{w}) = (\nabla\mathbf{u}, \nabla\mathbf{w}),$$

i.e.,

$$\int_{\Omega} A\mathbf{u}(\mathbf{x}) \cdot \mathbf{w}(\mathbf{x}) \, d\mathbf{x} = \int_{\Omega} \sum_{i=1}^d \frac{\partial \mathbf{u}}{\partial x_i} \frac{\partial \mathbf{w}}{\partial x_i} \, d\mathbf{x}.$$

The Stokes operator  $A$  is an unbounded, linear, self-adjoint, positive definite operator in  $L$ . In addition to these nice properties, its inverse  $A^{-1}$  is a compact, self-adjoint operator [3; 4]. Thus, there exists an orthonormal basis  $\{\mathbf{u}_k\}_{k=1}^\infty$  in  $L$  and a set of positive eigenvalues  $\{\delta_k\}_{k=1}^\infty$  accumulating at zero such that

$$A^{-1} \mathbf{u}_k = \delta_k \mathbf{u}_k,$$

for  $k = 1, 2, 3, \dots$ . Therefore,  $\{\mathbf{u}_k\}_{k=1}^\infty \in U$  with corresponding eigenvalues  $\{\lambda_k\}_{k=1}^\infty$  with  $\lambda_k = 1/\delta_k$  are such that

$$(2) \quad A\mathbf{u}_k = \lambda_k \mathbf{u}_k.$$

The eigenvalues (including multiplicities) satisfy

$$(3) \quad 0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_k \leq \dots \rightarrow \infty \quad \text{as } k \rightarrow \infty.$$

Observe that by taking the scalar product with orthonormal  $\mathbf{u}_k$  we have

$$(4) \quad \lambda_k = \|\nabla \mathbf{u}_k\|^2 = \int_{\Omega} \sum_{i=1}^d \frac{\partial \mathbf{u}_k}{\partial x_i} \cdot \frac{\partial \mathbf{u}_k}{\partial x_i} dx.$$

The eigenvalues also satisfy the Weyl asymptotic formulas [4; 8; 17] for  $d \geq 2$ :

$$(5) \quad \lambda_k \sim 4\pi \left( \frac{\Gamma(1 + \frac{d}{2})}{(d-1)|\Omega|} \right)^{\frac{2}{d}} k^{\frac{2}{d}},$$

where  $\Gamma(x)$  denotes the gamma function  $\Gamma(x) = \int_0^\infty t^{x-1} e^{-t} dt$  for  $x > 0$  and  $|\Omega|$  denotes the Lebesgue measure of  $\Omega$ .

The eigenvalue bounds that will be established in this paper closely resemble those associated with the Dirichlet Laplacian. A substantial body of literature exists regarding inequalities for the eigenvalues of the Dirichlet Laplacian; notable references include the survey articles cited in [1; 2; 7]. In his renowned paper [18], G. Pólya demonstrated that for small values of  $k$ , the ratio  $4\pi k/|\Omega|$  serves as a lower bound for the eigenvalues  $\mu_k$  of the Dirichlet Laplacian in tiling domains in  $\mathbb{R}^2$ . Pólya conjectured that this result could be extended to any bounded domain in  $\mathbb{R}^d$ . This conjecture remains unresolved. A significant advance in establishing a lower bound was achieved by P. Li and S.-T. Yau [14], who demonstrated the following inequality related to the sums of the eigenvalues  $\mu_j$  of the Dirichlet Laplacian on the domain  $\Omega$ :

$$(6) \quad \sum_{j=1}^k \mu_j \geq 4\pi \frac{d}{d+2} \left( \frac{\Gamma(1 + \frac{d}{2})}{|\Omega|} \right)^{\frac{2}{d}} k^{1+\frac{2}{d}},$$

Recent research has focused significantly on these types of bounds, along with their extensions and enhancements, particularly in relation to other operators. For instance, A. Melas [16] provided an improvement on (6) that incorporates moments of inertia. In two dimensions, H. Kovařík, S. Vugalter, and T. Weidl [13] further advanced Melas’s findings by introducing a positive correction term that involves the size of the boundary. T. Weidl [21] achieved an enhancement of the sharp Berezin-type bounds concerning the Riesz means  $\sum_k (z - \mu_k)_+^\sigma$  of the eigenvalues associated with the Dirichlet Laplacian operator within a specified domain for  $\sigma \geq 3/2$ . Regarding other operators, Harrell and Yıldırım Yolcu [5] along with Yolcu [29] proved inequalities of Berezin–Li–Yau type applicable to the eigenvalues of Klein–Gordon operators. Yıldırım Yolcu and Yolcu [25; 28; 27] established

Melas-type improvements for the eigenvalues of fractional Laplacian operators  $(-\Delta)^{\alpha/2}$ ,  $\alpha \in (0, 2]$ .

It is noteworthy that the coefficient of  $k^{1+\frac{2}{d}}$  in (6) is slightly larger for the tiling domains [18]. See [23] for more extensive research and a generalized conjecture.

Lower bounds on the sums of eigenvalues are important for the theoretical framework of attractors associated with the Navier–Stokes equations [9; 10]. The eigenvalue bounds of the Stokes operator have garnered attention following the work by A. A. Ilyin [9], who proved an inequality of Berezin–Li–Yau type for the Stokes operator:

$$(7) \quad \sum_{j=1}^k \lambda_j \geq 4\pi \frac{d}{d+2} \left( \frac{\Gamma(1+\frac{d}{2})}{(d-1)|\Omega|} \right)^{\frac{2}{d}} k^{1+\frac{2}{d}},$$

where  $k \geq 1$ . In this article, we first build upon (7) by establishing upper bounds for the sums of negative powers and lower bounds for the sums of positive powers of the eigenvalues of the Stokes operator as follows:

**Theorem 1.** *For  $k \geq 1$ ,  $0 < b < d/2$ , and  $d \geq 2$ , the sums of negative powers of eigenvalues  $\{\lambda_i\}_{i=1}^{\infty}$  of the Stokes operator on  $\Omega$  in (1) satisfy*

$$(8) \quad \sum_{j=1}^k \lambda_j^{-b} \leq (4\pi)^{-b} \frac{d}{d-2b} \left( \frac{(d-1)|\Omega|}{\Gamma(1+\frac{d}{2})} \right)^{\frac{2b}{d}} k^{1-\frac{2b}{d}}.$$

**Theorem 2.** *For  $k \geq 1$ ,  $0 < a \leq 1$ , and  $d \geq 2$ , the sums of positive powers of eigenvalues  $\{\lambda_i\}_{i=1}^{\infty}$  of the Stokes operator on  $\Omega$  in (1) satisfy*

$$(9) \quad \sum_{j=1}^k \lambda_j^a \geq (4\pi)^a \frac{d}{d+2a} \left( \frac{\Gamma(1+\frac{d}{2})}{(d-1)|\Omega|} \right)^{\frac{2a}{d}} k^{1+\frac{2a}{d}}.$$

By appropriately translating the open set  $\Omega$  if needed, we can assume that the second moment of inertia  $I(\Omega)$  is defined as

$$I(\Omega) = \int_{\Omega} |\mathbf{x}|^2 d\mathbf{x}.$$

Ilyin [10] was able to improve the Berezin–Li–Yau inequality in (7) for the Stokes operator by adding a lower-order term involving the moment of inertia  $I(\Omega)$  in dimensions 2, 3, and 4 as follows:

$$(10) \quad \sum_{j=1}^k \lambda_j \geq 4\pi \frac{d}{d+2} \left( \frac{\Gamma(1+\frac{d}{2})}{(d-1)|\Omega|} \right)^{\frac{2}{d}} k^{1+\frac{2}{d}} + \frac{d-1}{48} \beta_d^S \frac{|\Omega|}{I(\Omega)} k,$$

where in the two-dimensional case  $\beta_2^S = \frac{239}{240}$ , while for  $d = 3, 4$  it suffices to take  $\beta_3^S = 0.986$  and  $\beta_4^S = 0.978$ .

Let

$$B_R(\mathbf{x}) := \{\mathbf{y} \in \mathbb{R}^d : |\mathbf{y} - \mathbf{x}| \leq R\}$$

be the ball of radius  $R$  centered at  $\mathbf{x}$  in  $\mathbb{R}^d$ . Let

$$(11) \quad \omega_d = \frac{\pi^{\frac{d}{2}}}{\Gamma(1 + \frac{d}{2})}$$

be the volume of the unit ball  $B_1(\mathbf{x})$  in  $\mathbb{R}^d$ .

Building on the improvements in (10), Yıldırım Yolcu and Yolcu [24] introduced additional correction terms beyond those dependent on  $k^{1+\frac{2}{d}}$  and  $k \geq 1$ , along with their explicit coefficients, applicable for any  $d \geq 2$ :

$$(12) \quad \sum_{j=1}^k \lambda_j \geq 4\pi \frac{d}{d+2} \left( \frac{\Gamma(1 + \frac{d}{2})}{(d-1)|\Omega|} \right)^{\frac{2}{d}} k^{1+\frac{2}{d}} + \frac{d-1}{24(d^2+2d)} \frac{|\Omega|}{I(\Omega)} k \\ + \frac{((d-1)|\Omega|)^{\frac{3d+2}{2d}}}{144\sqrt{\pi} d^{\frac{3}{2}} (d+2) I(\Omega)^{\frac{3}{2}} \Gamma(1 + \frac{d}{2})^{\frac{1}{d}}} k^{1-\frac{1}{d}}.$$

In this paper we establish several effective lower bounds for the sums of the eigenvalues of the Stokes problem, including the following improved estimates:

**Theorem 3** (refinement of the Berezin–Li–Yau inequality for the Stokes operator). *Let  $\Omega \subset \mathbb{R}^d$ . For any  $d \geq m+1 \geq 2$ , the eigenvalues  $\{\lambda_i\}_{i=1}^\infty$  of the Stokes operator in (1) satisfy*

$$\sum_{i=1}^k \lambda_i \geq 4\pi \left( \frac{\Gamma(1 + \frac{d}{2})}{(d-1)|\Omega|} \right)^{\frac{2}{d}} k^{1+\frac{2}{d}} - \frac{2\omega_d^{\frac{m-1}{d}} A_{m+2} \varrho^{\frac{(m+1)d+m-1}{d}}}{(d+2)M^{m+1}} k^{1-\frac{m-1}{d}} \\ + c \frac{2\omega_d^{\frac{m}{d}} (m+1) A_{m+3} \varrho^{\frac{(m+2)d+m}{d}}}{(d+2)(m+3)M^{m+2}} k^{1-\frac{m}{d}},$$

where  $k \geq 1$ ,  $a$  is defined by the equality in (39),

$$(13) \quad \varrho = \frac{d-1}{(2\pi)^d} |\Omega|, \quad m_s = \frac{2\sqrt{(d^2-d)|\Omega|I(\Omega)}}{(2\pi)^d}, \quad M = \frac{\sqrt{d^2-d}}{(2\pi)^d} |\Omega|^{\frac{d+1}{d}} \omega_d^{-\frac{1}{d}},$$

and the remaining notation is defined by

$$A_r = (a+1)^r - a^r, \quad c = \min\{1, \max\{a_1, a_2\}\}, \\ a_1 = \frac{(d+2)(m+3)}{(m+1)((m+2)d+m)} \frac{(2\pi)^{m+2}}{A_{m+3}} d^{\frac{m+2}{2}} (d-1)^{-\frac{m(d+2)+2m}{2d}},$$

and

$$a_2 = \frac{\sqrt{2}d^{\frac{1}{d}}}{2(d-1)^{\frac{d+2}{2d}}} \frac{[(m+1)d+m-1](m+3)}{[(m+2)d+m](m+1)} \frac{A_{m+2}}{A_{m+3}}.$$

Theorem 3 yields no improvement when  $d = 2$ , but it quickly yields the following enhanced lower bound for any dimension  $d \geq 3$ :

**Corollary 1.** *For any bounded domain  $\Omega \subseteq \mathbb{R}^d$ , with  $d \geq 3$ , and any  $k \geq 1$ , the eigenvalues  $\{\lambda_i\}_{i=1}^\infty$  of the Stokes operator in (1) satisfy*

$$\begin{aligned} \sum_{i=1}^k \lambda_i \geq & \frac{4\pi d}{d+2} \left( \frac{\Gamma(1+\frac{d}{2})}{(d-1)|\Omega|} \right)^{\frac{2}{d}} k^{1+\frac{2}{d}} + \frac{\varrho^2}{6(d+2)m_s^2} k + \frac{\omega_d^{-\frac{1}{d}} \varrho^{\frac{3d+1}{d}}}{9(d+2)m_s^3} k^{\frac{d-1}{d}} \\ & + \frac{3\omega_d^{-\frac{1}{d}} \varrho^{\frac{4d+2}{d}}}{80(n+2)m_s^4} k^{\frac{d-2}{d}}, \end{aligned}$$

where  $\varrho$  and  $m_s$  are as in (13).

When  $d = 3$ , we have an improvement:

**Theorem 4.** *Let  $\Omega \subset \mathbb{R}^d$  with  $d = 3$ . Then the eigenvalues  $\{\lambda_i\}_{i=1}^\infty$  of the Stokes operator in (1) satisfy*

$$(14) \quad \sum_{i=1}^k \lambda_i \geq \frac{2\pi\varrho^6}{15m_s^5} ((t+1)^6 - t^6),$$

where  $m_s$  and  $\varrho$  are defined in (13) and

$$t = \frac{1}{2} \left( \left( T + \sqrt{T^2 + \frac{1}{27}} \right)^{\frac{1}{3}} - \left( -T + \sqrt{T^2 + \frac{1}{27}} \right)^{\frac{1}{3}} - 1 \right), \quad T = \frac{(d+1)m_s^d}{\omega_d \varrho^{d+1}} k.$$

Furthermore,

$$\sum_{i=1}^k \lambda_i \geq \frac{3}{5} \left( \frac{2\pi^3}{2\omega_3|\Omega|} \right)^{\frac{2}{3}} k^{\frac{5}{3}} + \frac{1}{24} \frac{|\Omega|}{I(\Omega)} k - \frac{11\pi\omega_3^{-\frac{1}{3}} \varrho^{\frac{14}{3}}}{180 m_s^4} k^{\frac{1}{3}} + \frac{659\pi\omega_3^{\frac{1}{3}} \varrho^{\frac{22}{3}}}{75000 \cdot 4^{\frac{1}{3}} m_s^6} k^{-\frac{1}{3}}.$$

Inspired by Melas's research [16], we adopted a similar methodology using fundamental techniques from prior studies [10; 22; 24; 27; 28; 29; 26]. While preserving the core strategy of these works, we introduced significant modifications to achieve stronger lower bounds.

**Outline.** Section 2 presents results needed for proving the theorems discussed in this work. The main content is found in Sections 3–6, which respectively establish Theorems 1, 2, 3 and 4, proving intermediate results as needed.

**2. Preliminaries**

The set of eigenfunctions  $\{\mathbf{u}_j\}_{j=1}^\infty$  of the Stokes operator is orthonormal in  $L^2(\Omega)$ , and so the set of Fourier transforms  $\{\hat{\mathbf{u}}_j\}_{j=1}^\infty$  also forms an orthonormal set in  $L^2(\mathbb{R}^d)$ , by Plancherel’s theorem. Let us set

$$(15) \quad U_k(\xi) = \sum_{j=1}^k |\hat{\mathbf{u}}_j(\xi)|^2 = \sum_{j=1}^k \left| \frac{1}{(2\pi)^{\frac{d}{2}}} \int_{\Omega} e^{-ix \cdot \xi} \mathbf{u}_j(\mathbf{x}) \, d\mathbf{x} \right|^2 \geq 0.$$

The integral is taken over  $\Omega$  instead of  $\mathbb{R}^d$  because the support of  $\mathbf{u}_j$  is  $\Omega$ .

The following result, proven by A. A. Ilyin [10, Section 2], is essential to our calculations.

**Lemma 1** [10]. *The functions  $U_k$  defined by (15) satisfy*

$$(16) \quad \int_{\mathbb{R}^d} U_k(\xi) \, d\xi = k,$$

$$(17) \quad U_k(\xi) \leq \frac{d-1}{(2\pi)^d} |\Omega|, \quad \xi \in \mathbb{R}^d,$$

$$(18) \quad \int_{\mathbb{R}^d} |\xi|^2 U_k(\xi) \, d\xi = \sum_{j=1}^k \lambda_j.$$

Note that equations (16)–(18) bear a striking resemblance to their Laplacian counterparts. However, the proof of Lemma 1 employs suborthonormal functions [10].

We highlight some important properties of the function  $U$ . Let  $U^*(\xi)$  designate the radial decreasing rearrangement of  $U(\xi)$ . By approximating  $U$  if needed, we may assume that there exists a real-valued absolutely continuous function

$$\zeta : [0, \infty) \rightarrow \left[0, \frac{d-1}{(2\pi)^d} |\Omega|\right]$$

such that  $U^*(\xi) = \zeta(|\xi|)$ . Then

$$(19) \quad 0 \leq -\zeta' \leq m_s,$$

where  $m_s$  depends only on  $d$  and  $\Omega$  and is explicitly defined in (13). To see (19), let  $\mu$  be the distribution function defined by

$$\mu(s) = |\{U(\xi) > s\}| = |\{U^*(\xi) > s\}|.$$

Then, we observe that

$$\mu(\zeta(t)) = |\{U^*(\xi) > \zeta(t)\}| = |\{\xi : |\xi| < t\}| = |B_t(0)| = w_d t^d.$$

Invoking the coarea formula in view of (17), we have

$$\mu(s) = \int_s^\infty \int_{\{U^{-1}(t)\}} \frac{1}{|\nabla U|} d\mathcal{H} dt = \int_s^{\frac{d-1}{(2\pi)^d} |\Omega|} \int_{\{U=t\}} \frac{1}{|\nabla U|} d\mathcal{H} dt,$$

where  $\mathcal{H}$  is the  $(d-1)$ -dimensional Hausdorff measure. Let us consider  $t > 0$  values such that  $\zeta'(t) < 0$ .

Let  $\bar{K}$  denote the closure of  $K \subset \mathbb{R}^d$  and  $\partial K$  the boundary of  $K$ . The isoperimetric inequality,

$$\mathcal{H}(\partial K) \geq dw_d^{\frac{1}{d}} |\bar{K}|^{\frac{d-1}{d}},$$

enables us to deduce

$$\begin{aligned} \frac{dw_d t^{d-1}}{\zeta'(t)} &= \mu'(\zeta(t)) = - \int_{\{U=\zeta(t)\}} \frac{1}{|\nabla U|} d\mathcal{H} \leq -\frac{1}{m_S} \mathcal{H}(\{U = \zeta(t)\}) \\ &\leq -\frac{1}{m_S} dw_d^{\frac{1}{d}} \mu(\zeta(t))^{\frac{d-1}{d}} = -\frac{1}{m_S} dw_d t^{d-1}. \end{aligned}$$

This inequality together with  $\zeta' \leq 0$  easily yields (19). Note that (19) essentially states that if the gradient vector of the original function is bounded, then the gradient of the rearrangement retains the same bound. (An alternative derivation is possible, using the Pólya–Szegő inequality.)

Now, let  $r$  represent the real number such that  $|\Omega| = w_d r^d$ . We can get a lower bound for  $I(\Omega)$ :

$$I(\Omega) \geq \int_{B_r(0)} |\mathbf{x}|^2 d\mathbf{x} = \frac{dw_d}{d+2} r^{d+2} = \frac{d}{d+2} w_d^{-\frac{2}{d}} |\Omega|^{\frac{d+2}{d}},$$

which quickly results in a lower bound for  $m_S$  as follows:

$$(20) \quad m_S = \frac{2\sqrt{(d^2-d)|\Omega|I(\Omega)}}{(2\pi)^d} \geq \frac{\sqrt{d^2-d}}{(2\pi)^d} |\Omega|^{\frac{d+1}{d}} \omega_d^{-\frac{1}{d}} =: M.$$

### 3. Proof of Theorem 1

The following proof is inspired from the proof of the Berezin–Li–Yau inequality in [5; 14; 28; 29]. An analogous proof is also given in [20] by means of the bathtub principle [15].

*Proof of Theorem 1.* Assume the properties (16)–(18). Since  $|\hat{\mathbf{u}}_j(\xi)|^2 d\xi$  is a probability measure on  $\mathbb{R}^d$  and  $t \mapsto t^{-b}$  is convex for  $t > 0$  and  $b > 0$ , employing Jensen’s inequality and (15), we obtain

$$(21) \quad \sum_{j=1}^k \lambda_j^{-b} = \sum_{j=1}^k \left( \int_{\mathbb{R}^d} |\xi|^2 |\hat{\mathbf{u}}_j(\xi)|^2 d\xi \right)^{-b} \leq \int_{\mathbb{R}^d} \frac{U_k(\xi)}{|\xi|^{2b}} d\xi.$$

Let  $\mathbf{1}_\Omega(t)$  denote the characteristic function of the set  $\Omega$ . Define

$$(22) \quad \begin{aligned} \Psi_k(\xi) &= \frac{d-1}{(2\pi)^d} |\Omega| \mathbf{1}_{B_{S_k}(0)}(\xi) \quad \text{and} \\ S_k &= \left( \frac{(d-2b)(2\pi)^d (\sum_{j=1}^k \lambda_j^{-b})}{d(d-1)\omega_d |\Omega|} \right)^{\frac{1}{d-2b}}, \end{aligned}$$

so that

$$(23) \quad \begin{aligned} \int_{\mathbb{R}^d} \frac{\Psi_k(\xi)}{|\xi|^{2b}} d\xi &= \frac{d-1}{(2\pi)^d} |\Omega| d\omega_d \int_0^{S_k} \frac{1}{t^{2b}} t^{d-1} dt \\ &= \frac{d-1}{(2\pi)^d} |\Omega| \frac{d\omega_d}{d-2b} S_k^{d-2b} = \sum_{j=1}^k \lambda_j^{-b}. \end{aligned}$$

Notice that

$$(24) \quad \left( \frac{1}{|\xi|^{2b}} - \frac{1}{S_k^{2b}} \right) (U_k(\xi) - \Psi_k(\xi)) \leq 0.$$

Integrating (24) on  $\mathbb{R}^d$  and using (23) we arrive at

$$(25) \quad \frac{1}{S_k^{2b}} \int_{\mathbb{R}^d} (U_k(\xi) - \Psi_k(\xi)) d\xi \geq \int_{\mathbb{R}^d} \frac{U_k(\xi) - \Psi_k(\xi)}{|\xi|^{2b}} d\xi \geq 0$$

from which it follows that

$$\int_{\mathbb{R}^d} U_k(\mu) d\mu \geq \int_{\mathbb{R}^d} \Psi_k(\mu) d\mu.$$

Thus, by (16), we obtain

$$(26) \quad k \geq \int_{\mathbb{R}^d} \Psi_k(\mu) d\mu = \frac{d-1}{(2\pi)^d} |\Omega| d\omega_d \left( \frac{S_k^d}{d} \right).$$

Substituting  $\omega_d$  given by (11) and  $S_k$  given by (22) into (26) and rearranging the terms, we obtain (8).  $\square$

#### 4. Proof of Theorem 2

We sketch a direct proof of (9) for the sake of completeness.

*Proof of Theorem 2.* Recall that  $U_k$  satisfies (16)–(18). Since  $|\hat{\mathbf{u}}_j(\xi)|^2 d\xi$  is a probability measure on  $\mathbb{R}^d$  and  $r \mapsto r^a$  is concave for  $r > 0$  and  $0 < a \leq 1$ , we can use Jensen's inequality to derive that

$$(27) \quad \sum_{j=1}^k \lambda_j^a = \sum_{j=1}^k \left( \int_{\mathbb{R}^d} |\xi|^2 |\hat{\mathbf{u}}_j(\xi)|^2 d\xi \right)^a \geq \int_{\mathbb{R}^d} |\xi|^{2a} U_k(\xi) d\xi.$$

Setting

$$(28) \quad \Theta_k(\xi) = \frac{d-1}{(2\pi)^d} |\Omega| \mathbf{1}_{B_{T_k}(0)}(\xi), \quad T_k = \left( \frac{(d+2a)(2\pi)^d (\sum_{j=1}^k \lambda_j^a)}{d(d-1)\omega_d |\Omega|} \right)^{\frac{1}{d+2a}},$$

we have

$$(29) \quad \begin{aligned} \int_{\mathbb{R}^d} |\xi|^{2a} \Theta_k(\xi) d\xi &= \frac{d-1}{(2\pi)^d} |\Omega| d \omega_d \int_0^{T_k} r^{2a} r^{d-1} dr \\ &= \frac{d-1}{(2\pi)^d} |\Omega| \frac{d \omega_d}{d+2a} T_k^{d+2a} = \sum_{j=1}^k \lambda_j^a. \end{aligned}$$

Now observe that

$$(30) \quad (|\xi|^{2a} - T_k^{2a})(U_k(\xi) - \Theta_k(\xi)) \geq 0.$$

Integrating (30) on  $\mathbb{R}^d$  and using (27) and (29) we conclude that

$$(31) \quad T_k^{2a} \int_{\mathbb{R}^d} (U_k(\xi) - \Theta_k(\xi)) d\xi \leq \int_{\mathbb{R}^d} |\xi|^{2a} (U_k(\xi) - \Theta_k(\xi)) d\xi \leq 0,$$

which yields

$$\int_{\mathbb{R}^d} U_k(\xi) d\xi \leq \int_{\mathbb{R}^d} \Theta_k(\xi) d\xi.$$

Now we use (16) to obtain

$$(32) \quad k \leq \int_{\mathbb{R}^d} \Theta_k(\xi) d\xi = \frac{d-1}{(2\pi)^d} |\Omega| d \omega_d \left( \frac{T_k^d}{d} \right).$$

Substituting the values of  $\omega_d$  and  $T_k$  given in (11) and (28) into (32) and simplifying, we deduce (9).  $\square$

### 5. Proof of Theorem 3

Before the proof of Theorem 3, we give some elementary results from [11; 24]. To connect two key integrals in our proof, we introduce an important quantity,  $a$ , using the idea in [24]:

**Lemma 2.** For any function  $\alpha : [0, \infty) \rightarrow [0, 1]$  and integer  $d \geq 1$  with

$$\int_0^\infty \phi(t) dt = 1, \quad \int_0^\infty t^d \phi(t) dt < \infty, \quad \int_0^\infty t^{d+2} \phi(t) dt \leq \infty,$$

there exists an  $a \geq 0$  such that

$$\int_a^{a+1} t^d dt = \int_0^\infty t^d \phi(t) dt \quad \text{and} \quad \int_a^{a+1} t^{d+2} dt \leq \int_0^\infty t^{d+2} \phi(t) dt.$$

The following sharp inequality is key in estimating the lower bounds for  $\sum_{i=1}^k \lambda_i$ .

**Lemma 3** [11]. *For an integer  $d \geq m + 1 \geq 2$  and positive real numbers  $t$  and  $\tau$  we have*

$$(33) \quad dt^{d+2} - (d + 2)\tau^2 t^d + 2\tau^{d+2} - \sum_{k=1}^{m+1} 2kt^{k-1}\tau^{n-k+1}(\tau - t)^2 \geq 0.$$

Considering all of these findings, we arrive at the following intermediate estimate.

**Proposition 1.** *If  $d \geq m + 1 \geq 2$ , we have*

$$(34) \quad \sum_{i=1}^k \lambda_i \geq \omega_d (dA)^{\frac{d+2}{d}} (\zeta(0))^{-\frac{2}{d}} - \frac{2\omega_d A_{m+2} (dA)^{\frac{d-m+1}{d}} (\zeta(0))^{\frac{(m+1)d+m-1}{d}}}{(d + 2)m_s^{m+1}} + \frac{2\omega_d (m + 1) A_{m+3} (dA)^{\frac{d-m}{d}} (\zeta(0))^{\frac{(m+2)d+m}{d}}}{(d + 2)(m + 3)m_s^{m+2}},$$

where  $m_s$  is defined in (13) and

$$A_r := (a + 1)^r - a^r \geq 1, \quad A := \frac{1}{d} \int_a^{a+1} t^d dt, \quad 0 \leq \zeta(0) \leq \frac{d - 1}{(2\pi)^d} |\Omega|,$$

the quantity  $a$  being defined by

$$\int_a^{a+1} t^d dt = \frac{k}{\omega_d}.$$

*Proof.* We use a strategy similar to that of [11; 24]. Since the map  $\xi \mapsto |\xi|^2$  is radial and increasing, Lemma 1 implies that

$$(35) \quad k = \int_{\mathbb{R}^d} U(\xi) d\xi = \int_{\mathbb{R}^d} U^*(\xi) d\xi = d\omega_d \int_0^\infty t^{d-1} \zeta(t) dt,$$

$$(36) \quad \sum_{i=1}^k \lambda_i = \int_{\mathbb{R}^d} |\xi|^2 U(\xi) d\xi \geq \int_{\mathbb{R}^d} |\xi|^2 U^*(\xi) d\xi = d\omega_d \int_0^\infty t^{d+1} \zeta(t) dt.$$

For convenience, we rescale  $\zeta$  as

$$S(t) = \frac{1}{\zeta(0)} \zeta\left(\frac{\zeta(0)}{m_s} t\right).$$

Obviously,  $S(t)$  is a positive function with  $S(0) = 1$  and  $0 \leq -S'(t) \leq 1$ . Let  $\phi(t) := -S'(t)$  and define

$$(37) \quad A = \int_0^\infty t^{d-1} S(t) dt, \quad B = \int_0^\infty t^{d+1} S(t) dt.$$

Integrating by parts, we get

$$(38) \quad \int_0^\infty t^d \phi(t) dt = dA, \quad \int_0^\infty t^{d+2} \phi(t) dt = (d+2)B.$$

According to (38) and Lemma 2, there exists  $a \geq 0$  such that

$$(39) \quad \int_a^{a+1} t^d dt = dA \quad \text{and} \quad \int_a^{a+1} t^{d+2} dt \leq (d+2)B.$$

For  $i \geq 0$ ,  $\tau \geq \frac{1}{2}$  and  $a \geq 0$ , we have

$$(40) \quad \int_a^{a+1} t^i (\tau - t)^2 dt = \frac{A_{i+3}}{i+3} - \frac{2A_{i+2}}{i+2} \tau + \frac{A_{i+1}}{i+1} \tau^2,$$

where

$$A_r = (a+1)^r - a^r \geq 1.$$

Taking into account Lemma 3 and (40), we may integrate (33) in  $t$  from  $a$  to  $a+1$  to get

$$(41) \quad d(d+2)B \geq d(d+2)A\tau^2 - 2\tau^{d+2} + \sum_{i=1}^{m+1} 2i\tau^{d-i+1} \left( \frac{A_i}{i} \tau^2 - \frac{2A_{i+1}}{i+1} \tau + \frac{A_{i+2}}{i+2} \right).$$

The summation on the second row of (41) can be rewritten as

$$\begin{aligned} &= 2\tau^{d+2} + 2 \sum_{i=1}^m A_{i+1} \tau^{d-i+2} - 2 \sum_{i=1}^{m+1} \frac{2iA_{i+1}}{i+1} \tau^{d-i+2} + 2 \sum_{i=1}^{m+1} \frac{iA_{i+2}}{i+2} \tau^{d-i+1} \\ &= 2\tau^{d+2} + 2A_2\tau^{d+1} + \frac{2mA_{m+2}}{m+2} \tau^{d-m+1} + \frac{2(m+1)A_{m+3}}{m+3} \tau^{d-m} \\ &\quad - 2A_2\tau^{d+1} - \frac{4(m+1)A_{m+2}}{m+2} \tau^{d-m+1} + 2 \sum_{i=2}^m \left( 1 + \frac{i-1}{i+1} - \frac{2i}{i+1} \right) A_{i+1} \tau^{d-i+2}. \end{aligned}$$

The parenthetical factor in this last summation vanishes, so the right-hand side of the preceding display boils down to

$$\frac{2(m+1)A_{m+3}}{m+3} \tau^{d-m} + 2\tau^{d+2} - 2A_{m+2}\tau^{d-m+1},$$

and we deduce that

$$(42) \quad d(d+2)B - (d+2)\tau^2 dA \geq \frac{2(m+1)A_{m+3}}{m+3} \tau^{d-m} - 2A_{m+2}\tau^{d-m+1}.$$

Using Jensen's inequality, we obtain  $(dA)^{\frac{1}{d}} \geq \int_\delta^{\delta+1} t dt \geq \frac{1}{2}$  for any  $\delta \geq 0$ . Hence,

putting  $\tau = (dA)^{\frac{1}{d}}$  in (42), we arrive at

$$(43) \quad B \geq \frac{(dA)^{\frac{d+2}{d}}}{d} - \frac{2A_{m+2}(dA)^{\frac{d-m+1}{n}}}{d(d+2)} + \frac{2(m+1)A_{m+3}(nA)^{\frac{d-m}{d}}}{d(d+2)(m+3)}.$$

Taking the rescaling of  $\zeta(t)$  into consideration, we obtain from (43) the integral inequality

$$\int_0^\infty t^{s+1} \zeta(t) dt \geq \frac{(dA)^{\frac{d+2}{n}} (\zeta(0))^{-\frac{2}{d}}}{d} - \frac{2S_{m+2}(dA)^{\frac{d-m+1}{d}} (\zeta(0))^{\frac{(m+1)d+m-1}{d}}}{d(d+2)m_s^{m+1}} + \frac{2(m+1)S_{m+3}(dA)^{\frac{d-m}{d}} (\zeta(0))^{\frac{(m+2)n+m}{d}}}{d(d+2)(m+3)m_s^{m+2}}.$$

Notice that  $\inf_t \zeta(t) \leq \zeta(0) \leq \sup_t \zeta(t)$ . Due to (36), the estimate of  $\int_0^\infty t^{s+1} \zeta(t) dt$  provides a lower bound

$$\sum_{i=1}^k \lambda_i \geq \omega_d (dA)^{\frac{d+2}{d}} (\zeta(0))^{-\frac{2}{d}} - \frac{2\omega_d A_{m+2}(dA)^{\frac{d-m+1}{d}} (\zeta(0))^{\frac{(m+1)d+m-1}{d}}}{(d+2)m_s^{m+1}} + \frac{2\omega_d(m+1)A_{m+3}(dA)^{\frac{d-m}{d}} (\zeta(0))^{\frac{(m+2)d+m}{d}}}{(d+2)(m+3)m_s^{m+2}}.$$

This is what we required in Proposition 1.  $\square$

The right-hand side of the intermediate lower bound (34) depends on  $\zeta(0)$ . Our goal is to derive a final eigenvalue inequality that depends solely on  $d$  and  $k$ . To achieve this, we adopt a strategy of minimizing (34) with respect to  $\zeta(0)$  over the interval  $[0, \varrho]$ , where  $\varrho = \frac{d-1}{(2\pi)^d} |\Omega|$  as defined in (13).

*Proof of Theorem 3.* Using (35) and (38) in combination with integration by parts, we obtain

$$\frac{k}{\omega_n} = d \int_0^\infty t^{d-1} \zeta(t) dt = \int_0^\infty -t^d \zeta'(t) dt = \int_0^\infty t^d \phi(t) dt = dA.$$

Using the fact that  $dA = k/\omega_d$ , we define  $G_1(x)$ ,  $G_2(x)$  and  $G(x)$  for  $x \in [0, \varrho]$  by

$$G_1(x) = \omega_d^{-\frac{2}{d}} x^{-\frac{2}{d}} k^{\frac{d+2}{d}} + c_2 \frac{2\omega_d^{\frac{m}{d}} (m+1)A_{m+3} x^{\frac{(m+2)d+m}{d}}}{(d+2)(m+3)m_s^{m+2}} k^{\frac{d-m}{d}},$$

$$G_2(x) = -\frac{2\omega_d^{\frac{m-1}{d}} A_{m+2} x^{\frac{(m+1)d+m-1}{d}}}{(d+2)m_s^{m+1}} k^{\frac{d-m+1}{d}},$$

$$G(x) = G_1(x) - G_2(x).$$

Here  $0 < c_2 \leq 1$  is a positive number to be determined. Since  $m \geq 1$ , we conclude that  $G_2(x)$  is decreasing on  $[0, \varrho]$ . By direct calculation, we have

$$\frac{d}{2} \frac{\omega_d^{\frac{2}{d}} G_1'(x)}{k^{\frac{d-m}{d}}} = -k^{\frac{2+d}{d}} x^{-\frac{d+2}{d}} + c_2 \omega_d^{\frac{m+2}{d}} \frac{(m+1)((m+2)d+m)}{(d+2)(m+3)} \frac{A_{m+3}}{m_s^{m+2}} x^{\frac{(m+1)d+m}{d}}.$$

Therefore, when

$$c_2 \leq \frac{k^{\frac{d+2}{d}}}{\omega_d^{\frac{m+2}{d}}} \frac{(d+2)(m+3)}{(m+1)[(m+2)d+m]} \frac{m_s^{m+2}}{A_{m+3}} \varrho^{-\frac{(m+2)(d+1)}{d}},$$

we conclude that  $G_1'(x) \leq 0$  on  $[0, \varrho]$ . Since

$$(44) \quad m_s \geq M = \frac{\sqrt{d^2 - d}}{(2\pi)^d \omega_d^{\frac{1}{d}}} |\Omega|^{\frac{d+1}{d}}, \quad \varrho = \frac{d-1}{(2\pi)^d} |\Omega|,$$

we can choose  $c_1 = \min\{1, a_1\}$ , where  $a_1$  is defined by

$$(45) \quad a_1 = \frac{(d+2)(m+3)}{(m+1)[(m+2)d+m]} \frac{(2\pi)^{m+2}}{A_{m+3}} \frac{d^{\frac{m+2}{2}}}{(d-1)^{\frac{m(d+2)+2m}{2d}}},$$

to obtain

$$(46) \quad \sum_{i=1}^k \lambda_i \geq \omega_d^{-\frac{2}{d}} \varrho^{-\frac{2}{d}} k^{\frac{d+2}{d}} - \frac{2\omega_d^{\frac{m-1}{d}} A_{m+2} \varrho^{\frac{(m+1)d+m-1}{d}}}{(d+2)M^{m+1}} k^{\frac{d-m+1}{d}} + c_1 \frac{2\omega_d^{\frac{m}{d}} (m+1) A_{m+3} \varrho^{\frac{(m+2)d+m}{d}}}{(d+2)(m+3)M^{m+2}} k^{\frac{d-m}{d}},$$

where  $\varrho$  and  $M$  are defined in (13). Similarly, we can split  $G(x)$  as a sum of two functions:

$$G(x) = g_1(x) + g_2(x), \quad x \in [0, \varrho],$$

where  $g_1$  and  $g_2$  are defined by  $g_1(x) = \omega_d^{-\frac{2}{d}} x^{-\frac{2}{d}} k^{\frac{d+2}{d}}$  and

$$g_2(x) = c_2 \frac{2\omega_d^{\frac{m}{d}} (m+1) A_{m+3} x^{\frac{(m+2)d+m}{d}}}{(d+2)(m+3)m_s^{m+2}} k^{\frac{d-m}{d}} - \frac{2\omega_d^{\frac{m-1}{d}} A_{m+2} x^{\frac{(m+1)d+m-1}{d}}}{(d+2)m_s^{m+1}} k^{\frac{d-m+1}{d}}.$$

Obviously,  $g_1(x)$  is decreasing on  $[0, \varrho]$ . Rewriting  $g_2(x)$  as

$$\frac{d(d+2)}{2} \frac{m_s}{\omega_d^{\frac{m}{d}} k^{\frac{d-m}{d}}} g_2(x) = c_2 \frac{[(m+2)d+m](m+1)A_{m+3}}{(m+3)m_s} x^{\frac{(m+1)d+m}{d}} - ((m+1)d+m-1)\omega_d^{-\frac{1}{d}} A_{m+2} k^{\frac{1}{d}} x^{\frac{(d+1)m-1}{d}}$$

allows us to observe that if

$$c_2 \leq \frac{[(m+1)d+m-1](m+3)}{[(m+2)d+m](m+1)} \frac{A_{m+2}}{A_{m+3}} \frac{m_s}{\omega_d^{1/d}} \varrho^{-\frac{d+1}{d}},$$

then  $g'_2(x) \leq 0$  on  $[0, \varrho]$ . Using (44) again, we choose  $c_3 = \min\{1, a_2\}$ , where  $a_2$  is defined by

$$(47) \quad a_2 = \frac{\sqrt{2}d^{\frac{1}{d}}}{2(d-1)^{\frac{d+2}{2d}}} \frac{[(m+1)d+m-1](m+3)}{[(m+2)d+m](m+1)} \frac{A_{m+2}}{A_{m+3}},$$

to derive that

$$(48) \quad \sum_{i=1}^k \lambda_i \geq \omega_d^{-\frac{2}{d}} \varrho^{-\frac{2}{d}} k^{\frac{d+2}{d}} - \frac{2\omega_d^{\frac{m-1}{d}} A_{m+2} \varrho^{\frac{(m+1)d+m-1}{d}}}{(d+2)M^{m+1}} k^{\frac{d-m+1}{d}} + c_3 \frac{2\omega_d^{\frac{m}{d}} (m+1) A_{m+3} \varrho^{\frac{(m+2)d+m}{d}}}{(d+2)(m+3)M^{m+2}} k^{\frac{d-m}{d}}.$$

In light of equations (46) and (48), we can use  $c = \min\{1, \max\{a_1, a_2\}\}$ , where  $a_1$  and  $a_2$  are defined in (45) and (47), to get

$$(49) \quad \sum_{i=1}^k \lambda_i \geq \omega_d^{-\frac{2}{d}} \varrho^{-\frac{2}{d}} k^{\frac{d+2}{d}} - \frac{2\omega_d^{\frac{m-1}{d}} A_{m+2} \varrho^{\frac{(m+1)d+m-1}{d}}}{(d+2)M^{m+1}} k^{\frac{d-m+1}{d}} + c \frac{2\omega_d^{\frac{m}{d}} (m+1) A_{m+3} \varrho^{\frac{(m+2)d+m}{d}}}{(d+2)(m+3)M^{m+2}} k^{\frac{d-m}{d}}.$$

Finally, we use

$$(50) \quad \omega_d^{-\frac{2}{d}} \varrho^{-\frac{2}{d}} k^{\frac{d+2}{d}} = 4\pi \left( \frac{\Gamma(1 + \frac{d}{2})}{(d-1)|\Omega|} \right)^{\frac{2}{d}} k^{1 + \frac{2}{d}}$$

to complete the proof of Theorem 3. □

*Proof of Corollary 1.* An argument similar to that of [11] gives

$$d(d+2)B - d(d+2)A\tau^2 + 2\tau^{d+2} \geq \frac{1}{6}(\tau^d) + \frac{1}{9}(\tau^{d-1}) + \frac{3}{80}\tau^{d-2}.$$

Hence, we get

$$\sum_{i=1}^k \lambda_i \geq \frac{d\omega_d^{-\frac{2}{d}} \zeta(0)^{-\frac{2}{d}}}{d+2} k^{\frac{d+2}{d}} + \frac{\zeta(0)^2}{6(d+2)m_s^2} k + \frac{c_4 \omega_d^{-\frac{1}{d}} \zeta(0)^{\frac{3d+1}{d}}}{(d+2)m_s^3} k^{\frac{d-1}{d}} + \frac{c_5 \omega_d^{-\frac{1}{d}} \zeta(0)^{\frac{4d+2}{d}}}{(n+2)m_s^4} k^{\frac{d-2}{d}},$$

where  $0 < c_4 \leq \frac{1}{9}$  and  $0 < c_5 \leq \frac{3}{80}$  are two constants to be chosen. Following an approach analogous to the discussion in the proof of Theorem 3, we can choose  $c_4 = \frac{1}{9}$  and  $c_5 = \frac{3}{80}$ . Using the equality in (50) completes the proof.  $\square$

## 6. Proof of Theorem 4

The following lemma from [10] plays a pivotal role in establishing our eigenvalue inequality and makes use of the piecewise continuous function

$$\Phi_s(r) = \begin{cases} m_s & \text{if } 0 \leq r \leq s, \\ m_s - \varrho(r - s) & \text{if } s \leq r \leq s + m_s/\varrho, \\ 0 & \text{if } s + m_s/\varrho \leq r, \end{cases}$$

where  $m_s$  and  $\varrho$  are defined in (13).

**Lemma 4** [10]. *Let  $\alpha > 0$  be a real number and  $\Psi(r)$  be a decreasing and absolutely continuous function such that*

$$\int_0^\infty r^\alpha \Psi(r) dr = \int_0^\infty r^\alpha \Phi_s(r) dr, \quad 0 \leq \Psi \leq m_s, \quad -L < \Psi' < 0.$$

Then, for any  $\beta \geq \alpha$ , the following integral inequality holds:

$$\int_0^\infty r^\beta \Psi(r) dr \geq \int_0^\infty r^\beta \Phi_s(r) dr.$$

Moreover, for any  $\gamma \geq 0$ , we have

$$\int_0^\infty r^\gamma \Phi_s(r) dr = \frac{\varrho^{\gamma+2}}{(\gamma+1)(\gamma+2)m_s^{\gamma+1}} ((t+1)^{\gamma+2} - t^{\gamma+2}), \quad s = \frac{tm_s}{\varrho}.$$

*Proof of Theorem 4.* Let  $m_s$  be as in (13). Applying Lemma 4 to  $\Psi(x) = \zeta(x)$  and setting  $\alpha = d - 1$ , we obtain

$$\int_0^\infty t^{d-1} \zeta(t) dt = \frac{k}{d\omega_d} = \frac{\varrho^{d+1}}{d(d+1)m_s^d} ((t+1)^{d+1} - t^{d+1}),$$

which in turn implies that  $t$  is the unique root of the equation

$$(t+1)^{d+1} - t^{d+1} = T := \frac{(d+1)m_s^d}{\omega_d \varrho^{d+1}} k.$$

Then Lemma 4 implies

$$(51) \quad \int_0^\infty t^{d+1} \zeta(t) dt \geq \frac{\varrho^{d+3}}{(d+2)(d+3)m_s^{d+2}} ((t+1)^{d+3} - t^{d+3}).$$

One can easily check that  $T$  satisfies

$$T \geq 1, \quad T \geq \frac{d+1}{d-1} \frac{(4\pi)^d}{\omega_d^2} \left( \frac{d^2}{(d-1)(d+2)} \right)^{\frac{d}{2}}.$$

In particular, when  $d = 3$ , we consider the equation

$$(t + 1)^4 - t^4 = T = \frac{4\varrho^3}{\omega_3 m_s^4} k,$$

whose positive root is

$$t = \frac{1}{2} \left( \left( T + \sqrt{T^2 + \frac{1}{27}} \right)^{\frac{1}{3}} - \left( -T + \sqrt{T^2 + \frac{1}{27}} \right)^{\frac{1}{3}} - 1 \right).$$

From (36), we obtain our desired result as

$$(52) \quad \sum_{i=1}^k \lambda_i \geq \frac{2\pi\varrho^6}{15m_s^5} ((t + 1)^6 - t^6).$$

Considering the Taylor series of  $t$ , we arrive at

$$t \geq -\frac{1}{2} + 4^{-\frac{1}{3}} T^{\frac{1}{3}} - \frac{1}{6} 2^{-\frac{1}{3}} T^{-\frac{1}{3}} + \frac{1}{324} 4^{-\frac{1}{3}} T^{-\frac{5}{3}} + \frac{1}{1944} 2^{-\frac{1}{3}} T^{-\frac{7}{3}} - \frac{1}{26244} 4^{-\frac{1}{3}} T^{-\frac{11}{3}} - \frac{1}{629856} 2^{-\frac{1}{3}} T^{-\frac{13}{3}},$$

where we used the fact that  $T \geq 1$ . Hence, we obtain

$$(t + 1)^6 - t^6 \geq \frac{3}{4} 2^{-\frac{1}{3}} T^{\frac{5}{3}} + \frac{5}{8} T - \frac{11}{24} 2^{\frac{2}{3}} T^{\frac{1}{3}} + \frac{659}{10000} T^{-\frac{1}{3}}.$$

Putting the lower bound of  $(t + 1)^6 - t^6$  into (52), we get

$$\sum_{i=1}^k \lambda_i \geq \frac{3}{5} \left( \frac{2\pi^3}{2\omega_3 |\Omega|} \right)^{\frac{2}{3}} k^{\frac{5}{3}} + \frac{1}{24} \frac{|\Omega|}{I(\Omega)} k - \frac{11\pi}{180} \frac{\varrho^{\frac{14}{3}}}{\omega_3^{\frac{1}{3}} m_s^4} k^{\frac{1}{3}} + \frac{659\pi\omega_3^{\frac{1}{3}}}{75000 \cdot 4^{\frac{1}{3}}} \frac{\varrho^{\frac{22}{3}}}{m_s^6} k^{-\frac{1}{3}}.$$

This concludes the proof of Theorem 4. □

### Acknowledgements

The authors wish to thank the referees for carefully reading this paper and making numerous valuable suggestions and corrections. Z. Ji’s research was supported by the National Natural Science Foundation of China, Grant no. 12501068, and Zhejiang Provincial Natural Science Foundation of China, Grant no. LQN25A010001. T. Yolcu acknowledges support from the Caterpillar Fellowship (no. 2511045) during the preparation of this article. Ji expresses his deep gratitude to Prof. Hongwei Xu, whose guidance and mentorship in mathematics have profoundly shaped his understanding and growth as a researcher. It is with great respect and appreciation that we dedicate this paper to him.

## References

- [1] M. S. Ashbaugh, “The universal eigenvalue bounds of Payne–Pólya–Weinberger, Hile–Protter, and H. C. Yang”, *Proc. Indian Acad. Sci. Math. Sci.* **112**:1 (2002), 3–30. MR
- [2] M. S. Ashbaugh and R. D. Benguria, “Isoperimetric inequalities for eigenvalues of the Laplacian”, pp. 105–139 in *Spectral theory and mathematical physics: a Festschrift in honor of Barry Simon’s 60th birthday*, vol. 1, Proc. Sympos. Pure Math. **76**:1, Amer. Math. Soc., 2007. MR
- [3] P. Constantin and C. Foias, *Navier–Stokes equations*, University of Chicago Press, 1988. MR
- [4] C. Foias, O. Manley, R. Rosa, and R. Temam, *Navier–Stokes equations and turbulence*, Encyclopedia of Mathematics and its Applications **83**, Cambridge University Press, 2001. MR
- [5] E. M. Harrell, II and S. Yıldırım Yolcu, “Eigenvalue inequalities for Klein–Gordon operators”, *J. Funct. Anal.* **256**:12 (2009), 3977–3995. MR
- [6] J. Hu and Y. Huang, “Lower bounds for eigenvalues of the Stokes operator”, *Adv. Appl. Math. Mech.* **5**:1 (2013), 1–18. MR
- [7] D. Hundertmark, “Some bound state problems in quantum mechanics”, pp. 463–496 in *Spectral theory and mathematical physics: a Festschrift in honor of Barry Simon’s 60th birthday*, vol. 1, Proc. Sympos. Pure Math. **76**:1, Amer. Math. Soc., 2007. MR
- [8] A. A. Ilyin, “Attractors for Navier–Stokes equations in domains with finite measure”, *Nonlinear Anal.* **27**:5 (1996), 605–616. MR
- [9] A. A. Ilyin, “On the spectrum of the Stokes operator”, *Funktsional. Anal. i Prilozhen.* **43**:4 (2009), 14–25. In Russian; translated in *Funct. Anal. Appl.* **43**:4, 254–263 (2009). MR
- [10] A. A. Ilyin, “Lower bounds for the spectrum of the Laplace and Stokes operators”, *Discrete Contin. Dyn. Syst.* **28**:1 (2010), 131–146. MR
- [11] Z. Ji and H. Xu, “Lower bounds for eigenvalues of Laplacian operator and the clamped plate problem”, *Calc. Var. Partial Differential Equations* **62**:6 (2023), art. id. 175, 27 pp. MR
- [12] J. P. Kelliher, “Eigenvalues of the Stokes operator versus the Dirichlet Laplacian in the plane”, *Pacific J. Math.* **244**:1 (2010), 99–132. MR
- [13] H. Kovařík, S. Vugalter, and T. Weidl, “Two-dimensional Berezin–Li–Yau inequalities with a correction term”, *Comm. Math. Phys.* **287**:3 (2009), 959–981. MR
- [14] P. Li and S. T. Yau, “On the Schrödinger equation and the eigenvalue problem”, *Comm. Math. Phys.* **88**:3 (1983), 309–318. MR
- [15] E. H. Lieb and M. Loss, *Analysis*, 2nd ed., Graduate Studies in Mathematics **14**, Amer. Math. Soc., 2001. MR
- [16] A. D. Melas, “A lower bound for sums of eigenvalues of the Laplacian”, *Proc. Amer. Math. Soc.* **131**:2 (2003), 631–636. MR
- [17] G. Métivier, “Valeurs propres d’opérateurs définis par la restriction de systèmes variationnels à des sous-espaces”, *J. Math. Pures Appl.* (9) **57**:2 (1978), 133–156. MR
- [18] G. Pólya, “On the eigenvalues of vibrating membranes”, *Proc. London Math. Soc.* (3) **11** (1961), 419–433. MR
- [19] R. Temam, *Navier–Stokes equations: Theory and numerical analysis*, 3rd ed., Studies in Mathematics and its Applications **2**, North-Holland, Amsterdam, 1984. MR
- [20] V. Vougalter, “Sharp semiclassical bounds for the moments of eigenvalues for some Schrödinger type operators with unbounded potentials”, *Math. Model. Nat. Phenom.* **8**:1 (2013), 237–245. MR

- [21] T. Weidl, “Improved Berezin–Li–Yau inequalities with a remainder term”, pp. 253–263 in *Spectral theory of differential operators*, Amer. Math. Soc. Transl. Ser. 2 **225**, Amer. Math. Soc., 2008. MR
- [22] S. Yıldırım Yolcu, “An improvement to a Berezin–Li–Yau type inequality”, *Proc. Amer. Math. Soc.* **138**:11 (2010), 4059–4066. MR
- [23] S. Yıldırım Yolcu and T. Yolcu, “Bounds for the eigenvalues of the fractional Laplacian”, *Rev. Math. Phys.* **24**:3 (2012), art. id. 1250003, 18 pp. MR
- [24] S. Yıldırım Yolcu and T. Yolcu, “Multidimensional lower bounds for the eigenvalues of Stokes and Dirichlet Laplacian operators”, *J. Math. Phys.* **53**:4 (2012), art. id. 043508, 17 pp. MR
- [25] S. Yıldırım Yolcu and T. Yolcu, “Estimates for the sums of eigenvalues of the fractional Laplacian on a bounded domain”, *Commun. Contemp. Math.* **15**:3 (2013), art. id. 1250048, 15 pp. MR
- [26] S. Yıldırım Yolcu and T. Yolcu, “Estimates on the eigenvalues of the clamped plate problem on domains in Euclidean spaces”, *J. Math. Phys.* **54**:8 (2013), art. id. 043515. MR
- [27] S. Yıldırım Yolcu and T. Yolcu, “Refined eigenvalue bounds on the Dirichlet fractional Laplacian”, *J. Math. Phys.* **56**:7 (2015), art. id. 073506, 12 pp. MR
- [28] S. Yıldırım Yolcu and T. Yolcu, “Sharper estimates on the eigenvalues of Dirichlet fractional Laplacian”, *Discrete Contin. Dyn. Syst.* **35**:5 (2015), 2209–2225. MR
- [29] T. Yolcu, “Refined bounds for the eigenvalues of the Klein–Gordon operator”, *Proc. Amer. Math. Soc.* **141**:12 (2013), 4305–4315. MR

Received July 12, 2025. Revised February 11, 2026.

ZHENGCHAO JI  
 DEPARTMENT OF MATHEMATICS  
 CHINA JILIANG UNIVERSITY  
 HANGZHOU  
 CHINA  
 jizhengchao@zju.edu.cn

TÜRKAY YOLCU  
 DEPARTMENT OF MATHEMATICS  
 BRADLEY UNIVERSITY  
 PEORIA, IL  
 UNITED STATES  
 tyolcu@bradley.edu



## THE CLASS $Q$ AND MIXTURE DISTRIBUTIONS WITH DOMINATED CONTINUOUS SINGULAR PARTS

ALEXEY A. KHARTOV

We consider a new class  $Q$  of distribution functions  $F$  that have the property of rational-infinite divisibility: there exist some infinitely divisible distribution functions  $F_1$  and  $F_2$  such that  $F_1 = F * F_2$ . A distribution function of class  $Q$  is quasi-infinite divisible in the sense that its characteristic function admits the Lévy-type representation with a “signed spectral measure”. This class is a wide natural extension of the fundamental class of infinitely divisible distribution functions and it is being actively studied now. We are interested in conditions for a distribution function  $F$  to belong to class  $Q$  for the unexplored case, where  $F$  may have a continuous singular part. We propose a criterion under the assumption that the continuous singular part of  $F$  is dominated by the discrete part in a certain sense. The criterion generalizes the previous results by Alexeev and Khartov for discrete probability laws and the results by Berger and Kutlu for the mixtures of discrete and absolutely continuous laws. In addition, we describe the characteristic triplet of the corresponding Lévy-type representation, which may contain a continuous singular part. We also show that the assumption of the dominated continuous singular part cannot be omitted or even slightly extended (without some special assumptions). We apply the general criterion to some interesting particular examples. We also positively solve the decomposition problem stated by Lindner, Pan and Sato within the case being considered.

### 1. Introduction

This paper is devoted to the study of a new class of probability laws that naturally extends the fundamental class of infinitely divisible distributions.

Let  $F$  be a distribution function on the real line  $\mathbb{R}$ . Recall that  $F$  and the corresponding probability law are called *infinitely divisible* if for every positive integer  $n$  there exists a distribution function  $F_{1/n}$  such that  $F = (F_{1/n})^{*n}$ , where

---

*MSC2020:* 60E05, 60E07, 60E10.

*Keywords:* distribution function, characteristic function, continuous singular part, infinite divisibility, rational-infinite divisibility, quasi-infinite divisibility, Lévy-type representation.

“ $*$ ” denotes the convolution operation, i.e.,  $F$  is the  $n$ -fold convolution power of  $F_{1/n}$ . Let  $f$  be the characteristic function of  $F$ , i.e.,

$$f(t) := \int_{\mathbb{R}} e^{itx} dF(x), \quad t \in \mathbb{R}.$$

It is well-known (see [12; 31; 37]) that  $F$  is infinitely divisible if and only if  $f$  admits the *Lévy representation*

$$(1) \quad f(t) = \exp \left\{ it\gamma - \frac{\sigma^2 t^2}{2} + \int_{\mathbb{R} \setminus \{0\}} (e^{itx} - 1 - it \sin(x)) dL(x) \right\}, \quad t \in \mathbb{R},$$

with a *shift parameter*  $\gamma \in \mathbb{R}$ , a *Gaussian variance*  $\sigma^2 \geq 0$ , and a *Lévy spectral function*  $L : \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}$ , which is nondecreasing on each of the intervals  $(-\infty, 0)$  and  $(0, +\infty)$ , and satisfies

$$(2) \quad \lim_{x \rightarrow -\infty} L(x) = \lim_{x \rightarrow +\infty} L(x) = 0,$$

and also

$$\int_{O_\delta} x^2 dL(x) < \infty \quad \text{for any } \delta > 0,$$

where  $O_\delta := (-\delta, 0) \cup (0, \delta)$ . The function  $L$  is assumed to be right-continuous at every point of the real line. Importantly, the *characteristic triplet*  $(\gamma, \sigma^2, L)$  is uniquely determined by  $f$  and hence by  $F$ . Due to this representation, the class of infinitely divisible probability laws has found a lot of applications through Lévy processes (see [31]), the stochastic calculus (see [4]), teletraffic models (see [25]), and actuarial mathematics (see [32]).

Let  $\mathbf{I}$  denote the class of all infinitely divisible distribution functions on the real line. This class is naturally extended in the following way. We call a distribution function  $F$  (and the corresponding probability law) *rational-infinitely divisible* if there exist some infinitely divisible distribution functions  $F_1$  and  $F_2$  such that  $F_1 = F * F_2$ . In terms of characteristic functions, this definition is equivalent to the formula  $f(t) = f_1(t)/f_2(t)$ ,  $t \in \mathbb{R}$ , for the characteristic function  $f$  of  $F$ , where  $f_1$  and  $f_2$  are the characteristic functions of some infinitely divisible distribution functions  $F_1$  and  $F_2$ . We denote by  $\mathbf{Q}$  the class of all rational-infinitely divisible distribution functions. Since  $F_2$  may be chosen as degenerate at some point  $a$  (i.e.,  $f_2(t) = e^{ita}$ ,  $t \in \mathbb{R}$ ), it is clear that, indeed,  $\mathbf{I} \subset \mathbf{Q}$ . Moreover, from the definition, it is seen that the characteristic function  $f$  of any  $F \in \mathbf{Q}$  admits a *Lévy-type representation*. Namely, if  $F_1$  and  $F_2$  have characteristic triplets  $(\gamma_1, \sigma_1^2, L_1)$  and  $(\gamma_2, \sigma_2^2, L_2)$ , then formula (1) holds with the *shift parameter*  $\gamma = \gamma_1 - \gamma_2 \in \mathbb{R}$ , the *Gaussian variance*  $\sigma^2 = \sigma_1^2 - \sigma_2^2$ , and the *spectral function*  $L = L_1 - L_2$ . In that case,  $L$  has a bounded total variation on  $\mathbb{R} \setminus O_\delta$  for every  $\delta > 0$ , and, in general, it is nonmonotonic on the intervals  $(-\infty, 0)$  and  $(0, +\infty)$ . The function  $L$  also

inherits from  $L_1$  and  $L_2$  right-continuity on  $\mathbb{R}$  and property (2). Moreover,

$$\int_{O_\delta} x^2 d|L|(x) < \infty \quad \text{for any } \delta > 0,$$

where we integrate over the variation of  $L$ . We now suppose that, conversely,  $f$  admits representation (1) with some  $\gamma \in \mathbb{R}$ ,  $\sigma^2 \geq 0$ , and  $L$  satisfying the above conditions. Following Lindner and Sato [26], the corresponding distribution function  $F$  and probability law are called *quasi-infinitely divisible*. Let us fix any real  $\gamma_1$  and  $\gamma_2$  such that  $\gamma = \gamma_1 - \gamma_2$ . Let us fix any nonnegative  $\sigma_1^2$  and  $\sigma_2^2$  such that  $\sigma_1^2 \geq \sigma_2^2$  and  $\sigma^2 = \sigma_1^2 - \sigma_2^2$ . Due to the Hahn–Jordan decomposition, it is not difficult to show that there exist some canonical Lévy spectral functions  $L_1$  and  $L_2$  satisfying the usual conditions with monotonicity such that  $L = L_1 - L_2$ . Then  $f(t) = f_1(t)/f_2(t)$ ,  $t \in \mathbb{R}$ , where  $f_1$  and  $f_2$  are represented by the canonical Lévy formula (1) with characteristic triplets  $(\gamma_1, \sigma_1^2, L_1)$  and  $(\gamma_2, \sigma_2^2, L_2)$ . So  $f_1$  and  $f_2$  are the characteristic functions for some infinitely divisible distribution functions and hence the distribution function  $F$  corresponding to such an  $f$  is rational-infinately divisible. Thus  $F \in \mathcal{Q}$  if and only if  $f$  admits representation (1) with some  $(\gamma, \sigma^2, L)$  satisfying the above conditions. Additionally, *the characteristic triplet*  $(\gamma, \sigma^2, L)$  is uniquely determined by  $f$  and hence by  $F$  as for infinitely divisible laws (this can be concluded from the assertion in [12, p. 80]). It is also clear that for any rational-infinately divisible  $F$  its characteristic function  $f$  has no zeroes on the real line, i.e.,  $f(t) \neq 0$ ,  $t \in \mathbb{R}$ .

The class  $\mathcal{Q}$  and its multivariate analog are objects of active study (see [2; 8; 26]) and they find interesting applications in probability limit and compactness theorems (see Sections 4 and 8 in [26], Section 3 in [3], the paper [19], and also [1; 17]), and in other areas (see, for instance, [10; 29; 30]). But, actually, nondegenerate representatives of  $\mathcal{Q} \setminus \mathcal{I}$  appeared even earlier in the theory of decompositions of probability laws as components of certain infinitely divisible distribution functions (see [12, pp. 81–83; 27, p. 165]).

The class  $\mathcal{Q}$  is seen to be rather wide. For instance, it contains the distribution function of every probability law that has a mass  $> 1/2$  at some point. Hence the class  $\mathcal{Q}$  contains nondegenerate distribution functions of some probability laws with bounded supports (see examples in [26]), which are “far” from the infinite divisibility property in a known sense (see [5]). So it is interesting and important to obtain criteria for belonging to the class  $\mathcal{Q}$ . The existing results in this direction usually have simple and nice formulations in terms of characteristic functions. The first quite general result of such type was obtained by Lindner, Pan, and Sato in [26] (see Theorem 8.1, p. 30). It states that a lattice distribution function  $F$  belongs to the class  $\mathcal{Q}$  if and only if its characteristic function  $f$  does not have zeroes on the real line, i.e.,  $f(t) \neq 0$  for any  $t \in \mathbb{R}$ .

In [3] and [18], this result was generalized to the class of arbitrary discrete probability laws. Namely, a discrete distribution function  $F$  belongs to  $\mathcal{Q}$  if and only if its characteristic function  $f$  is separated from zero, i.e.,  $|f(t)| \geq \mu$  for any  $t \in \mathbb{R}$  and for some constant  $\mu > 0$ . Moreover, in that case, the components of the characteristic triplet are fully described. This result is a generalization of the previous one for discrete lattice distributions, because the absolute value  $|f(\cdot)|$  of the characteristic function  $f$  of a discrete lattice distribution is a periodic continuous function on  $\mathbb{R}$ . Therefore such  $f$  is zero-free on the period segment (and hence on  $\mathbb{R}$ ) if and only if it is separated from zero.

In [6], Berger considered mixtures of a degenerate law (with a nonzero coefficient) and absolutely continuous distributions. According to his result, a distribution function  $F$  of such type belongs to  $\mathcal{Q}$  if and only if  $f(t) \neq 0$ ,  $t \in \mathbb{R}$ . Moreover, the result describes the structure of the components of the characteristic triplet in that case. The author also formulated more general criterion for the case, when the degenerate law from the previous one is replaced by a discrete lattice distribution with characteristic function, which has no zeroes on the real line. At present, however, the most general criterion (among those that use assumptions about the type of distribution) is the following result for the mixtures of discrete and absolutely continuous probability laws, which was obtained by Berger and Kutlu in the paper [7]. Let us formulate it with more details here. Namely, assume that  $F(x) = c_d F_d(x) + c_a F_a(x)$ ,  $x \in \mathbb{R}$ , where  $F_d$  is a discrete distribution function,  $F_a$  is an absolutely continuous distribution function,  $c_d > 0$ ,  $c_a \geq 0$ , and  $c_d + c_a = 1$ . We write the characteristic function  $f$  in the corresponding form:  $f(t) = c_d f_d(t) + c_a f_a(t)$ ,  $t \in \mathbb{R}$ . Then  $F \in \mathcal{Q}$  if and only if  $f(t) \neq 0$  and  $|f_d(t)| \geq \mu$  for any  $t \in \mathbb{R}$  with some constant  $\mu > 0$ . It is equivalent to the condition  $|f(t)| \geq \mu'$  for any  $t \in \mathbb{R}$  and for some constant  $\mu' > 0$ . Moreover, Berger and Kutlu showed the existence of some discrete part in the spectral function and they fully described its absolutely continuous part for this case. It should be noted that we are not aware of any similar results for purely absolutely continuous distribution functions  $F$ . However, for some cases the problem of membership in class  $\mathcal{Q}$  for a given distribution function of such type is not difficult to solve by the general criteria proposed in [20] with some additional analysis.

This article is devoted to generalizing and complementing all the mentioned results (except [20]) for the case, when  $F$  may have a continuous singular part. Namely, we propose a criterion for a distribution function  $F$  to belong to class  $\mathcal{Q}$  under the assumption that its continuous singular part is dominated by its discrete part in a certain sense. In fact, we show that the conditions on  $f$  from the results [6] and [7] are carried over to this case. In addition, we describe the characteristic triplet of the corresponding Lévy-type representation, which may contain some continuous singular part.

We next show that the assumption of a dominated continuous singular part cannot be omitted or even slightly extended without some additional assumptions. In addition, for any  $F \in \mathcal{Q}$  we solve the decomposition problem, which was stated by Lindner, Pan, Sato in [26] (see Open Question 8.4), within the considered case. Here we obtain a positive solution generalizing similar results from [6] and [7].

The article has the following structure. Section 2 contains necessary preliminaries, more detailed statements of some preexisting results mentioned above and the formulations of the new results of the paper. In Section 3, we formulate some important known theorems and useful lemmata, which will be auxiliary tools needed for the proofs of our results. In Section 4, we first prove a key auxiliary lemma and we next propose the proofs of the main results of the article.

Throughout the paper, we use the following notation. We denote by  $\mathbb{N}$  the set of positive integers and let  $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$ . The symbols  $\mathbb{Z}$  and  $\mathbb{Q}$  denote as usual the sets of all integers and rational numbers, respectively. Next,  $\mathbb{C}$  is the set of all complex numbers. For any  $z \in \mathbb{C}$  we denote by  $\text{Im}\{z\}$  and  $\text{arg}(z)$  its imaginary part and the principal value of the argument of  $z$ , respectively. If  $\psi$  is a complex-valued continuous function on  $\mathbb{R}$  satisfying  $\psi(0) = c \in \mathbb{C} \setminus \{0\}$  and  $\psi(t) \neq 0$  for any  $t \in \mathbb{R}$ , then the *distinguished logarithm*  $\text{Ln } \psi$  is defined by the formula  $\text{Ln } \psi(t) := \ln |\psi(t)| + i \text{Arg } \psi(t)$ ,  $t \in \mathbb{R}$ , where  $\text{Arg } \psi(t)$  is the argument of  $\psi(t)$  uniquely defined on  $\mathbb{R}$  by the continuity with the condition  $\text{Arg } \psi(0) = \text{arg}(c) \in (-\pi, \pi]$ . The symbol  $\mathbb{1}_a$  with fixed  $a \in \mathbb{R}$  denotes the distribution function of the degenerate law concentrated at the point  $a$ , i.e.,  $\mathbb{1}_a(x) = 1$  for  $x \geq a$  and  $\mathbb{1}_a(x) = 0$  for  $x < a$ . For any set  $A$  we denote by  $\mathbb{1}_A$  the indicator function of  $A$ , i.e.,  $\mathbb{1}_A(x) = 1$  for any  $x \in A$  and  $\mathbb{1}_A(x) = 0$  for any  $x \notin A$ . The signum function is denoted by  $\text{sgn}(\cdot)$ , i.e.,  $\text{sgn}(x) = +1$  for  $x > 0$ ,  $\text{sgn}(x) = -1$  for  $x < 0$ , and  $\text{sgn}(0) = 0$ . For any finite set  $A$  the symbol  $|A|$  denotes the number of elements of  $A$ . We always set  $\sum_{k \in A} a_k = 0$  and  $\prod_{k \in A} a_k = 1$  in the case  $A = \emptyset$ . For any two vectors  $x$  and  $y$  from  $\mathbb{R}^n$  the standard scalar product is denoted by  $\langle x, y \rangle$ .

For any function  $G$  defined on  $\mathbb{R}$  the limits  $\lim_{x \rightarrow \pm\infty} G(x)$  are denoted by  $G(\pm\infty)$ , respectively, if these limits exist. The class of all functions  $G : \mathbb{R} \rightarrow \mathbb{R}$  of bounded total variation on  $\mathbb{R}$  (nonmonotonic in general), which are right-continuous at every point and  $G(-\infty) = 0$ , is denoted by  $\mathbf{V}$ . We denote by  $\mathbf{V}_{\mathbb{C}}$  the set of all functions  $G : \mathbb{R} \rightarrow \mathbb{C}$  of the form  $G(x) = G_1(x) + iG_2(x)$ ,  $x \in \mathbb{R}$ , where  $G_1$  and  $G_2$  are from  $\mathbf{V}$ . For every  $G \in \mathbf{V}$  (or  $\mathbf{V}_{\mathbb{C}}$ ) its total variation on  $\mathbb{R}$  will be denoted by  $\|G\|$  and the total variation on  $(-\infty, x]$  by  $|G|(x)$ ,  $x \in \mathbb{R}$ . So we have  $|G(x)| \leq |G|(x) \leq \|G\|$ ,  $x \in \mathbb{R}$ , and  $|G|(+\infty) = \|G\|$ . Next, we adopt the following convention. Let  $G$  be a function from  $\mathbf{V}$  with the Fourier–Stieltjes transform  $g$ , i.e.,  $g(t) = \int_{\mathbb{R}} e^{itx} dG(x)$ ,  $t \in \mathbb{R}$ . In view of the uniqueness theorem for functions from  $\mathbf{V}$ , we set  $\|g\| := \|G\|$ . So  $\|g\| = 0$  if and only if  $g(t) = 0$ ,  $t \in \mathbb{R}$ , and  $\|c \cdot g\| = |c| \cdot \|g\|$  for any  $c \in \mathbb{R}$ . Let  $G_1$  and  $G_2$  be functions from  $\mathbf{V}$  with the

Fourier–Stieltjes transforms  $g_1$  and  $g_2$ , correspondingly. The known inequalities  $\|G_1 + G_2\| \leq \|G_1\| + \|G_2\|$  and  $\|G_1 * G_2\| \leq \|G_1\| \cdot \|G_2\|$  are correspondingly written as  $\|g_1 + g_2\| \leq \|g_1\| + \|g_2\|$  and  $\|g_1 \cdot g_2\| \leq \|g_1\| \cdot \|g_2\|$ . We recall that both  $V$  and the corresponding space of functions  $g$  with norm  $\|\cdot\|$  will be complete normed spaces (see [11, p. 165]).

## 2. Criteria for belonging to class $\mathcal{Q}$

Let  $F$  be an arbitrary distribution function on the real line. According to the Lebesgue decomposition theorem,  $F$  admits the representation

$$(3) \quad F(x) = c_d F_d(x) + c_a F_a(x) + c_s F_s(x), \quad x \in \mathbb{R},$$

where  $F_d$ ,  $F_a$ , and  $F_s$  are discrete, absolutely continuous and continuous singular distribution functions, respectively. Here the coefficients  $c_d$ ,  $c_a$ , and  $c_s$  are nonnegative constants such that  $c_d + c_a + c_s = 1$ . Let  $f$  be the characteristic function of  $F$ . It is represented in a similar way:

$$(4) \quad f(t) = c_d f_d(t) + c_a f_a(t) + c_s f_s(t), \quad t \in \mathbb{R},$$

where  $f_d$ ,  $f_a$ , and  $f_s$  are the characteristic functions corresponding to  $F_d$ ,  $F_a$ , and  $F_s$ , respectively. It is well known that the summands in (3) or (4) are uniquely determined. So if any of the terms is not identically zero, then the corresponding coefficient, distribution function, and characteristic function are uniquely determined.

We will consider only the case that  $F$  has nonzero discrete part, i.e.,  $c_d > 0$  in (3). We write the distribution function  $F_d$  in the form

$$(5) \quad F_d(x) = \sum_{\substack{k \in \mathbb{N}_0 \\ x_k \leq x}} p_k, \quad x \in \mathbb{R},$$

where  $x_k$  are distinct reals associated with weights  $p_k \geq 0$ ,  $k \in \mathbb{N}_0$ ,  $\sum_{k=0}^{\infty} p_k = 1$ . Hence  $f_d$  has the form

$$(6) \quad f_d(t) = \sum_{k \in \mathbb{N}_0} p_k e^{itx_k}, \quad t \in \mathbb{R}.$$

We define the carrier of the distribution corresponding to  $F_d$ :

$$\mathcal{X} := \{x_k : p_k > 0, k \in \mathbb{N}_0\}.$$

Obviously,  $\mathcal{X} \neq \emptyset$ . We also need the set of all finite  $\mathbb{Z}$ -linear combinations of elements from the set  $\mathcal{X}$ :

$$\langle \mathcal{X} \rangle := \left\{ \sum_{k=1}^m a_k z_k : a_k \in \mathbb{Z}, z_k \in \mathcal{X}, m \in \mathbb{N} \right\}.$$

So  $\langle \mathcal{X} \rangle$  is the module over the ring  $\mathbb{Z}$  with the generating set  $\mathcal{X}$ . It easily seen that, in particular,  $\mathcal{X} \subset \langle \mathcal{X} \rangle$  and  $0 \in \langle \mathcal{X} \rangle$ . If  $\mathcal{X} \neq \{0\}$ , then  $\langle \mathcal{X} \rangle$  is an infinite countable set.

Now we are ready to formulate in detail the most general existing results on criteria for belonging to class  $\mathcal{Q}$ . We start with the result obtained in [3] and [18] by Alexeev and Khartov for the case of discrete  $F$ .

**Theorem 1.** *Suppose that  $F$  is a discrete distribution function,  $c_d = 1$  and  $c_a = c_s = 0$  in (3) ( $F$  and  $F_d$  are identical and hence  $f$  and  $f_d$  are too;  $F$  has the form (5),  $f$  is represented by (6)). Then  $F \in \mathcal{Q}$  if and only if  $\inf_{t \in \mathbb{R}} |f(t)| > 0$ . In that case,  $f$  admits the representation*

$$(7) \quad f(t) = \exp \left\{ it\gamma_0 + \sum_{u \in \langle \mathcal{X} \rangle \setminus \{0\}} \lambda_u (e^{itu} - 1) \right\}, \quad t \in \mathbb{R},$$

with some  $\gamma_0 \in \langle \mathcal{X} \rangle$  and  $\lambda_u \in \mathbb{R}$  for all  $u \in \langle \mathcal{X} \rangle \setminus \{0\}$ , and  $\sum_{u \in \langle \mathcal{X} \rangle \setminus \{0\}} |\lambda_u| < \infty$ .

It is not difficult to rewrite the representation (7) in integral form (1) with some  $\gamma \in \mathbb{R}$ ,  $\sigma^2 = 0$ , and some discrete  $L$ , which will satisfy all the conditions for a spectral function for the quasi-infinitely divisibility. We will do this below for a more general case. Hence, if the characteristic function  $f$  is represented by (7), then  $F \in \mathcal{Q}$ .

It is clear that if  $F \in \mathcal{Q}$  and  $\lambda_u \geq 0$  for all  $u \in \langle \mathcal{X} \rangle \setminus \{0\}$  in representation (7), then  $F$  is infinitely divisible, i.e.,  $F \in \mathcal{I}$ . If there exists  $\lambda_v < 0$  with some  $v \in \langle \mathcal{X} \rangle \setminus \{0\}$ , then  $F \in \mathcal{Q} \setminus \mathcal{I}$ , because the (uniquely defined) function  $L$  will be decreasing in the neighborhood of  $v$ . For examples of the latter case see [26, p. 10] and [27, p. 165].

We now formulate Berger and Kutlu's result in [7] for the important case  $c_a \geq 0$ . As mentioned, at present, this is the most general criterion using information about the type of the distribution function  $F$ .

**Theorem 2.** *Suppose that  $F$  has the decomposition (3) with some  $c_d > 0$ ,  $c_a \geq 0$ , and  $c_s = 0$ . Then the following statements are equivalent:*

- (i)  $F \in \mathcal{Q}$ .
- (ii)  $\inf_{t \in \mathbb{R}} |f(t)| > 0$ .
- (iii)  $f(t) \neq 0$  for any  $t \in \mathbb{R}$ , and  $\inf_{t \in \mathbb{R}} |f_d(t)| > 0$ .

If one (hence all) of the conditions is satisfied, then  $f$  admits the representation

$$(8) \quad f(t) = \exp \left\{ it\gamma_0 + \sum_{u \in \mathcal{U}} \lambda_u (e^{itu} - 1) + \int_{\mathbb{R}} (e^{itx} - 1) \left( v(x) + \operatorname{sgn}(x) \frac{\mathbf{m} \cdot e^{-|x|}}{|x|} \right) dx \right\},$$

$t \in \mathbb{R},$

where  $\gamma_0 \in \mathbb{R}$ ,  $\mathbf{m} \in \mathbb{Z}$ ,  $\mathcal{U}$  is a discrete set,  $\lambda_u \in \mathbb{R}$  for all  $u \in \mathcal{U}$ , with  $\sum_{u \in \mathcal{U}} |\lambda_u| < \infty$ , and  $v$  is a real-valued function from  $L_1(\mathbb{R})$ .

In addition to this theorem, the paper [7] contains a number of interesting assertions concerning decomposition of  $F$  and characterization of the existence of the  $H$ -moment of  $F$  for certain positive functions  $H$ . Additionally, there is a rather general theorem similar to Theorem 2 for the function  $f$ , but without the assumption that it is the characteristic function of some probability law (Theorem 2.1 in [7]).

Theorem 2 generalizes the results by Berger ([6, Theorems 4.5 and 4.12]) mentioned in the introduction, where  $F_d$  was assumed to be a discrete lattice. It is also seen that, in fact, Theorem 2 is a strengthening of Theorem 1. However, we note that  $\gamma_0$  and the discrete part in (7) are described in greater detail than in formula (8).

We now propose a generalization of the previous criteria for the case when  $F$  may have a continuous singular part, i.e., when  $c_s \geq 0$  in (3). The following theorem is the main result of this article.

For convenience, we preliminarily select the following property of distributions. Let us define  $\mu_d := \inf_{t \in \mathbb{R}} |f_d(t)|$ . We say that a distribution function  $F$  has a *dominated continuous singular part* if  $c_s < c_d \mu_d$  for the case  $\mu_d > 0$  and if  $c_s = 0$  for the case  $\mu_d = 0$ .

**Theorem 3.** *Suppose that  $F$  has a decomposition (3) with  $c_d > 0$ ,  $c_a \geq 0$ ,  $c_s \geq 0$ , and that  $F$  has a dominated continuous singular part. Then the following statements are equivalent:*

- (i)  $F \in \mathcal{Q}$ .
- (ii)  $\inf_{t \in \mathbb{R}} |f(t)| > 0$ .
- (iii)  $f(t) \neq 0$  for any  $t \in \mathbb{R}$ , and  $\inf_{t \in \mathbb{R}} |f_d(t)| > 0$ .

If one (hence all) of the conditions is satisfied,  $f$  admits the representation

$$(9) \quad f(t) = \exp \left\{ it\gamma_0 + \sum_{u \in \langle \mathcal{X} \rangle \setminus \{0\}} \lambda_u (e^{itu} - 1) + \int_{\mathbb{R} \setminus \{0\}} (e^{itx} - 1) \left( v_a(x) + \operatorname{sgn}(x) \frac{\mathfrak{m}_a \cdot e^{-|x|}}{|x|} \right) dx + \int_{\mathbb{R} \setminus \{0\}} (e^{itx} - 1) dW(x) \right\}, \quad t \in \mathbb{R}.$$

Here  $\gamma_0 \in \langle \mathcal{X} \rangle$ ,  $\lambda_u \in \mathbb{R}$  for all  $u \in \langle \mathcal{X} \rangle \setminus \{0\}$ , and  $\sum_{u \in \langle \mathcal{X} \rangle \setminus \{0\}} |\lambda_u| < \infty$ . Next, the function  $v_a : \mathbb{R} \mapsto \mathbb{R}$  satisfies  $\int_{\mathbb{R}} |v_a(x)| dx < \infty$ , and, in the case  $c_a = 0$ ,  $v_a$  is identically 0; the constant  $\mathfrak{m}_a \in \mathbb{Z}$  is defined by the formula

$$(10) \quad \mathfrak{m}_a := \frac{1}{2\pi} \left( \lim_{t \rightarrow \infty} \operatorname{Arg} R_a(t) - \lim_{t \rightarrow -\infty} \operatorname{Arg} R_a(t) \right),$$

with

$$R_a(t) := 1 + \frac{c_a f_a(t)}{c_d f_d(t) + c_s f_s(t)}, \quad t \in \mathbb{R},$$

where, in particular,  $m_a = 0$  for the case  $c_a = 0$ . Next, the function  $W : \mathbb{R} \rightarrow \mathbb{R}$  belongs to  $\mathcal{V}$  and it is always continuous on  $\mathbb{R}$ . If  $c_s = 0$  then  $W$  is identically 0. If  $c_s \neq 0$  then  $W$  is not absolutely continuous on  $\mathbb{R}$ , i.e., it always contains some continuous singular part. In addition, if all the functions  $F_s^{*k}, k \in \mathbb{N}$ , are continuous singular, then the function  $W$  is (purely) continuous singular.

Note that the discrete part in the exponent in (9) depends only on discrete part of  $f$ , i.e., on  $f_d$ . More precisely, we have the following remark, which will be seen from the proof of Theorem 3.

**Remark 1.** In the representation (9),

$$\text{Ln } f_d(t) = it\gamma_0 + \sum_{u \in \langle \mathcal{X} \rangle \setminus \{0\}} \lambda_u (e^{itu} - 1), \quad t \in \mathbb{R}.$$

It is easily seen that if we set  $c_a := 0$  and  $c_s := 0$  in the assumptions of this theorem, then we come to Theorem 1. If we set only  $c_s := 0$ , then we conclude the statement of Theorem 2, but with a full description of the discrete part in (8). In the unexplored case  $c_s > 0$ , the new function  $W$  appears and it always has some continuous singular part.

Using Theorem 3, it is easy to construct a lot of particular examples of  $F \in \mathcal{Q}$  with nonzero continuous singular parts.

**Example 1.** Suppose that

$$F(x) := c_d \mathbb{1}_0(x) + c_a F_a(x) + c_s F_s(x), \quad x \in \mathbb{R},$$

with  $c_d > c_s > 0, c_a \geq 0$ , and  $c_d + c_a + c_s = 1$ . Let  $F_s$  be an arbitrary continuous singular function, but  $F_a$  be an absolutely continuous distribution function, whose characteristic function  $f_a$  is real and nonnegative (for instance,  $f_a$  is a Pólya-type characteristic function, or  $f_a$  corresponds to a symmetric continuous stable distribution). Then  $F \in \mathcal{Q}$ . Indeed,  $F$  has dominated continuous singular part, because  $f_d$  is identically 1 here, i.e.,  $\mu_d = 1$ , and hence  $c_d \mu_d = c_d > c_s$ . Let us check (ii) from Theorem 3. For this, we consider the characteristic function of  $F$ ,

$$f(t) = c_d + c_a f_a(t) + c_s f_s(t), \quad t \in \mathbb{R}.$$

Under the assumptions, we observe that

$$|f(t)| \geq |c_d + c_a f_a(t)| - c_s |f_s(t)| = c_d + c_a f_a(t) - c_s |f_s(t)|, \quad t \in \mathbb{R}.$$

Since  $f_a(t) \geq 0$  and  $|f_s(t)| \leq 1$  for any  $t \in \mathbb{R}$ , we have  $|f(t)| \geq c_d - c_s > 0$  for any  $t \in \mathbb{R}$ , i.e., (ii) holds. Thus  $F \in \mathcal{Q}$  by Theorem 3.

In the general case, of course, checking (ii) or (iii) of Theorem 3 may require more subtle analysis. We illustrate this with the following special example, which is interesting also because there is a (purely) continuous singular function  $W$  in (9).

**Example 2.** Let us consider the mixture of the degenerate law, the uniform distribution on  $[0, 1]$  and the classical Cantor distribution. Namely, we set

$$F(x) := c_d \mathbb{1}_0(x) + c_a U(x) + c_s S(x), \quad x \in \mathbb{R},$$

where  $U$  is an absolutely continuous distribution function with density  $\mathbb{1}_{[0,1]}$ , and  $S$  is the cumulative function of the classical Cantor distribution supported on the Cantor set  $\mathcal{C} \subset [0, 1]$ ;  $c_d > c_s > 0$ ,  $c_a \geq 0$ , and  $c_d + c_a + c_s = 1$ .

We first observe that  $f_d$  is identically 1 here and, as in Example 1,  $F$  has dominated continuous singular part. We next check condition (iii) of Theorem 3. Since  $\inf_{t \in \mathbb{R}} |f_d(t)| = 1$ , it remains to show that  $f(t) \neq 0$  for any  $t \in \mathbb{R}$ . The function  $f$  is expressed by the formula

$$f(t) = c_d + c_a \frac{e^{it} - 1}{it} + c_s e^{it/2} \prod_{k=1}^{\infty} \cos(t/3^k), \quad t \in \mathbb{R}.$$

We set

$$\hat{f}(t) := f(t)e^{-it/2} = c_d e^{-it/2} + c_a \frac{\sin(t/2)}{t/2} + c_s \prod_{k=1}^{\infty} \cos(t/3^k), \quad t \in \mathbb{R}.$$

The functions  $f$  and  $\hat{f}$  have the same set of zeroes on  $\mathbb{R}$ , a subset of the set of zeroes of the function  $\text{Im}\{\hat{f}(t)\} = -c_d \sin(t/2)$ ,  $t \in \mathbb{R}$ . The last set is exactly  $\{2\pi m : m \in \mathbb{Z}\}$ . Therefore it is sufficient to show that  $f(2\pi m) \neq 0$  for any  $m \in \mathbb{Z}$ . Obviously, we can exclude  $m = 0$ . For any  $m \in \mathbb{Z} \setminus \{0\}$  we write

$$\begin{aligned} |f(2\pi m)| &= |\hat{f}(2\pi m)| = \left| c_d \cos(\pi m) + c_a \frac{\sin(\pi m)}{\pi m} + c_s \prod_{k=1}^{\infty} \cos(2\pi m/3^k) \right| \\ &= \left| c_d (-1)^m + c_s \prod_{k=1}^{\infty} \cos(2\pi m/3^k) \right|. \end{aligned}$$

It follows that

$$|f(2\pi m)| \geq c_d - c_s \prod_{k=1}^{\infty} |\cos(2\pi m/3^k)| \geq c_d - c_s > 0.$$

Thus  $f(t) \neq 0$  for any  $t \in \mathbb{R}$  and condition (iii) of Theorem 3 is satisfied. So, by the theorem,  $F \in \mathcal{Q}$ .

Let us consider the representation (9) for  $f$ . Since  $f_d(t) = 1$  for any  $t \in \mathbb{R}$ , the sum

$$it\gamma_0 + \sum_u \lambda_u (e^{itu} - 1)$$

is identically 0 according to Remark 1. There are the function  $v_a \in L_1(\mathbb{R})$  and the constant  $m_a \in \mathbb{Z}$ , but we will not determine these here. Next, we turn to the function

$W$ . It is known that all convolution powers  $S^{*n}$ ,  $n \in \mathbb{N}$ , of the Cantor distribution function  $S$  are continuous singular; this follows from the famous Jessen–Wintner theorem (see [16], Theorem 35) and the representation of  $S(x + \frac{1}{2})$ ,  $x \in \mathbb{R}$ , as the infinite symmetric Bernoulli convolution, and it was also explicitly shown in [36] (see p. 520). According to Theorem 3, this implies that  $W$  is purely continuous singular. Moreover, it will be seen from the proof of Theorem 3 that  $W$  is the limit of the sums

$$W_n(x) := \sum_{k=1}^n \frac{(-1)^{k-1}}{k} \left(\frac{c_s}{c_d}\right)^k S^{*k}(x), \quad x \in \mathbb{R}, \quad n \in \mathbb{N},$$

as  $n \rightarrow \infty$  with respect to convergence in total variation on  $\mathbb{R}$ .

So Theorem 3 yields a sufficient condition for  $W$  to be (purely) continuous singular. Is it true that  $W$  is always a (purely) continuous singular function in the case  $c_s \neq 0$ ?

**Proposition 1.** *Suppose that  $F \in \mathcal{Q}$  satisfies the assumptions of Theorem 3 with  $c_s > 0$  and the characteristic function  $f$  is represented by (9) with some  $W$ . Suppose there is an integer  $n_a \geq 2$  such that the function  $F_s^{*(n_a-1)}$  is (purely) continuous singular, but the function  $F_s^{*n_a}$  is not, i.e.,  $F_s^{*n_a}(x) = \alpha H_a(x) + (1-\alpha)H_s(x)$ ,  $x \in \mathbb{R}$ , where  $\alpha$  is a number in  $(0, 1]$ ,  $H_a$  is an absolutely continuous distribution function, and  $H_s$  is a continuous singular distribution function. If*

$$\alpha \geq \frac{n_a}{n_a+1} \cdot \frac{c_s}{c_d}$$

*then the function  $W$  is not (purely) continuous singular. In particular, this is always true for the case  $\alpha = 1$ .*

It should be recalled here that there exist examples of continuous singular distribution functions (say of  $F_s$ ), whose convolution squares (say  $F_s^{*2}$ ) are absolutely continuous (see [14; 15; 36]). Therefore we answer the question asked before Proposition 1 in the negative.

The following remark yields the characteristic triplet for  $F \in \mathcal{Q}$  in the explicit form under the assumptions of Theorem 3.

**Remark 2.** The representation (9) can be written in the form

$$\begin{aligned} f(t) &= \exp \left\{ it\gamma_0 + \int_{\mathbb{R} \setminus \{0\}} (e^{itx} - 1) dL(x) \right\} \\ &= \exp \left\{ it\gamma + \int_{\mathbb{R} \setminus \{0\}} (e^{itx} - 1 - it \sin(x)) dL(x) \right\}, \quad t \in \mathbb{R}, \end{aligned}$$

where

$$\gamma := \gamma_0 + \int_{\mathbb{R} \setminus \{0\}} \sin(x) dL(x)$$

and  $L(x) := L_d(x) + L_a(x) + L_s(x)$  for any  $x \in \mathbb{R} \setminus \{0\}$  with

$$(11) \quad L_d(x) := \begin{cases} \sum_{\substack{u \in \langle \mathcal{X} \rangle \setminus \{0\} \\ u \leq x}} \lambda_u & \text{if } x < 0, \\ - \sum_{\substack{u \in \langle \mathcal{X} \rangle \setminus \{0\} \\ u > x}} \lambda_u & \text{if } x > 0, \end{cases}$$

$$(12) \quad L_a(x) := \begin{cases} \int_{u \leq x} \left( v_a(u) + \operatorname{sgn}(u) \frac{m_a \cdot e^{-|u|}}{|u|} \right) du & \text{if } x < 0, \\ - \int_{u > x} \left( v_a(u) + \operatorname{sgn}(u) \frac{m_a \cdot e^{-|u|}}{|u|} \right) du & \text{if } x > 0, \end{cases}$$

$$(13) \quad L_s(x) := \begin{cases} W(x) & \text{if } x < 0, \\ W(x) - W(+\infty) & \text{if } x > 0. \end{cases}$$

It is seen that  $L$  satisfies all admissible conditions for a spectral function in the Lévy type representation. Thus, under the assumptions of Theorem 3,  $(\gamma, 0, L)$  is the characteristic triplet for  $F$  satisfying one of the conditions (i)–(iii). On the other hand, if we know that (9) represents the characteristic function of some probability law, then its distribution function  $F$  is quasi-infinitely divisible by the definition and hence  $F \in \mathcal{Q}$ .

The following remark shows that the property of dominated singular part is not a necessary condition for belonging to class  $\mathcal{Q}$ .

**Remark 3.** Let  $F$  satisfy the assumptions of Theorem 3 with  $c_s > 0$ . Suppose that  $F \in \mathcal{Q}$ . Then  $F^{*n} \in \mathcal{Q}$  for any  $n \in \mathbb{N}$ , but  $F^{*n}$  does not have dominated singular part for all sufficiently large  $n$ .

Indeed, it is seen from the definition of the rational-infinite divisibility that the convolution of two any distribution functions from  $\mathcal{Q}$  belongs to  $\mathcal{Q}$ . Therefore  $F^{*n} \in \mathcal{Q}$  for any  $n \in \mathbb{N}$ . Next, we consider the characteristic function  $f$  of  $F$ . It admits the decomposition (4) with  $c_d > 0$ ,  $c_a \geq 0$ ,  $c_s > 0$ , and  $c_s < c_d \mu_d$ , where  $\mu_d = \inf_{t \in \mathbb{R}} |f_d(t)|$  as before. Then every  $F^{*n}$  has the characteristic function

$$f(t)^n = (c_d f_d(t) + c_a f_a(t) + c_s f_s(t))^n, \quad t \in \mathbb{R}.$$

From this it is easily seen that  $t \mapsto c_d^n f_d(t)^n$ ,  $t \in \mathbb{R}$ , is the Fourier–Stieltjes transform of the discrete part of  $F^{*n}$ . Hence  $c_d^n$  is the weight of this part and  $\inf_{t \in \mathbb{R}} |f_d(t)^n| = \mu_d^n$ . Next, there are  $n$  terms  $c_s c_d^{n-1} f_s(t) f_d(t)^{n-1}$  that appear in the Fourier–Stieltjes transform of the continuous singular part of  $F^{*n}$ . This is because a convolution of continuous singular function with a discrete function is continuous singular (see [28, p. 190] or [34, p. 319]). So the weight of the continuous singular part of  $F^{*n}$  is not less than  $nc_s c_d^{n-1}$ . For any integer  $n \geq c_d/c_s$  we have  $nc_s c_d^{n-1} \geq c_d^n \geq c_d^n \mu_d^n$ . Therefore the exact weight of the continuous singular part of  $F^{*n}$  is not less than

$c_d^n \mu_d^n$  too; i.e., the condition of dominated singular part doesn't hold for  $F^{*n}$  for  $n \geq c_d/c_s$ .

The following interesting example shows that the condition of a dominated singular part cannot be simply omitted, or even extended to the case  $c_s = c_d \mu_d$  with  $\mu_d > 0$  without certain additional assumptions.

**Example 3.** Let  $B$  denote the distribution function of the Bernoulli law on the points  $\pm 1$  with equal probabilities, i.e.,  $B(x) = \frac{1}{2} \mathbb{1}_{-1}(x) + \frac{1}{2} \mathbb{1}_1(x)$ ,  $x \in \mathbb{R}$ . We set

$$(14) \quad F_*(x) := \frac{1}{2} \mathbb{1}_0(x) + \frac{1}{2} F_s(x), \quad x \in \mathbb{R},$$

where  $F_s$  is the following infinite symmetric Bernoulli convolution

$$F_s = B_1 * B_2 * \dots * B_n * \dots$$

with  $B_k(x) := B(k!x)$ ,  $x \in \mathbb{R}$ ,  $k \in \mathbb{N}$ . Let  $f_*$  and  $f_s$  denote the characteristic functions of  $F_*$  and  $F_s$ . Then  $f_*(t) = \frac{1}{2} + \frac{1}{2} f_s(t)$ ,  $t \in \mathbb{R}$ , and

$$(15) \quad f_s(t) = \prod_{k=1}^{\infty} \cos(t/k!), \quad t \in \mathbb{R}.$$

It is known (see [28, pp. 20, 67]) that the function  $F_s$  is continuous singular. Let us consider Lebesgue's decomposition (4) for  $f = f_*$ . We have  $c_d = \frac{1}{2}$ ,  $c_a = 0$ , and  $c_s = \frac{1}{2}$ . Observe that the component  $f_d$  is identically 1 for  $f_*$  and hence  $\mu_d = \inf_{t \in \mathbb{R}} |f_d(t)| = 1$ . So  $c_d \mu_d = c_s$ , i.e., the condition of dominated continuous singular part, doesn't hold. Next,  $f_*(t) \neq 0$  for any  $t \in \mathbb{R}$ , since otherwise  $f_s(t_0) = -1$  for some  $t_0 \neq 0$  and hence  $|f_s(t_0)| = 1$ , which would mean that  $F_s$  is a discrete lattice distribution function (see [28, Theorem 2.1.4]), a contradiction. Thus  $F_*$  satisfies condition (iii) of Theorem 3 and the assumption  $c_d > 0$ , but  $F_*$  doesn't have dominated continuous singular part.

**Proposition 2.** *The function  $F_*$  doesn't belong to  $\mathcal{Q}$ .*

Thus, in general, conditions (i) and (iii) from Theorem 3 are not equivalent even if  $c_s = c_d \mu_d > 0$ .

By the way, we recall that distribution functions of discrete laws with a point mass  $\frac{1}{2}$ , which have characteristic functions without zeroes on the real line, don't always belong to class  $\mathcal{Q}$  (see [22] for more details). However, if the distribution is an atom of mass  $\frac{1}{2}$  plus a continuous part, which is not purely singular, then the answer is definite.

**Example 4.** Suppose that

$$F(x) := \frac{1}{2} \mathbb{1}_{\gamma_0}(x) + c_d F_a(x) + c_s F_s(x), \quad x \in \mathbb{R},$$

where  $F_a$  is an absolutely continuous distribution function,  $F_s$  is a continuous

singular distribution function,  $\gamma_0 \in \mathbb{R}$ ,  $c_a > 0$ ,  $c_s \geq 0$ , and  $c_a + c_s = \frac{1}{2}$ . Such a function  $F$  always belongs to class  $\mathcal{Q}$ . Let us check it. We consider the characteristic function of  $F$ :

$$f(t) = \frac{1}{2} e^{it\gamma_0} + c_a f_a(t) + c_s f_s(t), \quad t \in \mathbb{R}.$$

Here  $c_d = \frac{1}{2}$  and  $f_d(t) = e^{it\gamma_0}$ ,  $t \in \mathbb{R}$ . Hence  $\mu_d = \inf_{t \in \mathbb{R}} |f_d(t)| = 1$  and  $c_s = \frac{1}{2} - c_a < \frac{1}{2} = c_d \mu_d$ , i.e.,  $F$  has dominated continuous singular part. Next,  $f(t) \neq 0$  for any  $t \in \mathbb{R}$ . For otherwise  $2c_a f_a(t_0) + 2c_s f_s(t_0) = -e^{it_0\gamma_0}$  for some  $t_0 \neq 0$  and so  $|2c_a f_a(t_0) + 2c_s f_s(t_0)| = 1$ . This equality implies that the distribution function with the characteristic function  $2c_a f_a(\cdot) + 2c_s f_s(\cdot)$  is discrete lattice, but this is actually continuous by the assumptions, a contradiction.

So  $F$  satisfies condition (iii) and the other assumptions of Theorems 3. It follows that  $F \in \mathcal{Q}$ .

The next assertion solves the decomposition problem for any distribution function  $F \in \mathcal{Q}$  satisfying the assumptions of Theorem 3.

**Proposition 3.** *Let  $F$  be a distribution function with decomposition (3) with some  $c_d > 0$ ,  $c_a \geq 0$ ,  $c_s \geq 0$ , and a dominated continuous singular part. Suppose that  $F \in \mathcal{Q}$  and  $F = F_1 * F_2$  with some distribution functions  $F_1$  and  $F_2$ . Then  $F_1$  and  $F_2$  belong to  $\mathcal{Q}$ .*

This proposition generalizes a similar result from [7] (see Corollary 2.3 to Theorem 2.2), which solves the problem for  $F \in \mathcal{Q}$  satisfying the assumptions of Theorem 2. We note that there is a general decomposition problem for arbitrary  $F \in \mathcal{Q}$ , which was stated by Lindner, Pan and Sato in [26] (see Open Question 8.4): *Is it true that if  $F \in \mathcal{Q}$  and  $F = F_1 * F_2$  ( $F_1$  and  $F_2$  being distribution functions on  $\mathbb{R}$ ), then  $F_1 \in \mathcal{Q}$  and  $F_2 \in \mathcal{Q}$ ?* So Proposition 3 answers this question in the affirmative for any  $F \in \mathcal{Q}$  satisfying the assumptions of Theorem 3. However, there is a result in [21] that asserts that the general answer is negative.

### 3. Auxiliary theorems and lemmata

The proof of the main result (Theorem 3) of the article essentially uses the following Wiener–Pitt theorem [35] (see also [33; 11, p. 191]).

**Theorem 4.** *Let  $H$  be a function in  $V_{\mathbb{C}}$  with Lebesgue decomposition*

$$H(x) = H_d(x) + H_a(x) + H_s(x), \quad x \in \mathbb{R},$$

where  $H_d$ ,  $H_a$ ,  $H_s$  are the discrete, absolutely continuous and singular parts of  $H$ , each of which belongs to  $V_{\mathbb{C}}$ . Let

$$h(t) := \int_{\mathbb{R}} e^{itx} dH(x), \quad \text{and} \quad h_d(t) := \int_{\mathbb{R}} e^{itx} dH_d(x), \quad t \in \mathbb{R}.$$

Suppose that  $\inf_{t \in \mathbb{R}} |h(t)| > 0$  and  $\|H_S\| < \inf_{t \in \mathbb{R}} |h_d(t)|$ . Then there exists a function  $K$  in  $V_{\mathbb{C}}$  such that

$$\frac{1}{h(t)} = \int_{\mathbb{R}} e^{itx} dK(x), \quad t \in \mathbb{R}.$$

The following theorem was proposed by Berger [6]. In fact, it is a modification of one of Krein’s results (see [23, Theorem L, p. 15]).

**Theorem 5.** Suppose that a function  $h : \mathbb{R} \rightarrow \mathbb{C}$  is defined by the formula

$$(16) \quad h(t) := c + \int_{\mathbb{R}} e^{itx} u(x) dx, \quad t \in \mathbb{R},$$

where  $c \in \mathbb{C} \setminus \{0\}$  is a constant,  $u : \mathbb{R} \rightarrow \mathbb{C}$  is a function satisfying the condition  $\int_{\mathbb{R}} |u(x)| dx < \infty$ . Assume that  $h(0) = 1$  and  $h(t) \neq 0$  for any  $t \in \mathbb{R}$ . Then

$$h(t) = \exp \left\{ \int_{\mathbb{R} \setminus \{0\}} (e^{itx} - 1) \left( v(x) + \operatorname{sgn}(x) \frac{m \cdot e^{-|x|}}{|x|} \right) dx \right\}, \quad t \in \mathbb{R},$$

where  $m$  is a constant defined by the formula

$$m := \frac{1}{2\pi} \left( \lim_{t \rightarrow \infty} \operatorname{Arg} h(t) - \lim_{t \rightarrow -\infty} \operatorname{Arg} h(t) \right),$$

and the function  $v : \mathbb{R} \rightarrow \mathbb{C}$  satisfies the condition  $\int_{\mathbb{R}} |v(x)| dx < \infty$ .

Let us consider the quantity  $m$  from the theorem. It is the *index* of  $h$  (see [23]). By the Riemann–Lebesgue lemma, for the function  $h$  defined by (16) we have  $h(t) \rightarrow c$  as  $t \rightarrow \pm\infty$ . Therefore it is not difficult to conclude the following fact.

**Remark 4.** The quantity  $m$  from Theorem 5 is well-defined and it is an integer.

We next formulate a very useful result obtained by Berger [6].

**Theorem 6.** Let  $F$  be a distribution function on  $\mathbb{R}$  with the characteristic function  $f$ . Suppose that  $f$  admits the representation

$$f(t) = \exp \left\{ it\gamma_1 - \frac{\sigma^2 t^2}{2} + \int_{\mathbb{R} \setminus \{0\}} (e^{itx} - 1 - it \mathbb{1}_{[-1,1]}(x)) dL(x) \right\}, \quad t \in \mathbb{R},$$

with some  $\gamma_1 \in \mathbb{C}$ ,  $\sigma^2 \in \mathbb{C}$ , and the function  $L(x) = L_1(x) + iL_2(x)$ ,  $x \in \mathbb{R} \setminus \{0\}$ . Here for every  $j \in \{1, 2\}$  the function  $L_j$  has bounded variation on  $\mathbb{R} \setminus O_\delta$  for any  $\delta > 0$  (in general,  $L_j$  may be nonmonotonic on the intervals  $(-\infty, 0)$  and  $(0, +\infty)$ ),  $L_j$  is right-continuous at every point of  $\mathbb{R}$ ,  $L_j(-\infty) = L_j(+\infty) = 0$ , and

$$\int_{O_\delta} x^2 d|L_j|(x) < \infty \quad \text{for any } \delta > 0,$$

where  $O_\delta := (-\delta, 0) \cup (0, \delta)$ . Then  $\gamma_1 \in \mathbb{R}$ ,  $\sigma^2 \geq 0$ , and  $\operatorname{Im}\{L(x)\} = L_2(x) = 0$  for any  $x \in \mathbb{R} \setminus \{0\}$ . In addition,  $F \in \mathcal{Q}$ .

The next theorem yields one special property of characteristic functions of rational-infinitely divisible laws. Its proof can be found in [18, p. 3] or [19, p. 360].

**Theorem 7.** *Let  $F$  be a distribution function on  $\mathbb{R}$  with the characteristic function  $f$ . If  $F \in \mathcal{Q}$  then for any  $\tau > 0$  there exists  $C_\tau > 0$  such that*

$$\sup_{t \in \mathbb{R}} \left| \frac{f(t - \tau)f(t + \tau)}{f(t)^2} \right| \leq C_\tau.$$

**Lemma 1** (Bochner [9]). *Let  $(W_n)_{n \in \mathbb{N}}$  be a sequence in  $\mathcal{V}$  such that  $\sup_{n \in \mathbb{N}} \|W_n\| = B < \infty$ . Let*

$$w_n(t) := \int_{\mathbb{R}} e^{itx} dW_n(x), \quad t \in \mathbb{R}, \quad n \in \mathbb{N}.$$

*Suppose that  $w_n(t) \rightarrow w(t)$  as  $n \rightarrow \infty$  uniformly on every bounded interval. Then there exists  $W \in \mathcal{V}$  with  $\|W\| \leq B$  such that*

$$w(t) = \int_{\mathbb{R}} e^{itx} dW(x), \quad t \in \mathbb{R}.$$

We will need the following multivariate version of the Leibniz rule for partial derivatives of a product of two functions (see [13, p. 10]).

**Lemma 2.** *Suppose that the functions  $f : \mathbb{R}^d \rightarrow \mathbb{C}$  and  $g : \mathbb{R}^d \rightarrow \mathbb{C}$  have continuous mixed partial derivatives up to and including order  $d$  in some open set  $T \subset \mathbb{R}^d$ . Then, for any  $t = (t_1, \dots, t_d) \in T$ ,*

$$\frac{\partial^d (f(t) \cdot g(t))}{\partial t_1 \cdots \partial t_d} = \sum_{u \subset D} \left( \frac{\partial^{|u|} f(t)}{\prod_{j \in u} \partial t_j} \cdot \frac{\partial^{d-|u|} g(t)}{\prod_{j \in D \setminus u} \partial t_j} \right),$$

*where  $D = \{1, \dots, d\}$ , and the parameter  $u$  of the sum ranges over all the subsets of  $D$ .*

We will also need the multivariate form of Faà di Bruno's formula for the partial derivatives of a composition of two functions (see [13, p. 4]).

**Lemma 3.** *Suppose that a function  $g : \mathbb{R}^d \rightarrow \mathbb{C}$  has continuous mixed partial derivatives up to and including order  $d$  in some open set  $T \subset \mathbb{R}^d$ . Suppose that  $f : \mathbb{C} \rightarrow \mathbb{C}$  is a holomorphic function in some domain in  $\mathbb{C}$  that which include the values of  $g(t)$ ,  $t \in T$ . Suppose that  $D := \{1, \dots, d\}$ ,  $u$  is nonempty subset of  $D$ ,  $\mathcal{P}(u)$  is the set of all partitions of the set  $u$ . Then, for any  $t = (t_1, \dots, t_d) \in T$ ,*

$$\frac{\partial^{|u|} f(g(t))}{\prod_{j \in u} \partial t_j} = \sum_{P \in \mathcal{P}(u)} \left( \frac{d^{|P|} f(z)}{dz^{|P|}} \Big|_{z=g(t)} \cdot \prod_{s \in P} \frac{\partial^{|s|} g(t)}{\prod_{j \in s} \partial t_j} \right).$$

We now formulate a lemma that will be a key tool in proving of Theorem 3. Its proof is given at the start of the next section.

**Lemma 4.** *Suppose a function  $\varphi : \mathbb{R}^d \rightarrow \mathbb{C}$ , where  $d \in \mathbb{N}$ , admits the representation*

$$\varphi(t) = \sum_{m=1}^N q_m e^{i\langle t, c_m \rangle}, \quad t = (t_1, \dots, t_d) \in \mathbb{R}^d,$$

with  $N \in \mathbb{N}$ ,  $q_m \in \mathbb{C}$ , and  $c_m = (c_{m,1}, \dots, c_{m,d}) \in \mathbb{Z}^d$ ,  $m = 1, \dots, N$ . Then, for any  $k \in \mathbb{N}$ ,

$$(17) \quad \|\varphi^k\| \leq \left(\frac{\pi}{\sqrt{6}}\right)^d \sum_{u \subset D} \left( \prod_{j \in D \setminus u} |k\alpha_j - 1| \cdot \sum_{\substack{P \in \mathcal{P}(u) \\ |P| \leq k}} \left( \prod_{l=1}^{|P|} (k-l+1) \cdot S_\varphi^{k-|P|} \cdot R_\varphi(P) \right) \right),$$

where  $D = \{1, \dots, d\}$ ,

$$(18) \quad \alpha_j := \min_{m=1, \dots, N} c_{m,j}, \quad j = 1, \dots, d, \\ S_\varphi := \sup_{t \in [-\pi, \pi]^d} |\varphi(t)|, \quad R_\varphi(P) := \sup_{t \in [-\pi, \pi]^d} \prod_{s \in P} \left| \frac{\partial^{|s|} \varphi(t)}{\prod_{j \in s} \partial t_j} \right|.$$

In particular,

$$(19) \quad \|\varphi^k\| \leq A_\varphi k^d S_\varphi^k \quad \text{for any } k \in \mathbb{N},$$

with

$$(20) \quad A_\varphi := \left(\frac{\pi}{\sqrt{6}}\right)^d \sum_{u \subset D} \left( \prod_{j \in D \setminus u} (|\alpha_j| + 1) \cdot \sum_{P \in \mathcal{P}(u)} S_\varphi^{-|P|} R_\varphi(P) \right),$$

which is independent of  $k$ .

#### 4. Proofs

*Proof of Lemma 4.* We will first prove inequality (17) for  $k = 1$ . We define a vector  $a := (a_1, \dots, a_d)$  with  $a_j := 1 - \alpha_j$ ,  $j = 1, \dots, d$ , and we introduce the function

$$\hat{\varphi}_a(t) := \varphi(t) e^{i\langle t, a \rangle} = \sum_{m=1}^N q_m e^{i\langle t, c_m + a \rangle}, \quad t \in \mathbb{R}^d.$$

Note that  $c_m + a \in \mathbb{N}^d$  for any  $m = 1, \dots, N$ . Without loss of generality we can assume that  $c_m$  are distinct if  $N > 1$ . Then we have

$$(21) \quad \|\varphi\| = \sum_{m=1}^N |q_m| = \sum_{m=1}^N \left( |q_m| \cdot \prod_{j=1}^d (c_{m,j} + a_j) \cdot \prod_{j=1}^d \frac{1}{c_{m,j} + a_j} \right) \\ \leq \left( \sum_{m=1}^N \left( |q_m|^2 \cdot \prod_{j=1}^d (c_{m,j} + a_j)^2 \right) \right)^{1/2} \cdot \left( \sum_{m=1}^N \prod_{j=1}^d \frac{1}{(c_{m,j} + a_j)^2} \right)^{1/2}.$$

Since  $c_m + a = (c_{m,1} + a_1, \dots, c_{m,d} + a_d) \in \mathbb{N}^d$  are distinct vectors for different  $m$ , the following estimate holds:

$$\sum_{m=1}^N \prod_{j=1}^d \frac{1}{(c_{m,j} + a_j)^2} \leq \prod_{j=1}^d \left(1 + \frac{1}{2^2} + \dots + \frac{1}{n^2} + \dots\right) = \left(\frac{\pi^2}{6}\right)^d.$$

We next write the function  $\hat{\varphi}_a$  in expanded form as

$$\hat{\varphi}_a(t) = \sum_{m=1}^N q_m \exp \left\{ i \sum_{j=1}^d t_j (c_{m,j} + a_j) \right\}, \quad t = (t_1, \dots, t_d) \in \mathbb{R}^d.$$

So it is easily seen that

$$\frac{\partial^d \hat{\varphi}_a(t)}{\partial t_1 \cdots \partial t_d} = \sum_{m=1}^N \left( q_m \cdot i^d \cdot \prod_{j=1}^d (c_{m,j} + a_j) \cdot \exp \left\{ i \sum_{j=1}^d t_j (c_{m,j} + a_j) \right\} \right),$$

$$t = (t_1, \dots, t_d) \in \mathbb{R}^d.$$

We consider the written exponential functions (additionally normed by  $(2\pi)^d$ ) as a rank- $N$  orthonormal system of  $L_2([-\pi, \pi]^d)$  and, by Parseval's identity, we get

$$\sum_{m=1}^N \left( |q_m|^2 \cdot \prod_{j=1}^d (c_{m,j} + a_j)^2 \right) = \frac{1}{(2\pi)^d} \int_{[-\pi, \pi]^d} \left| \frac{\partial^d \hat{\varphi}_a(t)}{\partial t_1 \cdots \partial t_d} \right|^2 dt_1 \cdots dt_d.$$

Hence

$$\sum_{m=1}^N \left( |q_m|^2 \cdot \prod_{j=1}^d (c_{m,j} + a_j)^2 \right) \leq \sup_{t \in [-\pi, \pi]^d} \left| \frac{\partial^d \hat{\varphi}_a(t)}{\partial t_1 \cdots \partial t_d} \right|^2.$$

Using these estimates in (21), we come to the inequality

$$\|\varphi\| \leq \left( \frac{\pi}{\sqrt{6}} \right)^d \sup_{t \in [-\pi, \pi]^d} \left| \frac{\partial^d \hat{\varphi}_a(t)}{\partial t_1 \cdots \partial t_d} \right|.$$

We next find an upper estimate for the latter supremum in terms of  $\varphi$  and  $a$ . We apply Lemma 2 for the function  $\hat{\varphi}_a(t) = \varphi(t)e^{i\langle t, a \rangle}$  with  $f(t) := \varphi(t)$  and  $g(t) := e^{i\langle t, a \rangle}$ ,  $t \in \mathbb{R}^d$ :

$$\frac{\partial^d \hat{\varphi}_a(t)}{\partial t_1 \cdots \partial t_d} = \sum_{u \subset D} \left( \frac{\partial^{|u|} \varphi(t)}{\prod_{j \in u} \partial t_j} \cdot \frac{\partial^{d-|u|} e^{i\langle t, a \rangle}}{\prod_{j \in D \setminus u} \partial t_j} \right) = \sum_{u \subset D} \left( \frac{\partial^{|u|} \varphi(t)}{\prod_{j \in u} \partial t_j} e^{i\langle t, a \rangle} i^{d-|u|} \prod_{j \in D \setminus u} a_j \right),$$

$$t \in \mathbb{R}^d.$$

Here  $D := \{1, \dots, d\}$  and the index  $u$  of the sum ranges over all subsets of  $D$ .

Hence we get the estimate

$$\sup_{t \in [-\pi, \pi]^d} \left| \frac{\partial^d \hat{\varphi}_a(t)}{\partial t_1 \cdots \partial t_d} \right| \leq \sum_{u \subset D} \left( \prod_{j \in D \setminus u} |a_j| \cdot \sup_{t \in [-\pi, \pi]^d} \left| \frac{\partial^{|u|} \varphi(t)}{\prod_{j \in u} \partial t_j} \right| \right).$$

Thus (on account of the equalities  $a_j = 1 - \alpha_j$ ,  $j = 1, \dots, d$ ) we have

$$(22) \quad \|\varphi\| \leq \left( \frac{\pi}{\sqrt{6}} \right)^d \sum_{u \subset D} \left( \prod_{j \in D \setminus u} |\alpha_j - 1| \cdot \sup_{t \in [-\pi, \pi]^d} \left| \frac{\partial^{|u|} \varphi(t)}{\prod_{j \in u} \partial t_j} \right| \right).$$

The right-hand side of this inequality coincides with that of (17) for  $k = 1$ . Indeed, in the inner sum in (17), the index  $P$  can be equal to  $\{u\}$ , i.e.,  $|P| = 1$ , and, in the product in  $R_\varphi(P)$ , the index  $s$  assumes only the value  $u$ . Therefore  $\prod_{l=1}^{|P|} (k - l + 1) = 1$ ,  $S_\varphi^{k-|P|} = S_\varphi^0 = 1$ , and

$$R_\varphi(P) = \sup_{t \in [-\pi, \pi]^d} \left| \frac{\partial^{|u|} \varphi(t)}{\prod_{j \in u} \partial t_j} \right|.$$

Substituting these values in (17), we see the match with (22).

We now prove that (17) is true for any  $k \in \mathbb{N}$ . Let us fix an arbitrary  $k \in \mathbb{N}$ . We consider the function  $\varphi^k$  given by

$$\varphi^k(t) = \left( \sum_{m=1}^N q_m e^{i\langle t, c_m \rangle} \right)^k, \quad t \in \mathbb{R}^d.$$

Clearly, it can be written in the form

$$\varphi^k(t) = \sum_{m=1}^M Q_m e^{i\langle t, C_m \rangle}, \quad t \in \mathbb{R}^d,$$

with some  $M \in \mathbb{N}$ ,  $Q_m \in \mathbb{C}$ , and distinct  $C_m = (C_{m,1}, \dots, C_{m,d}) \in \mathbb{Z}^d$ ,  $m = 1, \dots, M$ . Observe that

$$C_m \in \left\{ \sum_{j=1}^k c_{m_j} : m_1, \dots, m_k \in \{1, \dots, N\} \right\}, \quad m = 1, \dots, M.$$

Therefore

$$\min_{m=1, \dots, M} C_{m,j} = k \min_{m=1, \dots, N} c_{m,j} = k\alpha_j, \quad j = 1, \dots, d.$$

Taking this into account, we use inequality (22) for the function  $\varphi^k$ :

$$\|\varphi^k\| \leq \left( \frac{\pi}{\sqrt{6}} \right)^d \sum_{u \subset D} \left( \prod_{j \in D \setminus u} |k\alpha_j - 1| \cdot \sup_{t \in [-\pi, \pi]^d} \left| \frac{\partial^{|u|} \varphi^k(t)}{\prod_{j \in u} \partial t_j} \right| \right).$$

We now write expressions for the mixed partial derivatives from the right-hand side. We apply Lemma 3 with  $g(t) := \varphi(t)$ ,  $t \in \mathbb{R}^d$ , and  $f(z) := z^k$ ,  $z \in \mathbb{C}$ :

$$\begin{aligned} \frac{\partial^{|\mathbf{u}|} \varphi^k(t)}{\prod_{j \in \mathbf{u}} \partial t_j} &= \sum_{P \in \mathcal{P}(\mathbf{u})} \left( \frac{d^{|P|} z^k}{dz^{|P|}} \Big|_{z=\varphi(t)} \cdot \prod_{s \in P} \frac{\partial^{|\mathbf{s}|} \varphi(t)}{\prod_{j \in \mathbf{s}} \partial t_j} \right) \\ &= \sum_{\substack{P \in \mathcal{P}(\mathbf{u}) \\ |P| \leq k}} \left( \prod_{l=1}^{|P|} (k-l+1) \cdot \varphi(t)^{k-|P|} \cdot \prod_{s \in P} \frac{\partial^{|\mathbf{s}|} \varphi(t)}{\prod_{j \in \mathbf{s}} \partial t_j} \right). \end{aligned}$$

This formula holds even in the case  $\mathbf{u} = \emptyset$ , where  $P$  must be  $\emptyset$ , i.e.,  $|P| = 0$  and  $\mathcal{P}(\mathbf{u}) = \{\emptyset\}$ . Indeed, there is only one summand in the latter sum, in which the written products are equal to 1 (because the indexes formally belong to the empty set). Hence the right-hand side equals  $\varphi(t)^k$ , as does the left-hand side, since  $|\mathbf{u}| = 0$ .

We have the estimate

$$\sup_{t \in [-\pi, \pi]^d} \left| \frac{\partial^{|\mathbf{u}|} \varphi^k(t)}{\prod_{j \in \mathbf{u}} \partial t_j} \right| \leq \sum_{\substack{P \in \mathcal{P}(\mathbf{u}) \\ |P| \leq k}} \left( \prod_{l=1}^{|P|} (k-l+1) \cdot S_\varphi^{k-|P|} \cdot R_\varphi(P) \right),$$

where  $S_\varphi$  and  $R_\varphi(P)$  are defined by (18). Thus we obtain the required inequality (17).

We next show that (19) holds. Let us obtain an upper estimate for the right-hand side of inequality (17). We fix  $k \geq 1$ . Observe that

$$\prod_{j \in D \setminus \mathbf{u}} |k\alpha_j - 1| \leq \prod_{j \in D \setminus \mathbf{u}} (k|\alpha_j| + 1) \leq \prod_{j \in D \setminus \mathbf{u}} (k|\alpha_j| + k) \leq k^{d-|\mathbf{u}|} \cdot \prod_{j \in D \setminus \mathbf{u}} (|\alpha_j| + 1).$$

Next, since  $|P| \leq |\mathbf{u}|$  for any  $P \in \mathcal{P}(\mathbf{u})$ , we have

$$\prod_{l=1}^{|P|} (k-l+1) \leq \prod_{l=1}^{|P|} (k-1+1) = k^{|P|} \leq k^{|\mathbf{u}|}.$$

We apply these inequalities to (17), obtaining

$$\begin{aligned} \|\varphi^k\| &\leq \left( \frac{\pi}{\sqrt{6}} \right)^d \sum_{\mathbf{u} \subset D} \left( k^{d-|\mathbf{u}|} \prod_{j \in D \setminus \mathbf{u}} (|\alpha_j| + 1) \cdot \sum_{\substack{P \in \mathcal{P}(\mathbf{u}) \\ |P| \leq k}} k^{|\mathbf{u}|} \cdot S_\varphi^{k-|P|} \cdot R_\varphi(P) \right) \\ &= k^d S_\varphi^k \cdot \left( \frac{\pi}{\sqrt{6}} \right)^d \sum_{\mathbf{u} \subset D} \left( \prod_{j \in D \setminus \mathbf{u}} (|\alpha_j| + 1) \cdot \sum_{\substack{P \in \mathcal{P}(\mathbf{u}) \\ |P| \leq k}} S_\varphi^{-|P|} R_\varphi(P) \right). \end{aligned}$$

Since  $S_\varphi$  and  $R_\varphi(P)$  are always nonnegative, we come to the needed inequality

$$\|\varphi^k\| \leq k^d S_\varphi^k \left( \frac{\pi}{\sqrt{6}} \right)^d \sum_{\mathbf{u} \subset D} \left( \prod_{j \in D \setminus \mathbf{u}} (|\alpha_j| + 1) \cdot \sum_{P \in \mathcal{P}(\mathbf{u})} S_\varphi^{-|P|} R_\varphi(P) \right) = A_\varphi k^d S_\varphi^k. \quad \square$$

*Proof of Theorem 3.* We assume that  $F$  admits the decomposition (3) with some  $c_d \in (0, 1]$ , and that  $F$  has a dominated singular part. We set  $\mu_d := \inf_{t \in \mathbb{R}} |f_d(t)|$ .

(i)  $\Rightarrow$  (ii). Suppose that  $F \in \mathcal{Q}$ . If  $\mu_d = 0$  then  $c_s = 0$  by the assumption of a dominated singular part, and hence we can apply Theorem 2. According to that theorem, (i) implies (ii). Next, if  $\mu_d > 0$  then we know that  $c_s < c_d \mu_d$ , i.e.,  $c_d \mu_d - c_s > 0$ . Observe that, for any  $t \in \mathbb{R}$ ,

$$\begin{aligned} |f(t)| &= |c_d f_d(t) + c_s f_s(t) + c_a f_a(t)| \geq c_d |f_d(t)| - c_s |f_s(t)| - c_a |f_a(t)| \\ &\geq c_d \mu_d - c_s - |f_a(t)|. \end{aligned}$$

Since  $f_a(t) \rightarrow 0$  as  $t \rightarrow \pm\infty$ , there exists  $T > 0$  such that  $|f_a(t)| < \frac{1}{2}(c_d \mu_d - c_s)$  as  $|t| > T$ . Hence  $|f(t)| > \frac{1}{2}(c_d \mu_d - c_s) > 0$  as  $|t| > T$ . Next, let us consider the function  $t \mapsto |f(t)|$  on the segment  $[-T, T]$ . Since it is continuous, there exists  $t_{\min} \in [-T, T]$  such that  $|f(t)| \geq |f(t_{\min})|$  for any  $t \in [-T, T]$ . Since  $F \in \mathcal{Q}$ , we know that  $f(t) \neq 0$  for any  $t \in \mathbb{R}$  (see the introduction) and thus  $C_T := |f(t_{\min})| > 0$ . So we get  $|f(t)| \geq C_T > 0$  for any  $t \in [-T, T]$ . Thus, for any  $t \in \mathbb{R}$ ,

$$|f(t)| \geq \min\left\{\frac{1}{2}(c_d \mu_d - c_s), C_T\right\} > 0,$$

i.e.,  $\inf_{t \in \mathbb{R}} |f(t)| > 0$ . So we come to (ii).

(ii)  $\Rightarrow$  (iii). Obviously, (ii) yields that  $f(t) \neq 0$  for any  $t \in \mathbb{R}$ . To obtain a contradiction, suppose that  $\mu_d = \inf_{t \in \mathbb{R}} |f_d(t)| = 0$ . Then  $c_s = 0$  by the condition of dominated singular part. According to Theorem 2, (iii) follows from (ii), i.e., in particular, we have  $\mu_d > 0$ , a contradiction. Thus we conclude that  $\inf_{t \in \mathbb{R}} |f_d(t)| > 0$  and (iii) holds.

(iii)  $\Rightarrow$  (i). Let us consider  $f$  represented by formula (4) with  $c_d > 0$ . Here we assume that  $f(t) \neq 0$  for any  $t \in \mathbb{R}$ , that

$$(23) \quad \mu_d = \inf_{t \in \mathbb{R}} |f_d(t)| > 0,$$

and that  $c_s < c_d \mu_d$ . Due to these assumptions, for any  $t \in \mathbb{R}$  we have

$$|c_d f_d(t) + c_s f_s(t)| \geq c_d |f_d(t)| - c_s |f_s(t)| \geq c_d \mu_d - c_s > 0,$$

i.e.,

$$(24) \quad \inf_{t \in \mathbb{R}} |c_d f_d(t) + c_s f_s(t)| > 0.$$

Then for any  $t \in \mathbb{R}$  we can write

$$\begin{aligned} f(t) &= c_d f_d(t) + c_s f_s(t) + c_a f_a(t) \\ &= f_d(t) \cdot \frac{c_d f_d(t) + c_s f_s(t) + c_a f_a(t)}{c_d f_d(t) + c_s f_s(t)} \cdot \frac{c_d f_d(t) + c_s f_s(t)}{f_d(t)} \\ &= f_d(t) \cdot \left(1 + \frac{c_a f_a(t)}{c_d f_d(t) + c_s f_s(t)}\right) \cdot \frac{c_d f_d(t) + c_s f_s(t)}{f_d(t)}. \end{aligned}$$

So it is convenient to represent  $f$  as

$$(25) \quad f(t) = f_d(t) f_{a,ds}(t) f_{s,d}(t), \quad t \in \mathbb{R},$$

where

$$f_{a,ds}(t) := c_d + c_s + \frac{c_a(c_d + c_s) f_a(t)}{c_d f_d(t) + c_s f_s(t)}, \quad f_{s,d}(t) := \frac{c_d f_d(t) + c_s f_s(t)}{(c_d + c_s) f_d(t)}, \quad t \in \mathbb{R}.$$

Let us consider  $f_d$  represented by (6). Due to (23), by Theorem 1, it admits the representation

$$(26) \quad f_d(t) = \exp \left\{ it\gamma_0 + \sum_{u \in \langle \mathcal{X} \rangle \setminus \{0\}} \lambda_u (e^{itu} - 1) \right\}, \quad t \in \mathbb{R},$$

with some  $\gamma_0 \in \langle \mathcal{X} \rangle$  and  $\lambda_u \in \mathbb{R}$  for all  $u \in \langle \mathcal{X} \rangle \setminus \{0\}$ , and  $\sum_{u \in \langle \mathcal{X} \rangle \setminus \{0\}} |\lambda_u| < \infty$ .

We now consider the function  $f_{a,ds}$ . If  $c_a = 0$  then  $f_{a,ds}(t) = c_d + c_s = 1 - c_a = 1$ . We next suppose that  $c_a > 0$ . The function  $t \mapsto c_d f_d(t) + c_s f_s(t)$ ,  $t \in \mathbb{R}$ , is separated from 0 according to (24) and for its continuous singular part we have

$$\|c_s f_s\| = c_s < c_d \mu_d = c_d \inf_{t \in \mathbb{R}} |f_d(t)|.$$

By Theorem 4, there exists a function  $I_{ds} : \mathbb{R} \rightarrow \mathbb{C}$  in  $V_{\mathbb{C}}$  such that

$$\frac{1}{c_d f_d(t) + c_s f_s(t)} = \int_{\mathbb{R}} e^{itx} dI_{ds}(x), \quad t \in \mathbb{R}.$$

We next observe that

$$\frac{f_a(t)}{c_d f_d(t) + c_s f_s(t)} = \int_{\mathbb{R}} e^{itx} dF_a(x) \cdot \int_{\mathbb{R}} e^{itx} dI_{ds}(x) = \int_{\mathbb{R}} e^{itx} d(F_a * I_{ds})(x), \quad t \in \mathbb{R},$$

and we write

$$\frac{c_a(c_d + c_s) f_a(t)}{c_d f_d(t) + c_s f_s(t)} = \int_{\mathbb{R}} e^{itx} d\tilde{F}_a(x), \quad t \in \mathbb{R},$$

where  $\tilde{F}_a(x) := c_a(c_d + c_s)(F_a * I_{ds})(x)$ ,  $x \in \mathbb{R}$ . The function  $\tilde{F}_a \in V_{\mathbb{C}}$  inherits absolute continuity from  $F_a$ , i.e., there exists a function  $\tilde{p}_a : \mathbb{R} \rightarrow \mathbb{C}$  such that  $\int_{\mathbb{R}} |\tilde{p}_a(x)| dx < \infty$  and  $\tilde{F}_a(x) = \int_{y \leq x} \tilde{p}_a(y) dy$ ,  $x \in \mathbb{R}$ . Hence

$$\int_{\mathbb{R}} e^{itx} d\tilde{F}_a(x) = \int_{\mathbb{R}} e^{itx} \tilde{p}_a(x) dx, \quad t \in \mathbb{R},$$

and we have

$$f_{a,ds}(t) = c_d + c_s + \frac{c_a(c_d + c_s) f_a(t)}{c_d f_d(t) + c_s f_s(t)} = c_d + c_s + \int_{\mathbb{R}} e^{itx} \tilde{p}_a(x) dx, \quad t \in \mathbb{R},$$

where  $c_d + c_s \geq c_d > 0$ . Note that  $f_{a,ds}(0) = 1$ . Since  $f(t) \neq 0$  for any  $t \in \mathbb{R}$  and the function  $t \mapsto c_d f_d(t) + c_s f_s(t)$ ,  $t \in \mathbb{R}$ , is bounded, we conclude that  $f_{a,ds}(t) \neq 0$

for any  $t \in \mathbb{R}$ . By Theorem 5,  $f_{a,d_s}$  admits the representation

$$(27) \quad f_{a,d_s}(t) = \exp \left\{ \int_{\mathbb{R} \setminus \{0\}} (e^{itx} - 1) \left( v_a(x) + \operatorname{sgn}(x) \frac{m_a \cdot e^{-|x|}}{|x|} \right) dx \right\}, \quad t \in \mathbb{R},$$

where  $v_a : \mathbb{R} \rightarrow \mathbb{C}$  is a function satisfying the condition  $\int_{\mathbb{R}} |v_a(x)| dx < \infty$ ,  $m_a$  is a constant defined by the formula

$$m_a := \frac{1}{2\pi} \left( \lim_{t \rightarrow \infty} \operatorname{Arg} f_{a,d_s}(t) - \lim_{t \rightarrow -\infty} \operatorname{Arg} f_{a,d_s}(t) \right).$$

Since  $c_d + c_s$  is positive, formula (10) gives an equivalent definition of  $m_a$ . By the way, it is easily seen that  $m_a = 0$  if  $c_a = 0$ . In general,  $m_a \in \mathbb{Z}$  (see Remark 4). Therefore, in the case  $c_a = 0$ , we set  $v_a(x) := 0$  for every  $x \in \mathbb{R}$  and the formula (27) remains valid.

We now turn to the function

$$f_{s,d}(t) = \frac{c_d f_d(t) + c_s f_s(t)}{(c_d + c_s) f_d(t)} = \left( 1 + \frac{c_s f_s(t)}{c_d f_d(t)} \right) \cdot \frac{c_d}{c_d + c_s}, \quad t \in \mathbb{R}.$$

From (24) we know that  $f_{s,d}(t) \neq 0$  for any  $t \in \mathbb{R}$  and we see that  $f_{s,d}(0) = 1$ . Hence the distinguished logarithm  $\operatorname{Ln} f_{s,d}$  is uniquely defined on  $\mathbb{R}$  with condition  $\operatorname{Ln} f_{s,d}(0) = 0$ . Due to the assumptions,

$$(28) \quad \left| \frac{c_s f_s(t)}{c_d f_d(t)} \right| \leq \frac{c_s}{c_d \mu_d} < 1 \quad \text{for any } t \in \mathbb{R},$$

and we can write

$$(29) \quad \operatorname{Ln} f_{s,d}(t) = \ln \left( 1 + \frac{c_s f_s(t)}{c_d f_d(t)} \right) - \ln \left( 1 + \frac{c_s}{c_d} \right), \quad t \in \mathbb{R},$$

where  $\ln(\cdot)$  returns the principal value of the logarithm. Let us consider the first logarithm in the right-hand side. Due to (28), we have the expansion

$$(30) \quad \ln \left( 1 + \frac{c_s f_s(t)}{c_d f_d(t)} \right) = \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k} \left( \frac{c_s f_s(t)}{c_d f_d(t)} \right)^k = \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k} \left( \frac{c_s}{c_d} \right)^k f_s(t)^k g_d(t)^k, \quad t \in \mathbb{R},$$

where we have set  $g_d(t) := 1/f_d(t)$ ,  $t \in \mathbb{R}$ . The series converges uniformly in  $\mathbb{R}$  by the Weierstrass M-test, because, due to (28), for any  $t \in \mathbb{R}$  the absolute values of its terms are majorized by the terms of the convergent numerical series

$$\sum_{k=1}^{\infty} \frac{1}{k} \left( \frac{c_s}{c_d \mu_d} \right)^k.$$

Let us consider the function  $g_d$  and show that it is actually a Fourier–Stieltjes transform of some function from class  $\mathbf{V}$ . Indeed, according to (26),

$$g_d(t) = e^{-it\gamma_0} \cdot e^{-\lambda_0} \cdot \exp\left\{-\sum_{u \in \langle \mathcal{X} \rangle \setminus \{0\}} \lambda_u e^{itu}\right\}, \quad t \in \mathbb{R},$$

where we have set

$$\lambda_0 := -\sum_{u \in \langle \mathcal{X} \rangle \setminus \{0\}} \lambda_u.$$

There is an absolutely convergent Fourier series with real coefficients in the last exponential function. Hence this function itself expands into an absolutely convergent Fourier series with coefficients in  $\mathbb{R}$  and exponents in  $\langle \mathcal{X} \rangle$ . This remains true after multiplying this series by  $e^{-\lambda_0}$  and  $e^{-it\gamma_0}$  (which leads to  $g_d$ ), because  $\lambda_0 \in \mathbb{R}$  and  $\gamma_0 \in \langle \mathcal{X} \rangle$ . Thus we have

$$(31) \quad g_d(t) = \sum_{y \in \langle \mathcal{X} \rangle} q_y e^{ity}, \quad t \in \mathbb{R},$$

where  $q_y \in \mathbb{R}$  for every  $y \in \langle \mathcal{X} \rangle$  and  $\sum_{y \in \langle \mathcal{X} \rangle} |q_y| < \infty$ . Obviously, the series (31) can be written as a Fourier–Stieltjes transform,

$$g_d(t) = \int_{\mathbb{R}} e^{itx} dI_d(x), \quad t \in \mathbb{R},$$

with the discrete function  $I_d \in \mathbf{V}$  given by

$$(32) \quad I_d(x) := \sum_{\substack{y \in \langle \mathcal{X} \rangle \\ y \leq x}} q_y, \quad x \in \mathbb{R}.$$

Let us return to the formula (30). We observe that for any  $k \in \mathbb{N}$  the functions  $f_s^k, g_d^k, f_s^k \cdot g^k$  are the Fourier–Stieltjes transforms of functions from  $\mathbf{V}$ :

$$(33) \quad \begin{aligned} f_s(t)^k &= \int_{\mathbb{R}} e^{itx} dF_s^{*k}(x), & g_d(t)^k &= \int_{\mathbb{R}} e^{itx} dI_d^{*k}(x), \\ f_s(t)^k g_d(t)^k &= \int_{\mathbb{R}} e^{itx} d(F_s^{*k} * I_d^{*k})(x), & t \in \mathbb{R}. \end{aligned}$$

Consequently, the partial sums of the series (30) are the Fourier–Stieltjes transforms of the functions

$$(34) \quad W_n(x) := \sum_{k=1}^n \frac{(-1)^{k-1}}{k} \left(\frac{c_s}{c_d}\right)^k (F_s^{*k} * I_d^{*k})(x), \quad x \in \mathbb{R}, \quad n \in \mathbb{N}.$$

Let us show that the sum of the whole series (30) admits the representation

$$(35) \quad \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k} \left(\frac{c_s}{c_d}\right)^k f_s(t)^k g_d(t)^k = \int_{\mathbb{R}} e^{itx} dW(x), \quad t \in \mathbb{R},$$

with some function  $W \in \mathbf{V}$ . According to Lemma 1, for this it is sufficient that (1) the partial sums of (30) uniformly converge to the infinite sum on every bounded interval and (2)  $\sup_{n \in \mathbb{N}} \|W_n\| < \infty$ . The uniform convergence of the partial sums (even on whole real line) was shown above, i.e., (1) holds. Let us prove (2). For any  $n \in \mathbb{N}$  we have the estimate

$$\|W_n\| \leq \sum_{k=1}^n \frac{1}{k} \left(\frac{c_s}{c_d}\right)^k \|F_s^{*k} * I_d^{*k}\| \leq \sum_{k=1}^n \left\{ \frac{1}{k} \left(\frac{c_s}{c_d}\right)^k \|F_s^{*k}\| \cdot \|I_d^{*k}\| \right\}.$$

Here  $\|F_s^{*k}\| = 1$  for every  $k \in \mathbb{N}$ , because all the  $F_s^{*k}$  are distribution functions. Also, it is convenient for us to set  $\|g_d^k\| = \|I_d^{*k}\|$ ,  $k \in \mathbb{N}$ . Thus we come to the inequalities

$$(36) \quad \|W_n\| \leq \sum_{k=1}^n \frac{1}{k} \left(\frac{c_s}{c_d}\right)^k \|g_d^k\|, \quad n \in \mathbb{N}.$$

We now estimate  $\|g_d^k\|$  for every  $k \in \mathbb{N}$ . Let us return to decomposition (31) for the function  $g_d$ . The set  $\langle \mathcal{X} \rangle$  is countable, and  $\sum_{y \in \langle \mathcal{X} \rangle} |q_y| < \infty$ . So we fix an arbitrary  $\varepsilon \in (0, 1)$  and choose  $N_\varepsilon \in \mathbb{N}$  and distinct points  $y_1, \dots, y_{N_\varepsilon} \in \langle \mathcal{X} \rangle$  such that

$$\sum_{y \in \langle \mathcal{X} \rangle \setminus \{y_1, \dots, y_{N_\varepsilon}\}} |q_y| < \varepsilon.$$

We introduce the polynomial

$$(37) \quad \tilde{g}_\varepsilon(t) := \sum_{m=1}^{N_\varepsilon} q_{y_m} e^{it y_m}, \quad t \in \mathbb{R}.$$

So we have  $\|g_d - \tilde{g}_\varepsilon\| < \varepsilon$ . We also observe that

$$(38) \quad \left\| \frac{g_d - \tilde{g}_\varepsilon}{g_d} \right\| \leq \|g_d - \tilde{g}_\varepsilon\| \cdot \left\| \frac{1}{g_d} \right\| = \|g_d - \tilde{g}_\varepsilon\| \cdot \|f_d\| = \|g_d - \tilde{g}_\varepsilon\| < \varepsilon.$$

Let us fix  $k \in \mathbb{N}$  and write

$$\|g_d^k\| = \left\| \tilde{g}_\varepsilon^k \cdot \left(\frac{g_d}{\tilde{g}_\varepsilon}\right)^k \right\| \leq \|\tilde{g}_\varepsilon^k\| \cdot \left\| \frac{g_d}{\tilde{g}_\varepsilon} \right\|^k.$$

In order to get a convenient representation for  $g_d/\tilde{g}_\varepsilon$ , we observe that for any  $n \in \mathbb{N}$

$$\begin{aligned} 1 - \left(\frac{g_d(t) - \tilde{g}_\varepsilon(t)}{g_d(t)}\right)^n &= \left(1 - \frac{g_d(t) - \tilde{g}_\varepsilon(t)}{g_d(t)}\right) \cdot \sum_{j=0}^{n-1} \left(\frac{g_d(t) - \tilde{g}_\varepsilon(t)}{g_d(t)}\right)^j \\ &= \frac{\tilde{g}_\varepsilon(t)}{g_d(t)} \cdot \sum_{j=0}^{n-1} \left(\frac{g_d(t) - \tilde{g}_\varepsilon(t)}{g_d(t)}\right)^j, \quad t \in \mathbb{R}. \end{aligned}$$

So we have

$$\frac{g_d(t)}{\tilde{g}_\varepsilon(t)} = \sum_{j=0}^{n-1} \left( \frac{g_d(t) - \tilde{g}_\varepsilon(t)}{g(t)} \right)^j + \left( \frac{g_d(t) - \tilde{g}_\varepsilon(t)}{g_d(t)} \right)^n \cdot \frac{g_d(t)}{\tilde{g}_\varepsilon(t)}, \quad t \in \mathbb{R}, \quad n \in \mathbb{N}.$$

Hence, due to (38), we get

$$\left\| \frac{g_d}{\tilde{g}_\varepsilon} \right\| \leq \sum_{j=0}^{n-1} \left\| \frac{g_d - \tilde{g}_\varepsilon}{g_d} \right\|^j + \left\| \frac{g_d - \tilde{g}_\varepsilon}{g_d} \right\|^n \cdot \left\| \frac{g_d}{\tilde{g}_\varepsilon} \right\| < \sum_{j=0}^{n-1} \varepsilon^j + \varepsilon^n \cdot \left\| \frac{g_d}{\tilde{g}_\varepsilon} \right\|, \quad n \in \mathbb{N}.$$

Since  $\varepsilon \in (0, 1)$ , letting  $n \rightarrow \infty$  yields

$$\left\| \frac{g_d}{\tilde{g}_\varepsilon} \right\| \leq \sum_{j=0}^{\infty} \varepsilon^j = \frac{1}{1 - \varepsilon}.$$

Thus we have

$$(39) \quad \|g_d^k\| \leq \frac{\|\tilde{g}_\varepsilon^k\|}{(1 - \varepsilon)^k}.$$

So the estimation of  $\|I_d^{*k}\| = \|g_d^k\|$  is reduced to finding an upper bound for  $\|\tilde{g}_\varepsilon^k\|$ .

Let us consider  $\tilde{g}_\varepsilon$  defined by formula (37). Let  $Y_\varepsilon := \{y_1, y_2, \dots, y_{N_\varepsilon}\}$ . Suppose that  $Y_\varepsilon = \{0\}$ , i.e.,  $\sum_{y \in \langle \mathcal{X} \rangle \setminus \{0\}} |q_y| < \varepsilon$  and  $\tilde{g}_\varepsilon(t) = q_0$  for any  $t \in \mathbb{R}$ . Then  $\|\tilde{g}_\varepsilon^k\| = |q_0|^k$  for any  $k \in \mathbb{N}$ . Observe that

$$|q_0| = |\tilde{g}_\varepsilon(0)| \leq |g_d(0)| + |\tilde{g}_\varepsilon(0) - g_d(0)|,$$

where  $|g_d(0)| = |1/f_d(0)| = 1$  and

$$|\tilde{g}_\varepsilon(0) - g_d(0)| = \left| \sum_{y \in \langle \mathcal{X} \rangle \setminus \{0\}} q_y \right| \leq \sum_{y \in \langle \mathcal{X} \rangle \setminus \{0\}} |q_y| < \varepsilon.$$

Thus  $|q_0| < 1 + \varepsilon$  and hence  $\|\tilde{g}_\varepsilon^k\| < (1 + \varepsilon)^k$ .

We next assume that  $Y_\varepsilon \neq \{0\}$ , i.e.,  $Y_\varepsilon$  contains nonzero elements. We select a basis over  $\mathbb{Q}$  in the set  $Y_\varepsilon$  (see [24, p. 67–68]), i.e., we choose nonzero elements  $\beta_1, \dots, \beta_d \in Y_\varepsilon$  with some  $d \in \{1, \dots, N_\varepsilon\}$ , which are linearly independent over  $\mathbb{Q}$  and for any  $m \in \{1, \dots, N_\varepsilon\}$  there exist some  $r_{m,1}, \dots, r_{m,d} \in \mathbb{Q}$  such that  $y_m = \sum_{l=1}^d r_{m,l} \beta_l$ . Linear independence over  $\mathbb{Q}$  means that with  $r_1, \dots, r_d \in \mathbb{Q}$ , vanishes only in the case  $r_1 = r_2 = \dots = r_d = 0$ . It is clear that the coefficients  $r_{m,l}$  of the decomposition of  $y_m$  are uniquely determined. Let  $\varkappa$  be the minimal positive integer such that  $\bar{r}_{m,l} := \varkappa \cdot r_{m,l} \in \mathbb{Z}$  for any admissible  $m$  and  $l$ . We set  $\bar{\beta}_l := \beta_l / \varkappa$  for every  $l \in \{1, \dots, d\}$ . Then we have  $y_m = \sum_{l=1}^d \bar{r}_{m,l} \bar{\beta}_l$  for any  $m \in \{1, \dots, N_\varepsilon\}$ . Here the coefficients  $\bar{r}_{m,l}$  are uniquely determined by  $y_m$  too. We define the vectors  $\bar{r}_m := (\bar{r}_{m,1}, \dots, \bar{r}_{m,d}) \in \mathbb{Z}^d$ ,  $m = 1, \dots, N_\varepsilon$ . Since  $y_1, \dots, y_{N_\varepsilon}$  are assumed to be distinct, all  $\bar{r}_m$  are distinct too. Let us introduce the function

$$\varphi_\varepsilon(t_1, \dots, t_d) := \sum_{m=1}^{N_\varepsilon} q_{y_m} \exp \left\{ i \sum_{l=1}^d \bar{r}_{m,l} t_l \right\}, \quad t_1, \dots, t_d \in \mathbb{R}.$$

It is easily seen that this function is continuous and  $2\pi$ -periodic over every variable. We will also use a shorthand for it:

$$\varphi_\varepsilon(t) = \sum_{m=1}^{N_\varepsilon} q_{y_m} e^{i\langle t, \bar{r}_m \rangle}, \quad t = (t_1, \dots, t_d) \in \mathbb{R}^d.$$

The functions  $\varphi_\varepsilon$  and  $\tilde{g}_\varepsilon$  are related:

$$\varphi_\varepsilon(\bar{\beta}_1 t, \dots, \bar{\beta}_d t) = \sum_{m=1}^{N_\varepsilon} q_{y_m} \exp \left\{ i t \sum_{l=1}^d \bar{r}_{m,l} \bar{\beta}_l \right\} = \sum_{m=1}^{N_\varepsilon} q_{y_m} e^{i t y_m} = \tilde{g}_\varepsilon(t), \quad t \in \mathbb{R},$$

i.e.,  $\tilde{g}_\varepsilon$  is diagonal for the function  $(t_1, \dots, t_d) \mapsto \varphi_\varepsilon(\bar{\beta}_1 t_1, \dots, \bar{\beta}_d t_d)$ . Hence the image of the first function is dense in the image of the second one (see [24, p. 116, Theorem 2.4.1]) and, consequently, in the image of  $\varphi_\varepsilon$ . This yields

$$(40) \quad \sup_{t \in \mathbb{R}} |\tilde{g}_\varepsilon(t)| = \sup_{t \in [-\pi, \pi]^d} |\varphi_\varepsilon(t)|.$$

Let us show that

$$(41) \quad \|\tilde{g}_\varepsilon^k\| = \|\varphi_\varepsilon^k\| \quad \text{for any } k \in \mathbb{N}.$$

We fix  $k \in \mathbb{N}$  and write

$$\begin{aligned} \tilde{g}_\varepsilon^k(t) &= \left( \sum_{m=1}^{N_\varepsilon} q_{y_m} e^{i t y_m} \right)^k = \sum_{m_1=1}^{N_\varepsilon} \cdots \sum_{m_k=1}^{N_\varepsilon} (q_{y_{m_1}} \cdots q_{y_{m_k}} e^{i t (y_{m_1} + \cdots + y_{m_k})}), \quad t \in \mathbb{R}, \\ \varphi_\varepsilon^k(t) &= \left( \sum_{m=1}^{N_\varepsilon} q_{y_m} e^{i \langle t, \bar{r}_m \rangle} \right)^k = \sum_{m_1=1}^{N_\varepsilon} \cdots \sum_{m_k=1}^{N_\varepsilon} (q_{y_{m_1}} \cdots q_{y_{m_k}} e^{i \langle t, \bar{r}_{m_1} + \cdots + \bar{r}_{m_k} \rangle}), \quad t \in \mathbb{R}^d. \end{aligned}$$

We next have the representation

$$\begin{aligned} \tilde{g}_\varepsilon^k(t) &= \sum_{z \in \mathcal{Y}_\varepsilon^{(k)}} \left( \sum_{\substack{1 \leq m_1, \dots, m_k \leq N_\varepsilon \\ y_{m_1} + \cdots + y_{m_k} = z}} q_{y_{m_1}} \cdots q_{y_{m_k}} \right) e^{i t z}, \quad t \in \mathbb{R}, \\ \varphi_\varepsilon^k(t) &= \sum_{s \in \mathcal{R}_\varepsilon^{(k)}} \left( \sum_{\substack{1 \leq m_1, \dots, m_k \leq N_\varepsilon \\ \bar{r}_{m_1} + \cdots + \bar{r}_{m_k} = s}} q_{\bar{r}_{m_1}} \cdots q_{\bar{r}_{m_k}} \right) e^{i \langle t, s \rangle}, \quad t \in \mathbb{R}^d, \end{aligned}$$

where

$$\begin{aligned} \mathcal{Y}_\varepsilon^{(k)} &:= \{y_{m_1} + \cdots + y_{m_k} : m_1, \dots, m_k \in \{1, \dots, N_\varepsilon\}\}, \\ \mathcal{R}_\varepsilon^{(k)} &:= \{\bar{r}_{m_1} + \cdots + \bar{r}_{m_k} : m_1, \dots, m_k \in \{1, \dots, N_\varepsilon\}\}. \end{aligned}$$

Thus we have

$$(42) \quad \|\tilde{g}_\varepsilon^k\| = \sum_{z \in \mathcal{Y}_\varepsilon^{(k)}} \left| \sum_{\substack{m_1=1, \dots, N_\varepsilon, \\ \dots \\ m_k=1, \dots, N_\varepsilon \\ y_{m_1} + \dots + y_{m_k} = z}} q_{y_{m_1}} \cdots q_{y_{m_k}} \right|,$$

$$(43) \quad \|\varphi_\varepsilon^k\| = \sum_{s \in \mathcal{R}_\varepsilon^{(k)}} \left| \sum_{\substack{m_1=1, \dots, N_\varepsilon, \\ \dots \\ m_k=1, \dots, N_\varepsilon \\ \bar{r}_{m_1} + \dots + \bar{r}_{m_k} = s}} q_{y_{m_1}} \cdots q_{y_{m_k}} \right|.$$

There is a natural map between the two finite sets  $\mathcal{Y}_\varepsilon^{(k)}$  and  $\mathcal{R}_\varepsilon^{(k)}$ : every number  $z = \sum_{j=1}^k y_{m_j}$  from the first set is paired with a vector  $s = \sum_{j=1}^k \bar{r}_{m_j}$  from the second set. Here  $s$  is the vector of coefficients of the decomposition for  $z$  with the basis  $\bar{\beta}_1, \dots, \bar{\beta}_d$ . This map is actually a bijection. Indeed, it is injective, because distinct numbers from  $\mathcal{Y}_\varepsilon^{(k)}$  have the distinct decompositions, i.e., they are paired with the distinct vectors from the set  $\mathcal{R}_\varepsilon^{(k)}$ . The map is surjective, because any vector  $v \in \mathcal{R}_\varepsilon^{(k)}$  is a sum  $\bar{r}_{m'_1} + \dots + \bar{r}_{m'_k}$  with some indices  $m'_1, \dots, m'_k \in \{1, \dots, N_\varepsilon\}$ , and this sum corresponds to the number  $z = y_{m'_1} + \dots + y_{m'_k}$  from the set  $\mathcal{Y}_\varepsilon^{(k)}$  by construction. Next, it is clear from the basis decompositions that for any fixed pair of corresponding elements  $z \in \mathcal{Y}_\varepsilon^{(k)}$  and  $s \in \mathcal{R}_\varepsilon^{(k)}$  the equalities  $y_{m_1} + \dots + y_{m_k} = z$  and  $\bar{r}_{m_1} + \dots + \bar{r}_{m_k} = s$  are equivalent for varying index vectors  $(m_1, \dots, m_k)$ . Since the inner sums in (42) and (43) add the same weight  $q_{y_{m_1}} \cdots q_{y_{m_k}}$  for every index vector  $(m_1, \dots, m_k)$ , we conclude that these inner sums are equal. Thus we come to the equality  $\|\tilde{g}_\varepsilon^k\| = \|\varphi_\varepsilon^k\|$ .

We now apply Lemma 4 to estimate  $\|\varphi_\varepsilon^k\|$  for any  $k \in \mathbb{N}$  (we set  $\varphi := \varphi_\varepsilon$ ,  $N := N_\varepsilon$ ,  $q_m := q_{y_m}$  and  $c_m := \bar{r}_m$  for every  $m = 1, \dots, N_\varepsilon$ ). Using inequality (19), we get

$$\|\varphi_\varepsilon^k\| \leq A_{\varphi_\varepsilon} k^d \sup_{t \in [-\pi, \pi]^d} |\varphi_\varepsilon(t)|^k, \quad k \in \mathbb{N},$$

where  $A_{\varphi_\varepsilon}$  is a constant defined by (20), which doesn't depend on  $k$ . Applying (40) and (41), we come to the inequality

$$\|\tilde{g}_\varepsilon^k\| \leq A_{\varphi_\varepsilon} k^d \sup_{t \in \mathbb{R}} |\tilde{g}_\varepsilon(t)|^k, \quad k \in \mathbb{N}.$$

Observe that

$$\sup_{t \in \mathbb{R}} |\tilde{g}_\varepsilon(t)| \leq \sup_{t \in \mathbb{R}} |g_d(t)| + \sup_{t \in \mathbb{R}} |g_d(t) - \tilde{g}_\varepsilon(t)|.$$

On account of (38), we have

$$\frac{\sup_{t \in \mathbb{R}} |g_d(t) - \tilde{g}_\varepsilon(t)|}{\sup_{t \in \mathbb{R}} |g_d(t)|} \leq \sup_{t \in \mathbb{R}} \left| \frac{g_d(t) - \tilde{g}_\varepsilon(t)}{g_d(t)} \right| \leq \left\| \frac{g_d - \tilde{g}_\varepsilon}{g_d} \right\| < \varepsilon,$$

i.e.,  $\sup_{t \in \mathbb{R}} |g_d(t) - \tilde{g}_\varepsilon(t)| < \varepsilon \sup_{t \in \mathbb{R}} |g_d(t)|$ . Therefore

$$\sup_{t \in \mathbb{R}} |\tilde{g}_\varepsilon(t)| \leq \sup_{t \in \mathbb{R}} |g_d(t)| + \varepsilon \sup_{t \in \mathbb{R}} |g_d(t)| = (1 + \varepsilon) \sup_{t \in \mathbb{R}} |g_d(t)|,$$

where

$$\sup_{t \in \mathbb{R}} |g_d(t)| = \sup_{t \in \mathbb{R}} \left| \frac{1}{f_d(t)} \right| = \frac{1}{\inf_{t \in \mathbb{R}} |f_d(t)|} = \frac{1}{\mu_d}.$$

So we obtain the estimate

$$\|\tilde{g}_\varepsilon^k\| \leq A_{\varphi_\varepsilon} k^d \cdot \frac{(1 + \varepsilon)^k}{\mu_d^k}, \quad k \in \mathbb{N}.$$

We note that here  $k^d \geq 1$  and  $1/\mu_d^k \geq 1$  for any  $k \in \mathbb{N}$ . Therefore the estimates of  $\|\tilde{g}_\varepsilon^k\|$  in the cases  $Y_\varepsilon = \{0\}$  and  $Y_\varepsilon \neq \{0\}$  can be unified as

$$\|\tilde{g}_\varepsilon^k\| \leq C_\varepsilon k^d \cdot \frac{(1 + \varepsilon)^k}{\mu_d^k} \quad \text{for any } k \in \mathbb{N},$$

with some constant  $C_\varepsilon > 0$ . We use this in (39), obtaining:

$$\|g_d^k\| \leq \frac{\|\tilde{g}_\varepsilon^k\|}{(1 - \varepsilon)^k} \leq C_\varepsilon k^d \cdot \frac{1}{\mu_d^k} \cdot \left( \frac{1 + \varepsilon}{1 - \varepsilon} \right)^k, \quad k \in \mathbb{N}.$$

Let us return to (36). Due to the last estimate, for any  $\varepsilon \in (0, 1)$  and  $n \in \mathbb{N}$  we have

$$(44) \quad \|W_n\| \leq \sum_{k=1}^n \left( \frac{1}{k} \left( \frac{c_s}{c_d} \right)^k C_\varepsilon k^d \cdot \frac{1}{\mu_d^k} \cdot \left( \frac{1 + \varepsilon}{1 - \varepsilon} \right)^k \right) = C_\varepsilon \sum_{k=1}^n k^{d-1} \left( \frac{c_s}{c_d \mu_d} \cdot \frac{1 + \varepsilon}{1 - \varepsilon} \right)^k.$$

Since  $c_s < c_d \mu_d$  by assumption, the fixed number  $\varepsilon \in (0, 1)$  can be specified by

$$0 < \frac{c_s}{c_d \mu_d} \cdot \frac{1 + \varepsilon}{1 - \varepsilon} < 1.$$

Then we obtain

$$\sup_{n \in \mathbb{N}} \|W_n\| \leq C_\varepsilon \sum_{k=1}^{\infty} k^{d-1} \left( \frac{c_s}{c_d \mu_d} \cdot \frac{1 + \varepsilon}{1 - \varepsilon} \right)^k < \infty,$$

i.e., we showed that the sufficient condition (2) for (35) is satisfied (condition (1) was proved above). So, according to (30) and (35), we have

$$(45) \quad \ln \left( 1 + \frac{c_s f_s(t)}{c_d f_d(t)} \right) = \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k} \left( \frac{c_s}{c_d} \right)^k f_s(t)^k g_d(t)^k = \int_{\mathbb{R}} e^{itx} dW(x), \quad t \in \mathbb{R},$$

with some function  $W \in \mathcal{V}$ . Hence (29) takes the form

$$\text{Ln } f_{s,d}(t) = \int_{\mathbb{R}} e^{itx} dW(x) - \ln \left( 1 + \frac{c_s}{c_d} \right), \quad t \in \mathbb{R}.$$

The equality  $\text{Ln } f_{s,d}(0) = 0$ , which was mentioned above, implies

$$\ln \left( 1 + \frac{c_s}{c_d} \right) = \int_{\mathbb{R}} dW(x).$$

Then

$$\text{Ln } f_{s,d}(t) = \int_{\mathbb{R}} (e^{itx} - 1) dW(x), \quad t \in \mathbb{R},$$

i.e., we come to the representation

$$(46) \quad \begin{aligned} f_{s,d}(t) &= \exp \left\{ \int_{\mathbb{R}} (e^{itx} - 1) dW(x) \right\} \\ &= \exp \left\{ \int_{\mathbb{R} \setminus \{0\}} (e^{itx} - 1) dW(x) \right\}, \quad t \in \mathbb{R}. \end{aligned}$$

Let us investigate the function  $W$ . If  $c_s = 0$ , then  $W$  is identically 0, which is seen from (45). We next focus on the case  $c_s \neq 0$ . We will prove that  $W$  is continuous on  $\mathbb{R}$ . For this we first observe that the  $W_n$ ,  $n \in \mathbb{N}$ , are continuous functions on  $\mathbb{R}$ . Indeed, according to (34),  $W_n$  is a finite linear combination of the functions  $F_s^{*k} * I_d^{*k}$ ,  $k \in \mathbb{N}$ , which are continuous on  $\mathbb{R}$ , being convolutions with continuous  $F_s$ . Next, we observe that, similarly to (44), we have the estimate

$$\|W_{n_2} - W_{n_1}\| \leq C_\varepsilon \sum_{k=n_1+1}^{n_2} k^{d-1} \left( \frac{c_s}{c_d \mu_d} \cdot \frac{1+\varepsilon}{1-\varepsilon} \right)^k,$$

with the same  $\varepsilon$ ,  $C_\varepsilon$  and for any positive integers  $n_1$  and  $n_2$  such that  $n_1 \leq n_2$ . The sum on the right can be made arbitrarily small for all sufficiently large  $n_1$  and  $n_2$ , because its terms (for  $k \in \mathbb{N}$ ) form a convergent series. This means that  $(W_n)_{n \in \mathbb{N}}$  is a fundamental sequence in the space  $V$ . Since  $V$  is a complete norm space (see the comments at the end of the introduction), there exists  $W_* \in V$  such that  $\|W_n - W_*\| \rightarrow 0$  as  $n \rightarrow \infty$ . Hence

$$\lim_{n \rightarrow \infty} \int_{\mathbb{R}} e^{itx} dW_n(x) = \int_{\mathbb{R}} e^{itx} dW_*(x), \quad t \in \mathbb{R}.$$

On the other hand, we know that

$$\begin{aligned} \lim_{n \rightarrow \infty} \int_{\mathbb{R}} e^{itx} dW_n(x) &= \lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{(-1)^{k-1}}{k} \left( \frac{c_s}{c_d} \right)^k f_s(t)^k g_d(t)^k \\ &= \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k} \left( \frac{c_s}{c_d} \right)^k f_s(t)^k g_d(t)^k = \int_{\mathbb{R}} e^{itx} dW(x), \quad t \in \mathbb{R}. \end{aligned}$$

Then  $\int_{\mathbb{R}} e^{itx} dW_*(x) = \int_{\mathbb{R}} e^{itx} dW(x)$  for any  $t \in \mathbb{N}$ , and we conclude that  $W_* = W$ . So we have proved that

$$(47) \quad \|W_n - W\| \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

In general, from the well-known relation  $\sup_{x \in \mathbb{R}} |U(x)| \leq \|U\|$  for any  $U \in V$ , we have uniform convergence:

$$\sup_{x \in \mathbb{R}} |W_n(x) - W(x)| \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Due to the already proved continuity of  $W_n, n \in \mathbb{N}$ , we conclude that  $W$  is continuous on  $\mathbb{R}$  too.

We now show that  $W$  cannot be purely absolutely continuous (in the case  $c_s \neq 0$ ). For suppose that  $W$  is absolutely continuous. Let

$$w(t) := \int_{\mathbb{R}} e^{itx} dW(x), \quad t \in \mathbb{R}.$$

Then, according to (45), we have

$$1 + \frac{c_s f_s(t)}{c_d f_d(t)} = \exp \left\{ \int_{\mathbb{R}} e^{itx} dW(x) \right\} = \exp\{w(t)\} = 1 + \sum_{k=1}^{\infty} \frac{w(t)^k}{k!}, \quad t \in \mathbb{R}.$$

Hence, on the one hand,

$$\begin{aligned} \frac{c_s f_s(t)}{c_d f_d(t)} &= \lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{w(t)^k}{k!} = \lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{1}{k!} \int_{\mathbb{R}} e^{itx} dW^{*k}(x) \\ &= \lim_{n \rightarrow \infty} \int_{\mathbb{R}} e^{itx} d \left( \sum_{k=1}^n \frac{1}{k!} W^{*k}(x) \right), \quad t \in \mathbb{R}. \end{aligned}$$

On the other hand, on account of (33), we have

$$\frac{c_s f_s(t)}{c_d f_d(t)} = \frac{c_s}{c_d} \cdot f_s(t) \cdot g_d(t) = \frac{c_s}{c_d} \cdot \int_{\mathbb{R}} e^{itx} d(F_s * I_d)(x) = \int_{\mathbb{R}} e^{itx} d \left( \frac{c_s}{c_d} \cdot (F_s * I_d)(x) \right),$$

for  $t \in \mathbb{R}$ . Since the variance

$$\left\| \sum_{k=n_1+1}^{n_2} \frac{1}{k!} W^{*k} \right\| \leq \sum_{k=n_1+1}^{n_2} \frac{\|W^{*k}\|}{k!} \leq \sum_{k=n_1+1}^{n_2} \frac{\|W\|^k}{k!}$$

can be made arbitrarily small for all sufficiently large  $n_1$  and  $n_2$ , the sequence of sums  $\sum_{k=1}^n W^{*k}/k!, n \in \mathbb{N}$ , is fundamental in the space  $\mathbf{V}$ . Due to the completeness of  $\mathbf{V}$ , these sums converge in variation to some function from  $\mathbf{V}$ , namely, to  $(c_s/c_d) \cdot (F_s * I_d)$  by the uniqueness of the Fourier–Stieltjes transform:

$$\left\| \sum_{k=1}^n \frac{1}{k!} W^{*k} - \frac{c_s}{c_d} \cdot (F_s * I_d) \right\| \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

But this is impossible, because  $\sum_{k=1}^n W^{*k}/k!, n \in \mathbb{N}$ , are absolutely continuous as linear combinations of convolution powers of absolutely continuous  $W$  by the assumption, but  $(c_s/c_d) \cdot (F_s * I_d)$  is continuous singular as the convolution of the continuous singular function  $F_s$  and the discrete function  $I_d$  (see the comments after Remark 3). Thus  $W$  is not a (purely) absolutely continuous function from  $\mathbf{V}$ , i.e., it always has some continuous singular part.

Let us return to (47) and prove the sufficient condition (from the statement of the theorem) for  $W$  to be (purely) continuous singular. Suppose that all the functions  $F_s^{*k}, k \in \mathbb{N}$ , are continuous singular. Hence all  $W_n, n \in \mathbb{N}$  are too. Let  $W_a$  and  $W_s$

be the absolutely continuous part and the continuous singular part of the Lebesgue decomposition for  $W$ :  $W = W_a + W_s$ . Then

$$\|W_n - W\| = \|W_n - W_a - W_s\| = \|W_n - W_s\| + \|W_a\| \geq \|W_a\| \geq 0.$$

Due to (47), we conclude that  $\|W_a\| = 0$ , i.e.,  $W_a(x) = 0$  for any  $x \in \mathbb{R}$ . Thus  $W = W_s$ .

We now combine the representations (26), (27) and (46) with formula (25):

$$\begin{aligned} f(t) &= \exp \left\{ it\gamma_0 + \sum_{u \in (\mathcal{X}) \setminus \{0\}} \lambda_u (e^{itu} - 1) \right. \\ &\quad \left. + \int_{\mathbb{R} \setminus \{0\}} (e^{itx} - 1) \left( v_a(x) + \operatorname{sgn}(x) \frac{m_a e^{-|x|}}{|x|} \right) dx + \int_{\mathbb{R} \setminus \{0\}} (e^{itx} - 1) dW(x) \right\} \\ &= \exp \left\{ it\gamma_0 + \int_{\mathbb{R} \setminus \{0\}} (e^{itx} - 1) dL(x) \right\}, \quad t \in \mathbb{R}, \end{aligned}$$

with  $L(x) := L_d(x) + L_a(x) + L_s(x)$ ,  $x \in \mathbb{R} \setminus \{0\}$ , where  $L_d$ ,  $L_a$ , and  $L_s$  are defined by formulas (11), (12), and (13). It is not difficult to check that  $L_d$ ,  $L_a$ ,  $L_s$ , and, consequently,  $L$  satisfy all conditions the for a spectral function of the Lévy type representation (see introduction or Theorem 6). However, we recall that the function  $v_a$  is potentially complex-valued and hence  $L$  is too. Next, it is seen that

$$\int_{S_1} |x| d|L_*(x)| < \infty$$

with  $S_1 := [-1, 1] \setminus \{0\}$  and for any  $L_* \in \{L_d, L_a, L_s\}$ . Hence it is also true for  $L_* = L$ , and we can write

$$f(t) = \exp \left\{ it\gamma_1 + \int_{\mathbb{R} \setminus \{0\}} (e^{itx} - 1 - itx \mathbb{1}_{[-1,1]}(x)) dL(x) \right\}, \quad t \in \mathbb{R},$$

with  $\gamma_1 := \gamma_0 + \int_{\mathbb{R} \setminus \{0\}} x \mathbb{1}_{[-1,1]}(x) dL(x)$ , which is potentially complex.

We now apply Theorem 6 to derive that  $L$  is actually real-valued, which means that  $v_a$  is a real-valued function. Moreover, we have  $F \in \mathcal{Q}$ , i.e., (i) is proved.  $\square$

*Proof of Proposition 1.* To obtain a contradiction, we assume that  $W$  is continuous singular. For convenience, we introduce the operator  $[\cdot]_s$ , which acts from  $\mathbf{V}$  to  $\mathbf{V}$ , and for any  $U \in \mathbf{V}$  it returns the continuous singular part of  $U$  as  $[U]_s$  (which can be identically zero). We start with (47) from the proof of Theorem 3, which holds under the assumptions of the proposition. For every  $n \in \mathbb{N}$  we write

$$\|W_n - W\| = \|(W_n - [W_n]_s) + ([W_n]_s - W)\| = \|W_n - [W_n]_s\| + \|[W_n]_s - W\|,$$

where the latter equality is valid because  $W_n - [W_n]_s$  is absolutely continuous (or identically zero) and  $[W_n]_s - W$  is continuous singular by assumption. Therefore  $\|W_n - W\| \geq \|[W_n]_s - W\| \geq 0$  and, due to (47), we conclude that

$$(48) \quad \|[W_n]_s - W\| \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Next, according to (34), we have

$$[W_n]_s = \left( \sum_{k=1}^n \frac{(-1)^{k-1}}{k} \left( \frac{c_s}{c_d} \right)^k F_s^{*k} * I_d^{*k} \right)_s = \sum_{k=1}^n \frac{(-1)^{k-1}}{k} \left( \frac{c_s}{c_d} \right)^k [F_s^{*k} * I_d^{*k}]_s$$

for  $n \in \mathbb{N}$ . Since  $I_d$  is discrete (see (32)), the functions  $I_d^{*k}$  are discrete and hence  $[F_s^{*k} * I_d^{*k}]_s = [F_s^{*k}]_s * I_d^{*k}$  for any  $k \in \mathbb{N}$ . Thus we come to the equalities

$$(49) \quad [W_n]_s = \sum_{k=1}^n \frac{(-1)^{k-1}}{k} \left( \frac{c_s}{c_d} \right)^k [F_s^{*k}]_s * I_d^{*k}, \quad n \in \mathbb{N}.$$

Let us consider the functions  $[F_s^{*k}]_s$  and numbers  $[F_s^{*k}]_s(\infty)$ . By the definition of  $n_a$ , for any  $k < n_a$  we have  $[F_s^{*k}]_s = F_s^{*k}$  and, in particular,  $[F_s^{*k}]_s(\infty) = 1$ . For the case  $k = n_a$ , by the assumption, we have  $[F_s^{*n_a}]_s(x) = (1 - \alpha)H_s(x)$ ,  $x \in \mathbb{R}$ , and hence  $[F_s^{*n_a}]_s(\infty) = 1 - \alpha$ . Next, for any integer  $k \geq n_a$  we observe that

$$[F_s^{*(k+1)}]_s(x) = [F_s^{*k} * F_s]_s(x) = [[F_s^{*k}]_s * F_s]_s(x) \leq ([F_s^{*k}]_s * F_s)(x), \quad x \in \mathbb{R}.$$

By the Lebesgue dominated convergence theorem, we conclude that

$$([F_s^{*k}]_s * F_s)(\infty) = \lim_{z \rightarrow \infty} \int_{\mathbb{R}} F_s(z-x) d[F_s^{*k}]_s(x) = \int_{\mathbb{R}} d[F_s^{*k}]_s(x) = [F_s^{*k}]_s(\infty).$$

Thus  $[F_s^{*(k+1)}]_s(\infty) \leq [F_s^{*k}]_s(\infty)$  for any integer  $k \geq n_a$ . Let us introduce the sequence  $A_k := [F_s^{*k}]_s(\infty)$ ,  $k \in \mathbb{N}$ . So we have

$$(50) \quad \begin{aligned} A_1 = \dots = A_{n_a-1} = 1, \quad A_{n_a} = 1 - \alpha, \\ A_k \geq A_{k+1} \geq 0 \quad \text{for any } k \geq n_a. \end{aligned}$$

Due to (48), we next conclude that

$$\int_{\mathbb{R}} e^{itx} dW(x) = \lim_{n \rightarrow \infty} \int_{\mathbb{R}} e^{itx} d[W_n]_s(x), \quad t \in \mathbb{R}.$$

Then, according to (45) and (49), we have

$$\ln \left( 1 + \frac{c_s f_s(t)}{c_d f_d(t)} \right) = \lim_{n \rightarrow \infty} \sum_{k=1}^n \frac{(-1)^{k-1}}{k} \left( \frac{c_s}{c_d} \right)^k f_{k,s}(t) g_d(t)^k \quad \text{for any } t \in \mathbb{R},$$

where

$$f_{k,s}(t) := \int_{\mathbb{R}} e^{itx} d[F_s^{*k}]_s(x), \quad t \in \mathbb{R}.$$

In particular, we write

$$\ln \left( 1 + \frac{c_s f_s(0)}{c_d f_d(0)} \right) = \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k} \left( \frac{c_s}{c_d} \right)^k f_{k,s}(0) g_d(0)^k.$$

Recall that  $f_s(0) = f_d(0) = g_d(0) = 1$  and  $f_{k,s}(0) = [F_s^{*k}]_s(\infty) = A_k$  for any  $k \in \mathbb{N}$ . Then, on the one hand,

$$\ln\left(1 + \frac{c_s}{c_d}\right) = \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k} \left(\frac{c_s}{c_d}\right)^k A_k.$$

On the other hand, since  $c_s < c_d \mu_d \leq c_d$ , we have the expansion

$$\ln\left(1 + \frac{c_s}{c_d}\right) = \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k} \left(\frac{c_s}{c_d}\right)^k.$$

Thus we come to the equality

$$\sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k} \left(\frac{c_s}{c_d}\right)^k A_k = \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k} \left(\frac{c_s}{c_d}\right)^k.$$

Due to (50), it is reduced to

$$(51) \quad \sum_{k=n_a}^{\infty} \frac{(-1)^{k-n_a}}{k} \left(\frac{c_s}{c_d}\right)^k A_k = \sum_{k=n_a}^{\infty} \frac{(-1)^{k-n_a}}{k} \left(\frac{c_s}{c_d}\right)^k.$$

The series on the left is alternating with nonincreasing absolute values of the terms. Therefore

$$\sum_{k=n_a}^{\infty} \frac{(-1)^{k-n_a}}{k} \left(\frac{c_s}{c_d}\right)^k A_k \leq \frac{1}{n_a} \left(\frac{c_s}{c_d}\right)^{n_a} A_{n_a} = \frac{1}{n_a} \left(\frac{c_s}{c_d}\right)^{n_a} (1 - \alpha).$$

Since the series on the right in (51) is alternating with strictly decreasing absolute values, the following inequality holds:

$$\sum_{k=n_a}^{\infty} \frac{(-1)^{k-n_a}}{k} \left(\frac{c_s}{c_d}\right)^k > \frac{1}{n_a} \left(\frac{c_s}{c_d}\right)^{n_a} - \frac{1}{n_a+1} \left(\frac{c_s}{c_d}\right)^{n_a+1} = \frac{1}{n_a} \left(\frac{c_s}{c_d}\right)^{n_a} \left(1 - \frac{n_a}{n_a+1} \cdot \frac{c_s}{c_d}\right).$$

We see that the assumption  $\alpha \geq \frac{n_a}{n_a+1} \cdot \frac{c_s}{c_d}$  implies the inequality

$$\sum_{k=n_a}^{\infty} \frac{(-1)^{k-n_a}}{k} \left(\frac{c_s}{c_d}\right)^k A_k < \sum_{k=n_a}^{\infty} \frac{(-1)^{k-n_a}}{k} \left(\frac{c_s}{c_d}\right)^k.$$

Thus we come to a contradiction with (51). So the assumption that  $W$  is continuous singular is false.

Due to the condition of dominated continuous singular part, we have  $c_s < c_d$ . Since  $\frac{n_a}{n_a+1} < 1$ , the inequality  $\alpha \geq \frac{n_a}{n_a+1} \cdot \frac{c_s}{c_d}$  always holds in the case  $\alpha = 1$ .  $\square$

*Proof of Proposition 2.* Let us consider the function  $f_s$  defined by formula (15). We first find the limit of the sequence  $f_s(t_n)$ ,  $n \in \mathbb{N}$ , with  $t_n := \pi(2n)!$  as  $n \rightarrow \infty$ . We write  $f_s(t_n) = M_n \cdot R_n$ ,  $n \in \mathbb{N}$ , where

$$M_n := \prod_{k=1}^{2n} \cos(t_n/k!) \quad \text{and} \quad R_n := \prod_{k=2n+1}^{\infty} \cos(t_n/k!), \quad n \in \mathbb{N}.$$

For any  $n \in \mathbb{N}$

$$\begin{aligned} M_n &= \cos\left(\frac{\pi(2n)!}{(2n)!}\right) \cdot \cos\left(\frac{\pi(2n)!}{(2n-1)!}\right) \cdot \cos\left(\frac{\pi(2n)!}{(2n-2)!}\right) \cdot \dots \cdot \cos\left(\frac{\pi(2n)!}{1!}\right) \\ &= \cos(\pi) \cdot \cos(\pi \cdot 2n) \cdot \cos(\pi \cdot 2n(2n-1)) \cdot \dots \cdot \cos(\pi \cdot 2n(2n-1) \cdot \dots \cdot 1) \\ &= (-1) \cdot 1 \cdot 1 \cdot \dots \cdot 1 = -1. \end{aligned}$$

Next, for every  $n \in \mathbb{N}$  and for every integer  $k \geq 2n+1$  we have

$$0 < \frac{t_n}{k!} = \frac{\pi(2n)!}{k \cdot (k-1)!} \leq \frac{\pi}{k} \leq \frac{\pi}{3},$$

and, by the well-known inequality  $\cos(x) \geq 1 - x^2/2$ ,  $x \in \mathbb{R}$ , we get

$$\cos(t_n/k!) \geq 1 - \frac{(t_n/k!)^2}{2} \geq 1 - \frac{\pi^2}{2k^2} \geq 1 - \frac{\pi^2}{18} > 0.$$

Therefore, on the one hand, it is clear that for any  $n \in \mathbb{N}$  we have  $\cos(t_n/k!) < 1$  for any  $k \geq 2n+1$  and hence  $R_n < 1$ . On the other hand,

$$R_n \geq \prod_{k=2n+1}^{\infty} \left(1 - \frac{\pi^2}{2k^2}\right) = \exp\left\{\sum_{k=2n+1}^{\infty} \ln\left(1 - \frac{\pi^2}{2k^2}\right)\right\}, \quad n \in \mathbb{N},$$

where the sum in the exponent tends to 0 as  $n \rightarrow \infty$ . So we conclude that  $R_n \rightarrow 1$  as  $n \rightarrow \infty$ . Thus

$$f_s(t_n) = M_n \cdot R_n = -R_n \rightarrow -1 \quad \text{as} \quad n \rightarrow \infty,$$

and we know that  $f_s(t_n) > -1$  for any  $n \in \mathbb{N}$ .

We next observe that

$$f_s(t_n \pm \pi) = \prod_{k=1}^{\infty} \cos\left(\frac{t_n \pm \pi}{k!}\right) = 0.$$

Indeed,

$$\cos\left(\frac{t_n \pm \pi}{k!}\right) \Big|_{k=2} = \cos\left(\frac{\pi(2n)! \pm \pi}{2!}\right) = \cos\left(\pi \cdot 3 \cdot \dots \cdot (2n) \pm \frac{\pi}{2}\right) = 0.$$

We now return to the characteristic function  $f_*(t) = \frac{1}{2} + \frac{1}{2} f_s(t)$ ,  $t \in \mathbb{R}$ , and we consider the quantities

$$\frac{f_*(t_n - \pi) f_*(t_n + \pi)}{f_*(t_n)^2} = \frac{(\frac{1}{2} + \frac{1}{2} f_s(t_n - \pi))(\frac{1}{2} + \frac{1}{2} f_s(t_n + \pi))}{(\frac{1}{2} + \frac{1}{2} f_s(t_n))^2}, \quad n \in \mathbb{N}.$$

By the above,

$$\frac{f_*(t_n - \pi) f_*(t_n + \pi)}{f_*(t_n)^2} = \frac{1}{(1 + f_s(t_n))^2} \rightarrow \infty \quad \text{as } n \rightarrow \infty.$$

Therefore, by Theorem 7,  $f_*$  cannot be the characteristic function of a distribution function from class  $\mathcal{Q}$ . Thus  $F_* \notin \mathcal{Q}$ .  $\square$

*Proof of Proposition 3.* We have decompositions (3) and (4) for the distribution function  $F$  and its characteristic function  $f$ , respectively. Since  $F \in \mathcal{Q}$ , we know that  $f(t) \neq 0$  for any  $t \in \mathbb{R}$  and  $\mu_d = \inf_{t \in \mathbb{R}} |f_d(t)| > 0$  according to equation (9). Due to the assumption of dominated singular part and  $c_d > 0$ , we have  $c_s < c_d \mu_d$ .

Let  $f_1$  and  $f_2$  denote the characteristic functions of  $F_1$  and  $F_2$ . The assumed decomposition  $F = F_1 * F_2$  means that  $f(t) = f_1(t) f_2(t)$ ,  $t \in \mathbb{R}$ . This means that  $f_1(t) \neq 0$  and  $f_2(t) \neq 0$  for all  $t \in \mathbb{R}$ .

Let us write the Lebesgue decompositions for  $F_1$  and  $F_2$ :

$$F_j(x) = c_{j,d} F_{j,d}(x) + c_{j,a} F_{j,a}(x) + c_{j,s} F_{j,s}(x), \quad x \in \mathbb{R}, \quad j \in \{1, 2\}.$$

We also write the corresponding decompositions for  $f_1$  and  $f_2$ :

$$f_j(t) = c_{j,d} f_{j,d}(t) + c_{j,a} f_{j,a}(t) + c_{j,s} f_{j,s}(t), \quad t \in \mathbb{R}, \quad j \in \{1, 2\}.$$

Here  $c_{j,d}$ ,  $c_{j,a}$ ,  $c_{j,s}$  are nonnegative and  $c_{j,d} + c_{j,a} + c_{j,s} = 1$  for  $j = 1$  and  $j = 2$ . For clarity, we write the equality  $F = F_1 * F_2$  in the expanded form:

$$(52) \quad c_d F_d + c_a F_a + c_s F_s \\ = (c_{1,d} F_{1,d} + c_{1,a} F_{1,a} + c_{1,s} F_{1,s}) * (c_{2,d} F_{2,d} + c_{2,a} F_{2,a} + c_{2,s} F_{2,s}).$$

Since  $F$  has nonzero discrete part ( $c_d > 0$ ), the functions  $F_1$  and  $F_2$  have nonzero discrete parts too, i.e.,  $c_{1,d} > 0$  and  $c_{2,d} > 0$ . Since a convolution of any two distribution functions is discrete if and only if these functions are discrete, we conclude that  $c_d F_d(x) = c_{1,d} c_{2,d} (F_{1,d} * F_{2,d})(x)$ ,  $x \in \mathbb{R}$ , i.e.,  $c_d = c_{1,d} c_{2,d}$  and  $F_d = F_{1,d} * F_{2,d}$ . Thus we have  $f_d(t) = f_{1,d}(t) f_{2,d}(t)$ ,  $t \in \mathbb{R}$ . Since  $|f_{1,d}(t)| \leq 1$  and  $|f_{2,d}(t)| \leq 1$  for any  $t \in \mathbb{R}$ , we conclude that

$$\mu_{1,d} := \inf_{t \in \mathbb{R}} |f_{1,d}(t)| \geq \inf_{t \in \mathbb{R}} |f_d(t)| = \mu_d > 0,$$

and, analogously,

$$\mu_{2,d} := \inf_{t \in \mathbb{R}} |f_{2,d}(t)| \geq \inf_{t \in \mathbb{R}} |f_d(t)| = \mu_d > 0.$$

We next observe that  $F_{1,s} * F_{2,d}$  and  $F_{2,s} * F_{1,d}$  are continuous singular. Therefore the corresponding summands from (52) are included in the continuous part of  $F$ , i.e.,  $c_s F_s(x) \geq c_{1,s} c_{2,d} (F_{1,s} * F_{2,d})(x) + c_{2,s} c_{1,d} (F_{2,s} * F_{1,d})(x)$ ,  $x \in \mathbb{R}$ . Consequently,

$$\begin{aligned} c_s &= c_s F_s(\infty) \geq c_{1,s} c_{2,d} (F_{1,s} * F_{2,d})(\infty) + c_{2,s} c_{1,d} (F_{2,s} * F_{1,d})(\infty) \\ &= c_{1,s} c_{2,d} + c_{2,s} c_{1,d}. \end{aligned}$$

Then we get

$$c_{1,s} c_{2,d} \leq c_s < c_d \mu_d = c_{1,d} c_{2,d} \mu_d \leq c_{1,d} c_{2,d} \mu_{1,d},$$

i.e.,  $c_{1,s} < c_{1,d} \mu_{1,d}$ . Analogously,

$$c_{2,s} c_{1,d} \leq c_s < c_d \mu_d = c_{1,d} c_{2,d} \mu_d \leq c_{1,d} c_{2,d} \mu_{2,d},$$

i.e.,  $c_{2,s} < c_{2,d} \mu_{2,d}$ . Thus  $F_1$  and  $F_2$  have dominated continuous singular parts.

We have shown that  $F_1$  and  $F_2$  satisfy the assumptions of Theorem 3 and condition (iii) from it. Thus, according to that theorem,  $F_1$  and  $F_2$  belong to class  $\mathcal{Q}$ .  $\square$

## References

- [1] I. A. Alexeev and A. A. Khartov, “On convergence and compactness in variation with a shift of discrete probability laws”, *Vestn. St.-Peterbg. Univ. Mat. Mekh. Astron.* **8**:3 (2021), 385–393. In Russian; translated in *Vestn. St. Petersburg. Univ. Mat.* **54**:3 (2021), 221–226. MR
- [2] I. A. Alexeev and A. A. Khartov, “A criterion and a Cramér–Wold device for quasi-infinite divisibility for discrete multivariate probability laws”, *Electron. J. Probab.* **28** (2023), art. id. 130, 17 pp. MR
- [3] I. A. Alexeev and A. A. Khartov, “Spectral representations of characteristic functions of discrete probability laws”, *Bernoulli* **29**:2 (2023), 1392–1409. MR
- [4] D. Applebaum, *Lévy processes and stochastic calculus*, 2nd ed., Cambridge Studies in Advanced Mathematics **116**, Cambridge University Press, 2009. MR
- [5] G. Baxter and J. M. Shapiro, “On bounded infinitely divisible random variables”, *Sankhyā* **22**:3–4 (1960), 253–260. MR
- [6] D. Berger, “On quasi-ininitely divisible distributions with a point mass”, *Math. Nachr.* **292**:8 (2019), 1674–1684. MR
- [7] D. Berger and M. Kutlu, “Quasi-infinite divisibility of a class of distributions with discrete part”, *Proc. Amer. Math. Soc.* **151**:5 (2023), 2211–2224. MR
- [8] D. Berger, M. Kutlu, and A. Lindner, “On multivariate quasi-ininitely divisible distributions”, pp. 87–120 in *A lifetime of excursions through random walks and Lévy processes: a volume in honour of Ron Doney’s 80th birthday*, Progr. Probab. **78**, Birkhäuser, 2021. MR
- [9] S. Bochner, “A theorem on Fourier–Stieltjes integrals”, *Bull. Amer. Math. Soc.* **40**:4 (1934), 271–276. MR
- [10] H. Chhaiba, N. Demni, and Z. Mouayn, “Analysis of generalized negative binomial distributions attached to hyperbolic Landau levels”, *J. Math. Phys.* **57**:7 (2016), art. id. 072103, 14 pp. MR
- [11] I. Gelfand, D. Raikov, and G. Shilov, *Commutative normed rings*, Chelsea, New York, 1964. MR

- [12] B. V. Gnedenko and A. N. Kolmogorov, *Limit distributions for sums of independent random variables*, Addison-Wesley, Cambridge, MA, 1954. MR
- [13] M. Hardy, “Combinatorics of partial derivatives”, *Electron. J. Combin.* **13**:1 (2006), art. id. 1, 13 pp. MR
- [14] E. Hewitt and H. S. Zuckerman, “Singular measures with absolutely continuous convolution squares”, *Proc. Cambridge Philos. Soc.* **62** (1966), 399–420. MR
- [15] O. S. Ivashchev-Musatov, “On coefficients of trigonometric null-series”, *Izv. Akad. Nauk SSSR Ser. Mat.* **21** (1957), 559–578. In Russian. MR
- [16] B. Jessen and A. Wintner, “Distribution functions and the Riemann zeta function”, *Trans. Amer. Math. Soc.* **38**:1 (1935), 48–88. MR
- [17] A. A. Khartov, “Compactness criteria for quasi-infinitely divisible distributions on the integers”, *Statist. Probab. Lett.* **153** (2019), 1–6. MR
- [18] A. A. Khartov, “A criterion of quasi-infinite divisibility for discrete laws”, *Statist. Probab. Lett.* **185** (2022), art. id. 109436, 4 pp. MR
- [19] A. A. Khartov, “On weak convergence of quasi-infinitely divisible laws”, *Pacific J. Math.* **322**:2 (2023), 341–367. MR
- [20] A. A. Khartov, “Some criteria of rational-infinite divisibility for probability laws”, *Electron. J. Probab.* **30** (2025), art. id. 111, 29 pp. MR
- [21] A. A. Khartov, “On decomposition problem for distribution functions of class  $\mathcal{Q}$ ”, *Proc. Amer. Math. Soc.* **154**:1 (2026), 405–419. MR
- [22] A. A. Khartov and I. A. Alekseev, “Quasi-infinite divisibility and three-point probability laws”, *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)* **495** (2020), 305–316. In Russian; translated in *J. Math. Sci.* **268** (2022), 731–738. MR
- [23] M. G. Krein, “Integral equations on the half-line with a kernel depending on the difference of the arguments”, *Uspehi Mat. Nauk* **13**:83 (1958), 3–120. In Russian. MR
- [24] B. M. Levitan, Почти-периодические функции, Gosudar Izdat. Tehn.-Teor. Lit., Moscow, 1953. MR
- [25] M. A. Lifshits, *Random processes by example*, World Scientific, Hackensack, NJ, 2014. MR
- [26] A. Lindner, L. Pan, and K.-I. Sato, “On quasi-infinitely divisible distributions”, *Trans. Amer. Math. Soc.* **370**:12 (2018), 8483–8520. MR
- [27] Y. V. Linnik and I. V. Ostrovskii, *Decomposition of random variables and vectors*, Translations of Mathematical Monographs **48**, Amer. Math. Soc., 1977. MR
- [28] E. Lukacs, *Characteristic functions*, 2nd ed., Griffin, London, 1970. MR
- [29] T. Nakamura, “A complete Riemann zeta distribution and the Riemann hypothesis”, *Bernoulli* **21**:1 (2015), 604–617. MR
- [30] R. Passeggeri, “On quasi-infinitely divisible random measures”, *Bayesian Anal.* **18**:1 (2023), 253–286. MR
- [31] K.-i. Sato, *Lévy processes and infinitely divisible distributions*, Cambridge Studies in Advanced Mathematics **68**, Cambridge University Press, 1999. MR
- [32] W. Schoutens, *Lévy processes in finance: pricing financial derivatives*, Wiley, Chichester, 2003.
- [33] Y. A. Shreider, “The structure of maximal ideals in rings of measures with convolution”, *Mat. Sbornik N.S.* **69** (1950), 297–318. MR
- [34] H. G. Tucker, “On a necessary and sufficient condition that an infinitely divisible distribution be absolutely continuous”, *Trans. Amer. Math. Soc.* **118** (1965), 316–330. MR

- [35] N. Wiener and H. R. Pitt, “On absolutely convergent Fourier–Stieltjes transforms”, *Duke Math. J.* **4**:2 (1938), 420–436. MR
- [36] N. Wiener and A. Wintner, “Fourier–Stieltjes transforms and singular infinite convolutions”, *Amer. J. Math.* **60**:3 (1938), 513–522. MR
- [37] V. M. Zolotarev, *Modern theory of summation of random variables*, VSP, Utrecht, 1997. MR

Received June 17, 2025. Revised March 18, 2026.

ALEXEY A. KHARTOV

INSTITUTE FOR INFORMATION TRANSMISSION PROBLEMS RAS (KHARKEVICH INSTITUTE) OF  
THE RUSSIAN ACADEMY OF SCIENCES

MOSCOW

RUSSIA

and

SMOLENSK STATE UNIVERSITY

SMOLENSK

RUSSIA

khartov.a@iitp.ru



# DATA FOR SHIMURA VARIETIES INTERSECTING THE TORELLI LOCUS

WANLIN LI, ELENA MANTOVAN AND RACHEL PRIES

**For infinitely many Hurwitz spaces parametrizing cyclic covers of the projective line, we provide a method to determine the integral PEL datum of the Shimura variety that contains the image of the Hurwitz space under the Torelli morphism.**

## 1. Introduction

**1.1. Overview.** In [15], Shimura studied unitary groups associated with Hermitian spaces over algebraic number fields and their maximal lattices. In [14], he developed this theory to study isomorphism classes of polarized abelian varieties and Riemann forms. Using this, in [16], Shimura determined the lattice and Hermitian matrix associated with each of six families of cyclic covers of the projective line  $\mathbb{P}^1$ . The lattice and Hermitian matrix determine *the integral PEL datum* of the family, as defined in Section 2.5.

For  $m$  an odd prime such that  $\mathbb{Q}(\zeta_m)$  has class number one, we provide in Theorem 5.5 a method to determine the lattice and Hermitian matrix, and thus the integral PEL datum, for all positive-dimensional families of degree  $m$  cyclic covers of  $\mathbb{P}^1$ .

Our results compute the integral PEL datum of certain unitary Shimura varieties that contain Hurwitz spaces of cyclic covers of the projective line.

---

We would like to thank the American Institute of Mathematics for their support through the Square program. Li was partially supported by the Simons Collaboration on Arithmetic Geometry, Number Theory and Computation and by NSF grant DMS-2302511. Mantovan was partially supported by NSF grant DMS-22-00694. Pries was partially supported by NSF grants DMS-19-01819 and DMS-22-00418.

We are thankful to Yunqing Tang for her support and advice on this paper. We also thank Eran Assaf, Bjorn Poonen, and John Voight for helpful conversations about class groups. We thank the referee for helpful comments.

*MSC2020:* primary 11G15, 11G18, 14H10, 14K10, 14K22; secondary 11G10, 11G30, 11R18, 14G35, 14H40.

*Keywords:* abelian variety, curve, Jacobian, complex multiplication, moduli space, Hurwitz space, Shimura variety, Torelli locus, PEL type, lattice, Hermitian form, cyclotomic field, class group.

**1.2. Results.** Consider a family  $H_\gamma$  of cyclic covers of the projective line  $\mathbb{P}^1$ , indexed by the monodromy datum  $\gamma = (m, N, a)$ , where  $m$  is the degree,  $N$  is the number of branch points, and  $a$  is the inertia type (Section 2.1). Let  $g = g_\gamma$  be the genus of the curves in this family; see (2-1).

For  $g \geq 1$ , let  $\mathcal{A}_g$  be the moduli space of principally polarized abelian varieties of dimension  $g$ . Let  $Z_\gamma$  be the image of the family  $H_\gamma$  in  $\mathcal{A}_g$  under the Torelli morphism. In [12, Section 3], Moonen describes the smallest PEL type Shimura substack  $S_\gamma$  of  $\mathcal{A}_g$  containing  $Z_\gamma$ . Our goal is to compute the Shimura datum of  $S_\gamma$  or, more precisely, the *integral PEL datum* of  $S_\gamma$  as defined in Definition 2.2. We use [17, Proposition 4.1], which shows that the integral PEL datum of  $S_\gamma$  can be computed at any point  $P$  of  $Z_\gamma$ .

Suppose  $m$  is an odd prime such that  $\mathbb{Q}(\zeta_m)$  has class number 1. In other words,  $m \in \{3, 5, 7, 11, 13, 17, 19\}$ . In Theorem 5.5, for every monodromy datum  $\gamma$  for degree  $m$  covers, we provide a method that determines the integral PEL datum of  $S_\gamma$ . Our approach is to find a point  $P \in Z_\gamma$  for which we can analyze the Jacobian  $J_P$  represented by  $P$  and the lattice  $\Lambda = H^1(J_P, \mathbb{Z})$ . From this, we can

- (1) express  $V = \mathbb{Q}^{2g}$  as a vector space over  $F = \mathbb{Q}(\zeta_m)$ , find the  $\mathcal{O}_F$ -lattice  $\Lambda \subset V$ , and
- (2) explicitly find the Hermitian form  $\langle \cdot, \cdot \rangle$  on  $V$ , which takes integral values on  $\Lambda$ .

Our technique also applies to infinitely many monodromy data  $\gamma$  when  $m$  is a composite number such that  $\mathbb{Q}(\zeta_m)$  has class number 1; see Section 7.

Specifically, we choose  $P$  to be a *distinguished point* of  $Z_\gamma$ , meaning that the Jacobian  $J_P$  represented by  $P$  is a product of principally polarized abelian varieties each having complex multiplication, see Definition 5.1. For every monodromy datum  $\gamma$  with  $m$  an odd prime, we prove in Proposition 5.2 that  $Z_\gamma$  has a distinguished point  $P$  such that each of the abelian varieties in the product has complex multiplication by  $\mathbb{Z}[\zeta_m]$ . In fact,  $J_P$  is the Jacobian of a singular curve in the boundary of the Hurwitz family.

**1.3. Comparison of methods.** Our approach to this problem is different from Shimura's and may be more accessible to people with background in the areas of algebraic number theory and moduli spaces of curves (Hurwitz spaces).

One advantage of our approach is that it provides a straightforward method to determine the lattice and the Hermitian form explicitly. We provide many examples of this in Section 6 (for  $m$  prime) and Section 7 (for  $m$  composite). The reason is that the decomposition of  $J_P$  provides a basis for the lattice as a free  $\mathbb{Z}[\zeta_m]$ -module. The Hermitian form is diagonal with respect to that basis and is determined by the CM-types of the factors of  $J_P$ , which can be explicitly computed from  $\gamma$ . In Remark 6.6 we give more careful attention to the subtleties caused by the choice of primitive  $m$ -th root of unity.

With Shimura's approach, it is necessary to find a Witt decomposition of the signature and lattice; this is straightforward for Shimura's six examples, five of which are of the form  $y^m = f(x)$  for  $m$  an odd prime and  $f(x)$  a separable polynomial, but could cause subtleties for more complicated families.

One drawback to our approach is that it only works when the CM-types produced in the method are simple. The simple property guarantees that the principal polarization on each of the abelian varieties is unique up to isomorphism; this guarantees that the Hermitian form we compute is correct. We use several methods to address this issue.

For composite  $m$ , there are several complications with both approaches.

**1.4. Outline.** Section 2 contains information about families of cyclic covers of  $\mathbb{P}^1$  and information about Shimura varieties and Shimura data.

Section 3 contains some results in algebraic number theory about the narrow class group and units of independent signs that we need for this paper and later papers.

In Section 4, we prove a result about uniqueness of principal polarizations on abelian varieties with complex multiplication (Propositions 4.4 and 4.5).

Section 5 contains the main result of the paper, Theorem 5.5, which determines the integral PEL data of the unitary Shimura varieties  $S_\gamma$  for all families  $S_\gamma$  when  $m$  is an odd prime such that  $\mathbb{Q}(\zeta_m)$  has class number 1.

Section 6 contains examples of Theorem 5.5: for all  $\gamma$  when  $m = 3$ ; and for all  $\gamma$  with  $N = 4$  branch points when  $m = 5$  and  $m = 7$ . In Section 6.4, we apply the technique of Theorem 5.5 to determine the integral PEL datum using a different kind of distinguished point that represents the Jacobian of a curve with extra automorphisms.

In Section 7, we provide some examples when  $m = 4, 6, 10$  to illustrate that the technique of Theorem 5.5 sometimes works when  $m$  is composite.

**Remark 1.1.** In [12, Theorem 3.6], Moonen proved that there are exactly 20 equivalence classes of  $\gamma$  for which  $Z_\gamma$  is open and dense in  $S_\gamma$ ; in this situation, the family is called *special*. The six families in Shimura's paper are  $M[n]$  when  $n = 6, 10, 8, 11, 16, 17$ , where  $M[n]$  denotes the  $n$ -th row in [12, Table 1]. As an application of Theorem 5.5, we determine the integral PEL datum for 12 of these 20 families, including the 6 families from [16]. We emphasize that the special property is not necessary for either approach.

We would like to thank the referee for this observation: when  $\dim(Z_\gamma) < \dim(S_\gamma)$ , the Torelli locus gives a cycle on  $S_\gamma$  which is not obviously associated to a Shimura subvariety.

## 2. Hurwitz families and Shimura varieties

**2.1. Families of cyclic covers of the projective line.** This section is a shortened version of [8, Section 2.2]. Consider the universal family of  $\mu_m$ -covers  $\psi : \mathcal{C} \rightarrow \mathbb{P}^1$ , branched at  $N$  points, with inertia type  $a = (a(1), \dots, a(N))$ ; the data  $\gamma = (m, N, a)$  is called a *monodromy datum*. Over  $\mathbb{Q}(\zeta_m)$ , such a cover  $\psi$  has an equation of the form  $y^m = \prod_{i=1}^N (x - t(i))^{a(i)}$ , and a chosen automorphism  $h$  of order  $m$  given by  $(x, y) \mapsto (x, \zeta_m y)$ . By the Riemann–Hurwitz formula, the genus of the fibers of  $\mathcal{C}$  is

$$(2-1) \quad g = g_\gamma = 1 + \frac{1}{2} \left( (N - 2)m - \sum_{i=1}^N \gcd(a(i), m) \right)$$

For  $0 \leq n < m$ , let  $\tau_n : \mathbb{Q}[\mu_m] \rightarrow \mathbb{C}$  be given by  $\tau_n(\zeta_m) = e^{2\pi i n/m}$ . We identify  $\mathbb{Z}/m\mathbb{Z} = \text{Hom}(\mathbb{Q}[\mu_m], \mathbb{C})$  by  $n \mapsto \tau_n$ . The *signature type* of  $\gamma$  is a function  $\mathfrak{f} : \text{Hom}(\mathbb{Q}[\mu_m], \mathbb{C}) \rightarrow \mathbb{Z}_{\geq 0}$  which we denote by  $\mathfrak{f} = (\mathfrak{f}(\tau_1), \dots, \mathfrak{f}(\tau_{m-1}))$ , where  $\mathfrak{f}(\tau_n)$  is the dimension of the eigenspace of  $H^0(\mathcal{C}(\mathbb{C}), \Omega^1)$  on which  $h$  acts by multiplication by  $\zeta_m^n$ .

For any  $x \in \mathbb{Q}$ , let  $\langle x \rangle$  denote the fractional part of  $x$ . By [12, Lemma 2.7, §3.2],

$$(2-2) \quad \mathfrak{f}(\tau_n) = \begin{cases} -1 + \sum_{i=1}^N \left\langle \frac{-na(i)}{m} \right\rangle & \text{if } n \not\equiv 0 \pmod{m}, \\ 0 & \text{if } n \equiv 0 \pmod{m}. \end{cases}$$

We describe how the inertia type  $a$  and signature  $\mathfrak{f}$  change under the action of  $\text{Aut}(\mu_m) \simeq (\mathbb{Z}/m\mathbb{Z})^*$ . Let  $\sigma_i \in \text{Aut}(\mu_m)$  denote the automorphism such that  $\sigma_i(\zeta_m) = \zeta_m^i$ .

**Lemma 2.1.** *The automorphism  $\sigma_i \in \text{Aut}(\mu_m)$  takes the inertia type*

$$a = (a_1, a_2, \dots, a_N) \quad \text{to} \quad a' = i^{-1} \cdot a = (i^{-1} \cdot a_1, i^{-1} \cdot a_2, \dots, i^{-1} \cdot a_N)$$

*and the signature*

$$\mathfrak{f} \quad \text{to} \quad \mathfrak{f}', \quad \text{where } \mathfrak{f}'(\tau_n) = \mathfrak{f}(\tau_{ni^{-1}}).$$

*Proof.* The  $j$ -th entry  $a_j$  of the inertia type signifies that the canonical generator of inertia above the  $j$ -th branch point is the  $a_j$ -th power of the generator  $\zeta_m$  of  $\mu_m$ . The canonical generator of inertia is the  $i^{-1}a_j$ -th power of the new generator  $\zeta_m^i$  of  $\mu_m$ , so the  $j$ -th entry of the inertia type  $a'$  is  $i^{-1}a_j$ .

The automorphism  $\sigma_i$  permutes the eigenspaces for the action of  $h$  on  $H^0(\mathcal{C}(\mathbb{C}), \Omega^1)$ , taking the eigenspace indexed by  $n$  to the one indexed by  $ni^{-1}$ . This yields the formula  $\mathfrak{f}'(\tau_n) = \mathfrak{f}(\tau_{ni^{-1}})$ , which can also be deduced from the formula for  $a'$  and (2-2). □

Consider the Hurwitz family of  $\mu_m$ -covers of  $\mathbb{P}^1$  with monodromy datum  $\gamma = (m, N, a)$ . As in [8, Definition 2.1], let  $Z^0 = Z_\gamma^0$  denote the image of this family in  $\mathcal{A}_g$ . Let  $Z = Z_\gamma$  denote the closure of  $Z_\gamma^0$  in  $\mathcal{A}_g$ .

**2.2. Complex abelian varieties.** Let  $g \geq 1$  and let  $V = \mathbb{R}^{2g}$ . Suppose  $V$  has a complex structure and  $\Lambda \subset V$  is a lattice of rank  $2g$ . A Riemann form on a pair  $(V, \Lambda)$  is an alternating pairing  $\Psi : \Lambda \times \Lambda \rightarrow \mathbb{Z}$  such that the pairing  $\Psi_{\mathbb{R}}(\cdot, \sqrt{-1}\cdot) : V \times V \rightarrow \mathbb{R}$  is symmetric and positive definite. Every complex abelian variety of dimension  $g$  is isomorphic to  $V/\Lambda$  for some pair  $(V, \Lambda)$  that admits a Riemann form.

By Riemann’s theorem, the category of polarized abelian varieties of dimension  $g$  over  $\mathbb{C}$  is equivalent to the category of pairs  $(V, \Lambda)$ , where  $V$  is a nontrivial  $\mathbb{C}$ -vector space of dimension  $g$  and  $\Lambda$  is a lattice in  $V$  together with a Riemann form integral on  $\Lambda$ .

**2.3. Siegel modular varieties.** For  $g \geq 1$ , let  $V = \mathbb{Q}^{2g}$  together with the standard lattice  $\Lambda = \mathbb{Z}^{2g}$ . Let  $\Psi : V \times V \rightarrow \mathbb{Q}$  be the standard symplectic form, which is integral on  $\Lambda$ . The points of the Siegel upper half space  $\mathcal{H}_g$  are complex structures  $J$  on  $V_{\mathbb{R}}$  such that  $\Psi_J = \Psi_{\mathbb{R}}(\cdot, J\cdot)$  is a Riemann form. These points parametrize principally polarized complex abelian varieties  $A$  of dimension  $g$ , equipped with a trivialization  $\Lambda \simeq H^1(A, \mathbb{Z})$ . The Shimura datum for  $\mathcal{A}_g$  arises from the symplectic  $\mathbb{Q}$ -vector space  $(V, \Psi)$ ; its integral datum given by the self-dual  $\mathbb{Z}$ -lattice  $\Lambda$ . The associated algebraic group over  $\mathbb{Q}$  is  $G_g := \mathrm{GSp}(V, \Psi)$ , the algebraic group of symplectic similitudes; see [11, Example 11.12].

**2.4. Shimura data of PEL type.** A Shimura datum of PEL type is a Shimura datum arising from a PEL datum  $(B, *, V, \langle \cdot, \cdot \rangle, h)$ , where (see [5, Section 4])

- $B$  is a semisimple finite dimensional  $\mathbb{Q}$ -algebra;
- $*$  is a positive involution on  $B$ ;
- $V$  is a finitely generated  $B$ -module;
- $\langle \cdot, \cdot \rangle$  is a nondegenerate skew-symmetric  $*$ -Hermitian form on  $V$ ; and
- $h : \mathbb{C} \rightarrow \mathrm{End}_{B \otimes \mathbb{R}}(V_{\mathbb{R}})$  is an  $\mathbb{R}$ -algebra homomorphism satisfying  $\langle h(z)\cdot, \cdot \rangle = \langle \cdot, h(\bar{z})\cdot \rangle$ , for all  $z \in \mathbb{C}$ ; and such that the symmetric bilinear form  $\langle \cdot, h(i)\cdot \rangle$  on  $V$  is positive definite.

The associated algebraic group  $H$  over  $\mathbb{Q}$  is  $H = \mathrm{GL}_B(V) \cap \mathrm{GSp}(V, \langle \cdot, \cdot \rangle)$ . By definition,  $H \subset G_g$  for  $g = (\dim_{\mathbb{Q}} V)/2$ .

Given a PEL datum  $(B, *, V, \langle \cdot, \cdot \rangle, h)$ , there is an associated PEL moduli space  $\mathrm{Sh} = \mathrm{Sh}(B, *, V, \langle \cdot, \cdot \rangle, h)$ , which is a subspace of  $\mathcal{A}_g$ . The space  $\mathrm{Sh}$  is defined over a finite extension of  $\mathbb{Q}$ , called the reflex field [5, Section 5]. The manifold  $\mathrm{Sh}(\mathbb{C})^{\mathrm{an}}$  is the disjoint union of finitely many connected Shimura varieties [5, Sections 7–8].

**2.5. Integral PEL data.** Let  $(B, *, V, \langle \cdot, \cdot \rangle, h)$  be a PEL datum.

**Definition 2.2.** An *integral PEL datum* of the PEL datum  $(B, *, V, \langle \cdot, \cdot \rangle, h)$  consists of a tuple  $(\mathcal{O}_B, *, \Lambda, \langle \cdot, \cdot \rangle, h)$  where

- $\mathcal{O}_B$  is an order of  $B$  that is  $*$ -stable, and
- $\Lambda$  is a lattice of  $V$  that is  $\mathcal{O}_B$ -stable, such that the  $*$ -Hermitian form  $\langle \cdot, \cdot \rangle$  is integral on  $\Lambda$ .

A rational prime  $p$  is *good* for an integral PEL datum if  $\mathcal{O}_B$  is maximal at  $p$  and  $\Lambda$  is self-dual at  $p$ . All but finitely many primes are good, [6, Definition 1.4.1.1].

To an integral PEL datum  $(\mathcal{O}_B, *, \Lambda, \langle \cdot, \cdot \rangle, h)$ , there is an associated (canonical) integral model  $\text{Sh}(\mathcal{O}_B, *, \Lambda, \langle \cdot, \cdot \rangle, h)$  of  $\text{Sh}(B, *, V, \langle \cdot, \cdot \rangle, h)$ , defined over the ring of integers of the reflex field localized at good primes, [6, Theorem 1.4.1.1]. If  $p$  is a good prime for the integral PEL datum, then  $\text{Sh}(\mathcal{O}_B, *, \Lambda, \langle \cdot, \cdot \rangle, h)$  has good reduction at  $p$ .

**2.6. Shimura varieties given by the action of the covering group.** Let  $\gamma = (m, N, a)$  be a monodromy datum, and let  $Z_\gamma$  be as in Section 2.1. Then  $Z_\gamma$  is contained in a subspace of PEL type  $S_\gamma$  of  $\mathcal{A}_g$  [12, §3]. The PEL datum  $(B, *, V, \langle \cdot, \cdot \rangle, h)$  of  $S_\gamma$  (defined as in Section 2.4) satisfies the following conditions:

- $B = \mathbb{Q}[\mu_m]$  is the group algebra of  $\mu_m$  over  $\mathbb{Q}$ .
- $*$  is the involution on  $\mathbb{Q}[\mu_m]$  induced by the inverse map on  $\mu_m$ .
- $V = \mathbb{Q}^{2g}$  for  $g = g_\gamma$  as given in (2-1).

The structure of  $V$  as a  $B$ -module, and the complex structure on  $V_{\mathbb{R}}$  (namely,  $h$ ) are uniquely determined by the signature type  $\mathfrak{f} = \mathfrak{f}_\gamma$  given in (2-2).

Our goal is to compute the integral PEL datum  $(\mathcal{O}_B, *, \Lambda, \langle \cdot, \cdot \rangle, h)$  of  $S_\gamma$ .

More precisely, we set  $\mathcal{O}_B = \mathbb{Z}[\mu_m]$ , which is a  $*$ -stable order of  $B$ , and compute an  $\mathcal{O}_B$ -stable lattice  $\Lambda$  of  $V$  and a skew Hermitian form  $\langle \cdot, \cdot \rangle$  on  $V$  which is integral on  $\Lambda$ . By construction, the order  $\mathcal{O}_B$  is maximal away from  $m$  and the lattice  $\Lambda$  is self-dual away from  $m$ ; hence  $S_\gamma$  has good reduction at every prime  $p$  not dividing  $m$ .

**2.7. Complex multiplication.** Let  $L$  be a CM-field and let  $L_0$  be the maximal totally real subfield of  $L$ . Let  $n = [L_0 : \mathbb{Q}]$ ; hence,  $[L : \mathbb{Q}] = 2n$ . A CM-type of  $L$  is an ordered set  $\Phi$  of distinct embeddings  $\phi_i : L \hookrightarrow \mathbb{C}$ , for  $1 \leq i \leq n$ , no two of which are complex conjugate. A CM-type of  $L$  is called *simple* if it is not induced from the CM-type of a proper CM-subfield of  $L$ .

Let  $A$  be a complex torus such that  $L \subset \text{End}(A) \otimes \mathbb{Q}$ . We say that  $A$  is of CM-type  $(L, \Phi)$  if  $\dim(A) = n$  and the complex representation of  $L$  on  $\text{Lie}(A)$  is isomorphic to  $\sum_{\phi \in \Phi} \phi$ . If, in addition,  $\text{End}(A) \simeq \mathcal{O}_L$ , we say  $A$  has type  $(\mathcal{O}_L, \Phi)$ .

The following lemma is well-known. See [9, Lemma 3.1] for some explanation.

**Lemma 2.3.** *Suppose  $\psi : \mathcal{C} \rightarrow \mathbb{P}^1$  is a  $\mu_m$ -cover, branched at  $N = 3$  points, with inertia type  $a = (a_1, a_2, a_3)$  and signature type  $\mathfrak{f}$ . Then the Jacobian of  $\mathcal{C}$  has complex multiplication by  $\prod_d \mathbb{Q}(\zeta_d)$ , where  $d$  satisfies  $1 < d \mid m$ , and  $d \nmid a(i)$  for  $1 \leq i \leq 3$ . Its CM-type  $\Phi$  satisfies*

$$(2-3) \quad \phi \in \Phi \iff \mathfrak{f}(\phi) > 0.$$

The maximal order in  $\prod_d \mathbb{Q}(\zeta_d)$  is  $\prod_d \mathbb{Z}[\zeta_d]$ . The image of the group ring  $\mathbb{Z}[\mu_m]$  in  $\prod_d \mathbb{Q}(\zeta_d)$  has finite index inside the maximal order. This index is 1 if and only if there is a unique  $d$  satisfying  $1 < d \mid m$ , and  $d \nmid a(i)$  for  $1 \leq i \leq 3$ . This is true if  $m$  is prime.

The next result is immediate from Lemma 2.1 and (2-3).

**Lemma 2.4.** *If  $N = 3$ , then  $\sigma_i \in \text{Aut}(\mu_m)$  takes the CM-type  $\Phi$  to  $\Phi' = i \cdot \Phi = \{\tau_{ni} \mid \tau_n \in \Phi\}$ .*

For any CM-type  $(L, \Phi)$ , there exists a unique 0-dimensional PEL type moduli space  $\text{Sh} = \text{Sh}(L, \Phi)$  whose points represent abelian varieties with complex multiplication of type  $(\mathcal{O}_L, \Phi)$ . (When  $L = \mathbb{Q}(\zeta_m)$ , the points of  $\text{Sh}$  represent the Jacobians of curves for which there is a  $\mu_m$ -cover of  $\mathbb{P}^1$  branched at  $N = 3$  points with CM-type  $\Phi$ .) The signature type  $\mathfrak{f}$  of  $\text{Sh}$  is given by

$$(2-4) \quad \text{for each } \phi : L \rightarrow \mathbb{C}, \quad \mathfrak{f}(\phi) = \begin{cases} 1 & \text{if } \phi \in \Phi, \\ 0 & \text{if } \phi \notin \Phi. \end{cases}$$

For any signature type  $\mathfrak{f}$  on  $L$  taking values in  $\{0, 1\}$ , there is a unique CM-type  $\Phi$  of  $L$  compatible with  $\mathfrak{f}$ .

### 3. Some background from algebraic number theory

**3.1. Class group and narrow class group.** For a number field  $L$ : let  $\mathcal{O}_L$  denote its ring of integers; let  $\mathcal{U}_L$  denote the units in  $\mathcal{O}_L$ ; and let  $D_{L/\mathbb{Q}}$  denote the different of  $L$  over  $\mathbb{Q}$ .

Let  $\text{cl}_L$  be the class group of  $L$ . Recall that  $\text{cl}_L = I_L/P_L$ , where  $I_L$  is the group of nonzero fractional ideals and  $P_L$  is the subgroup of nonzero principal fractional ideals. Let  $h_L = |\text{cl}_L|$  be the class number of  $L$ .

An element  $\alpha \in L$  is totally positive if it is positive under every real embedding of  $L$ . Let  $\mathcal{U}_L^+ \subset \mathcal{U}_L$  be the subgroup of totally positive units. Let  $P_L^+ \subset P_L$  be the subgroup of principal ideals generated by a totally positive element. The narrow class group of  $L$  is  $\text{cl}_L^+ = I_L/P_L^+$  and the narrow class number is  $h_L^+ = |\text{cl}_L^+|$ . There is a surjection

$$\nu_L : \text{cl}_L^+ \rightarrow \text{cl}_L.$$

**3.2. Units of independent signs.** Let  $L$  be a CM-field and  $L_0$  its maximal totally real subfield. Let  $n = [L_0 : \mathbb{Q}]$ ; hence,  $[L : \mathbb{Q}] = 2n$ . By [19, Theorem 4.10],  $h_{L_0}$  divides  $h_L$ .

We fix an ordering  $\tau_1, \dots, \tau_n$  of the  $n$  real embeddings  $L_0 \hookrightarrow \mathbb{R}$ . If  $L_0/\mathbb{Q}$  is Galois, we identify these with the elements of  $\text{Gal}(L_0/\mathbb{Q})$ . Consider the group homomorphism

$$(3-1) \quad \rho_{L_0} : \mathcal{U}_{L_0} \rightarrow \{\pm 1\}^n, \quad \rho_{L_0}(u) = (\tau_i(u)/|\tau_i(u)|)_{1 \leq i \leq n} \text{ for } u \in \mathcal{U}_{L_0}.$$

We say that  $L_0$  has *units of independent signs* if, for each real embedding  $\tau$ , there is a unit which is negative under  $\tau$  but positive under all other real embeddings [2, Definition 12.1]. This is equivalent to saying that  $\rho_{L_0}$  is surjective or that  $\nu_{L_0} : \text{cl}_{L_0}^+ \rightarrow \text{cl}_{L_0}$  is an isomorphism [2, Lemma 11.2].

We say that  $L_0$  has *units of almost independent signs* if every unit in  $\mathcal{U}_{L_0}$  is negative under an even number of real embeddings and, for every pair of real embeddings, there is a unit which is negative under exactly the two embeddings in that pair [2, Definition 12.13]. This condition is equivalent to  $|\ker(\nu_{L_0})| = 2$  or  $|\text{coker}(\rho_{L_0})| = 2$ , by [2, Lemma 11.2].

**3.3. Norms and the Hasse unit index.** Consider the norm map  $N : \mathcal{U}_L \rightarrow \mathcal{U}_{L_0}$  given by  $N(y) = N_{L/L_0}(y)$  for  $y \in \mathcal{U}_L$ . By Dirichlet's unit theorem,  $[\mathcal{U}_{L_0} : \mathcal{U}_{L_0}^2] = 2^n$ .

**Lemma 3.1.** *Suppose  $L_0$  is a totally real field, and  $L/L_0$  is a CM-extension. Then*

$$(3-2) \quad \mathcal{U}_{L_0}^2 \subseteq N(\mathcal{U}_L) \subseteq \mathcal{U}_{L_0}^+ \subseteq \mathcal{U}_{L_0}.$$

*Proof.* By definition,  $\mathcal{U}_{L_0}^+ \subseteq \mathcal{U}_{L_0}$ . All elements in  $N(\mathcal{U}_L)$  are totally positive units because  $L$  is a CM-field, quadratic over its totally real subfield  $L_0$ . So  $N(\mathcal{U}_L) \subseteq \mathcal{U}_{L_0}^+$ . Also,  $\mathcal{U}_{L_0}^2 = N(\mathcal{U}_{L_0})$ , hence  $\mathcal{U}_{L_0}^2 \subset N(\mathcal{U}_L)$ .  $\square$

Let  $\mu_L$  be the group of roots of unity of  $L$ .

**Definition 3.2.** The *Hasse unit index* of a CM-extension  $L/L_0$  is

$$Q(L) = [\mathcal{U}_L : \mu_L \mathcal{U}_{L_0}].$$

By [19, Theorem 4.12],  $Q(L) = 1$  or  $Q(L) = 2$ . Since  $\text{Ker}(N) = \mu_L$ , it follows that  $\mathcal{U}_L = \mu_L \mathcal{U}_{L_0}$  if and only if  $N(\mathcal{U}_L) = N(\mathcal{U}_{L_0})$ . Also  $N(\mathcal{U}_{L_0}) = \mathcal{U}_{L_0}^2$ . Thus

$$(3-3) \quad Q(L) = [N(\mathcal{U}_L) : \mathcal{U}_{L_0}^2].$$

Let  $t$  be the number of finite primes ramified in  $L/L_0$ .

**Lemma 3.3.** *Suppose  $L$  has odd class number. Let  $L_0$  be a totally real field, and  $L/L_0$  a CM-extension. Then  $Q(L) = 1$  if and only if  $t = 1$ ; and  $Q(L) = 2$  if and only if  $t = 0$ .*

*Proof.* The material in [2, Chapter 13] is stated in terms of the type of the CM-extension  $L/L_0$ . By a theorem of Kummer [2, Theorem 13.4]  $L/L_0$  has type I (resp. II) if and only if  $Q(L) = 1$  (resp.  $Q(L) = 2$ ). The result is then immediate from [2, page 73].  $\square$

We summarize the result from this section that we need in this and later papers.

**Lemma 3.4.** *Suppose  $L$  is a CM-field with maximal totally real subfield  $L_0$ .*

- (1)  $L_0$  has units of independent signs if and only if  $Q(L)[\mathcal{U}_{L_0}^+ : N(\mathcal{U}_L)] = 1$ .
- (2)  $L_0$  has units of almost independent signs if and only if  $Q(L)[\mathcal{U}_{L_0}^+ : N(\mathcal{U}_L)] = 2$ .

*Proof.* (1) By [2, Lemma 12.2],  $L_0$  has units of independent signs if and only if every element of  $\mathcal{U}_{L_0}^+$  is a square. The result then follows from Lemma 3.1.

(2) It follows from the definitions that  $L_0$  has units of almost independent signs if and only if  $[\mathcal{U}_{L_0} : \mathcal{U}_{L_0}^+] = 2^{n-1}$ . Since  $[\mathcal{U}_{L_0} : \mathcal{U}_{L_0}^2] = 2^n$ , this is equivalent to  $[\mathcal{U}_{L_0}^+ : \mathcal{U}_{L_0}^2] = 2$ .  $\square$

**3.4. Cyclotomic fields.** Let  $m$  be a positive integer and let  $\zeta_m = e^{2\pi i/m} \in \mathbb{C}$ . Let  $\mu_m \subset \mathbb{C}$  be the group of  $m$ -th roots of unity. Let  $\mathbb{Q}[\mu_m]$  be the group algebra of  $\mu_m$  over  $\mathbb{Q}$ .

The cyclotomic field  $F = \mathbb{Q}(\zeta_m)$  is a CM-field over  $\mathbb{Q}$  with maximal totally real subfield  $F_0 = \mathbb{Q}(\zeta_m + \zeta_m^{-1})$ . The degree of  $F_0$  over  $\mathbb{Q}$  is  $n = \phi(m)/2$ .

**Notation 3.5.** The choice of  $\zeta_m$  fixes an embedding  $\sigma_1 : F \hookrightarrow \mathbb{C}$ . For  $1 \leq i < m$ , with  $\gcd(i, m) = 1$ , let  $\sigma_i$  be the embedding  $F \hookrightarrow \mathbb{C}$  (or automorphism in  $\text{Gal}(F/\mathbb{Q})$ ) determined by  $\sigma_i(\zeta_m) = \zeta_m^i$ . For  $x \in F$ , let  $\bar{x} = \sigma_{m-1}(x)$  denote its complex conjugate.

If  $m = p^r$  is a prime power, then  $F/F_0$  is ramified at the unique prime of  $F_0$  above  $p$  and at the  $n$  infinite primes of  $F_0$  and is unramified at all other primes. See [19, Proposition 2.15].

**Lemma 3.6.** *Let  $F = \mathbb{Q}(\zeta_m)$  and  $F_0 = \mathbb{Q}(\zeta_m + \zeta_m^{-1})$ . Suppose  $F$  has class number 1.*

(1) *If  $m$  is a prime power (or twice a prime power), then  $F_0$  has narrow class number 1 and thus has units of independent signs. The complete list of these  $m$  is*

$$m = 1, 2, 4, 8, 16, 32;$$

$$m = 3, 5, 7, 9, 11, 13, 17, 19, 25, 27; \text{ or twice a value of } m \text{ on this row.}$$

(2) *If  $m$  is not a prime power (or twice a prime power), then  $F_0$  has narrow class number 2 and thus has units of almost independent signs. The complete list of these  $m$  is*

$$m = 12, 16, 20, 24, 28, 36, 40, 44, 48, 60, 84;$$

$$m = 15, 21, 33, 35, 45; \text{ or twice a value of } m \text{ on this row.}$$

*Proof.* The complete list of  $m$  such that  $h_F = 1$  is well-known; it is the union of the lists in parts (1) and (2). For these  $m$ , since  $h_F = 1$ , also  $h_{F_0} = 1$ .

(1) By a result of Hasse (see [2, Corollary 3.9]), since  $h_F$  and  $h_{F_0}$  are odd,  $F_0$  has units of independent signs if and only if  $F/F_0$  is ramified at exactly one finite prime. This happens if and only if  $m = p^r$  or  $m = 2 \cdot p^r$ , for some prime  $p$ .

(2) Since  $h_F$  and  $h_{F_0}$  are odd, then  $F_0$  has units of almost independent signs if and only if  $F/F_0$  is not ramified at any finite prime [2, Corollary 13.10]. This happens if  $m$  is not a prime power (or twice a prime power).  $\square$

**3.5. A generator for the different of cyclotomic fields.**

**Lemma 3.7.** *Let  $F = \mathbb{Q}(\zeta_m)$ . The element  $\beta_0$  below generates  $D_{F/\mathbb{Q}}$  and  $\beta_0 = -\bar{\beta}_0$ .*

(1) *If  $m$  is an odd prime, then*

$$(3-4) \quad \beta_0 = \frac{m}{\zeta_m^{(m+1)/2} - \zeta_m^{(m-1)/2}}.$$

(2) *If  $m = 2^k$  with  $k \geq 2$ , then  $\beta_0 = -2^{k-1}i$ .*

(3) *If  $m = 3^k$ , then  $\beta_0 = -3^{k-1}\sqrt{3}i$ .*

(4) *If  $m = pq$  with  $p, q$  distinct odd primes, then*

$$(3-5) \quad \beta_0 = m \frac{\zeta_m^{(m+1)/2} - \zeta_m^{(m-1)/2}}{(\zeta_m^{q(p+1)/2} - \zeta_m^{q(p-1)/2})(\zeta_m^{p(q+1)/2} - \zeta_m^{p(q-1)/2})}.$$

*Proof.* In each case,  $\beta_0$  is on the imaginary axis, so  $\beta_0 = -\bar{\beta}_0$ .

Let  $c_m(x)$  denote the  $m$ -th cyclotomic polynomial. The different  $D_{F/\mathbb{Q}}$  is generated by  $\langle c'_m(\zeta_m) \rangle$  by [13, III, Proposition 2.4]. To show that  $\beta_0$  generates  $D_{F/\mathbb{Q}}$ , it suffices to show that it is an associate of  $\langle c'_m(\zeta_m) \rangle$  in  $\mathcal{O}_F$ .

(1) When  $m$  is an odd prime, then  $c_m(x) = (x^m - 1)/(x - 1)$ . Then

$$c'_m(x) = \frac{(x - 1)mx^{m-1} - (x^m - 1)}{(x - 1)^2}.$$

Hence  $c'_m(\zeta_m) = m\zeta_m^{-1}/(\zeta_m - 1)$ , and  $\beta_0$  from (3-4) is an associate of  $c'_m(\zeta_m)$  in  $\mathcal{O}_F$ .

(2) If  $m = 2^k$ , then  $c_m(x) = x^{2^{k-1}} + 1$ . Thus  $c'_m(\zeta_m) = 2^{k-1}\zeta_m^{2^{k-1}-1} = -2^{k-1}/\zeta_m$ , and the corresponding  $\beta_0$  is an associate of  $c'_m(\zeta_m)$  in  $\mathcal{O}_F$ .

(3) If  $m = 3^k$ , then  $c_m(x) = x^{2 \cdot 3^{k-1}} + x^{3^{k-1}} + 1$ . Thus  $c'_m(\zeta_m) = 3^{k-1}\zeta_m^{3^{k-1}-1}\sqrt{-3}$ , and this  $\beta_0$  is an associate of  $c'_m(\zeta_m)$  in  $\mathcal{O}_F$ .

(4) In this case,  $c_m(x) = \frac{(x^m - 1)(x - 1)}{(x^p - 1)(x^q - 1)}$ , and  $\beta_0$  in (3-5) is an associate of  $c'_m(\zeta_m) = \frac{m(\zeta_m - 1)}{\zeta_m^p(\zeta_m^p - 1)(\zeta_m^q - 1)}$ .  $\square$

**3.6. Simple types.** Let  $F = \mathbb{Q}(\zeta_m)$  and recall that  $\text{Gal}(F/\mathbb{Q}) \simeq (\mathbb{Z}/m\mathbb{Z})^*$ . A CM-type  $\Phi$  of  $F$  is simple, if it is not induced from any CM-field  $F' \subset F$ .

Let  $H \subset \text{Gal}(F/\mathbb{Q})$  and let  $F^H$  be the fixed field of  $F$  under  $H$ . If  $F^H$  is a CM-field, then  $\Phi$  is induced from  $F^H$  if and only if it is a union of cosets of  $H$ .

**Lemma 3.8.** *Suppose  $m = 4$  or  $m$  is a Fermat prime or twice a Fermat prime. Then  $\Phi$  is simple.*

We will use Lemma 3.8 for  $m = 4$  and  $m = 3, 5, 17$  and  $m = 6, 10$ .

*Proof.* For these values of  $m$ , the field  $F$  contains no proper nontrivial CM-fields.  $\square$

**Lemma 3.9.** *Let  $m > 3$  be prime. Let  $\gamma = (m, 3, a)$  be a monodromy datum with inertia type  $a = (1, a_2, a_3)$ . Let  $\Phi$  be the CM-type of  $F$  corresponding to  $a$ . Then  $\Phi$  is simple if and only if  $a \neq (1, x, x^2)$  for some  $x \in (\mathbb{Z}/m\mathbb{Z})^*$  with order 3.*

*Proof.* This follows from [4, Theorem 2]. See also [7, Theorem 6.2].  $\square$

For example, when  $m = 7$  and  $a = (1, a_2, a_3)$ ,  $\Phi$  is simple unless  $a = (1, 2, 4)$ .

By a direct computation, one can check that  $\Phi$  is always simple when  $m = 25$  or  $m = 27$ .

#### 4. Principal polarizations of CM abelian varieties

Let  $L$  be a CM-field and  $L_0$  its maximal totally real subfield. We will now study principal polarizations on abelian varieties having complex multiplication by  $L$ .

Section 4.1 contains background about complex tori with complex multiplication. In Section 4.2, we study principal polarizations on CM-abelian varieties, following work of van Wamelen. In Section 4.3, we prove a result about existence and uniqueness of principal polarizations for abelian varieties with simple CM-type when  $L$  has class number 1. In Section 4.4, we specialize to the case that  $L$  is a cyclotomic field with class number 1.

**4.1. Complex tori.** We review some complex multiplication theory, following Lang in [7, Chapter 1]. Let  $A$  be a complex torus such that  $L \subset \text{End}(A) \otimes \mathbb{Q}$ . We say that  $A$  is of type  $(L, \Phi)$  if the complex representation of  $\text{End}(A) \otimes \mathbb{Q}$  is isomorphic to  $\sum_{\phi \in \Phi} \phi$ . If, in addition,  $\text{End}(A) \simeq \mathcal{O}_L$ , we say  $A$  has type  $(\mathcal{O}_L, \Phi)$ .

**Theorem 4.1.** [7, Chapter 1: Theorems 3.3, 3.5, 4.1, 4.2]

- (1) *If  $\mathfrak{a}$  is a lattice in  $L$  and  $\Phi$  is a CM-type of  $L$ , then  $\mathbb{C}^n / \Phi(\mathfrak{a})$  is a complex torus of type  $(L, \Phi)$ .*
- (2) *If  $A$  is a complex torus of type  $(L, \Phi)$ , then there exists a lattice  $\mathfrak{a}$  of  $L$  such that  $A \simeq \mathbb{C}^n / \Phi(\mathfrak{a})$ .*
- (3) *If  $\Phi$  is a simple type and  $\mathfrak{a}$  is a fractional ideal of  $L$ , then  $\text{End}(\mathbb{C}^n / \Phi(\mathfrak{a})) \simeq \mathcal{O}_L$ .*

(4) If  $\Phi$  is a simple type and  $\mathfrak{a}, \mathfrak{b}$  are fractional ideals of  $L$ , then  $\mathbb{C}^n/\Phi(\mathfrak{a}) \simeq \mathbb{C}^n/\Phi(\mathfrak{b})$  if and only if  $\mathfrak{a}$  and  $\mathfrak{b}$  are in the same ideal class.

In particular, if  $(L, \Phi)$  is a simple CM-type, then the set of isomorphism classes of complex tori of type  $(\mathcal{O}_L, \Phi)$  is in bijection with the class group of  $L$ .

Furthermore, by [7, Chapter 1, Theorem 4.5], every (admissible, nondegenerate) Riemann form on  $\mathbb{C}^n/\Phi(\mathfrak{a})$  is given by

$$(4-1) \quad \mathbb{E}(\Phi(x), \Phi(y)) = \mathrm{tr}_{L/\mathbb{Q}}(\xi x \bar{y}), \text{ for } x, y \in L,$$

for some  $\xi$  such that  $L = L_0(\xi)$ ,  $\xi^2 \in L_0$  is totally negative, and  $\mathrm{Im}(\phi(\xi)) > 0$  for  $\phi \in \Phi$ .

**4.2. Principal polarizations on CM abelian varieties.** In [18, p. 310], van Wamelen developed an algorithm to produce isomorphism classes of principally polarized abelian varieties of type  $(\mathcal{O}_L, \Phi)$  based on the following result.

**Theorem 4.2** (van Wamelen [18]). *Let  $(L, \Phi)$  be a CM-type.*

(1) (Theorem 4)<sup>1</sup> *Writing  $L = L_0(\sqrt{-c})$  for some  $c \in \mathcal{O}_{L_0}$ , there exist a fractional ideal  $\mathfrak{a} \subset L$  and an element  $b \in \mathcal{O}_{L_0}$  such that  $D_{L/\mathbb{Q}} \cdot \mathfrak{a}\bar{\mathfrak{a}} = \langle b\sqrt{-c} \rangle$ .*

(2) (Theorem 3) *Let  $\xi \in L$  be such that  $L = L_0(\xi)$ ,  $\xi^2 \in L_0$ , and  $D_{L/\mathbb{Q}} \cdot \mathfrak{a}\bar{\mathfrak{a}} = \langle \xi^{-1} \rangle$ , for some fractional ideal  $\mathfrak{a}$  of  $L$  (for example, with the notation of part (1),  $\xi = (b\sqrt{-c})^{-1}$ ). Define a Riemann form  $\mathbb{E} : \mathbb{C}^n \times \mathbb{C}^n \rightarrow \mathbb{C}$  by*

$$\mathbb{E}(z, w) = \sum_{i=1}^n \phi_i(\xi)(\bar{z}_i w_i - z_i \bar{w}_i),$$

for  $z, w \in \mathbb{C}^n$ . If  $\mathrm{Im}(\phi(\xi)) > 0$  for  $\phi \in \Phi$ , then  $\mathbb{E}$  defines a principal polarization on  $\mathbb{C}^n/\Phi(\mathfrak{a})$  of type  $(\mathcal{O}_L, \Phi)$ . Furthermore, if  $(L, \Phi)$  is a simple CM-type, then all principal polarizations on  $\mathbb{C}^n/\Phi(\mathfrak{a})$  of type  $(\mathcal{O}_L, \Phi)$  are given by such a  $\xi$ .<sup>2</sup>

(3) (Corollary 1) *Two principal polarizations of the same CM-type on  $\mathbb{C}^n/\Phi(\mathfrak{a})$  arising from  $\xi_1$  and  $\xi_2$  give isomorphic principally polarized abelian varieties if and only if there exists a unit  $u \in \mathcal{O}_L^*$  such that  $\xi_1 = u\bar{u}\xi_2$ .*

**Corollary 4.3.** *Let  $\Phi$  be a CM-type of  $L$ . An element  $\xi = \beta^{-1}$  for some  $\beta \in \mathcal{O}_L$  defines a principal polarization on  $A_\Phi = \mathbb{C}^n/\Phi(\mathcal{O}_L)$  of CM-type  $(\mathcal{O}_L, \Phi)$  if and only if*

- (1)  $\beta$  generates the different  $D_{L/\mathbb{Q}}$ ,
- (2)  $\beta = -\bar{\beta}$ , and
- (3)  $\mathrm{Im}(\phi(\beta)) < 0$ , for each  $\phi \in \Phi$ .

<sup>1</sup>We have made a small adjustment to the notation to be consistent with other parts of this theorem.

<sup>2</sup>Note that type  $(L, \Phi)$  and type  $(\mathcal{O}_L, \Phi)$  are equivalent in the last two statements.

Two elements  $\beta, \beta'$  satisfying the above conditions yield isomorphic principally polarized abelian varieties if and only if there exists a unit  $u \in \mathcal{U}_L$  such that  $\beta = u\bar{u}\beta'$ . If the CM-type  $\Phi$  is simple, then all principal polarizations of  $A_\Phi$  of CM-type  $(\mathcal{O}_L, \Phi)$  arise this way.

*Proof.* This follows from Theorem 4.2, replacing  $\xi, \xi_1, \xi_2$  with  $\beta^{-1}, \beta^{-1}, \beta'^{-1}$ .  $\square$

In Lemma 3.7, we determined an element  $\beta_0 \in \mathcal{O}_F$  satisfying conditions (1) and (2) in Corollary 4.3 when  $F = \mathbb{Q}(\zeta_m)$  for many values of  $m$ .

**4.3. Existence and uniqueness of principal polarizations.** In this section, we study principal polarizations on CM-abelian varieties. Under conditions on the class group and unit group of the field, we show such principal polarizations exist and are uniquely determined. Recall that  $L$  is a CM-field with maximal totally real subfield  $L_0$  and  $n = [L_0 : \mathbb{Q}]$ .

**Proposition 4.4.** *Suppose  $L_0$  has units of independent signs and  $(L, \Phi)$  is a CM-type.*

- (1) *There exists a principally polarized CM-abelian variety of type  $(\mathcal{O}_L, \Phi)$ .*
- (2) *If  $(L, \Phi)$  is simple, then any CM-abelian variety of type  $(\mathcal{O}_L, \Phi)$  has at most one principal polarization, up to isomorphism.*
- (3) *If  $(L, \Phi)$  is simple and  $L$  has class number 1, then there exists a unique principally polarized CM-abelian variety of type  $(\mathcal{O}_L, \Phi)$ , up to isomorphism.*

*Proof.* By Theorem 4.2(1), there exists a fractional ideal  $\mathfrak{a}$  in  $L$  and an element  $\beta_0 \in \mathcal{O}_L$  such that (1)  $\beta_0$  generates  $D_{L/\mathbb{Q}}\mathfrak{a}\bar{\mathfrak{a}}$  and (2)  $\beta_0 = -\bar{\beta}_0$ . An element  $\beta' \in \mathcal{O}_L$  satisfies conditions (1) and (2) if and only if  $\beta' = u_0\beta_0$  for a totally real unit  $u_0 \in \mathcal{U}_{L_0}$ .

By Theorem 4.2(2),  $\beta \in \mathcal{O}_L$  defines a principal polarization of type  $(\mathcal{O}_L, \Phi)$  on  $\mathbb{C}^n/\Phi(\mathfrak{a})$  if and only if it satisfies conditions (1) and (2), and also (3)  $\text{Im}(\phi(\beta)) < 0$ , for each  $\phi \in \Phi$ .

Hence, to finish the first claim, it suffices to check that there exists  $u_0 \in \mathcal{U}_{L_0}$  such that  $\beta = u_0\beta_0$  satisfies condition (3), for any CM-type  $\Phi$  of  $L$ . If  $L_0$  has units of independent signs, then  $\rho_{L_0}$  is surjective; this shows that the unit  $u_0$  described above exists.

By Theorem 4.2(2), if  $(L, \Phi)$  is simple, all principal polarizations of type  $\Phi$  on  $\mathbb{C}^n/\Phi(\mathfrak{a})$  arise from some  $\beta \in \mathcal{O}_L$  satisfying conditions (1)–(3). By Theorem 4.2(3),  $\beta, \beta' \in \mathcal{O}_L$  satisfying conditions (1)–(3) define isomorphic principally polarized abelian varieties if and only if  $\beta' = N(u)\beta$  for some unit  $u \in \mathcal{U}_L$ .

By definition, given a fractional ideal  $\mathfrak{a}$  of  $L$ , if  $\beta \in \mathcal{O}_L$  satisfying conditions (1)–(3) exists, then  $\beta' \in \mathcal{O}_L$  also satisfies conditions (1)–(3) if and only if  $\beta' = u^+\beta$  for some totally positive unit  $u^+ \in \mathcal{U}_{L_0}^+$ .

Hence, to finish the second claim, it suffices to check that for any  $u^+ \in \mathcal{U}_{L_0}^+$  there is  $u \in \mathcal{U}_L$  such that  $u^+ = N(u)$ . Since  $L_0$  has units of independent signs, this follows from Lemma 3.4. Finally, if  $L$  has class number 1, then by Theorem 4.2 and [18, Theorem 5], any CM-abelian variety of type  $(\mathcal{O}_L, \Phi)$  is isomorphic to  $A_\Phi = \mathbb{C}^n / \Phi(\mathcal{O}_L)$ .  $\square$

**Proposition 4.5.** *Suppose  $L_0$  has units of almost independent signs and  $Q(L) = 2$ , where  $Q(L)$  is defined in Definition 3.2. Suppose  $(L, \Phi)$  is a simple CM-type.*

- (1) *The number of isomorphism classes of principal polarizations on a CM-abelian variety of type  $(\mathcal{O}_L, \Phi)$  is at most one.*
- (2) *Suppose in addition that  $L$  has class number 1. If there is a principally polarized CM-abelian variety of type  $(\mathcal{O}_L, \Phi)$ , then it is unique up to isomorphism.*

*Proof.* The arguments in the proof of Proposition 4.4 still apply, after observing that  $U_{L_0}^+ = N(\mathcal{U}_L)$  when  $L_0$  has units of almost independent signs and  $Q(L) = 2$  by Lemma 3.4.  $\square$

**4.4. CM-types for cyclotomic fields.** We consider the case when  $L$  is the cyclotomic field  $F = \mathbb{Q}(\zeta_m)$ . By Section 3.2, the next result is a special case of Propositions 4.4 and 4.5.

**Corollary 4.6.** *Let  $m$  such that  $F = \mathbb{Q}(\zeta_m)$  has class number 1. Let  $\Phi$  be a CM-type of  $F$ .*

- (1) *If  $F_0$  has narrow class number 1, then there exists a principally polarized CM-abelian variety of type  $(\mathcal{O}_F, \Phi)$ . Furthermore, if  $(F, \Phi)$  is simple, then it is unique up to isomorphism.*
- (2) *Suppose  $F_0$  has narrow class number 2 and  $\Phi$  is simple. If there exists a principally polarized CM-abelian variety of type  $(\mathcal{O}_F, \Phi)$ , then it is unique up to isomorphism.*

We will now explicitly describe the data of the principal polarization.

**Corollary 4.7.** *Let  $m$ ,  $F = \mathbb{Q}(\zeta_m)$ , and  $\beta_0$  be as in Lemma 3.7. Let  $\Phi$  be a CM-type of  $F$ . Consider conditions (1)–(3) in Corollary 4.3.*

- (1) *If  $\Phi' = i\Phi$  and  $\beta$  satisfies conditions (1)–(3) for  $\Phi$ , then  $\beta' = \sigma_{i-1}(\beta)$  satisfies conditions (1)–(3) for  $\Phi'$ .*
- (2) *Suppose  $F_0$  has units of independent signs. Then, for any CM-type  $\Phi$  of  $F$ , there exists a unit  $u_0 \in \mathcal{U}_{F_0}$  such that  $\beta = u_0\beta_0$  satisfies conditions (1)–(3) for  $\Phi$ . Furthermore,  $\beta$  is the unique element satisfying conditions (1)–(3) for  $\Phi$ , up to multiplication by an element of  $N_{F/F_0}(\mathcal{U}_F)$ .*

The condition that  $F_0$  has units of independent signs is satisfied if the narrow class number of  $F_0$  is odd [2, bottom of page 58].

*Proof.* (1) This follows from Lemma 2.1 and Lemma 2.4.

(2) By Lemma 3.7,  $\beta_0$  satisfies conditions (1) and (2) of Corollary 4.3. Since  $F_0$  has units of independent signs, there exists  $u_0 \in \mathcal{U}_{F_0}$  such that  $\beta = u_0\beta_0$  satisfies condition (3). Since  $u_0 \in \mathcal{U}_{F_0}$  is a real unit,  $\beta$  also satisfies conditions (1) and (2) of Corollary 4.3. Note that  $\beta$  is unique up to multiplication by a totally positive unit of  $F_0$ . By Lemma 3.4,  $\mathcal{U}_{F_0}^+ = N(\mathcal{U}_F)$ , proving the uniqueness statement.  $\square$

See Examples 6.3 and 6.7 for illustrations of Corollary 4.7 when  $m = 5$  and  $m = 7$ , respectively. As another example, if  $m = 8$ , then  $\beta_0 = -4i$  from Lemma 3.7: if  $\mathfrak{f} = (0, 1, 0, 1)$ , set  $u_0 = -1$  and  $\beta = 4i$ ; if  $\mathfrak{f} = (1, 1, 0, 0)$ , set  $u_0 = \sqrt{2} - 1$  and  $\beta = 4(1 - \sqrt{2})i$ .

### 5. Shimura data

Consider a monodromy datum  $\gamma = (m, N, a)$  as in Section 2.1. Recall that  $H_\gamma$  is the Hurwitz space of  $\mu_m$ -cyclic covers of  $\mathbb{P}^1$  with monodromy datum  $\gamma$ . Recall also that  $Z_\gamma$  is the closure in  $\mathcal{A}_g$  of the image of  $H_\gamma$  under the Torelli morphism and  $S_\gamma$  is the smallest PEL type Shimura substack of  $\mathcal{A}_g$  containing  $Z_\gamma$ . In this section, under certain assumptions on  $m$ , we provide a method to determine the integral PEL datum of  $S_\gamma$ .

A point  $P$  in  $Z_\gamma$  represents the Jacobian of a stable curve  $C_P$  of compact type, where compact type means that the dual graph of  $C_P$  is a tree. Furthermore, the curve  $C_P$  is a  $\mu_m$ -cover of a tree of projective lines. The main idea of the proof is to find a point  $P$  in  $Z_\gamma$  such that the Jacobian of every irreducible component of  $C_P$  has complex multiplication by a CM-field.

**5.1. Admissible covers.** We briefly review some information about the boundary of  $H_\gamma$ , see [20, Chapter 4] or [3, Section 1]. The generic point of  $Z_\gamma$  represents the Jacobian of a smooth curve. More generally, a point  $P$  of  $Z_\gamma$  represents the Jacobian  $J_P$  of a stable curve  $C_P$  of compact type. There is a  $\mu_m$ -cover  $\psi : C_P \rightarrow T$  where  $T$  is a tree of projective lines. The nodes of  $C_P$  and  $T$  are ordinary double points.

Since  $P$  is in  $Z_\gamma$ , the closure of  $Z_\gamma^0$ , the cover  $\psi$  is *admissible* as defined below. If  $\eta$  is a node of  $C_P$ , consider the restrictions  $\phi_1$  and  $\phi_2$  of  $\phi$  to the two irreducible components of  $C_P$  that intersect at  $\eta$ . When  $m$  is prime, the compact type condition implies that  $\phi_1$  and  $\phi_2$  are each totally ramified at  $\eta$ . For  $i = 1, 2$ , let  $\alpha_i$  be the canonical generator of inertia of  $\phi_i$  at  $\eta$ . The admissible condition is that  $\alpha_1$  and  $\alpha_2$  are inverses in  $\mu_m$ .

**5.2. Distinguished points.** We prove that there exists a distinguished point  $P$  in  $Z_\gamma$  such that we can compute the lattice and Hermitian form for  $J_P$ .

**Definition 5.1.** A point  $P$  in  $Z_\gamma$  is a *distinguished point* if the Jacobian  $J_P$  is a principally polarized abelian variety with complex multiplication by a maximal order in a CM-field, or the direct sum of such together with the product polarization.

**Proposition 5.2.** *Let  $m$  be an odd prime. Let  $\gamma = (m, N, a)$  be a monodromy datum. Then  $Z_\gamma$  has a distinguished point  $P$ . More specifically, for  $r = N - 2$ :*

- (1) *In the family  $H_\gamma$  of  $\mu_m$ -covers of a genus 0 curve with monodromy datum  $\gamma$ , there is a point which represents an admissible  $\mu_m$ -cover  $\psi : C_P \rightarrow T$ , where  $T$  is a tree of  $r$  projective lines and  $C_P$  is a curve of compact type, with  $r$  irreducible components  $C_1, \dots, C_r$ , each of which is a curve of genus  $(m - 1)/2$  admitting a  $\mu_m$ -cover of  $\mathbb{P}^1$  branched at 3 points.*
- (2) *The Jacobian  $J_P$  of  $C_P$  is of the form  $A \simeq \bigoplus_{j=1}^r A_j$ , where each  $A_j$  is a principally polarized abelian variety of dimension  $(m - 1)/2$  having complex multiplication by  $\mathcal{O}_F = \mathbb{Z}[\zeta_m]$ , together with the product polarization.*

*Proof.* The fact that  $Z_\gamma$  has a distinguished point  $P$  is immediate from part (2), which we will show follows from part (1).

(1) Let  $a = (a(1), \dots, a(N))$  be the inertia type. If  $N \geq 4$  and  $m$  is an odd prime, then  $a$  has the following property: there is a pair  $(i, j)$  with  $1 \leq i < j \leq N$ , such that  $a(i) + a(j) \not\equiv 0 \pmod{m}$ . Without loss of generality, we can suppose  $i = 1$  and  $j = 2$ . Let  $\alpha_2 = a(1) + a(2)$  and  $\alpha_1 = -\alpha_2$ .

When  $N \geq 4$ , then  $Z_\gamma^0$  is affine. Consider a family of  $\mu_m$ -covers with monodromy datum  $\gamma$ , where the first branch point  $b_1$  approaches the second branch point  $b_2$ . When  $b_1 = b_2$ , the curve is singular and its normalization is a  $\mu_m$ -cover  $\phi$  of a tree of two projective lines. Let  $\phi_1$  (resp.  $\phi_2$ ) be the restriction of  $\phi$  over the first (resp. second) of these.

The cover  $\phi_1$  (resp.  $\phi_2$ ) is ramified at the specializations of 2 (resp.  $N - 2$ ) ramification points. For each of these, the canonical generator of inertia remains the same at the specialization. The values in the inertia type of  $\phi_1$  (or  $\phi_2$ ) sum to 0 modulo  $m$ . Also,  $\phi$  is admissible and ramified at the ordinary double point. Thus  $\phi_1$  is a  $\mu_m$ -cover of  $\mathbb{P}^1$  branched at 3 points with inertia type  $(a(1), a(2), \alpha_1)$ , and  $\phi_2$  is a  $\mu_m$ -cover of  $\mathbb{P}^1$  branched at  $N - 1$  points with inertia type  $(\alpha_2, a(3), \dots, a(N))$ . The cover  $\phi$  is called a *degeneration of compact type* in [8, Remark 5.2].

By induction on  $N$ , we see that  $H_\gamma$  contains a family of  $\mu_m$ -covers which degenerates completely to an admissible  $\mu_m$ -cover  $\psi : C_P \rightarrow T$ , where  $T$  is a tree of  $r$  projective lines and the restriction of  $\psi$  above each component of  $T$  is branched at 3 points. By (2-1), each irreducible component of  $C_P$  has genus  $(m - 1)/2$ .

(2) Choose a labeling  $C_1, \dots, C_r$  of the irreducible components of  $C_P$ . For  $1 \leq j \leq r$ , let  $A_j = \text{Jac}(C_j)$ . Note that  $A_j$  is a principally polarized abelian variety of

dimension  $(m - 1)/2$ . By [1, Section 9.2, Example 8, page 246],  $J_P \simeq \bigoplus_{j=1}^r A_j$ , and the principal polarization on  $J_P$  decomposes as the product polarization.

Consider the CM-field  $F = \mathbb{Q}(\zeta_m)$ . Then  $\mathcal{O}_F = \mathbb{Z}[\zeta_m] \subset \text{End}(A_j)$  since  $\mu_m$  acts on  $C_j$  for  $1 \leq j \leq r$ . Since  $\text{deg}(F/\mathbb{Q}) = 2 \cdot \dim(A_j)$ , the abelian variety  $A_j$  has complex multiplication by  $\mathcal{O}_F$  for  $1 \leq j \leq r$ .  $\square$

**5.3. Shimura datum for Hurwitz spaces.** We turn to the main result of the paper. Recall that  $\gamma = (m, N, a)$  and  $r = N - 2$ .

**Definition 5.3.** Let  $P \in Z_\gamma$  be a distinguished point as described in Proposition 5.2. Suppose  $\mathbb{Q}(\zeta_m + \zeta_m^{-1})$  has units of independent signs. (This is guaranteed by Lemma 3.6 when  $m$  is an odd prime such that  $F = \mathbb{Q}(\zeta_m)$  has class number 1.)

For  $1 \leq j \leq r$ , define  $\beta_j$  and  $\xi_j = \beta_j^{-1}$  as follows. Recall that  $C_j$  is an irreducible component of  $C_P$ . The inertia type of the  $\mu_m$ -cover  $C_j \rightarrow \mathbb{P}^1$  determines its signature  $\mathfrak{f}_j$  by (2-2), which determines its CM-type  $\Phi_j$  as in (2-4). Let  $\beta_0$  be as in Lemma 3.7. By Corollary 4.7, there exists  $u_j \in \mathcal{U}_{F_0}$  such that  $\beta_j := u_j \beta_0$  satisfies conditions (1)–(3) of Corollary 4.3 for  $\Phi_j$ . Furthermore,  $\beta_j$  is the unique element satisfying conditions (1)–(3) for  $\Phi_j$ , up to multiplication by an element of  $N_{F/F_0}(\mathcal{U}_F)$ .

**Remark 5.4.** The CM-type  $(F, \Phi_j)$  of  $A_j$  may not be simple. For example, for the special family  $M$ [17] with  $m = 7$ , we show in Section 6.3 that  $(F, \Phi_2)$  is not simple.

Let  $F = \mathbb{Q}(\zeta_m)$  and denote by  $\mathcal{O}_F = \mathbb{Z}[\zeta_m]$  the ring of integers of  $F$ .

**Theorem 5.5.** *Let  $m$  be an odd prime. Let  $\gamma = (m, N, a)$  be a monodromy datum with  $N \geq 4$ . Suppose  $F = \mathbb{Q}(\zeta_m)$  has class number 1. Let  $r = N - 2$ .*

*Let  $P \in Z_\gamma$  be a distinguished point as described in Proposition 5.2. For  $1 \leq j \leq r$ , this determines an abelian variety  $A_j$  with CM-type  $(\mathcal{O}_F, \Phi_j)$ ; let  $\xi_j = \beta_j^{-1}$  be as in Definition 5.3.*

*Then the integral PEL datum for  $S_\gamma$  is given by*

- *the  $F$ -vector space  $V = F^r$ , together with the standard  $\mathcal{O}_F$ -lattice  $\Lambda = (\mathcal{O}_F)^r \subseteq V$ ;*
- *the Hermitian form  $H_B = \langle \cdot, \cdot \rangle$  on  $V$  taking integral values on  $\Lambda$  and defined by*

$$(5-1) \quad \langle x, y \rangle = \text{tr}_{F/\mathbb{Q}}(x B \bar{y}^T) \quad \text{for } B = \text{diag}[\xi_1, \dots, \xi_r] \in \text{GL}_r(F) = \text{GL}(V).$$

It is straightforward to compute  $B$  from  $\gamma$ ; we give many examples in Section 6.

*Proof.* The variety  $Z_\gamma$  is an irreducible algebraic subvariety of  $\mathcal{A}_g$ . By [17, Proposition 4.1], the lattice and Hermitian form for the integral PEL datum of  $S_\gamma$  can be computed at any point of  $Z_\gamma$ . Since  $m$  is an odd prime, there is a distinguished point  $P \in Z_\gamma$  as described in Proposition 5.2. We compute the lattice and Hermitian form at  $P$ .

Let  $V = H^1(C_P(\mathbb{C}), \mathbb{Q})$ . Note that  $r = N - 2 = 2g/(m - 1)$ . From (2-2), we deduce that there is an isomorphism of  $\mathbb{Q}[\mu_m]$ -modules  $V \simeq F^r$ , where multiplication of  $\mathbb{Q}[\mu_m]$  on  $F^r$  is given by the natural homomorphism  $\mathbb{Q}[\mu_m] \rightarrow F$ . The complex structure on  $V_{\mathbb{R}}$  is given as  $V_{\mathbb{R}} \simeq (F \otimes_{\mathbb{Q}} \mathbb{R})^r \simeq \bigoplus_{n=1}^m \mathbb{C}^{f(\tau_n)}$ . By (2-2),  $f(\tau_n) + f(\tau_{-n}) = r$  for all  $n \not\equiv 0 \pmod{m}$ .

Let  $\Lambda = H^1(C_P(\mathbb{C}), \mathbb{Z})$  be the first Betti cohomology of the curve  $C_P$ . When  $F$  has class number 1, we prove that there is an isomorphism of  $\mathbb{Q}[\mu_m]$ -modules  $V \simeq F^r$  such that the  $\mathbb{Z}[\mu_m]$ -lattice  $\Lambda \subset V$  maps isomorphically onto  $(\mathcal{O}_F)^r \subset F^r$  and the Hermitian form  $\psi$  on  $V$  maps to the diagonal Hermitian form  $H_B$  on  $F^r$  in (5-1). Note  $(\mathcal{O}_F)^r \subset F^r$  is a  $\mathbb{Z}[\mu_m]$ -lattice, and  $H_B$  is integral on  $(\mathcal{O}_F)^r$ .

By Lemma 3.6, if  $m$  is prime and  $F$  has class number 1, then  $F_0 = \mathbb{Q}(\zeta_m + \zeta_m^{-1})$  has narrow class number 1 and has units of independent signs. For  $1 \leq j \leq r$ , by hypothesis,  $A_j$  has dimension  $(m - 1)/2$ ; it has complex multiplication by  $\mathcal{O}_F$  and has CM-type  $(\mathcal{O}_F, \Phi_j)$ . Furthermore,  $\beta_j$  defines a principal polarization on  $A_j$ ; the corresponding Hermitian form is given by  $\mathbb{E}(\Phi(x), \Phi(y)) = \text{tr}_{F/\mathbb{Q}}(\xi_j x \bar{y})$  for  $x, y \in L$  by (4-1).

Since  $F$  has class number 1, there is a unique complex torus of CM-type  $(\mathcal{O}_F, \Phi_j)$  up to isomorphism by Theorem 4.1; namely,  $A_j \simeq \mathbb{C}^g / \Phi_j(\mathcal{O}_F)$ . Since  $J_P \simeq \bigoplus_{j=1}^r A_j$ , we deduce that  $\Lambda \simeq (\mathcal{O}_F)^r$ . The product polarization on  $J_P$  defines an integral Hermitian form of the prescribed signature on  $\Lambda$ . By [16, Appendix, Proposition 8], there is a unique Hermitian form of given signature on  $\Lambda$ . Thus the Hermitian form on  $\Lambda$  is the one in (5-1).  $\square$

**Remark 5.6.** The values of  $m$  for which Theorem 5.5 applies are 3, 5, 7, 11, 13, 17, 19.

For  $m = 3, 5, 11, 17$ , it is not necessary to refer to [16, Appendix, Proposition 8]; the reason is that  $(F, \Phi_j)$  is automatically simple for  $1 \leq j \leq r$  by Lemmas 3.8 and 3.9. Thus by Corollary 4.6(1), there is a unique principally polarized CM-abelian variety of type  $(\mathcal{O}_F, \Phi_j)$  up to isomorphism. In Section 6.4, we give an alternative approach for  $m = 7$  that relies on Corollary 4.6(2) and does not refer to [16, Appendix, Proposition 8].

The strategy of Theorem 5.5 may apply even when  $m$  is composite; see Section 7.

## 6. Applications for $m$ prime

Theorem 5.5 gives a method to determine the integral PEL datum for  $S_\gamma$  for all monodromy data  $\gamma = (m, N, a)$  when  $m = 3, 5, 7, 11, 13, 17, 19$ , so that  $F = \mathbb{Q}(\zeta_m)$  has class number 1. In this section, we give examples of this. Specifically:

- In Section 6.1, when  $m = 3$ , we explicitly determine the Hermitian form for all  $\gamma$ .
- In Sections 6.2 and 6.3, when  $m = 5, 7$ , we give explicit examples for all equivalence classes of  $\gamma$  when  $N = 4$ .

- In Section 6.4, we follow an alternative approach for  $m = 7$  which avoids the possible occurrence of nonsimple CM-types at the distinguished point.

In particular, we determine the integral PEL datum for the 6 special families for which the degree  $m$  is an odd prime; these are denoted by  $M[n]$  as in [12, Table 1].

Given a signature  $\mathfrak{f}$ , let  $\Phi$  be the CM-type of  $F$  determined by  $\mathfrak{f}$ , as in (2-4). Given the  $F$ -vector space  $V = F^r$ , together with the standard  $\mathcal{O}_F$ -lattice  $\Lambda = (\mathcal{O}_F)^r \subseteq V$ , let  $H_B = \langle \cdot, \cdot \rangle$  denote the Hermitian form on  $V$  defined in (5-1).

**6.1. Integral PEL datum for all families when  $m = 3$ .**

**Example 6.1.** When  $m = 3$ , then  $\beta_0 = 3/(\zeta_3^2 - \zeta_3) = \sqrt{-3}$  by Lemma 3.7.

For  $\mathfrak{f}_1 = (1, 0)$ , set  $u_1 = -1$  and  $b_1 = u_1\beta_0$  so that  $b_1 = -\sqrt{-3}$ . Then  $\text{Im}(\sigma_1(b_1)) < 0$ .

For  $\mathfrak{f}_2 = (0, 1)$ , set  $u_2 = 1$  and  $b_2 = u_2\beta_0$  so that  $b_2 = \sqrt{-3}$ . Then  $\text{Im}(\sigma_2(b_2)) < 0$ .

The CM-type  $\Phi_i$  is simple by Lemma 3.8 and  $b_i$  satisfies Corollary 4.7 with respect to  $\Phi_i$ .

For  $m = 3$ , there are simple formulas relating the signature  $\mathfrak{f}$  and the inertia type  $a$ . If  $a$  has  $d_1$  entries of 1 and  $d_2$  entries of 2, then  $\mathfrak{f} = (f_1, f_2)$  with  $f_1 = (2d_1 + d_2 - 3)/3$  and  $f_2 = (d_1 + 2d_2 - 3)/3$ . Then  $d_1 = 2f_1 - f_2 + 1$  and  $d_2 = 2f_2 - f_1 + 1$ . One can check that  $0 \leq \max(f_1, f_2) \leq 2 \min(f_1, f_2) + 1$ .

**Corollary 6.2.** *Let  $m = 3$  and  $N \geq 4$ . Let  $\gamma = (3, N, a)$  be a monodromy datum with signature  $(f_1, f_2)$ . Let  $g = f_1 + f_2$ . Let  $S_\gamma$  be the component of the Shimura variety containing  $Z_\gamma$ .*

*Let  $F = \mathbb{Q}(\zeta_3)$  and  $\xi = -1/\sqrt{-3}$ . Then the integral PEL datum of  $S_\gamma$  has lattice  $(\mathcal{O}_F)^g$  with Hermitian form  $H_B$  where  $B \in \text{GL}_g(\mathcal{O}_F)$  is diagonal with  $f_1$  entries of  $\xi$  and  $f_2$  entries of  $-\xi$ .*

*Proof.* Consider the admissible  $\mu_3$ -cover  $\varphi: \mathcal{C}_P \rightarrow T$  represented by the distinguished point from Proposition 5.2. Here  $T$  is a tree of  $r = f_1 + f_2$  projective lines. Above  $f_1$  (resp.  $f_2$ ) components of  $T$ , the restriction of  $\varphi$  is a  $\mu_3$ -cover branched at 3 points with signature  $\mathfrak{f}_1 = (1, 0)$  (resp.  $\mathfrak{f}_2 = (0, 1)$ ). The result then follows from Theorem 5.5. □

In particular, Corollary 6.2 includes the three special families  $M[3]$ ,  $M[6]$ ,  $M[10]$ :

$M$	$(m, N, a)$	$B$
$M[3]$	$(3, 4, (1, 1, 2, 2))$	$\text{diag}[\xi, -\xi]$
$M[6]$	$(3, 5, (1, 1, 1, 1, 2))$	$\text{diag}[\xi, \xi, -\xi]$
$M[10]$	$(3, 6, (1, 1, 1, 1, 1, 1))$	$\text{diag}[\xi, \xi, \xi, -\xi]$

The cases  $M[6] : \xi[1, 1, -1]$  and  $M[10] : \xi[1, 1, 1, -1]$  match the table on [16, p. 1].

**6.2. Integral PEL datum when  $m = 5$ .** We illustrate Theorem 5.5 for several well-chosen examples when  $m = 5$ , including all families branched at  $N = 4$  points up to equivalence and one family branched at  $N = 5$  points. This includes the two special families when  $m = 5$ , namely  $M[11]$  and  $M[16]$ . A similar result can be obtained for any monodromy datum  $\gamma$  with  $N > 4$  by an inductive process.

**Example 6.3.** The following table summarizes the data for CM-types when  $m = 5$ .

$a$	$\mathfrak{f}$	$\Phi$	$\beta$
(4, 3, 3)	(0, 1, 0, 1)	{2, 4}	$\beta_1 = 5/(\zeta_5^3 - \zeta_5^2)$
(3, 1, 1)	(1, 1, 0, 0)	{1, 2}	$\beta_2 = 5/(\zeta_5 - \zeta_5^4)$
(1, 2, 2)	(1, 0, 1, 0)	{1, 3}	$\beta_3 = -\beta_1$
(2, 4, 4)	(0, 0, 1, 1)	{3, 4}	$\beta_4 = -\beta_2$

In the  $i$ -th line of the table, the CM-type  $\Phi_i$  is simple by Lemma 3.8 and  $\beta_i$  satisfies Corollary 4.7 with respect to  $\Phi_i$ . The automorphism  $\sigma_3$  permutes the rows via the cycle (1, 2, 3, 4) and its inverse  $\sigma_2$  permutes the fourth column via  $\beta_1 \rightarrow \beta_2 \rightarrow -\beta_1 \rightarrow -\beta_2 \rightarrow \beta_1$ .

*Proof.* When  $m = 5$ , then  $\beta_0 = 5/(\zeta_5^3 - \zeta_5^2)$  by Lemma 3.7. For  $\mathfrak{f}_1 = (0, 1, 0, 1)$ , set  $u_1 = 1$  and  $\beta_1 = u_1\beta_0$ . We compute that  $\text{Im}(\sigma_j(\beta_1)) < 0$  for  $j = 2, 4$ . For  $\mathfrak{f}_2 = (1, 1, 0, 0)$ , set  $u_2 = (\zeta_5^3 - \zeta_5^2)/(\zeta_5 - \zeta_5^4)$  and  $\beta_2 = u_2\beta_0$ . We compute that  $\text{Im}(\sigma_j(\beta_2)) < 0$  for  $j = 1, 2$ . The signature  $\mathfrak{f}_3 = (1, 0, 1, 0)$  (resp.  $\mathfrak{f}_4 = (0, 0, 1, 1)$ ) is the complex conjugate of  $\mathfrak{f}_1$  (resp.  $\mathfrak{f}_2$ ), which negates the value of  $\beta$ .  $\square$

**Corollary 6.4.** *Let  $m = 5$  and  $F = \mathbb{Q}(\zeta_5)$ . Every family of  $\mu_5$ -covers of  $\mathbb{P}^1$  with  $N = 4$  is equivalent to either (i), (ii) or  $M[11]$  in the table below. Recall  $\beta_1, \beta_2 \in \mathcal{O}_F$  from Example 6.3 and let  $\xi_i = \beta_i^{-1}$ . For the monodromy data  $\gamma = (5, N, a)$  and  $r = N - 2$  as below, the integral PEL datum of  $S_\gamma$  has lattice  $(\mathcal{O}_F)^r$  with Hermitian form  $H_B$  where  $B \in \text{GL}_r(\mathcal{O}_F)$  is as below.*

$M$	$(5, N, a)$	$B$
(i)	(5, 4, (1, 1, 4, 4))	$\text{diag}[\xi_2, -\xi_2]$
(ii)	(5, 4, (1, 2, 3, 4))	$\text{diag}[-\xi_1, \xi_1]$
$M[11]$	(5, 4, (1, 3, 3, 3))	$\text{diag}[\xi_1, \xi_2]$
$M[16]$	(5, 5, (2, 2, 2, 2, 2))	$\text{diag}[-\xi_1, -\xi_2, -\xi_1]$

*Proof.* Suppose  $m = 5$  and  $N = 4$  and let  $a$  be the inertia type of  $\gamma$ . If three of the values of  $a$  are the same, the family is equivalent to the one with  $a = (1, 3, 3, 3)$ , which is  $M[11]$ . If two of the values of  $a$  are the same, the family is equivalent to (i). If all values of  $a$  are distinct, the family is equivalent to (ii).

By Theorem 5.5, it suffices to find the CM-type  $(F, \Phi_i)$  for the abelian varieties  $A_i$  in the decomposition of the Jacobian of  $\mathcal{C}_P$ . We refer to [8, Remark 5.2,

Lemma 6.4] for information about the admissible degeneration, given in shorthand by: (i)  $(1, 1, 3) + (2, 4, 4)$ ; (ii)  $(1, 2, 2) + (3, 3, 4)$ ;  $M[11] (1, 3, 1) + (4, 3, 3)$ ; and  $M[16] (2, 2, 1) + (4, 2, 4) + (1, 2, 2)$ . Using the table in Example 6.3, we find the entries of the diagonal of  $B$ .  $\square$

**Remark 6.5.** The family (ii), with monodromy datum  $\gamma = (5, 4, (1, 2, 3, 4))$ , has a second degeneration of the form  $(1, 3, 1) + (4, 2, 4)$ , whose Hermitian form has matrix  $B' = \text{diag}[\xi_2, -\xi_2]$ . We give two reasons why the Hermitian forms determined by  $B'$  and  $B = \text{diag}[-\xi_1, \xi_1]$  are isomorphic.

First, by Lemma 2.1, the automorphism  $\sigma_2$  takes the inertia types in the first degeneration to those in the second by multiplying the entries by 3. By Corollary 4.7(1), the action on the entries of  $B$  is via  $\sigma_{2^{-1}} = \sigma_3$  and

$$\sigma_3(B) = \text{diag}[\sigma_3(-\xi_1), \sigma_3(\xi_1)] = \text{diag}[\xi_2, -\xi_2] = B'.$$

Second, for family (ii),  $S_\gamma$  has signature type  $(1, 1, 1, 1)$ ; hence its reflex field is  $\mathbb{Q}$  (which is smaller than  $F_0 \subset \mathbb{R}$ ). The matrices  $B$  and  $B'$  are conjugate under the action of  $\sigma_3 \in \text{Gal}(F_0/\mathbb{Q})$  and correspond to the two choices of a  $\mathbb{Q}$ -linear embedding  $F_0 \hookrightarrow \mathbb{R}$ .

**Remark 6.6.** To compare with Shimura’s work, write  $w = \zeta_5 + \zeta_5^4$ . Then we have  $w^2 + w - 1 = 0$ . So  $w = (-1 + \sqrt{5})/2$ . Then  $\xi_2/\xi_1 = -w - 1 = -(1 + \sqrt{5})/2$  so  $\xi_1/\xi_2 = (1 - \sqrt{5})/2$ .

Consider the family  $\gamma' = (5, 4, (1, 1, 1, 1))$ . A careful look at [16, Section 5] shows that Shimura replaced  $\zeta_5$  by  $\zeta_5^3$  in his computation for this family. This has the effect of switching to the family  $M[16]$  with  $\gamma = (5, 4, (2, 2, 2, 2, 2))$ ; indeed, Shimura computes that the signature is  $(2, 0, 3, 1)$ . By line 4 of the table in Corollary 6.4, the family  $\gamma$  has  $B = -\xi_1[1, 1, \xi_2/\xi_1] = -\xi_1[1, 1, -(1 + \sqrt{5})/2]$ . This does not exactly match what is written in line 5 of the table on [16, page 1], namely  $[1, 1, \xi_1/\xi_2] = [1, 1, (1 - \sqrt{5})/2]$ , but it has the same sign signature and thus yields an isomorphic Hermitian form.

Consider the family  $\gamma' = (5, 4, (2, 1, 1, 1))$ . The details for this family are not included in [16] but it appears that Shimura replaced  $\zeta_5$  by  $\zeta_5^3$  in his computation for this family also. This has the effect of switching to the family  $M[11]$  with  $\gamma = (5, 4, (1, 3, 3, 3))$ . By line 3 of the table in Corollary 6.4, the family  $\gamma$  has  $B = \xi_2[1, \xi_1/\xi_2] = \xi_2[1, (1 - \sqrt{5})/2]$ . This matches what is written in line 4 of the table on [16, page 1].

**6.3. Integral PEL datum when  $m = 7$ .** We illustrate Theorem 5.5 for several well-chosen examples when  $m = 7$ , including all families branched at  $N = 4$  points up to equivalence. This includes the special family  $M[17]$ . A similar result can be obtained for any monodromy datum  $\gamma$  with  $N > 4$  by an inductive process.

**Example 6.7.** The following table summarizes the cases when  $m = 7$ .

$a$	$\mathfrak{f}$	$\Phi$	$\beta$
(1, 1, 5)	(1, 1, 1, 0, 0, 0)	{1, 2, 3}	$\beta_1 = 7/(\zeta_7 - \zeta_7^6)$
(3, 3, 1)	(1, 0, 1, 0, 1, 0)	{1, 3, 5}	$\beta_2 = 7/(\zeta_7^3 - \zeta_7^4)$
(2, 2, 3)	(1, 0, 0, 1, 1, 0)	{1, 4, 5}	$\beta_3 = 7/(\zeta_7^2 - \zeta_7^5)$
(6, 6, 2)	(0, 0, 0, 1, 1, 1)	{4, 5, 6}	$\beta_4 = -\beta_1$
(4, 4, 6)	(0, 1, 0, 1, 0, 1)	{2, 4, 6}	$\beta_5 = -\beta_2$
(5, 5, 4)	(0, 1, 1, 0, 0, 1)	{2, 3, 6}	$\beta_6 = -\beta_3$
(1, 2, 4)	(1, 1, 0, 1, 0, 0)	{1, 2, 4}	$\beta = -\frac{7(\zeta_7^3 - \zeta_7^4)}{(\zeta_7 - \zeta_7^6)(\zeta_7^2 - \zeta_7^5)}$
(3, 1, 5)	(0, 0, 1, 0, 1, 1)	{3, 5, 6}	$\beta' = -\beta$

**Lemma 6.8.** *In the  $i$ -th line of the table in Example 6.7, for  $1 \leq i \leq 6$ , the CM-type  $\Phi_i$  is simple, and the element  $\beta_i$  satisfies Corollary 4.7 with respect to  $\Phi_i$ . The generator  $\sigma_3$  of the Galois group  $\text{Gal}(\mathbb{Q}(\zeta_7)/\mathbb{Q})$  permutes these via  $\sigma_3^{i-1}(\beta_1) = \beta_i$ , for  $i = 1, \dots, 6$ .*

*In the last two lines of the table, the element  $\beta$  (resp.  $\beta'$ ) satisfies Corollary 4.7 with respect to the CM-type  $\Phi = \{1, 2, 4\}$  (resp.  $\Phi' = \{3, 5, 6\}$ ). The CM types  $\Phi, \Phi'$  are not simple.*

*Proof.* When  $m = 7$ , then  $\beta_0 = 7/(\zeta_7^4 - \zeta_7^3)$  by Lemma 3.7.

For  $a_1 = (1, 1, 5)$  and  $\mathfrak{f}_1 = (1, 1, 1, 0, 0, 0)$ , set  $u_1 = (\zeta_7^4 - \zeta_7^3)/(\zeta_7 - \zeta_7^6)$  and  $\beta_1 = u_1\beta_0$ . We compute that  $\text{Im}(\sigma_j(\beta_1)) < 0$  for  $j = 1, 2, 3$ .

Let  $\sigma = \sigma_5$ , which is a generator of  $\text{Gal}(\mathbb{Q}(\zeta_7)/\mathbb{Q})$  and has inverse  $\sigma_3$ . Then, by Lemmas 2.1 and 2.4, for  $1 \leq i \leq 6$ , the action of  $\sigma_5^{i-1}$  changes the inertia type to  $a_i := (3^{i-1}) \cdot a_1$  and the CM-type to  $\Phi_i := 5^{i-1} \cdot \Phi$ , which also determines the signature  $\mathfrak{f}_i$ . Also, the element  $\beta_i = \sigma_3^{i-1}(\beta_1)$  satisfies Corollary 4.7 with respect to  $\Phi_i$ .

For  $a = (1, 2, 4)$  and  $\mathfrak{f} = (1, 1, 0, 1, 0, 0)$ , set

$$u = -\frac{(\zeta_7^4 - \zeta_7^3)^2}{(\zeta_7 - \zeta_7^6)(\zeta_7^2 - \zeta_7^5)} \quad \text{and} \quad \beta = u\beta_0 = -\frac{7(\zeta_7^3 - \zeta_7^4)}{(\zeta_7 - \zeta_7^6)(\zeta_7^2 - \zeta_7^5)}.$$

Then  $\text{Im}(\sigma_j(\beta_2)) < 0$  for  $j = 1, 2, 4$ . The last line follows from Lemma 2.1 by applying complex conjugation to the previous line.

By Lemma 3.9,  $\Phi_i$  is simple unless it is  $\{1, 2, 4\}$  or  $\{3, 5, 6\}$ . □

For  $m = 7$  and  $N = 4$ , every family is equivalent to one in the next result.

**Corollary 6.9.** *Let  $m = 7$  and  $F = \mathbb{Q}(\zeta_7)$ . Recall  $\beta_1, \beta_2, \beta_3, \beta \in \mathcal{O}_F$  from Example 6.7. Let  $\xi_i = \beta_i^{-1}$  for  $i = 1, 2, 3$  and  $\xi = \beta^{-1}$ . For the monodromy*

data  $\gamma = (7, 4, a)$  as below, the integral PEL datum of  $S_\gamma$  has lattice  $(\mathcal{O}_F)^2$  with Hermitian form  $H_B$  where  $B \in \text{GL}_2(\mathcal{O}_F)$  is as below.

$M$	$a$	$B$
(i)	(1, 1, 2, 3)	$\text{diag}[\xi_1, \xi_3]$
(ii)	(1, 1, 6, 6)	$\text{diag}[\xi_1, -\xi_1]$
(iii)	(1, 2, 5, 6)	$\text{diag}[\xi_1, -\xi_1]$
$M[17]$	(2, 4, 4, 4)	$\text{diag}[-\xi_2, \xi]$

*Proof.* We refer to [8, Remark 5.2, Lemma 6.4] for information about the admissible degeneration of each family, given in shorthand by (i)  $(1, 1, 5) + (2, 2, 3)$ ; (ii)  $(1, 1, 5) + (2, 6, 6)$ ; (iii)  $(1, 5, 1) + (6, 2, 6)$ ; and  $M[17]$   $(4, 4, 6) + (1, 4, 2)$ . For each family, consider the CM-types  $(F, \Phi_1)$  and  $(F, \Phi_2)$  for the abelian varieties  $A_i$  in the decomposition of the Jacobian at the distinguished point arising from the degeneration. We compute the associated integral PEL data by Theorem 5.5, using the table in Example 6.7.  $\square$

**Remark 6.10.** In cases (i), (ii), and (iii), both  $(F, \Phi_1)$  and  $(F, \Phi_2)$  are simple. For  $M[17]$ , the CM-type  $(F, \Phi_2)$  is not simple, but this is not a concern by [16, Appendix, Proposition 8] or Section 6.4. For  $M[17]$ , we note that  $B = \xi[1, -\xi_2/\xi]$  where  $-\xi_2/\xi = \zeta_7 + \zeta_7^6 - 1$ . In line 6 of the table on [16, page 1], for the related family  $\gamma = (7, 4, (4, 1, 1, 1))$ , Shimura writes the Hermitian form as  $H_B$  with  $B = \xi[1, -\sin(3\pi/7)/\sin(2\pi/7)]$ .

**6.4. Another approach for  $M[17]$ .** In this section, we compute the integral PEL datum for the family  $M[17]$  using another approach. This approach utilizes a different kind of distinguished point  $Q$  in the family, namely one that represents a curve  $C_Q$  with extra automorphisms. In this section, we see that the Jacobian  $J_Q$  has complex multiplication by a larger field and that its CM-type is simple.

In fact, we proved that every positive-dimensional family of  $\mu_7$ -covers of  $\mathbb{P}^1$  has a distinguished point representing a product of principally polarized abelian varieties, each of which has complex multiplication with a CM-type that is simple. In this way, one can avoid the use of [16, Appendix, Proposition 8] when  $m = 7$ . We omit the details of this.

We start by finding another distinguished point in the family  $Z_\gamma$ , under certain restrictive conditions on  $\gamma$ . Similarly to Notation 3.5, for  $0 \leq n \leq 3m$  with  $\text{gcd}(n, 3m) = 1$ , let  $\sigma_n$  be the embedding  $\mathbb{Q}(\zeta_{3m}) \hookrightarrow \mathbb{C}$  determined by  $\sigma_n(\zeta_{3m}) = \zeta_{3m}^n$ .

**Proposition 6.11.** *Let  $\gamma = (m, 4, a)$  where  $m > 3$  is prime and  $a = (1, 1, 1, m - 3)$ . Then  $Z_\gamma$  has a distinguished point  $Q$ . More specifically: in the family of  $\mu_m$ -covers with monodromy datum  $\gamma$ , there is a point which represents a  $\mu_m$ -cover*

$\psi : C_Q \rightarrow \mathbb{P}^1$ , where  $C_Q$  is a curve of genus  $m - 1$  having an automorphism of order 3.

The Jacobian of  $C_Q$  has complex multiplication by  $(\mathbb{Z}[\zeta_{3m}], \Phi_Q)$ , where, for any  $0 < n < 3m$  with  $(n, 3m) = 1$ , the embedding  $\sigma_n$  is in  $\Phi_Q$  if and only if

$$\begin{aligned} n \in [0, 2m/3] \cup [m, 5m/3] \cup [2m, 8m/3] & \text{ if } n \equiv 1 \pmod{3}, \text{ or} \\ n \in [0, m/3] \cup [m, 4m/3] \cup [2m, 7m/3] & \text{ if } n \equiv 2 \pmod{3}. \end{aligned}$$

It is possible to generalize Proposition 6.11, by replacing 3 by an odd prime  $\ell$  relatively prime to  $m$  and letting  $N = \ell + 1$  and  $a = (1, \dots, 1, m - \ell)$ . We omit this generalization.

*Proof of Proposition 6.11.* Let  $C_Q$  be the smooth projective curve with equation  $y^m = x^3 - 1$ . It admits a  $\mu_m$ -cover to  $\mathbb{P}^1$  branched at  $1, \zeta_3, \zeta_3^2, \infty$  with inertia type  $a = (1, 1, 1, m - 3)$ . The genus of  $C_Q$  is  $m - 1$  by (2-1). Also  $C_Q$  has an automorphism  $(x, y) \mapsto (\zeta_3 x, y)$  of order 3.

The Jacobian  $J_Q = \text{Jac}(C_Q)$  is a principally polarized abelian variety having dimension  $m - 1$ . Let  $F = \mathbb{Q}(\zeta_m)$ , and  $L = \mathbb{Q}(\zeta_{3m})$ . The field  $L$  is a CM-field of degree  $2 \cdot \dim(J_Q)$ . Then, the inclusion  $\mathcal{O}_F \subset \text{End}(J_Q)$  extends to an inclusion  $\mathcal{O}_L = \mathcal{O}_F[\zeta_3] \subset \text{End}(J_Q)$ . Thus  $J_Q$  has complex multiplication by  $\mathcal{O}_L = \mathbb{Z}[\zeta_{3m}]$ .

Consider the morphism  $\phi : C_Q \rightarrow \mathbb{P}^1$ , taking  $(x, y) \mapsto x^3$ . Then  $\phi$  is a  $\mu_{3m}$ -cover of  $\mathbb{P}^1$ , branched at  $0, 1, \infty$ . We compute the CM-type of  $J_Q$  by finding the inertia type and signature type of  $\phi$ . Let  $b_0$  (resp.  $b_1$ , resp.  $b_\infty$ ) denote the element of  $\mathbb{Z}/3m\mathbb{Z}$  that determines the canonical generator of inertia of  $\phi$  above  $0$ , (resp.  $1$ , resp.  $\infty$ ).

Note that  $b_1 = 3$ . This is because  $\phi$  is branched at the 3 points  $x = 1, \zeta_3, \zeta_3^2$  that lie above  $x^3 = 1$ ; the canonical generator of inertia of  $\phi$  at the points of  $C_Q$  above these is, by definition, the automorphism identified with  $\zeta_7^1 = \zeta_{21}^3$ .

Without loss of generality,  $b_0 = m$ . To see this, note there are  $m$  points of  $C_Q$  above  $x^3 = 0$ , so  $\gcd(b_0, 3m) = m$ ; this implies that  $b_0 = 2m$  or  $b_0 = m$ ; possibly after replacing the order 3 automorphism with its inverse, we can suppose that  $b_0 = m$ . Third,  $b_\infty = 3m - b_0 - b_1 = 2m - 3$ , so  $(b_0, b_1, b_\infty) = (m, 3, 2m - 3)$ .

We compute the signature  $\mathfrak{f}$  of  $\phi$ . Let  $0 < n < 3m$  and  $(n, 3m) = 1$ . By (2-2):

$$\mathfrak{f}(\sigma_n) = -1 + \langle -n/3 \rangle + \langle -n/m \rangle + \langle (-2n)/3 + (n/m) \rangle;$$

that is,

$$\mathfrak{f}(\sigma_n) = \begin{cases} -1 + \frac{2}{3} + \langle -n/m \rangle + \langle -\frac{2}{3} + n/m \rangle & \text{if } n \equiv 1 \pmod{3}, \\ -1 + \frac{1}{3} + \langle -n/m \rangle + \langle -\frac{1}{3} + n/m \rangle & \text{if } n \equiv 2 \pmod{3}, \end{cases}$$

We deduce that  $\mathfrak{f}(\sigma_n) = 1$  if  $n \equiv 1 \pmod{3}$  and  $n \in [0, 2m/3] \cup [m, 5m/3] \cup [2m, 8m/3]$  or if  $n \equiv 2 \pmod{3}$  and  $n \in [0, m/3] \cup [m, 4m/3] \cup [2m, 7m/3]$ ; otherwise  $\mathfrak{f}(\sigma_n) = 0$ .

Finally, by (2-3), the CM-type  $\Phi_Q$  of  $J_Q$  is as given in the statement.  $\square$

For  $m = 7$ , we verify that the CM-type  $\Phi_Q$  of the Jacobian of  $C_Q$  is simple.

**Lemma 6.12.** *When  $m = 7$ , then the CM-type  $\Phi_Q$  in Proposition 6.11 is simple.*

*Proof.* Let  $C_Q$  be the smooth projective curve with equation  $y^7 = x^3 - 1$ . From Proposition 6.11, its CM-type is  $\Phi = \{\tau_n \mid n = 1, 2, 4, 8, 10, 16\}$ . We check that  $\Phi$  is simple by showing that it is not induced from any proper CM-field. Let  $\sigma_i : \mathbb{Q}(\zeta_{3m}) \rightarrow \mathbb{C}$  be the embedding given by  $\sigma_i(\zeta_{3m}) = \zeta_{3m}^i$ , for  $1 \leq i \leq 3m$  with  $\gcd(i, 3m) = 1$ . Every subgroup of  $\text{Gal}(\mathbb{Q}(\zeta_{3m})/\mathbb{Q})$  not containing complex conjugation contains either  $\sigma_8$  (order 2),  $\sigma_{13}$  (order 2) or  $\sigma_4$  (order 3). It suffices to show that the subgroup generated by each of these has a coset not contained in  $\Phi$  but having nontrivial intersection with  $\Phi$ , so that  $\Phi$  is not a union of cosets. For  $\langle \sigma_8 \rangle$ , this is true because of the coset  $\{\sigma_4, \sigma_{11}\}$ . For  $\langle \sigma_{13} \rangle$ , this is true because of the coset  $\{\sigma_1, \sigma_{13}\}$ . For  $\langle \sigma_4 \rangle$ , this is true because of the coset  $\{\sigma_2, \sigma_8, \sigma_{11}\}$ .  $\square$

The following lemma will be helpful.

**Lemma 6.13.** *In the field  $L = \mathbb{Q}(\zeta_{21})$ , write  $\zeta = \zeta_{21}$ . Let  $\beta_3 = 7/(\zeta^6 - \zeta^{15})$ . Let*

$$(6-1) \quad \alpha = (\zeta^7 - \zeta^{14})(\zeta^2 - \zeta^{19}) = (\zeta^9 + \zeta^{12}) - (\zeta^5 + \zeta^{16}).$$

*With respect to the CM-type  $\Phi = \{1, 2, 4, 8, 10, 16\}$  of  $L$ , the following element  $z$  satisfies conditions (1)–(3) in Corollary 4.3:*

$$(6-2) \quad z = \beta_3 \alpha = 21(\zeta^2 - \zeta^{19})/((\zeta^{14} - \zeta^7)(\zeta^6 - \zeta^{15})).$$

*Proof.* We verify the first claim by computation using that  $\zeta^{14} - \zeta^7 = -\sqrt{-3}$ . We verify the second claim from Lemma 3.7, part (4). Note that  $z = -\sigma_4(\beta_0)$  for  $\beta_0$  as in loc. cit.  $\square$

We now compute the integral PEL datum of  $M[17]$  using Proposition 6.11.

**Proposition 6.14.** *Let  $m = 7$  and  $F = \mathbb{Q}(\zeta_7)$ . Let  $\xi_1 = \beta_1^{-1}$  where  $\beta_1 \in \mathcal{O}_F$  is defined in Example 6.7. Consider the units  $u_1 = \zeta_7^2 + \zeta_7^5$  and  $v = (1 + \zeta_7^3 + \zeta_7^4)^{-1}$  of  $\mathbb{Z}[\zeta_7]$ . For the family  $M[17]$ , with monodromy datum  $\gamma = (7, 4, (2, 4, 4, 4))$ , the integral PEL datum of  $S_\gamma$  has lattice  $\Lambda = (\mathcal{O}_F)^2$  with Hermitian form  $H_B$  where*

$$B = \xi_1 v \begin{bmatrix} u_1 & -\zeta_7^2 \\ -\zeta_7^5 & u_1 \end{bmatrix}.$$

*Proof.* Consider the family  $Z = Z_\gamma$  with monodromy datum  $\gamma = (7, 4, (1, 1, 1, 4))$ . Because  $(2, 4, 4, 4) = 4 \cdot (4, 1, 1, 1)$  and  $4 \equiv 2^{-1} \pmod{7}$ , the action of  $\sigma_2$  takes  $Z$  to the family  $M[17]$ .

Let

$$(6-3) \quad A = \frac{\zeta_7^2 - \zeta_7^5}{7(1 + \zeta_7 + \zeta_7^6)} \begin{bmatrix} \zeta_7^3 + \zeta_7^4 & -\zeta_7^4 \\ -\zeta_7^3 & \zeta_7^3 + \zeta_7^4 \end{bmatrix}.$$

One can check that  $A$  is integral away from 7. We claim that the integral PEL datum of  $Z$  is given by the lattice  $\Lambda = (\mathcal{O}_F)^2$  and the Hermitian form for the matrix  $A$ . This is sufficient to prove the proposition because, by the last statement of Corollary 4.3, the integral PEL datum of  $M[17]$  is then given by the lattice  $\Lambda = (\mathcal{O}_F)^2$  and the Hermitian form for the matrix  $B = \sigma_2^{-1}(A)$ .

We turn to computing the integral PEL datum of  $Z$ . Let  $L = \mathbb{Q}(\zeta_{21})$ . Consider the CM-type  $\Phi = \{1, 2, 4, 8, 10, 16\}$  for  $L$ . Consider the curve  $C_Q$  of genus 6 given by the equation  $y^7 = x^3 - 1$  and the associated  $\mu_7$ -cover  $C_Q \rightarrow \mathbb{P}^1$ . By Proposition 6.11: this cover is represented by a point  $Q$  of  $Z$ ; the Jacobian  $\text{Jac}(C_Q)$  is a principally polarized abelian variety with complex multiplication by  $(\mathcal{O}_L, \Phi_Q)$ ; and this CM-type is simple by Lemma 6.12. In other words,  $Q$  is a simple distinguished point of  $Z$ .

Note that  $L$  has class number 1 and  $L_0$  has narrow class number 2. By Corollary 4.6, there exists a unique principally polarized CM-abelian variety of type  $(\mathcal{O}_L, \Phi)$  up to isomorphism. By Lemma 6.13, the element  $z$  from (6-2) satisfies conditions (1)–(3) in Corollary 4.3 with respect to  $\Phi$ . Thus  $\text{Jac}(C_Q)$  is isomorphic to the torus  $\mathbb{C}^6/\Phi(\mathcal{O}_L)$ , with principal polarization given by  $\xi = z^{-1}$ . Furthermore, the integral PEL datum of  $Z$  is given by the lattice  $\Lambda = \mathcal{O}_L \subset V = L$  and the Hermitian form  $\langle x, y \rangle = \text{tr}_{L/\mathbb{Q}}(x\xi\bar{y}^T)$ . Note that  $\mathcal{O}_L = \mathbb{Z}[\zeta_{21}] = \mathcal{O}_F[\zeta_3]$ . With respect to the ordered basis  $1, \zeta_3$  for  $\mathcal{O}_L$  over  $\mathcal{O}_F$ , the Hermitian form is given by a matrix in  $\text{GL}_2(\mathcal{O}_F)$ . By Lemma 6.15, this matrix is  $A$ .  $\square$

**Lemma 6.15.** *With respect to the ordered basis  $1, \zeta_3$  for  $\mathcal{O}_L$  over  $\mathcal{O}_F$ , the Hermitian form  $\langle x, y \rangle = \text{tr}_{L/\mathbb{Q}}(x\xi\bar{y}^T)$  is given by the matrix  $A \in \text{GL}_2(\mathcal{O}_F)$ , where  $A$  is given in (6-3).*

*Proof.* Let  $x, y \in \Lambda$ . By Lemma 6.13, since  $\beta_3 = 7/(\zeta_7^2 - \zeta_7^5)$  is in  $\mathcal{O}_{F_0}$ , we have

$$\langle x, y \rangle = \text{tr}_{L/\mathbb{Q}}(xz\bar{y}) = \text{tr}_{F/\mathbb{Q}}(\beta_3^{-1} \text{tr}_{L/F}(x\alpha^{-1}\bar{y})),$$

where  $\alpha$  is as in (6-1).

Let  $\tau = \sigma_8 \in \text{Gal}(L/\mathbb{Q})$ . Then  $\tau$  is the generator of  $\text{Gal}(L/F)$ , and

$$\text{tr}_{L/\mathbb{Q}}(x\alpha^{-1}\bar{y}) = \text{tr}_{F/\mathbb{Q}}(x\alpha^{-1}\bar{y} + \tau(x)\tau(\alpha^{-1})\tau(\bar{y})).$$

Write  $x = x_1 + x_2\zeta_3$  and  $y = y_1 + y_2\zeta_3$ , for  $x_1, x_2, y_1, y_2 \in \mathcal{O}_F$ . We compute

$$\text{tr}_{L/\mathbb{Q}}(x\alpha^{-1}\bar{y}) = \text{tr}_{F/\mathbb{Q}}((x_1 + x_2\zeta_3)\alpha^{-1}(\bar{y}_1 + \bar{y}_2\zeta_3^2) + (x_1 + x_2\zeta_3^2)\tau(\alpha^{-1})(\bar{y}_1 + \bar{y}_2\zeta_3)).$$

Write  $a_{1,1} = \alpha^{-1} + \tau(\alpha^{-1})$ ,  $a_{1,2} = \zeta_3^2\alpha^{-1} + \zeta_3\tau(\alpha^{-1})$ , and  $a_{2,1} = \zeta_3\alpha^{-1} + \zeta_3^2\tau(\alpha^{-1})$ . Then  $\text{tr}_{L/\mathbb{Q}}(x\alpha^{-1}\bar{y}) = \text{tr}_{F/\mathbb{Q}}(x_1\bar{y}_1a_{1,1} + x_1\bar{y}_2a_{1,2} + x_2\bar{y}_1a_{2,1} + x_2\bar{y}_2a_{1,1})$ . We compute that  $a_{1,1} = \alpha^{-1} + \tau(\alpha^{-1}) = (\zeta_7^4 + \zeta_7^3)/(1 + \zeta_7 + \zeta_7^6)$ ,  $a_{1,2} = -\zeta_7^4/(1 + \zeta_7 + \zeta_7^6)$ , and  $a_{2,1} = -\zeta_7^3/(1 + \zeta_7 + \zeta_7^6)$ . Thus,  $\langle x, y \rangle = \text{tr}_{F/\mathbb{Q}}((x_1, x_2)A(\bar{y}_1, \bar{y}_2)^T)$ .  $\square$

## 7. Applications for composite $m$

We illustrate how to extend Theorem 5.5 by computing the integral PEL datum in some cases when  $m$  is composite. In particular, we determine the integral PEL datum for 6 of the Moonen special families where  $m = 4, 6, 10$ .

**Remark 7.1.** Our techniques do not apply well for symplectic Shimura varieties, so we exclude the modular curve  $M[1]$  with  $m = 2$ , the Picard surface  $M[2]$  with  $m = 2$ , the family  $M[7]$  with  $m = 4$ , and  $M[12]$  with  $m = 6$ . We exclude  $M[15]$  with  $m = 8$  and  $M[20]$  with  $m = 12$ , since the CM-types for biquadratic CM-fields are never simple. Our techniques are not sufficient to handle  $M[13]$  with  $m = 6$  or  $M[19]$  with  $m = 9$ .

**7.1. Difficulties when  $m$  is composite.** The key to Theorem 5.5 is the existence of a distinguished point  $P$  in  $Z_\gamma$  satisfying the properties in Proposition 5.2. When  $m$  is composite, this is not always possible for the following reasons.

**Remark 7.2.** One issue when  $m$  is composite is that the curve  $C_P$  might not have compact type. For example, the family  $M[12]$ , with  $\gamma = (6, 4, (1, 1, 1, 3))$ , has no admissible degenerations of compact type. The reason is that the two covers with inertia types  $a_1 = (1, 1, 4)$  and  $a_2 = (2, 1, 3)$  would need to be joined at two points, leading to a cycle in the dual graph of  $C_P$ .<sup>3</sup> The same is true for  $M[7]$ , with  $\gamma = (4, 4, (1, 1, 1, 1))$ .

**Remark 7.3.** Another issue when  $m$  is composite is that the integral group algebra  $\mathbb{Z}[\mu_m]$  has nontrivial index in  $\prod_{1 \leq d|m} \mathbb{Z}[\zeta_d]$ . Hence, the Jacobian of a  $\mu_m$ -cover of  $\mathbb{P}^1$  branched at 3 points might not have complex multiplication by a maximal order. See Section 7.5.

**7.2. Integral PEL data for two families with  $m = 4$ .** We find the integral PEL datum for two of the three special families with  $m = 4$ :  $M[4]$  and  $M[8]$ ; we exclude  $M[7]$  as it is not unitary. When  $m = 4$ , the hypotheses of Proposition 5.2 are not satisfied but we can find a distinguished point in the family by direct computation.

**Example 7.4.** If  $m = 4$ , then  $\beta_0 = -2i$  from Lemma 3.7. Set  $u_1 = -1$ , so  $\beta_1 = -\beta_0 = 2i$ . If  $\mathfrak{f}_1 = (0, 1)$ , then  $\beta_1$  satisfies the 3 conditions of Corollary 4.3 and all CM-types of  $F = \mathbb{Q}(i)$  are simple.

**Corollary 7.5.** Let  $m = 4$  and  $F = \mathbb{Q}(i)$ . Let  $\xi = \beta_1^{-1} = 1/2i$ . For the family  $M$  with monodromy datum  $\gamma$  as in the next table (top of page 206), the integral PEL datum for  $S_\gamma$  has lattice  $(\mathcal{O}_F)^r$  with Hermitian form  $H_B$ , where  $r$  and  $B \in \mathrm{GL}_r(\mathcal{O}_F)$

<sup>3</sup>There is a typo in this case in [8, Lemma 6.4].

are as follows:

$M$	$\gamma = (m, N, a)$	$r$	$B \in \text{GL}_r(\mathcal{O}_F)$
$M[4]$	$(4, 4, (1, 2, 2, 3))$	2	$\text{diag}[\xi, -\xi]$
$M[8]$	$(4, 5, (1, 1, 2, 2, 2))$	3	$\text{diag}[\xi, -\xi, \xi]$

*Proof.* When  $m = 4$ , then  $a = (3, 2, 3)$  has signature  $\mathfrak{f} = (0, 1)$  and  $\sigma_2(\beta_1) < 0$ . Also  $a = (1, 2, 1)$  has signature  $\mathfrak{f} = (1, 0)$  and  $\sigma_1(-\beta_1) < 0$ . By [8, Lemma 6.4], there is a distinguished point in the family  $M[4]$  and the family  $M[8]$ . The family  $M[4]$  has an admissible degeneration, expressed in short as  $(1, 2, 1) + (3, 2, 3)$ . The family  $M[8]$  has an admissible degeneration, expressed as  $(1, 2, 1) + (3, 2, 3) + (1, 2, 1)$ . The result then follows from Theorem 5.5.  $\square$

The case  $M[8] : \xi[1, -1, 1]$  matches line 3 of the table on [16, page 1].

**7.3. Remark when  $m = 9$ .** The techniques in this paper are not sufficient to find the integral PEL datum for the special family  $M[19]$  when  $m = 9$ . This family has inertia type  $a = (3, 5, 5, 5)$  and its only admissible degeneration can be expressed in short hand as  $(3, 5, 1) + (8, 5, 5)$ . For  $a_2 = (8, 5, 5)$ , the Jacobian has complex multiplication by  $\mathbb{Q}[\zeta_3] \times \mathbb{Q}[\zeta_9]$ . As the action of  $\mathbb{Z}[\mu_9]$  does not extend to the maximal order  $\mathbb{Z}[\zeta_3] \times \mathbb{Z}[\zeta_9]$ , we do not know how to compute its lattice.

We expect that our technique is sufficient to determine the integral PEL datum when  $m = 9$ ,  $N = 4$  and  $a = (3, 1, 8, 6)$ . We leave the details to the reader to check.

**7.4. Generalities for twice a prime.**

**Notation 7.6.** Let  $m = 2m'$  where  $m'$  is an odd prime. Let  $F = \mathbb{Q}(\zeta_m)$  and  $F' = \mathbb{Q}(\zeta_{m'})$ . We identify  $F = F'$  and  $\mathbb{Z}[\zeta_m] = \mathbb{Z}[\zeta_{m'}]$ , by  $\zeta_m = -\zeta_{m'}^{(m'+1)/2}$ . There is a bijection between CM types  $\Phi$  for  $F$  and CM types  $\Phi'$  for  $F'$  where  $\sigma_j \in \Phi$  for an odd integer  $1 \leq j \leq m$  if and only if  $\sigma_{j'} \in \Phi'$ , where  $j'$  is the reduction of  $j$  modulo  $m'$ .

If  $\beta$  satisfies Corollary 4.7 with respect to the CM type  $(\mathbb{Z}[\zeta_{m'}], \Phi')$  then it also satisfies Corollary 4.7 with respect to the CM type  $(\mathbb{Z}[\zeta_m], \Phi)$ . This can be verified in general but we only need a few cases from Example 6.1 when  $m' = 3$  and Example 6.3 when  $m' = 5$ .

**Example 7.7.**

$m$	$\beta$	$(\mathbb{Z}[\zeta_{m'}], \Phi')$	$(\mathbb{Z}[\zeta_m], \Phi)$
6	$b_2 = \sqrt{-3}$	$(\mathbb{Z}[\zeta_3], \{2\})$	$(\mathbb{Z}[\zeta_6], \{5\})$
10	$\beta_1 = 5/(\zeta_5^3 - \zeta_5^2)$	$(\mathbb{Z}[\zeta_5], \{2, 4\})$	$(\mathbb{Z}[\zeta_{10}], \{7, 9\})$
10	$\beta_2 = 5/(\zeta_5 - \zeta_5^4)$	$(\mathbb{Z}[\zeta_5], \{1, 2\})$	$(\mathbb{Z}[\zeta_{10}], \{1, 7\})$

**7.5. Integral PEL data for 4 families with  $m = 6, 10$ .** We find the integral PEL datum for three special families with  $m = 6$ , namely  $M[5]$ ,  $M[9]$  and  $M[14]$ , and for the special family with  $m = 10$ , namely  $M[18]$ . For  $m = 6$ , we exclude  $M[12]$  as it is not unitary and  $M[13]$  (Remark 7.10). For  $m$  composite, the hypotheses of Proposition 5.2 are not satisfied. In each case, we find a distinguished point  $P$  in the family for which the issues in Remarks 7.2 and 7.3 do not occur.

**Corollary 7.8.** (Recall Notation 7.6.) *For the special family  $M[18]$  with monodromy datum  $\gamma = (10, 4, (3, 5, 6, 6))$ , the integral PEL datum of  $S_\gamma$  has lattice*

$$\Lambda = (\mathcal{O}_{F'}) \oplus (\mathcal{O}_F)^2$$

with Hermitian form  $H_B$ , where  $B \in \text{GL}_1(\mathcal{O}_{F'}) \times \text{GL}_2(\mathcal{O}_F)$  is  $\text{diag}[\xi_2] \oplus \text{diag}[\xi_1, \xi_2]$  with  $\xi_i = \beta_i^{-1}$ .

*Proof.* Let  $m = 10$  and  $m' = 5$ . Let  $M = M[18]$  with  $\gamma = (10, 4, (3, 5, 6, 6))$  and signature type  $\mathfrak{f} = (1, 1, 0, 1, 0, 0, 2, 0, 1)$ . The proof is similar to that of Theorem 5.5. We produce a simple distinguished point  $P$  in the family similar to that in Proposition 5.2(1); it represents an admissible  $\mu_m$ -cover  $\psi : C_P \rightarrow T$ , where  $T$  is a tree of projective lines and  $C_P$  is a curve of compact type such that each of its irreducible components is a curve admitting a  $\mu_m$ -cover of  $\mathbb{P}^1$  branched at 3 points. We verify by direct computation that each irreducible component of  $C_P$  has complex multiplication by either  $\mathbb{Z}[\zeta_m]$  or  $\mathbb{Z}[\zeta_{m'}]$ .

Consider the  $\mu_5$ -cover  $\psi_2 : C_2 \rightarrow \mathbb{P}^1$  branched at three points, with inertia type  $a_2 = (4, 3, 3)$ , and signature  $\mathfrak{f}_2 = (0, 1, 0, 1)$ . Then  $A_2 = \text{Jac}(C_2)$  has complex multiplication by  $(\mathbb{Z}[\zeta_5], \{2, 4\})$ . Consider the induced curve  $\tilde{C}_2 = \text{Ind}_5^{10}(C_2)$ , which is the disconnected curve consisting of two copies of  $C_2$ , and the induced  $\mu_{10}$ -cover  $\Psi_2 : \tilde{C}_2 \rightarrow \mathbb{P}^1$ . Then  $\Psi_2$  is branched at three points and, somewhat imprecisely, we can say that it has inertia type  $(8, 6, 6) = \text{Ind}_5^{10}(4, 3, 3)$ . Above the first branch point, there are two points  $\eta_2$  and  $\eta'_2$  on  $\tilde{C}_2$ , and they are labeled by the two cosets of  $\mu_5 \subset \mu_{10}$ . Let  $\mathcal{A}_2 = \text{Jac}(\tilde{C}_2) \simeq A_2^2$ .

As explained in [10, Section 3.1], the signature type of  $\mathcal{A}_2$  is

$$\mathfrak{f}_2 = (0, 1, 0, 1, 0, 0, 1, 0, 1) = (0, 1, 0, 1, 0, 0, 0, 0, 0) + (0, 0, 0, 0, 0, 0, 1, 0, 1).$$

The action of  $\mathbb{Z}[\mu_{10}]$  on  $\mathcal{A}_2$  is the diagonal action of  $\mathbb{Z}[\zeta_5] \times \mathbb{Z}[\zeta_{10}]$  on  $A^2$ . The first (resp. second) copy of  $A_2$  has CM-type  $(\mathbb{Z}[\zeta_5], \{2, 4\})$  (resp.  $(\mathbb{Z}[\zeta_{10}], \{7, 9\})$ ); for this CM-type, the element  $\beta_1$  (resp.  $\beta_1$ ) satisfies Corollary 4.7, as seen in Example 7.7.

Consider the  $\mu_{10}$ -cover  $\psi_1 : C_1 \rightarrow \mathbb{P}^1$  branched at three points, with inertia type  $a_1 = (3, 5, 2)$ . Above the 3rd branch point, there are two points  $\eta_1$  and  $\eta'_1$  on  $C_1$  and they are labeled by the two cosets of  $\mu_5 \subset \mu_{10}$ . Let  $A_1 = \text{Jac}(C_1)$ ; it has signature type  $\mathfrak{f}_1 = (1, 0, 0, 0, 0, 0, 1, 0, 0)$ . By Lemma 2.3,  $A_1$  has complex multiplication

by  $(\mathbb{Z}[\zeta_{10}], \{1, 7\})$ . Since  $\{1, 7\} = \{3 \cdot 7, 3 \cdot 9\} \pmod{10}$ , the element  $\beta$  that satisfies Corollary 4.7 is  $\sigma_7(\beta_1) = \beta_2$ . All CM types of  $\mathbb{Q}(\zeta_5) = \mathbb{Q}(\zeta_{10})$  are simple.

Let  $\mathcal{C}$  denote the singular curve, whose components are  $C_1$  and  $\tilde{C}_2$ , formed by identifying  $\eta_1 \eta'_1$  with  $\eta_2$  and  $\eta'_2$ , so that the cosets are matched correctly, in two ordinary double points. The curve  $\mathcal{C}$  admits an admissible  $\mu_{10}$ -cover  $\Psi$  to a chain of two projective lines. By construction,  $\mathcal{C}$  has an action by  $\mathcal{O}_{F'} \oplus \mathcal{O}_F^2$ , with CM-type given by  $\text{diag}[\xi_2] \oplus \text{diag}[\xi_1, \xi_2]$ .

The last thing to check is that  $\Psi$  is represented by a point  $P \in Z_\gamma$ . Since  $\Psi$  is admissible, it can be deformed to a  $\mu_{10}$ -cover of  $\mathbb{P}^1$  with inertia type  $(3, 5, 6, 6)$ . This has signature type  $\mathfrak{f} = (1, 1, 0, 1, 0, 0, 2, 0, 1)$ . Hence,  $\Psi$  is represented by a point  $P \in Z_\gamma$  and by the preceding paragraph  $P$  is a simple distinguished point.  $\square$

**Corollary 7.9.** (Recall Notation 7.6.) *Let  $\xi = 1/\sqrt{-3}$ . For the special families  $M$  with monodromy datum  $\gamma = (6, N, a)$  as below, the integral PEL datum of  $S_\gamma$  has lattice  $\Lambda = (\mathcal{O}_{F'})^{r'} \oplus (\mathcal{O}_F)^r$  with Hermitian form  $H_B$  where  $r', r$ , and  $B$  are as follows:*

$M$	$\gamma = (m, N, a)$	$(r', r)$	$B \in \text{GL}_{r'}(\mathcal{O}_{F'}) \times \text{GL}_r(\mathcal{O}_F)$
$M[5]$	$(6, 4, (2, 3, 3, 4))$	$(0, 2)$	$\text{diag}[\xi, -\xi]$
$M[9]$	$(6, 4, (1, 3, 4, 4))$	$(1, 2)$	$\text{diag}[-\xi] \oplus \text{diag}[\xi, -\xi]$
$M[14]$	$(6, 5, (2, 2, 2, 3, 3))$	$(1, 3)$	$\text{diag}[\xi] \oplus \text{diag}[\xi, -\xi, \xi]$

*Proof.* The proof is very similar to that of Corollary 7.8 so we provide only a sketch.

(1) Let  $M = M[5]$  with  $\gamma = (6, 4, (2, 3, 3, 4))$  and  $\mathfrak{f} = (1, 0, 0, 0, 1)$ . Then  $C_P$  is the join of two  $\mu_6$ -covers with inertia types  $a_1 = (1, 2, 3)$  and  $a_2 = (3, 4, 5)$ . These have signatures  $\mathfrak{f}_1 = (1, 0, 0, 0, 0)$  and  $\mathfrak{f}_2 = (0, 0, 0, 0, 1)$ . By Lemma 2.3,  $A_1$  has CM by  $\mathcal{O}_F$  of type  $\Phi_1 = \{1\}$ , and  $A_2$  has CM by  $\mathcal{O}_F$  of type  $\Phi_2 = \{5\}$ . Let  $b_1 = -b_2 = -\sqrt{-3}$ . For  $i = 1, 2$ , the CM-type  $\Phi_i$  is simple, and the element  $b_i \in \mathcal{O}_F$  satisfies Corollary 4.7 with respect to  $\Phi_i$ .

(2) Let  $M = M[9]$  with  $\gamma = (6, 4, (1, 3, 4, 4))$  and  $\mathfrak{f} = (1, 1, 0, 0, 1)$ . Then  $C_P$  is the join of two covers with inertia types  $a_1 = (1, 2, 3)$  and  $a_2 = \text{Ind}_3^6(2, 2, 2)$ . These have signatures  $\mathfrak{f}_1 = (1, 0, 0, 0, 0)$  and  $\mathfrak{f}_2 = (0, 1, 0, 0, 1)$ . By Lemma 2.3,  $A_1 = \text{Jac}(C_1)$  has CM-type  $(\mathcal{O}_F, \{1\})$ , which has  $\beta = b_1 = -b_2$ . Also  $\mathcal{A}_2 \simeq A^2$  and the action of  $\mathbb{Z}[\mu_6]$  on  $\mathcal{A}_2$  is given by the diagonal action of  $\mathbb{Z}[\zeta_3] \times \mathbb{Z}[\zeta_6]$  on  $A^2$ . The first copy of  $A$  has CM-type  $(\mathbb{Z}[\zeta_3], \{2\})$  which has  $\beta = b_2$  and the second copy of  $A$  has CM-type  $(\mathbb{Z}[\zeta_6], \{5\})$  which has  $\beta = b_2$ .

(3) Let  $M = M[14]$  with  $\gamma = (6, 5, (2, 2, 2, 3, 3))$  and  $\mathfrak{f} = (2, 0, 0, 1, 1)$ . Then  $C_P$  is the join of three covers with inertia types  $a_1 = (1, 2, 3)$ ,  $a_2 = (3, 4, 5)$  and  $a_3 = \text{Ind}_3^6(1, 1, 1)$ . These have signatures  $\mathfrak{f}_1 = (1, 0, 0, 0, 0)$ ,  $\mathfrak{f}_2 = (0, 0, 0, 0, 1)$ , and  $\mathfrak{f}_3 = (1, 0, 0, 1, 0)$ . By Lemma 2.3,  $A_1$  has CM-type  $(\mathcal{O}_F, \{1\})$  which has  $\beta = b_1$

and  $A_2$  has CM-type  $(\mathcal{O}_F, \{5\})$  which has  $\beta = b_2$ . Also  $\mathcal{A}_3 \simeq A^2$  and the action of  $\mathbb{Z}[\mu_6]$  on  $\mathcal{A}_3$  is given by the diagonal action of  $\mathbb{Z}[\zeta_3] \times \mathbb{Z}[\zeta_6]$  on  $A^2$ . The first copy of  $A$  has CM-type  $(\mathbb{Z}[\zeta_3], \{1\})$  which has  $\beta = b_1$  and the second copy of  $A$  has CM-type  $(\mathbb{Z}[\zeta_6], \{4\})$  which has  $\beta = b_1$ .  $\square$

We checked Corollaries 7.8 and 7.9 independently, using the fact that the special families for  $m = 6, 10$  are subspaces of those for  $m' = 3, 5$ , up to a Galois twist.

**Remark 7.10.** Let  $M = M[13]$  be the special family with  $\gamma = (6, 4, (1, 1, 2, 2))$  and  $\mathfrak{f} = (2, 1, 0, 1, 0)$ . This has an admissible degeneration to the join of two covers with inertia types  $a_1 = (1, 1, 4)$  and  $a_2 = \text{Ind}_3^6(1, 1, 1)$ . The first one has signature  $\mathfrak{f}_1 = (1, 1, 0, 0, 0)$  and its Jacobian  $A_1$  has CM by an order of finite index in  $\mathcal{O}_{F'} \times \mathcal{O}_F$ . The index is nontrivial since  $\langle x^3 - 1, x^3 + 1 \rangle = \langle 2 \rangle$  in  $\mathbb{Z}[x]/\langle x^6 - 1 \rangle$ , so our technique does not apply. The other degeneration to  $a_1 = (1, 2, 3)$  and  $a_2 = (3, 1, 2)$  does not have compact type.

## References

- [1] S. Bosch, W. Lütkebohmert, and M. Raynaud, *Néron models*, Ergebnisse der Mathematik und ihrer Grenzgebiete **21**, Springer, 1990. MR
- [2] P. E. Conner and J. Hurrelbrink, *Class number parity*, Series in Pure Mathematics **8**, World Scientific, Singapore, 1988. MR
- [3] T. Ekedahl, “Boundary behaviour of Hurwitz schemes”, pp. 173–198 in *The moduli space of curves* (Texel, Netherlands, 1994), Progr. Math. **129**, Birkhäuser, Boston, 1995. MR
- [4] N. Koblitz and D. Rohrlich, “Simple factors in the Jacobian of a Fermat curve”, *Canadian J. Math.* **30**:6 (1978), 1183–1205. MR
- [5] R. E. Kottwitz, “Points on some Shimura varieties over finite fields”, *J. Amer. Math. Soc.* **5**:2 (1992), 373–444. MR
- [6] K.-W. Lan, *Arithmetic compactifications of PEL-type Shimura varieties*, London Mathematical Society Monographs Series **36**, Princeton University Press, 2013. MR
- [7] S. Lang, *Complex multiplication*, Grundlehr. Math. Wissen. **255**, Springer, 1983. MR
- [8] W. Li, E. Mantovan, R. Pries, and Y. Tang, “Newton polygons arising from special families of cyclic covers of the projective line”, *Res. Number Theory* **5**:1 (2019), art. id. 12, 31 pp. MR
- [9] W. Li, E. Mantovan, R. Pries, and Y. Tang, “Newton polygons of cyclic covers of the projective line branched at three points”, pp. 115–132 in *Research directions in number theory: women in numbers IV*, Assoc. Women Math. Ser. **19**, Springer, 2019. Preprint version available on arXiv:1805.04598. MR
- [10] W. Li, E. Mantovan, R. Pries, and Y. Tang, “Newton polygon stratification of the Torelli locus in unitary Shimura varieties”, *Int. Math. Res. Not.* **2022**:9 (2022), 6464–6511. MR
- [11] J. S. Milne, “Shimura varieties and moduli”, pp. 467–548 in *Handbook of moduli*, Adv. Lect. Math. (ALM) **25**, International Press, Somerville, MA, 2013. MR
- [12] B. Moonen, “Special subvarieties arising from families of cyclic covers of the projective line”, *Doc. Math.* **15** (2010), 793–819. MR
- [13] J. Neukirch, *Algebraic number theory*, Grundlehr. Math. Wissen. **322**, Springer, 1999. MR

- [14] G. Shimura, “On analytic families of polarized abelian varieties and automorphic functions”, *Ann. of Math. (2)* **78** (1963), 149–192. MR
- [15] G. Shimura, “Arithmetic of unitary groups”, *Ann. of Math. (2)* **79** (1964), 369–409. MR
- [16] G. Shimura, “On purely transcendental fields automorphic functions of several variable”, *Osaka Math. J.* **1**:1 (1964), 1–14. MR
- [17] G. Shimura, “Moduli and fibre systems of abelian varieties”, *Ann. of Math. (2)* **83** (1966), 294–338. MR
- [18] P. van Wamelen, “Examples of genus two CM curves defined over the rationals”, *Math. Comp.* **68**:225 (1999), 307–320. MR
- [19] L. C. Washington, *Introduction to cyclotomic fields*, 2nd ed., Graduate Texts in Mathematics **83**, Springer, 1997. MR
- [20] S. Wewers, *Construction of Hurwitz spaces*, dissertation, Universität Duisburg-Essen, 1998.

Received September 7, 2021. Revised December 22, 2024.

WANLIN LI  
DEPARTMENT OF MATHEMATICS  
VANDERBILT UNIVERSITY  
NASHVILLE, TN  
UNITED STATES  
wanlin.li@vanderbilt.edu

ELENA MANTOVAN  
DEPARTMENT OF MATHEMATICS  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
PASADENA, CA  
UNITED STATES  
mantovan@caltech.edu

RACHEL PRIES  
DEPARTMENT OF MATHEMATICS  
COLORADO STATE UNIVERSITY  
FORT COLLINS, CO  
UNITED STATES  
pries@colostate.edu

## THE DERIVED SERIES OF GGS GROUPS

J. MORITZ PETSCHICK

**We describe the structure and indices of the members of the derived series of a GGS group  $G$  defined over the  $p$ -adic tree. The values  $|G : G^{(n)}|$  exhibit a very limited dependence on the defining vector of the group. Furthermore, we establish that the derived and Frattini series of a GGS group defined by a nonconstant vector are identical.**

### 1. Introduction and statement of results

Groups of automorphisms of regular rooted trees provide examples with intriguing asymptotic and structural properties. One particularly well-studied case is the family of Grigorchuk–Gupta–Sidki groups (usually abbreviated as GGS groups), generalising the second Grigorchuk group and the Gupta–Sidki  $p$ -groups. This family encompasses at least one group of intermediate word growth, as shown in [6], and numerous finitely generated infinite periodic groups, as demonstrated in [10]. The GGS groups acting on the  $p$ -adic tree, where  $p$  denotes an odd prime, are best understood; hence, in the following, by a GGS group we shall mean specifically a GGS group acting on the  $p$ -adic tree.

GGS groups are defined by a nonzero element  $e$  of  $\mathbb{F}_p^{p-1}$  as “input data”. It is fortunate that many properties of interest are satisfied by the group defined by  $e$  if and only if  $e$  satisfies certain linear conditions. For instance, a GGS group is periodic and just-infinite if and only if the sum of the entries of  $e$  is zero [9; 17]. Similarly, it is a branch group (with the congruence subgroup property) if and only if not all entries of  $e$  are equal [7; 8]. Its Hausdorff dimension is governed by a function depending only on certain linear invariants of  $e$  [7]. Some of these results naturally extend to larger classes of groups [1; 2; 11; 14], but have been established first for GGS groups, making the class of GGS groups a fertile soil for establishing new techniques.

In this article, we provide a description of the derived series — i.e., the subgroups defined by  $G^{(0)} = G$  and  $G^{(n+1)} = [G^{(n)}, G^{(n)}]$  for all  $n \in \mathbb{N}$  — for every GGS

---

This research was funded by the Deutsche Forschungsgemeinschaft (DFG), no. 380258175.

*MSC2020:* primary 20E08; secondary 20F14, 20E15.

*Keywords:* derived series, GGS group, branch group, self-similar group, circulant matrix, automorphism group of regular rooted trees.

group  $G$ . Conventionally, we write  $G'$  and  $G''$  for  $G^{(1)}$  and  $G^{(2)}$ , respectively. Our description is formulated in terms of linear conditions on the defining vector  $\mathbf{e}$ .

A description of the derived series was obtained by Vieira [16] for the special case of the Gupta–Sidki 3-group, along with some results on its lower central series. However, the proof in [16] does not readily generalise to general GGS groups. We approach the problem by adopting methods developed by Fernández-Alcober and Zugadi-Reizabal in [7]. We note that the sequence of indices of the members of the derived series for the (first) Grigorchuk group was computed by Grigorchuk in [9].

Our main result is the following.

**Theorem 1.1.** *Let  $p$  be an odd prime and let  $G$  be a GGS group acting on the  $p$ -adic tree defined by a nonconstant vector  $\mathbf{e} \in \mathbb{F}_p^{p-1}$ . Let  $\mathbf{e}'$  be the tuple of differences between the entries of  $\mathbf{e}$  and let  $\mathbf{e}''$  be the tuple of differences of  $\mathbf{e}'$ . Then, for  $n \geq 2$ ,*

$$\log_p |G : G^{(n)}| = p^{n-2}(p + \text{con}(\mathbf{e}') + \text{sym}(\mathbf{e}'')) - \frac{p^{n-1} - 1}{p-1} \cdot \text{sym}(\mathbf{e}) + 1.$$

Here, a vector  $\mathbf{d}$  is called symmetric if its entries are the same when read from left to right or right to left, and  $\text{sym}(\mathbf{d})$  equals 1 if the vector  $\mathbf{d}$  is symmetric and 0 otherwise. Similarly,  $\text{con}(\mathbf{d})$  equals 1 if the vector  $\mathbf{d}$  is constant, i.e., if all its entries are equal, and 0 otherwise. In [8, Proposition 3.4], Fernández-Alcober, Garrido and Uria-Albizuri show that GGS groups with constant defining vector admit an infinite metabelian quotient, i.e., that  $|G : G''| = \infty$ . It is an elementary fact that  $|G : G'| = p^2$ , so Theorem 1.1 completes the description of all indices  $|G : G^{(n)}|$  for all GGS groups  $G$  and all integers  $n \in \mathbb{N}$ .

To arrive at Theorem 1.1, we need a description of the derived subgroups:

**Theorem 1.2.** *Let  $p$  be an odd prime and let  $G$  be a GGS group acting on the  $p$ -adic tree defined by a nonconstant vector  $\mathbf{e} \in \mathbb{F}_p^{p-1}$ . Let  $n \geq 3$ . Then*

$$\psi(G^{(n)}) = \chi_p G^{(n-1)}.$$

*If  $\text{con}(\mathbf{e}') + \text{sym}(\mathbf{e}'') = 0$ , the same holds for  $n = 2$ .*

Together with a description of the second derived subgroup, which we present in Proposition 3.2, this theorem yields a good account of the structure of the derived subgroups of  $G$ . One notable consequence is that the quotients  $G^{(n)}/G^{(n+1)}$  are elementary abelian  $p$ -groups, which immediately yields the following corollary.

**Corollary 1.3.** *Let  $G$  be a GGS group defined by a nonconstant vector. The Frattini series of  $G$  coincides with the derived series of  $G$ .*

Finally, we investigate the GGS groups defined by nonconstant vectors with the maximal possible indices  $|G : G^{(n)}|$ . We find that there are precisely two isomorphism classes of GGS groups with  $|G : G^{(n)}|$  maximal among GGS groups defined by nonconstant vectors (acting on a fixed  $p$ -adic tree); see Proposition 3.5.

## 2. On Grigorchuk–Gupta–Sidki-groups

**Notation.** The letter  $p$  refers to a fixed odd prime. Given integers  $m$  and  $n$  subject to  $m \leq n$ , the set  $\{m, m + 1, \dots, n - 1, n\}$  is denoted by  $[m, n]$ . The symbol  $X$  refers to the set  $[0, p - 1]$  underlying the field  $\mathbb{F}_p$ . For a group  $G$ , the  $p$ -fold direct product  $G \times \dots \times G$  is denoted by  $\chi_p G$ . For  $k \in \mathbb{N}$  distinct elements  $i_0, \dots, i_{k-1} \in X$  and  $p$  elements  $g_0, \dots, g_{p-1}$  of a group  $G$ , the expression

$$(i_0 : g_0, \dots, i_{k-1} : g_{k-1}, \diamond : g_\diamond) \in \chi_p G$$

denotes the tuple indexed by  $X$ , with  $g_j$  at position  $i_j$  for  $j \in [0, k - 1]$ , and with  $g_\diamond$  (possibly varying with  $\diamond$ ) at every other position  $\diamond \in X \setminus \{i_j \mid j \in [0, k - 1]\}$ . The symbol  $\diamond$  is reserved for this use.

Given a group  $G$  and two of its elements  $g$  and  $h$ , we use the following conventions for conjugation and the commutator:

$$g^h = h^{-1}gh \quad \text{and} \quad [g, h] = g^{-1}h^{-1}gh = g^{-1}g^h.$$

**Automorphisms of rooted trees.** The Cayley graph  $X^*$  of the free monoid on  $X$  is the rooted  $p$ -adic tree, i.e., a loop-free graph with a distinguished vertex (the “root”)  $\emptyset$  of valency  $p$  and all other vertices of valency  $p + 1$ . We write  $X^n$  for the set of all vertices of a given (geodesic) distance  $n$  to  $\emptyset$ . This set is called the  $n$ -th level of  $X^*$ .

Any (graph) automorphism  $g \in \text{Aut}(X^*)$  necessarily fixes  $\emptyset$  for its unique valency, and must consequently leave all levels  $X^n$  invariant. We write  $\text{St}(n)$  for the (pointwise) stabiliser of  $X^n$ , and  $\text{St}_G(n)$  for its intersection with a subgroup  $G \leq \text{Aut}(X^*)$ .

Let  $u$  and  $v$  be vertices of  $X^*$ . We write  $u^g$  for the image of  $u$  under  $g$ . Since every level is invariant under  $g$ , the equation

$$(uv)^g = u^g v^{g|_u}$$

uniquely defines a map  $|_u : \text{Aut}(X^*) \rightarrow \text{Aut}(X^*)$ , called the *section map at  $u$* . The image of  $g$  is called *the section of  $g$  at  $u$* . For  $u \in X^*$ , the restriction of  $|_u$  to the stabiliser  $\text{st}(u)$  of  $u$  is a group homomorphism, indeed, the map

$$\psi : \text{St}(1) \rightarrow \chi_p \text{Aut}(X^*), \quad g \mapsto (\diamond : g|_\diamond)$$

is an isomorphism. We record some equations for sections. Let  $u$  and  $v$  be vertices of  $X^*$  and  $g$  and  $h$  be automorphisms, then

$$(g|_u)|_v = g|_{uv}, \quad (gh)|_u = g|_u h|_{u^g}, \quad g^{-1}|_u = (g|_{u^{g^{-1}}})^{-1}.$$

The action of an automorphism  $g$  on the first level  $X = X^1$  is denoted  $g|^\emptyset \in \text{Sym}(X)$ . An element such that  $g|_u = \text{id}$  for all  $u \neq \emptyset$  is called *rooted*. We identify rooted

elements with the their images under  $|\varnothing$ . Let  $u \in X^n$  and  $g \in \text{Aut}(X^*)$ . The unique element  $h \in \text{St}(n)$  satisfying  $h|_u = g$  and  $h|_v = \text{id}$  for all  $v \in X^n \setminus \{u\}$  is denoted  $\text{ins}_u(g)$  and called the *insertion of  $g$  at  $u$* .

A group  $G \leq \text{Aut}(X^*)$  is called *self-similar*, if all sections  $g|_u$  are contained in  $G$  for all  $u \in X^*$  and  $g \in G$ . A group  $G \leq \text{Aut}(X^*)$  is called *fractal* if  $\text{st}_G(x)|_x = G$  for every  $x \in X$ . A group  $G \leq \text{Aut}(X^*)$  is called *spherically transitive* if it acts transitively on every level  $X^n$ . A self-similar group  $G \leq \text{Aut}(X^*)$  is called a *regular branch group* if it is spherically transitive and if there is a finite-index subgroup  $K \leq \text{St}_G(1)$  such that  $\chi_p K \leq \psi(K)$ . A standard technique to establish that a group is regular branch is the following:

**Proposition 2.1** [7, Proposition 2.18]. *Let  $G \leq \text{Aut}(X^*)$  be a spherically transitive fractal group, let  $H \leq G$  be a subgroup and let  $S \subseteq G$  be a subset. If  $\text{ins}_0(S) \subseteq H$ , then  $\chi_p \langle S \rangle^G \leq \psi(H^G)$ .*

**GGs groups and their defining vectors.** Let  $e = (e_1, \dots, e_{p-1}) \in \mathbb{F}_p^{p-1}$  be a nonzero vector. The group  $G$  generated by the rooted automorphism  $a = (0 \ 1 \ \dots \ p-1)$  induced by the addition of 1 in  $\mathbb{F}_p$ , and the automorphism defined by

$$b = \psi^{-1}(0 : b, \diamond : a^{e \diamond})$$

is called the *GGs group defined by  $e$* . The vector  $e$  is the *defining vector of  $G$* .

More generally, GGS groups acting on the (not necessarily prime)  $m$ -adic tree are defined in the same way, but using elements of  $(\mathbb{Z}/m\mathbb{Z})^{m-1}$  whose entries have no common divisor other than 1. In general, the structure of these groups is much less understood than the case considered here. Even for prime powers  $m = p^n$ , the situation is much more involved, see for example [5], where the branching structures for these groups have been evaluated. Various further generalisations of GGS groups have been studied, e.g. in [1; 3; 12; 14].

Distinct defining vectors may give rise to identical or isomorphic GGS groups. Nonzero multiples of a given vector  $e$  define the same subgroup of  $\text{Aut}(X^*)$ . Apart from that, certain reorderings of a given defining vector give rise to isomorphic (but not necessarily identical) GGS groups. We make use of the following characterisation. The group  $\mathbb{F}_p^\times \times \mathbb{F}_p^\times \cong \mathbb{C}_{p-1}^2$  acts on the set  $\mathbb{F}_p^{p-1} \setminus \{\mathbf{0}\}$  of defining vectors by

$$(2-1) \quad (e_1, \dots, e_{p-1}) * (\lambda, \mu) = (\lambda \cdot e_\mu, \lambda \cdot e_{2,\mu}, \dots, \lambda \cdot e_{(p-2),\mu}, \lambda \cdot e_{(p-1),\mu}).$$

**Theorem 2.2** [13, Corollary 4.5]. *Two GGS groups  $G$  and  $\tilde{G}$  defined by  $e$  and  $\tilde{e}$ , respectively, are isomorphic if and only if  $e$  and  $\tilde{e}$  share the same orbit under the action of  $\mathbb{F}_p^\times \times \mathbb{F}_p^\times$ .*

This allows us to choose defining vectors with desirable properties, as also done in [7, Theorem 2.16].

**Corollary 2.3.** *Let  $G$  be the GGS group defined by  $e$  and let  $\alpha \in \mathbb{F}_p^\times$ .*

- (i) *There exists a defining vector  $\tilde{e}$  with  $\tilde{e}_1 = \alpha$  such that the GGS group defined by  $\tilde{e}$  is isomorphic to  $G$ .*
- (ii) *If  $e$  is not constant, there exists a defining vector  $\tilde{e}$  with  $e_i - e_{i+1} = \alpha$ , for some  $i \in \{1, \dots, p-2\}$ , such that the GGS group defined by  $\tilde{e}$  is isomorphic to  $G$ .*

**Difference vectors.** Let  $\mathbf{d} = (d_1, \dots, d_n) \in \mathbb{F}_p^n$  be a vector. The *difference vector* of  $\mathbf{d}$  is the vector

$$\mathbf{d}' = (d'_1, d'_2, \dots, d'_{n-1}) = (d_1 - d_2, d_2 - d_3, \dots, d_{n-1} - d_n) \in \mathbb{F}_p^{n-1}.$$

A vector  $\mathbf{d}$  is called *constant* if  $\mathbf{d}' = \mathbf{0}$ , i.e., if all entries of  $\mathbf{d}$  are equal. We put  $\text{con}(\mathbf{d}) = 1$  if  $\mathbf{d}$  is constant, and  $\text{con}(\mathbf{d}) = 0$  otherwise.

The structure of the derived subgroups of the GGS group defined by  $e$  is influenced by the difference vector  $e'$  and by its difference vector  $e''$ , called, for convenience, the *second difference vector* of  $e$ . The significance of  $e'$  is suggested by the following computation. Every GGS group  $G$  is two-generated and its derived subgroup is normally generated by the commutator  $c = [a^{-1}, b]$ , whose section decomposition is closely related to  $e'$ ,

$$\begin{aligned} (2-2) \quad \psi(c) &= \psi([a^{-1}, b]) = \psi(b^{a^{-1}})^{-1} \psi(b) \\ &= (\diamond : a^{-e_{\diamond+1}}, p-1 : b^{-1})(0 : b, \diamond : a^{e_{\diamond}}) \\ &= (0 : a^{-e_1} b, \diamond : a^{e'_{\diamond}}, p-1 : b^{-1} a^{e_{p-1}}). \end{aligned}$$

The entries of  $e''$  appear in a similar way in the section decomposition of  $[a^{-1}, c]$ .

Perhaps surprisingly, the higher difference vectors of  $e$  do not affect the structure of the derived series.

**Symmetric vectors.** Let  $n \in \mathbb{N}$  be even, let  $K$  be a field not of characteristic 2, and let  $\mathbf{d} = (d_1, \dots, d_n) \in K^n$  be a vector. It is called *symmetric* if

$$d_i = d_{n-i+1}$$

for all  $i \in [1, n/2]$ . We put  $\text{sym}(\mathbf{d}) = 1$  if  $\mathbf{d}$  is symmetric, and  $\text{sym}(\mathbf{d}) = 0$  otherwise. Evidently, the set of all symmetric vectors constitutes a subspace  $S$  of  $K^n$ , being subject to the conditions  $d_i - d_{n-i+1} = 0$  for  $i \in [1, n/2]$ . If  $\mathbf{d}$  is symmetric, the second difference vector  $\mathbf{d}''$  is also symmetric; see Table 1 for an overview of the possible configurations of the values  $\text{con}(e)$ ,  $\text{sym}(e)$ ,  $\text{con}(e')$  and  $\text{sym}(e'')$  for a defining vector. More precisely, the second difference vector is symmetric if and only if

$$d_i - 2d_{i+1} + d_{i+2} = d''_i = d''_{n-i-1} = d_{n-1-i} - 2d_{n-i} + d_{n-i+1},$$

i.e., if and only if  $d''_i - d''_{n-i-1} = 0$  for all  $i \in [1, n/2 - 1]$ . It is apparent that

con( $\mathbf{e}$ )	sym( $\mathbf{e}$ )	con( $\mathbf{e}'$ )	sym( $\mathbf{e}''$ )
0	0	0	0
0	0	0	1
0	0	1	1
0	1	0	1
1	1	1	1

**Table 1.** Possible configurations of values for the invariants of  $\mathbf{e}$  influencing the indices of the members of the derived series of the corresponding GGS group. For  $p = 3$ , the first, second and fourth rows are nonexistent.

the set of all vectors such that  $\mathbf{d}''$  is symmetric is a subspace containing the space of symmetric vectors as a subspace of codimension 1. We need the following computational lemma.

**Lemma 2.4.** *Let  $\mathbf{e} \in \mathbb{F}_p^{p-1}$  be such that  $\text{sym}(\mathbf{e}'') = 1$ . Then*

$$2(e_{p-1} - e_1) + (e_2 - e_{p-2}) = 0.$$

*Proof.* Write  $s_i = e_i - e_{p-i}$  for  $i \in [1, p-1]$ . Note that

$$s'_i = s_i - s_{i+1} = e_i - e_{p-i} - e_{i+1} + e_{p-i-1} = e'_i - e'_{p-i-1},$$

and in the same way,  $s''_i = e''_i - e''_{p-i-2}$ . Moreover,  $s_i = -s_{p-i}$ , whence  $s'_{(p-1)/2} = s_{(p-1)/2} - s_{(p+1)/2} = 2 \cdot s_{(p-1)/2}$ . Two simple telescope sum computations yield

$$\sum_{i=1}^{(p-3)/2} s''_i = \sum_{i=1}^{(p-3)/2} (s'_i - s'_{i+1}) = s'_1 - s'_{(p-1)/2} = s_1 - s_2 - 2 \cdot s_{(p-1)/2}$$

and

$$\begin{aligned} \sum_{i=1}^{(p-3)/2} i \cdot s''_i &= \sum_{i=1}^{(p-3)/2} i \cdot (s'_i - s'_{i+1}) = \sum_{i=1}^{(p-3)/2} s'_i - ((p-3)/2) \cdot s'_{(p-1)/2} \\ &= s_1 - s_{(p-1)/2} + 3 \cdot s_{(p-1)/2} \\ &= s_1 + 2 \cdot s_{(p-1)/2}. \end{aligned}$$

Combining them, one finds that

$$- \sum_{i=1}^{(p-3)/2} (i+1) \cdot s''_i = -2s_1 + s_2.$$

Since  $\mathbf{e}''$  is symmetric,  $s''_i = 0$  for all  $i \in [1, p-3]$ . Thus the left-hand side of the equality above is zero, while the right-hand side evaluates to the desired expression  $2(e_{p-1} - e_1) + (e_2 - e_{p-2})$ . □

**Circulant spaces.** Let  $K$  be a field and let  $\mathbf{d} = (d_0, \dots, d_{\ell-1}) \in K^\ell$  be a vector. The *circulant matrix*  $\text{Circ}(\mathbf{d})$  associated to  $\mathbf{d}$  is the matrix whose rows are the cyclic shifts of  $\mathbf{d}$ , i.e.,

$$\begin{pmatrix} d_0 & d_1 & \dots & d_{\ell-2} & d_{\ell-1} \\ d_{\ell-1} & d_0 & \dots & d_{\ell-3} & d_{\ell-2} \\ \vdots & \vdots & & \vdots & \vdots \\ d_1 & d_2 & \dots & d_{\ell-1} & d_0 \end{pmatrix}.$$

The *semicirculant matrix*  $\text{SCirc}(\mathbf{d})$  associated to  $\mathbf{d}$  is the upper triangular matrix given by

$$\begin{pmatrix} d_0 & d_1 & \dots & d_{\ell-2} & d_{\ell-1} \\ 0 & d_0 & \dots & d_{\ell-3} & d_{\ell-2} \\ \vdots & \vdots & & \vdots & \vdots \\ 0 & 0 & \dots & 0 & d_0 \end{pmatrix}.$$

More generally, a *circulant subspace of  $K^\ell$*  is a subspace that is invariant under the automorphism induced by the cyclic permutation of the standard basis elements of  $K^\ell$ . Given a subset  $M \subseteq K^\ell$ , we write  $\text{Circ}(M)$  for the minimal circulant subspace containing  $M$ . In case of a singleton set,  $\text{Circ}(\{\mathbf{d}\})$  is spanned by the rows of the circulant matrix  $\text{Circ}(\mathbf{d})$ .

Ranks of circulant matrices, their computation and interpretation have long been studied; see for example [4] for the situation over the field of complex numbers. In positive characteristic  $p$ , the special case  $\ell = p^n$  allows an easy description of the circulant subspaces of  $K^\ell$ . Let  $\Pi \in K^{\ell \times \ell}$  be the Pascal matrix with entries  $\binom{i}{j}$  for  $i, j \in [0, \ell-1]$ , using the convention that  $\binom{i}{j} = 0$  for  $i < j$ , and write  $\Pi_i$  for the  $\ell$ -by- $i$  matrix consisting of the first  $i$  columns of  $\Pi$ . Note that  $\Pi$  is a lower unitriangular matrix; in particular, it is invertible.

**Proposition 2.5.** *Let  $K$  be a field of characteristic  $p > 0$  and let  $\ell = p^n$  be a power of the characteristic. There exists a complete flag*

$$\{\mathbf{0}\} = \text{Circ}_0 \subset \text{Circ}_1 \subset \dots \subset \text{Circ}_\ell = K^\ell$$

*containing all circulant subspaces of  $K^\ell$ , which are given by*

$$\text{Circ}_i = \ker \Pi_{\ell-i} = \{\mathbf{d} \in K^\ell \mid \text{rank } \text{Circ}(\mathbf{d}) \leq i\}.$$

*Proof.* Fix a vector  $\mathbf{d} = (d_0, \dots, d_{\ell-1}) \in K^\ell$ . Put

$$f_{\mathbf{d}}(X) = d_0 + d_1X + d_2X^2 + \dots + d_{\ell-1}X^{\ell-1}$$

and

$$P = \begin{pmatrix} 0_{\ell-1,1} & \text{I}_{\ell-1} \\ 1 & 0_{1,\ell-1} \end{pmatrix},$$

where  $0_{n,m}$  and  $I_n$  stand for the zero and unit matrices, respectively, of the indicated formats. The matrix  $P$  is the permutation matrix associated to the cyclic shift of the basis elements, and

$$\text{Circ}(\mathbf{d}) = d_0I + d_1P + d_2P^2 + \dots + d_{\ell-1}P^{\ell-1} = f_{\mathbf{d}}(P).$$

The characteristic polynomial of  $P$  is  $X^\ell - 1 = (X - 1)^\ell$  and splits over  $K$ ; here we use that  $\ell$  is a power of  $p$ . The geometric multiplicity of its unique eigenvalue 1 is 1, whence  $P$  is conjugate to the matrix  $J_\ell(1)$  consisting of a single Jordan block of eigenvalue 1. Consequently,  $\text{Circ}(\mathbf{d})$  is conjugate to  $f_{\mathbf{d}}(J_\ell(1))$ . For  $k \in \mathbb{N}$ , we have  $J_\ell(1)^k = \left( \binom{k}{j-i} \right)_{i,j \in [1,\ell]}$ , i.e.,  $J_\ell(1)^k = \text{SCirc} \left( \left( \binom{k}{0}, \binom{k}{1}, \dots, \binom{k}{\ell-1} \right) \right)$ . Thus  $f_{\mathbf{d}}(J_\ell(1))$  is the semicirculant matrix with respect to the vector  $\mathbf{b} = (b_0, \dots, b_{\ell-1})$ , with

$$b_i = \sum_{j=0}^{\ell-1} d_j \binom{j}{i}.$$

Then  $\mathbf{b} = \mathbf{d}\Pi$ . Since the rank of  $\text{SCirc}(\mathbf{b})$  is equal to  $\ell - \min\{j \in [0, \ell-1] \mid b_j \neq 0\}$ , we see that  $\text{rank Circ}(\mathbf{d}) \leq i$  if and only if  $\mathbf{d} \in \ker \Pi_{\ell-i}$ , with equality precisely if  $\mathbf{d} \notin \ker \Pi_{\ell-i+1}$ . Since  $\Pi$  is invertible, we see that  $\text{rank } \Pi_{\ell-i} = i$ . Thus  $\text{Circ}_i$  is indeed a circulant space, as  $\text{Circ}_i = \text{Circ}(\mathbf{d})$  for any  $\mathbf{d} \in \text{Circ}_i \setminus \text{Circ}_{i-1}$ . Evidently  $\text{Circ}_i \subset \text{Circ}_{i+1}$ . It remains to show that there are no further circulant spaces. Let  $C$  be a circulant subspace of dimension  $i$ . For every  $\mathbf{d} \in C$ , we necessarily find  $\text{rank Circ}(\mathbf{d}) \leq i$ . If there exists  $\mathbf{d}$  such that the rank of  $\text{Circ}(\mathbf{d})$  is  $i$ , naturally  $C = \text{Circ}(\mathbf{d}) = \text{Circ}_i$ . Thus assume that  $\text{Circ}(\mathbf{d}) < i$  for all  $\mathbf{d} \in C$ . But then  $C \subseteq \text{Circ}_{i-1}$ , an  $(i-1)$ -dimensional subspace, which is absurd.  $\square$

This description extends [7, Lemma 2.7]. Note that the crucial point is that the polynomial  $X^\ell - 1$  splits over  $K$ . Let  $\mathbf{e}$  be the defining vector of the GGS group  $G$  and put  $\bar{\mathbf{e}} = (0, e_1, \dots, e_{p-1})$ . The structure of  $G$  is heavily influenced by the *cyclic rank*  $\text{cr}(\mathbf{e})$  of  $\mathbf{e}$ , which is given by  $\text{cr}(\mathbf{e}) = \dim \text{Circ}(\bar{\mathbf{e}})$ . By Proposition 2.5,  $\text{cr}(\mathbf{e}) = p$  if and only if the sum of entries of  $\mathbf{e}$  is nonzero, whence by [9, Example 9.1], the group  $G$  is periodic if and only if  $\text{cr}(\mathbf{e}) < p$ . The Hausdorff dimension of  $G$  solely depends on whether  $\mathbf{e}$  is symmetric or constant and on the value of  $\text{cr}(\mathbf{e})$ , as demonstrated by Fernández-Alcober and Zugadi-Reizabal in [7, Theorem B].

**Corollary 2.6.** *Let  $K$  be a field of characteristic  $p > 0$ , let  $i \in [0, p]$  and let  $\text{Circ}_i$  be the members of the flag of circulant subspaces of  $K^p$ . A basis for  $\text{Circ}_i$  is given by the first  $i$  columns of the Pascal matrix  $\Pi \in K^{p \times p}$ .*

*Proof.* In view of Proposition 2.5 and the fact that the columns of  $\Pi$  are linearly independent, we have to show that

$$\sum_{n=j}^{p-1} \binom{n}{m} \binom{n}{j} \equiv_p 0$$

for every  $m \in [0, i-1]$  and  $j \in [0, p-i-1]$ . Put  $\phi(k, m, j) = \sum_{n=j}^{p-1} n^k \binom{n}{m} \binom{n}{j}$  and compute

$$\phi(k, m, j) = \sum_{n=j}^{p-1} n^k \binom{n}{m} \binom{n}{j-1} \frac{n-j+1}{j} = \frac{1}{j} \phi(k+1, m, j-1) + \frac{1-j}{j} \phi(k, m, j-1);$$

analogously,

$$\phi(k, m, j) = \frac{1}{m} \phi(k+1, m-1, j) + \frac{1-m}{m} \phi(k, m-1, j).$$

By iteration we find that  $\phi(0, m, j)$  is a linear combination of the values of  $\phi(k, 0, 0)$  for  $k \in [0, m+j]$ . But

$$\phi(k, 0, 0) = \sum_{n=0}^{p-1} n^k = \sum_{n=0}^{p-1} n \equiv_p 0$$

for every  $k$  that is not a multiple of  $p-1$ . Since  $m+j < p-1$ , we obtain the desired congruence. □

**Properties and structure of GGS groups.** Recall that, for a given GGS group  $G$ , the rooted generator is denoted  $a$ , the directed generator is denoted  $b$ , and we write  $c$  for the commutator  $[a^{-1}, b]$ , whose sections are given in (2-2). We shall use the following shorthand notation for the conjugates of  $c$ :

$$c_i = c^{a^i} = [a^{-1}, b^{a^i}].$$

In particular,  $c_0 = c$ . We collect some facts about GGS groups.

**Lemma 2.7.** *Every GGS group is fractal and self-similar.*

For a proof, see e.g. [8, Section 2] or [13, Section 2.3]. Next, we record some information on certain small quotients of GGS groups.

**Lemma 2.8.** *Let  $G$  be a GGS group. Then*

- (i)  $\log_p |G : G'| = 2$  and  $G/G'$  is elementary abelian,
- (ii)  $\log_p |G' : \gamma_3(G)| = 1$  and  $G/\gamma_3(G)$  is of exponent  $p$ , and
- (iii)  $\log_p |G' : \text{St}_G(1)'| = p-1$ .

For statements (i) and (ii), see Theorem 2.1(iii) of [7] and note that  $a^p = b^p = \text{id}$  and  $c^p \equiv_{\gamma_3(G)} \text{id}$ . Statement (iii) is proven as part of Theorem 2.14 of [7].

It is a result of Fernández-Alcober and Zugadi-Reizabal that every GGS group defined by a nonconstant vector is a regular branch group (see below). The same is not true for GGS groups defined by constant tuples, explaining their divergent behaviour.

**Theorem 2.9.** *Let  $G$  be a GGS group with nonconstant defining vector  $\mathbf{e}$ . Then:*

- (i)  $\psi(\gamma_3(\text{St}_G(1))) = \chi_p \gamma_3(G)$ .
- (ii)  $\psi(\text{St}_G(1)') \leq \chi_p G'$  and  $\log_p |\chi_p G' : \psi(\text{St}_G(1)')| = \text{sym}(\mathbf{e})$ .

*In particular, the group  $G$  is regular branch over  $\gamma_3(G)$ , and it is regular branch over  $G'$  if  $\mathbf{e}$  is not symmetric.*

These statements appear as Lemmas 3.2 and 3.4 of [7].

We need another fact concerning the subgroups of GGS groups.

**Proposition 2.10.** *Let  $G$  be a GGS group with a nonconstant defining vector. Then*

$$[\text{St}_G(1)', G'] = \gamma_3(\text{St}_G(1)).$$

*In particular,  $\gamma_3(\text{St}_G(1)) \leq G''$ .*

*Proof.* The inclusion  $[\text{St}_G(1)', G'] \leq \gamma_3(\text{St}_G(1))$  is a straightforward consequence of  $G' \leq \text{St}_G(1)$ , which itself follows from  $G/\text{St}_G(1)$  being cyclic. We have to establish the other inclusion. In view of Proposition 2.1, it is enough to prove that  $\text{ins}_0([c, b])$  and  $\text{ins}_0([c, a])$  are contained in  $[\text{St}_G(1)', G']$ , as  $\gamma_3(G)$  is normally generated by  $[c, b]$  and  $[c, a]$ . We distinguish two cases.

Case  $\text{sym}(\mathbf{e}) = 0$ : By Theorem 2.9(ii), the element  $\text{ins}_0(c)$  is contained in  $\text{St}_G(1)'$ . By Corollary 2.3(ii) we may assume  $e'_i = 1$  for some  $i \in \{1, \dots, p-2\}$ . By (2-2)  $c|_i = a$ , hence

$$c_{p-i}|_0 = c|_i = a, \quad (c_{p-i}^{e_1} c)|_0 = a^{e_1} a^{-e_1} b = b.$$

Since  $c_{p-i}$  and  $c$  are elements of  $G'$ , we obtain

$$[\text{ins}_0(c), c_{p-i}] = \text{ins}_0([c, a]) \quad \text{and} \quad [\text{ins}_0(c), c_{p-i}^{e_1} c] = \text{ins}_0([c, b])$$

are contained in  $[\text{St}_G(1)', G']$ .

Case  $\text{sym}(\mathbf{e}) = 1$ : Note that, since  $\mathbf{e}$  is by assumption not constant, the prime  $p$  is necessarily greater than 3. By Corollary 2.3(i), we may assume  $e_1 = e_{p-1} = -1$ . Observe that

$$\begin{aligned} \psi([b, b^a]) &= [\psi(b), \psi(b^a)] = (0 : [b, a^{-1}], 1 : [a^{-1}, b], \diamond : \text{id}) \\ &= (0 : c^{-1}, 1 : c, \diamond : \text{id}). \end{aligned}$$

We first show that there exists  $j \in \mathbb{F}_p^\times \setminus \{1, p-1\}$  such that  $e_{j-1} \neq e_{p-j-1}$ . Assume the converse for contradiction. Using that  $\mathbf{e}$  is symmetric, we find

$$e_j = e_{(1+j)-1} = e_{p-(1+j)-1} = e_{j+2}.$$

Thus  $e_2 = e_4 = \dots = e_{p-1}$  and  $e_3 = e_5 = \dots = e_{p-2}$ , using that  $p > 3$ . Since  $\mathbf{e}$  is symmetric,  $e_2 = e_{p-2}$  and  $e_1 = e_{p-1}$ , whence  $\mathbf{e}$  is constant, which is excluded.

Now let  $j$  be an element as described above. Compute

$$\psi([a^{2j}, b]) = (0 : a^{-e_{p-2j}}b, 2j : b^{-1}a^{e_{2j}}, \diamond : a^{e_{\diamond} - e_{\diamond-2j}}),$$

so  $[a^{2j}, b]|_j = a^{e_j - e_{p-j}} = \text{id}$ , since  $\mathbf{e}$  is symmetric. Put  $g = [a^{2j}, b]^{a^{1-j}}$  to find

$$g|_1 = [a^{2j}, b]|_j = \text{id} \quad \text{and} \quad g|_0 = [a^{2j}, b]|_{j-1} = a^{e_{j-1} - e_{p-j-1}},$$

since  $j \notin \{1, p-1\}$  forbids  $j-1 \in \{0, 2j\}$ . Let  $i = (e_{j-1} - e_{p-j-1})^{-1}$ , using that  $e_{j-1} \neq e_{p-j-1}$ . Observe that

$$\begin{aligned} [\text{St}_G(1)', G'] \ni [[b^a, b], g^i] &= [\psi^{-1}(0 : c, 1 : c^{-1}, \diamond : \text{id}), \psi^{-1}(0 : a, 1 : \text{id}, \diamond : g^i|_{\diamond})] \\ &= \text{ins}_0([c, a]). \end{aligned}$$

It remains to show  $\text{ins}_0([c, b]) \in [\text{St}_G(1)', G']$ . Consider  $h = g^{ie_{p-2}}[a^2, b]$ , which fulfils

$$h|_0 = a^{e_{p-2}}a^{-e_{p-2}}b = b \quad \text{and} \quad h|_1 = a^{e_1 - e_{p-1}} = \text{id}.$$

Then

$$[\text{St}_G(1)', G'] \ni [[b^a, b], h] = \text{ins}_0([c, b]). \quad \square$$

Using Proposition 2.10, we derive the following adjunct to Theorem 2.9.

**Proposition 2.11.** *Let  $G$  be a GGS group with nonconstant defining vector. Then  $G$  is branch over  $G''$ .*

*Proof.* Using Theorem 2.9(i) and Proposition 2.10 we find

$$\chi_p G'' \leq \chi_p \gamma_3(G) = \psi(\gamma_3(\text{St}_G(1))) \leq \psi(G''). \quad \square$$

### 3. The derived series of GGS groups

**The second derived subgroup.** By Lemma 2.8(ii), the quotient  $G'/\gamma_3(G)$  is a cyclic group of order  $p$ , generated by the element  $\bar{c} = c \cdot \gamma_3(G)$ . Thus the group  $V := \chi_p(G'/\gamma_3(G))$  is an elementary abelian  $p$ -group of rank  $p$ , which we write additively and treat as an  $\mathbb{F}_p$ -vector space with the natural basis


$$\{(\bar{c}, 0, \dots, 0), \dots, (0, \dots, 0, \bar{c})\}.$$

Write  $\phi: \text{St}_G(1)' \rightarrow V$  for the concatenation of  $\psi$  and the natural epimorphism  $\chi_p G' \rightarrow V$ .

**Lemma 3.1.** *Let  $G$  be a GGS group with nonconstant defining vector and let  $N \leq G$  be a subgroup satisfying*

$$\chi_p \gamma_3(G) \leq \psi(N) \leq \chi_p G'.$$

*Then  $N$  is normal in  $G$  if and only if  $\phi(N)$  is a circulant subspace of  $V$ , and as a consequence, there exist precisely  $p+1 - \text{sym}(\mathbf{e})$  such normal subgroups.*



**Figure 1.** Part of the top of the subgroup lattice of a GGS group, with some supergroups added. Passage from the left to the right side signifies the application of  $\psi$ . All indices are logarithmic. See [7] for the computation of the index of  $St_G(2)$ .

*Proof.* The conjugation action of  $a$  corresponds to a cyclic shift on  $\psi(N)$ , whence  $N$  is invariant under conjugation by  $a$  if and only if  $\phi(N)$  is circulant. It remains to notice that  $N$  is automatically invariant under conjugation by  $b$ : for every  $g \in \psi^{-1}(\chi_p G') \cap G$  and every  $h \in St_G(1)$

$$\psi(g^h) = (\diamond : g|_{\diamond}^{h|\diamond}) \equiv_{\chi_p, \gamma_3(G)} (\diamond : g|_{\diamond}) = g.$$

The last statement is a direct consequence of Proposition 2.5 and Theorem 2.9(ii).  $\square$

**Proposition 3.2.** *Let  $G$  be a GGS group with nonconstant defining vector. Then  $G'' = \phi^{-1}(\text{Circ}_{p-\text{con}(e')-\text{sym}(e'')}(V))$ . In particular,*

$$\log_p |\chi_p G' : \psi(G'')| = \text{con}(e') + \text{sym}(e'').$$

*Proof.* Since  $G' \leq St_G(1)$ , the second derived subgroup  $G''$  is contained in  $St_G(1)'$ , which, by Theorem 2.9(ii), is in turn contained in  $\psi^{-1}(\chi_p G')$ . On the other hand,

using Proposition 2.10 and Theorem 2.9(i), we find

$$\chi_p \gamma_3(G) = \psi(\gamma_3(\text{St}_G(1))) \leq \psi(G'').$$

Thus Lemma 3.1 applies and it remains to compute the dimension of  $\phi(G'')$ . To achieve this, we pick a subset  $S$  of  $G''$  normally generating  $G''$  and use Proposition 2.5 to compute the dimension of the circulant space generated by  $\phi(S)$ . Since the group  $G$  is 2-generated, a natural choice for  $S$  is

$$\{[c^g, c] \mid g \in G\}.$$

The kernel of  $\phi$  is  $\gamma_3(\text{St}_G(1)) = \psi^{-1}(\chi_p \gamma_3(G))$ , whence  $\phi([c^g, c]) = \phi([c_i, c])$  for  $a^i \equiv_{\text{St}_G(1)} g$ . Notice that

$$[c_i, c]^{-1} = [c, c_i] = [c, c^{a^i}] = [c_{p-i}, c]^{a^i}.$$

Thus the set  $\phi(\{[c_i, c] \mid i \in [1, (p-1)/2]\})$  generates  $\phi(G'')$  as a circulant space. The sections of the elements  $[c_i, c]$  are given by (compare (2-2))

$$\begin{aligned} \psi([c_1, c]) &= (0 : [b^{-1}a^{e_{p-1}}, a^{-e_1}b], 1 : [a^{-e_1}b, a^{e'_1}], \diamond : \text{id}, p-1 : [a^{e'_{p-2}}, b^{-1}a^{e_{p-1}}]) \\ &\equiv (0 : c^{e_1 - e_{p-1}}, 1 : c^{e'_1}, \diamond : \text{id}, p-1 : c^{e'_{p-2}}) \pmod{\chi_p \gamma_3(G)} \end{aligned}$$

and

$$\begin{aligned} \psi([c_i, c]) &= \begin{pmatrix} 0 : [a^{e'_{p-i}}, a^{-e_1}b], & p-1 : [a^{e'_{p-i-1}}, b^{-1}a^{e_{p-1}}] \\ i-1 : [b^{-1}a^{e_{p-1}}, a^{e'_{i-1}}], & i : [a^{-e_1}b, a^{e'_i}] \\ \diamond : \text{id} \end{pmatrix} \\ &\equiv \begin{pmatrix} 0 : c^{-e'_{p-i}}, & p-1 : c^{e'_{p-i-1}}, \\ i-1 : c^{-e'_{i-1}}, & i : c^{e'_i}, \\ \diamond : \text{id} \end{pmatrix} \pmod{\chi_p \gamma_3(G)} \end{aligned}$$

for  $i \in [2, (p-1)/2]$ . Thus the images  $\mathbf{d}_i = \phi([c_i, c])$  in  $V$  are given by

$$\mathbf{d}_1 = (e_1 - e_{p-1}, e'_1, 0, \dots, 0, e'_{p-2}),$$

and by

$$\mathbf{d}_i = (-e'_{p-i}, \underbrace{0, \dots, 0}_{i-2}, -e'_{i-1}, e'_i, \underbrace{0, \dots, 0}_{p-i-2}, e'_{p-i-1}),$$

for  $i \in [2, (p-1)/2]$ . By Proposition 2.5, the dimension of  $\phi(G'')$  is equal to the maximum value of  $\dim \text{Circ}(\mathbf{d}_i)$  for  $i \in [1, (p-1)/2]$ . To determine the dimension of the latter spaces, recall that for any  $\mathbf{f} = (f_0, \dots, f_{p-1})$  we find  $\mathbf{f} \Pi_3 = (\mathbf{f} \Xi_p, \mathbf{f} \Xi_{p-1}, \mathbf{f} \Xi_{p-2})$ ,

$$\mathbf{f} \Xi_p = \sum_{i=0}^{p-1} f_i, \quad \mathbf{f} \Xi_{p-1} = \sum_{i=0}^{p-1} i f_i, \quad \text{and} \quad \mathbf{f} \Xi_{p-2} = \sum_{i=0}^{p-1} \binom{i}{2} f_i$$

and compute

$$(3-1) \quad \mathbf{d}_1 \Xi_p = 2(e_1 - e_{p-1}) + e_{p-2} - e_2,$$

$$(3-2) \quad \mathbf{d}_1 \Xi_{p-1} = e'_1 - e'_{p-2}.$$

$$(3-3) \quad \mathbf{d}_1 \Xi_{p-2} = e'_{p-2},$$

using  $\binom{p-1}{2} \equiv_p 1$  in the last line. Then, for  $i \in [2, (p-1)/2]$ , we compute

$$(3-4) \quad \mathbf{d}_i \Xi_p = -e'_{p-i} - e'_{i-1} + e'_i + e'_{p-i-1} = e''_{p-i-1} - e''_{i-1},$$

and for  $i \in [2, (p-1)/2]$  we find

$$(3-5) \quad \begin{aligned} \mathbf{d}_i \Xi_{p-1} &= -0 \cdot e'_{p-i} - (i-1)e'_{i-1} + ie'_i + (p-1)e'_{p-i-1} \\ &= -i \cdot e''_{i-1} + e'_{i-1} - e'_{p-i-1} \\ &= -i \cdot e''_{i-1} + \sum_{j=i-1}^{p-i-2} e''_j. \end{aligned}$$

Note that for  $i = (p-1)/2$ , this is equal to

$$(3-6) \quad \mathbf{d}_{(p-1)/2} \Xi_{p-1} = (p+3)/2 \cdot e''_{(p-3)/2}.$$

Case  $\text{sym}(\mathbf{e}'') = 0$ . This implies  $\text{sym}(\mathbf{e}) = 0$ , and  $V = \phi(\text{St}_G(1)')$  by Theorem 2.9(ii). By definition, there exists  $i \in [1, (p-3)/2]$  such that  $e''_i \neq e''_{p-2-i}$ , hence  $\mathbf{d}_{i+1} \Xi_p \neq 0$  by (3-4) and  $\dim \text{Circ}(\mathbf{d}_{i+1}) = p$  by Proposition 2.5. Thus  $\phi(G'')$  is equal to  $V$ , i.e.,  $G'' = \text{St}_G(1)'$ . Note that  $\text{sym}(\mathbf{e}'') = 0$  implies  $\text{con}(\mathbf{e}') = 0$ , since the difference vector of a constant vector is zero and in particular symmetric. Thus we find

$$\dim \phi(G'') = p = p - (\text{sym}(\mathbf{e}'') + \text{con}(\mathbf{e}')).$$

Case  $\text{sym}(\mathbf{e}'') = 1$ . By (3-4),  $\mathbf{d}_i \Xi_p = 0$  for all  $i \in [2, (p-1)/2]$ , and, by Lemma 2.4 and (3-1), also  $\mathbf{d}_1 \Xi_p = 0$ ; hence Proposition 2.5 implies  $\dim \phi(G'') \leq p-1$ . Using (3-5), we see that

$$(3-7) \quad \mathbf{d}_i \Xi_{p-1} = -i \cdot e''_{i-1} + \sum_{j=i-1}^{p-i-2} e''_j = (1-i) \cdot e''_{i-1} + 2 \sum_{j=i}^{(p-3)/2} e''_j$$

for  $i \in [2, (p-3)/2]$ .

If  $\text{con}(\mathbf{e}') = 0$ , then  $p \neq 3$  and  $\mathbf{e}'' \neq 0$ . By symmetry, there exists  $i \in [1, (p-3)/2]$  such that  $e''_i \neq 0$ . If  $e''_{(p-3)/2} \neq 0$ , we find  $\mathbf{d}_{(p-1)/2} \Xi_{p-1} \neq 0$  by (3-6). Otherwise, the prime  $p$  is at least 7. Let  $i \in [1, (p-5)/2]$  be maximal such that  $e''_i \neq 0$ . Then  $\mathbf{d}_{i+1} \Xi_{p-1} \neq 0$  by (3-7). Thus, by Proposition 2.5,  $\dim \phi(G'') = p-1 = p - (\text{sym}(\mathbf{e}'') + \text{con}(\mathbf{e}'))$ .

On the other hand, if  $\text{con}(\mathbf{e}') = 1$ , we immediately find  $\mathbf{d}_1 \Xi_{p-1}$  by (3-2). Furthermore,  $\mathbf{e}'' = \mathbf{0}$ , whence we also find  $\mathbf{d}_i \Xi_{p-1} = 0$  for  $i \in [2, (p-1)/2]$ , using (3-5). But

by (3-3),  $d_1 \Xi_{p-2} = e'_{p-2} \neq 0$ ; otherwise,  $e' = \mathbf{0}$  since it is constant, which implies  $\text{con}(e) = 1$ , which was excluded. Thus  $\dim \phi(G'') = p-2 = p - (\text{sym}(e'') + \text{con}(e'))$ . □

**Lemma 3.3.** *Let  $G$  be a GGS group defined by the nonconstant vector  $e$ . Then*

$$\log_p |G' : G''| = p + \text{con}(e') + \text{sym}(e'') - \text{sym}(e) - 1.$$

*Proof.* This is an immediate consequence of Lemma 2.8(iii), Proposition 3.2 and Theorem 2.9(ii):

$$\begin{aligned} \log_p |G' : G''| &= \log_p |G' : \text{St}_G(1)'| + \log_p |\chi_p G' : \psi(G'')| - \log_p |\chi_p G' : \psi(\text{St}_G(1)')| \\ &= p + \text{con}(e') + \text{sym}(e'') - \text{sym}(e) - 1. \end{aligned} \quad \square$$

**Proofs of the main results.** We are now in the position to prove our theorems, which we state again for the convenience of the reader.

**Theorem 1.2.** *Let  $p$  be an odd prime and let  $G$  be a GGS group acting on the  $p$ -adic tree defined by a nonconstant vector  $e \in \mathbb{F}_p^{p-1}$ . Let  $n \geq 3$ . Then*

$$\psi(G^{(n)}) = \chi_p G^{(n-1)}.$$

*If  $\text{con}(e') + \text{sym}(e'') = 0$ , the same holds for  $n = 2$ .*

*Proof.* Assume that the given equation holds true for some  $n \geq 1$ . Then we find

$$\psi(G^{(n+1)}) = \psi(G^{(n)})' = (\chi_p G^{(n-1)})' = \chi_p G^{(n)},$$

since  $G^{(n)} \leq \text{St}_G(1)$ . Thus it is enough to consider the case  $n = 3$ , or the case  $n = 2$ , respectively.

First assume that  $\text{con}(e') = \text{sym}(e'') = 0$ , which also implies  $\text{sym}(e) = 0$ . By Lemma 3.3,  $\chi_p G' = \psi(G'')$ . Thus the equation holds for  $n = 2$ .

We forgo the above assumptions on the defining vector and prove the desired equation for  $n = 3$ . In view of Proposition 2.1, it is sufficient to prove that  $\text{ins}_0([c, c^g]) \in G^{(3)}$  for any  $g \in G$ , since  $G''$  is normally generated by the elements  $\{[c, c^g] \mid g \in G\}$ . By Proposition 2.10, the group  $G''$  contains  $\psi^{-1}(\chi_p \gamma_3(G))$ , in particular  $\text{ins}_0([c, g]) \leq G'$  for all  $g \in G$ . By Proposition 3.2 we find  $h \in G''$  with

$$\psi(h) = (0 : c, 1 : c^{-2}, 2 : c, \diamond : \text{id}).$$

Let  $g \in G$ . Then

$$[h, \text{ins}_0([c, g])] = \text{ins}_0([h]_0, [c, g]) = \text{ins}_0([c, [c, g]]) = \text{ins}_0([c, c^g]) \in G^{(3)}. \quad \square$$

Before we prove Theorem 1.1, we use Theorem 1.2 to derive some more results on the structure of  $G$ .

**Corollary 3.4.** *Let  $G$  be a GGS group defined by a nonconstant vector and let  $n \in \mathbb{N}$ . Then the quotient  $G^{(n)}/G^{(n+1)}$  is an elementary abelian  $p$ -group.*

*Proof.* It is sufficient to show that all  $p$ -th powers in  $G^{(n)}$  are contained in  $G^{(n+1)}$  for all  $n \in \mathbb{N}$ . For  $n = 0$ , this is the statement of Lemma 2.8(i). Since  $G$  is self-similar, we find  $\psi(G') \leq \chi_p G$ , and, as a consequence,  $\psi(G^{(n)}) \leq \chi_p G^{(n-1)}$  for all  $n \geq 1$ . Thus for  $n = 1$ , notice that

$$\psi((G')^p) \leq (\chi_p G)^p \leq \chi_p \gamma_3(G) = \psi(\gamma_3(\text{St}_G(1))) \leq \psi(G''),$$

using Lemma 2.8(ii), Theorem 2.9(i) and Proposition 2.10. For general  $n > 1$ , using induction and Theorem 1.2, we see that

$$\psi(G^{(n)})^p \leq (\chi_p G^{(n-1)})^p = \chi_p (G^{(n-1)})^p \leq \chi_p G^{(n)} = \psi(G^{(n+1)}). \quad \square$$

Corollary 1.3 follows immediately. It remains to prove Theorem 1.1.

**Theorem 1.1.** *Let  $p$  be an odd prime and let  $G$  be a GGS group acting on the  $p$ -adic tree defined by a nonconstant vector  $e \in \mathbb{F}_p^{p-1}$ . Let  $e'$  be the tuple of differences between the entries of  $e$  and let  $e''$  be the tuple of differences of  $e'$ . Then for  $n \geq 2$ ,*

$$\log_p |G : G^{(n)}| = p^{n-2}(p + \text{con}(e') + \text{sym}(e'')) - \frac{p^{n-1} - 1}{p-1} \cdot \text{sym}(e) + 1.$$

*Proof.* Using Theorem 1.2, we find for  $n \geq 3$

$$\log_p |G^{(n)} : G^{(n+1)}| = \log_p |\chi_p G^{(n-1)} : \chi_p G^{(n)}| = p \cdot \log_p |G^{(n-1)} : G^{(n)}|,$$

and consequently

$$\log_p |G'' : G^{(n)}| = \sum_{i=0}^{n-3} p^i \log_p |G'' : G^{(3)}| = \frac{p^{n-2} - 1}{p-1} \cdot \log_p |G'' : G^{(3)}|.$$

Employing our previous results we find

$$\begin{aligned} \log_p |G'' : G^{(3)}| &\stackrel{\text{Thm. 1.2}}{=} \log_p |\chi_p G' : \chi_p G''| - \log_p |\chi_p G' : \psi(G'')| \\ &\stackrel{\text{Prop. 3.2}}{=} p \cdot \log_p |G' : G''| - (\text{con}(e') + \text{sym}(e'')) \\ &\stackrel{\text{Lem. 3.3}}{=} p(p-1 + \text{con}(e') + \text{sym}(e'') - \text{sym}(e)) - (\text{con}(e') + \text{sym}(e'')) \\ &= (p-1)(p + \text{con}(e') + \text{sym}(e'')) - p \cdot \text{sym}(e). \end{aligned}$$

Putting everything together (using Lemma 2.8(i) and, again, Lemma 3.3), we find

$$\begin{aligned} \log_p |G : G^{(n)}| &= (p^{n-2} - 1)(p + \text{con}(e') + \text{sym}(e'')) \\ &\quad - \frac{p^{n-1} - p}{p-1} \cdot \text{sym}(e) + \log_p |G' : G''| + \log_p |G : G'| \\ &= p^{n-2}(p + \text{con}(e') + \text{sym}(e'')) - \frac{p^{n-1} - 1}{p-1} \cdot \text{sym}(e) + 1. \quad \square \end{aligned}$$

**GGs groups with differentially constant defining vector.** A vector  $e$  is called *differentially constant* if it satisfies  $\text{con}(e') = 1$  (and thus also  $\text{sym}(e'') = 1$ ). In view of Theorem 1.1, GGS groups with differentially constant defining vector display the largest indices  $|G : G^{(n)}|$  among GGS groups with nonconstant defining vector, as  $\text{con}(e') = 1$  implies  $\text{sym}(e) = 0$ . The condition  $\text{con}(e') = 1$  is a strong restriction on the defining vector, making it possible to determine the isomorphism classes of differentially constant GGS groups.

Resolving the definition, one finds that  $\text{con}(e') = 1$  implies that there exist  $k, m \in \mathbb{F}_p$  such that  $e = (k + m, k + 2m, \dots, k + (p-1)m)$ . We introduce the shorthand notation  $\text{dc}(k, m)$  for the vector given above. If  $m = 0$ , evidently  $\text{con}(e) = 1$ .

**Proposition 3.5.** *For any given odd prime  $p$  there are precisely three isomorphism classes of differentially constant GGS groups acting on the  $p$ -adic tree:*

- (i) *one consisting of the constant GGS group,*
- (ii) *one consisting of a single periodic GGS group,*
- (iii) *one containing precisely  $p-1$  nonperiodic GGS groups.*

*Proof.* Recall the isomorphism class preserving action  $*$  of  $(\mathbb{F}_p^\times)^2$  on the set of defining vectors given by (2-1). Let  $G$  be the GGS group defined by  $\text{dc}(k, m)$  and let  $\lambda, \mu \in \mathbb{F}_p^\times$ . Then

$$\text{dc}(k, m) * (\lambda, \mu) = \text{dc}(\lambda k, \lambda m) * (1, \mu) = \text{dc}(\lambda k, \lambda \mu m).$$

If  $m = 0$ , the vector  $\text{dc}(k, 0)$  is constant, and we are in case (i). If  $m \neq 0$ , we find

$$\text{dc}(0, m) * (1, m^{-1}) = \text{dc}(0, 1) \quad \text{and} \quad \text{dc}(k, m) * (k^{-1}, km^{-1}) = \text{dc}(1, 1)$$

for  $k \neq 0$ . At the same time,  $\text{dc}(0, m) * (\lambda, \mu) \neq \text{dc}(1, 1)$  for all  $\lambda, \mu \in \mathbb{F}_p^\times$ , whence  $\text{dc}(0, 1)$  and  $\text{dc}(1, 1)$  represent distinct isomorphism classes. The  $(\mathbb{F}_p^\times)^2$ -orbit of  $\text{dc}(0, 1)$  consists of the multiples of  $\text{dc}(0, 1)$  and all the associated GGS groups are identical. By [9, Example 9.1], the GGS group defined by  $e$  is periodic if and only if the sum of the entries of  $e$  vanishes, i.e., if  $\dim \text{Circ}(\bar{f}e) \leq p-1$ . For  $\text{dc}(0, 1) = (1, \dots, p-1)$  this is the case, but not for  $\text{dc}(1, 1) = (2, \dots, p-1, 0)$ . As seen above,  $(\mathbb{F}_p^\times)^2$  acts transitively on  $\{\text{dc}(k, m) \mid k, m \in \mathbb{F}_p^\times\}$ . Since only proportional vectors yield identical GGS groups, there are  $p-1$  distinct GGS groups isomorphic to the group defined by  $\text{dc}(1, 1)$ . □

In case of the prime  $p = 3$ , all GGS groups are differentially constant and the unique periodic GGS group is the Gupta–Sidki 3-group. It is interesting to see that other invariants take extremal values for the groups  $G_p$  defined by  $(1, 2, \dots, p-1)$  (for arbitrary  $p$ ): By [7, Theorem B], the Hausdorff dimension of the GGS group

defined by  $\mathbf{e}$  is given by

$$\frac{(p-1) \operatorname{cr}(\mathbf{e})}{p^2} - \frac{\operatorname{sym}(\mathbf{e})}{p^2} - \frac{\operatorname{con}(\mathbf{e})}{(p-1)p^2}.$$

By Corollary 2.6,  $\operatorname{cr}(\mathbf{e}) = 2$  if and only if  $\mathbf{e}$  is a nonzero multiple of  $\operatorname{dc}(0, 1)$ . In particular, a symmetric defining vector  $\mathbf{e}$  fulfils  $\operatorname{cr}(\mathbf{e}) > 2$ . Furthermore, a constant defining vector has  $\operatorname{cr}(\mathbf{e}) = p$ . Thus the group  $G_p$  is the unique GGS group with the minimal (among GGS groups acting on a fixed tree) possible Hausdorff dimension  $2(p-1)/p^2$ , while the group defined by  $\operatorname{dc}(1, 1)$  is among those with maximal Hausdorff dimension  $(p-1)/p$ .

The automorphism group of  $G_p$  is as large as possible; cf. [15, Example 6.2].

**Comparison with the congruence subgroups.** The members of the derived series of a GGS group share certain properties with the level stabilisers. They both form filtrations of the group, with  $\operatorname{St}_G(n) \geq G^{(n)}$  for all  $n \in \mathbb{N}$ ; for sufficiently high values of  $n$ , they satisfy  $\psi(G_n) = \chi_p G_{n-1}$  by Theorem 1.2 and [7, Lemma 3.3], respectively; furthermore, the quotients of respective consecutive members are elementary abelian  $p$ -groups. Using the formula for the index of the  $n$ -th level stabiliser provided by Fernández-Alcober and Zugadi-Reizabal in [7, Theorem A] one finds

$$\log_p |G : \operatorname{St}_G(2)| = \operatorname{cr}(\mathbf{e}) + 1,$$

hence, using the consequence  $\log_p |G : \operatorname{St}_G(1)'| = p + 1$  of Lemma 2.8, we find that  $\log_p |\operatorname{St}_G(2) : \operatorname{St}_G(1)'| = p - \operatorname{cr}(\mathbf{e})$ . A comparison with  $\log_p |G : G''| = p + \operatorname{con}(\mathbf{e}') + \operatorname{sym}(\mathbf{e}'') - \operatorname{sym}(\mathbf{e}) + 1$  makes it apparent that  $G'' = \operatorname{St}_G(2)$  if and only if  $\mathbf{e}'$  is nonconstant,  $\operatorname{cr}(\mathbf{e}) = p$ , and  $\operatorname{sym}(\mathbf{e}) = \operatorname{sym}(\mathbf{e}'')$ . Since  $\psi(G_n) = \chi_p G_{n-1}$  holds for both the derived series and the series of level stabilisers from  $n = 3$  onwards, the series only differ in their first term in this case, and otherwise have no equal terms at all. The largest difference is attained for the periodic groups  $G_p$  with the differentially constant defining vector  $\operatorname{dc}(0, 1)$ ; recall that  $\operatorname{cr}(\operatorname{dc}(0, 1)) = 2$ , and thus

$$\log_p |\operatorname{St}_{G_p}(n) : G_p^{(n)}| = p^{n-1}.$$

By Proposition 2.5,  $\operatorname{cr}(\mathbf{e}) = 2$  is the minimum possible value; however one finds that  $\operatorname{cr}(\mathbf{e})$  may take any value in  $[2, p]$ . Therefore the number of distinct sequences  $(|G : \operatorname{St}_G(n)|)_{n \in \mathbb{N}}$  obtained by any GGS group is between  $p$  and  $2p-1$  (note that symmetric defining vectors do not admit all values  $[2, p]$  under  $\operatorname{cr}$ ); in particular, the number of classes of GGS groups separated by the sequence of indices of their level stabilisers grows with  $p$ . In contrast, the sequences  $(|G : G^{(n)}|)_{n \in \mathbb{N}}$  yield a partition into five subsets for  $p \neq 3$ ; for  $p = 3$ , one obtains 2 classes.

### Acknowledgements

The results of this article have previously appeared as a part of the author's Ph.D. dissertation, written under the supervision of B. Klopsch at the Heinrich-Heine-Universität Düsseldorf. The author thanks K. Rajeev for helpful discussions and bringing his attention to the problem, and G. Fernández-Alcober, M. Garcia-rena Perez, M. Piccolo and D. Tijsma for their comments on preliminary versions of this paper. Finally, the author would like to thank the referees for their most valuable feedback and suggestions.

### References

- [1] T. Alexoudas, B. Klopsch, and A. Thillaisundaram, “Maximal subgroups of multi-edge spinal groups”, *Groups Geom. Dyn.* **10**:2 (2016), 619–648. MR
- [2] L. Bartholdi and Z. Šunić, “On the word and period growth of some groups of tree automorphisms”, *Comm. Algebra* **29**:11 (2001), 4923–4964. MR
- [3] L. Bartholdi, R. I. Grigorchuk, and Z. Šunić, “Branch groups”, pp. 989–1112 in *Handbook of algebra*, vol. 3, Elsevier, Amsterdam, 2003. MR
- [4] P. J. Davis, *Circulant matrices*, Wiley, New York, 1979. MR
- [5] E. Di Domenico, G. A. Fernández-Alcober, and N. Gavioli, “GGs-groups over primary trees: branch structures”, *Monatsh. Math.* **200**:4 (2023), 781–797. MR
- [6] J. Fabrykowski and N. Gupta, “On groups with sub-exponential growth functions”, *J. Indian Math. Soc. (N.S.)* **49**:3-4 (1985), 249–256. MR
- [7] G. A. Fernández-Alcober and A. Zugadi-Reizabal, “GGs-groups: order of congruence quotients and Hausdorff dimension”, *Trans. Amer. Math. Soc.* **366**:4 (2014), 1993–2017. MR
- [8] G. A. Fernández-Alcober, A. Garrido, and J. Uria-Albizuri, “On the congruence subgroup property for GGS-groups”, *Proc. Amer. Math. Soc.* **145**:8 (2017), 3311–3322. MR
- [9] R. I. Grigorchuk, “Just infinite branch groups”, pp. 121–179 in *New horizons in pro-p groups*, Progr. Math. **184**, Birkhäuser, Boston, 2000. MR
- [10] N. Gupta and S. Sidki, “On the Burnside problem for periodic groups”, *Math. Z.* **182**:3 (1983), 385–388. MR
- [11] B. Klopsch and A. Thillaisundaram, “Maximal subgroups and irreducible representations of generalized multi-edge spinal groups”, *Proc. Edinb. Math. Soc. (2)* **61**:3 (2018), 673–703. MR
- [12] E. Pervova, “Profinite completions of some groups acting on trees”, *J. Algebra* **310**:2 (2007), 858–879. MR
- [13] J. M. Petschick, “On conjugacy of GGS-groups”, *J. Group Theory* **22**:3 (2019), 347–358. MR
- [14] J. M. Petschick, “Two periodicity conditions for spinal groups”, *J. Algebra* **633** (2023), 242–269. MR
- [15] J. M. Petschick, “The automorphism group of a multi-GGS group”, *J. Algebra Appl.* **23**:14 (2024), art. id. 2450241, 24 pp.
- [16] A. C. Vieira, “On the lower central series and the derived series of the Gupta–Sidki 3-group”, *Comm. Algebra* **26**:4 (1998), 1319–1333. MR
- [17] T. Vovkivsky, “Infinite torsion groups arising as generalizations of the second Grigorchuk group”, pp. 357–377 in *Algebra (Moscow, 1998)*, de Gruyter, Berlin, 2000. MR

Received June 2, 2023. Revised March 15, 2026.

J. MORITZ PETSCHICK  
FAKULTÄT FÜR MATHEMATIK  
UNIVERSITÄT BIELEFELD  
BIELEFELD  
GERMANY  
[jpetschick@math.uni-bielefeld.de](mailto:jpetschick@math.uni-bielefeld.de)

# PROPERTY QT OF RELATIVELY HIERARCHICALLY HYPERBOLIC GROUPS

BINGXUE TAO

Using the projection complex machinery, a number of authors (Bestvina, Bromberg and Fujiwara; Hagen and Petyt; and Han, Nguyen and Yang) have proved that several classes of nonpositively curved groups admit equivariant quasi-isometric embeddings into finite products of quasitrees, i.e., having property QT. Here we unify and generalize those results by establishing a sufficient condition for relatively hierarchically hyperbolic groups to have property QT.

As applications, we show that a group has property QT if it is residually finite and belongs to one of the following classes of groups: admissible groups, hyperbolic-2-decomposable groups with no distorted elements, and Artin groups of large and hyperbolic type. We also introduce a slightly stronger version of property QT, called property  $QT_0$ , and show the invariance of property  $QT_0$  under graph products.

## 1. Introduction

Group actions on quasitrees have been studied intensively in recent years. A *quasitree* is a geodesic space quasi-isometric to a simplicial tree. We say that a finitely generated group  $G$  has *property QT* if  $G$  acts on a finite product of quasitrees (equipped with the  $\ell^1$ -metric) such that the orbit map is a quasi-isometric embedding. Such an embedding is called a *QT embedding* of  $G$ . Since a quasitree has asymptotic dimension at most 1, property QT is a stronger form of finite asymptotic dimension. Examples of groups with property QT include

- Coxeter groups [DJ99];
- residually finite hyperbolic groups [BBF21];
- mapping class groups of finite-type surfaces [BBF21];
- virtually colorable hierarchically hyperbolic groups whose associated hyperbolic spaces are all quasitrees [HP22] (including virtually compact special

---

*MSC2020*: primary 20F65, 20F67; secondary 20E06, 20E26, 20F36.

*Keywords*: hierarchically hyperbolic, quasitree, projection complex, residually finite.

groups [BHS17b], the genus 2 handlebody group [Che22], fundamental groups of nongeometric graph manifolds [HRSS24]);

- fundamental groups of compact orientable 3-manifolds whose sphere-disk decomposition does not support either Sol or Nil geometry [HNY25];

along with their undistorted subgroups.

The last four examples were proved to have property QT with the help of the projection complex techniques developed in [BBF15; BBFS19]. For mapping class groups, property QT strengthens Theorem C of [BBF15], which says that mapping class groups equivariantly quasi-isometrically embed in a finite product of hyperbolic graphs of finite asymptotic dimension. Counterexamples of property QT include certain special linear groups [Man06; Man08], generalized Baumslag–Solitar groups with infinite monodromy [But25] and groups with Property hereditary (NL) [BFG24]. For some basic corollaries of property QT, see [HNY25, §§2.1, 2.2]. Recently, Vergara [Ver24] proved that any finitely generated group with property QT has a proper uniformly Lipschitz affine action on  $\ell^1$  with quasi-isometrically embedded orbits.

As a generalization of the Masur–Minsky machinery [MM99; MM00], *hierarchically hyperbolic groups* [BHS17b; BHS19], abbreviated HHGs, have become an important bridge between mapping class groups, cubical groups, and many other nonpositively curved groups. A list of papers in this field can be found in [HRSS24]. Coarsely speaking, an HHG is a finitely generated group  $G$  whose geometry can be recovered from  $G$ -equivariant projections to a specified (possibly infinite) collection of hyperbolic spaces. For background on HHGs and relative HHGs, see Section 2.3. As shown in [BHS17a], HHGs have finite asymptotic dimension. This leads to a natural question:

**Question 1.1.** *Which HHGs have property QT?*

In this paper, we provide a sufficient condition for relative HHGs to have property QT, which reproves those of [BBF21; HP22; HNY25]. We also give a sufficient condition for the existence of a *quasimedial* QT embedding (for this notion, see [HP22]). This stronger property can be used to prove the existence of globally stable cylinders (see [PSZ25]), which connects to a long-standing question of Rips and Sela [RS95] about canonical representatives of elements in hyperbolic groups. The following is a collection of applications from Section 7. These results are new except for mapping class groups.

**Theorem 1.2.** *The following groups have property QT:*

- *mapping class groups of finite-type surfaces;*
- *residually finite admissible graphs of groups;*
- *residually finite hyperbolic-2-decomposable groups with no distorted elements;*

- *residually finite Artin groups of large and hyperbolic type.*

*The QT embeddings for all these groups are quasimedians.*

When studying the invariance of property QT in some cases, we want the group action on the product space to be diagonal. We say that a finitely generated group  $G$  has *property QT<sub>0</sub>* if  $G$  has property QT and the  $G$ -action on the finite product of quasitrees is diagonal. By [HNY25, Theorem 1.5], if a residually finite group  $G$  is hyperbolic relative to a collection of groups with property QT<sub>0</sub>, then  $G$  has property QT. Without ambiguity, we also say that a  $G$ -action on a metric space  $X$  has *property QT<sub>0</sub>* if  $X$  admits a  $G$ -equivariant quasi-isometric embedding into a finite product of quasitrees on which  $G$  acts diagonally. In particular, if  $X$  itself is a finite product of quasitrees, then any diagonal action on  $X$  has property QT<sub>0</sub>. We prove the following invariance of property QT<sub>0</sub> under graph products.

**Theorem 7.10.** *Any graph product of groups whose every vertex group has property QT<sub>0</sub> still has property QT<sub>0</sub>.*

Now we give the main definitions needed to state our main theorem.

**Definition 1.3.** Let  $(G, \mathfrak{S})$  be a relative HHG. For any  $U \in \mathfrak{S}$ , we write  $G_U < \text{Aut}(\mathfrak{S}_U)$  to mean the image of  $\text{Stab}_G(U)$  under the restriction homomorphism.

- (1) We say a domain  $U \in \mathfrak{S}$  is of *type I* if it has the following properties:
  - (*hyperbolicity*)  $\mathcal{C}U$  is hyperbolic.
  - (*acylindrical image*)  $G_U$  acts on  $\mathcal{C}U$  acylindrically.
  - (*cobounded nested region*)  $G_U$  acts on  $F_U$  coboundedly.
  - (*separable quasi-axes*) For any element  $g \in \text{Stab}_G(U)$  that acts loxodromically on  $\mathcal{C}U$ , the elementary closure  $EC(g)$  is separable in  $G$ .
- (2) We say a domain  $U \in \mathfrak{S}$  is of *type II* if the action  $G_U \curvearrowright \mathcal{C}U$  has property QT<sub>0</sub>.

For any  $U \in \mathfrak{S}$  of type II, property QT<sub>0</sub> provides quasitrees  $T_U^i$  along with  $G_U$ -equivariant maps  $t_U^i : \mathcal{C}U \rightarrow T_U^i$  for  $i = 1, \dots, n_U$  such that

$$\prod_{i=1}^{n_U} t_U^i : \mathcal{C}U \rightarrow \prod_{i=1}^{n_U} T_U^i$$

is a quasi-isometric embedding. Our main theorem is as follows.

**Theorem 1.4.** *Let  $(G, \mathfrak{S})$  be a relative HHG that is virtually colorable. If every  $U \in \mathfrak{S}$  is of type I or II, then  $G$  has property QT.*

*If, for any  $D \geq 1$ , there exists  $D' \geq 1$  such that for every  $U \in \mathfrak{S}$  of type II and each  $i = 1, \dots, n_U$  the map  $t_U^i : \mathcal{C}U \rightarrow T_U^i$  sends  $(D, D)$ -quasigeodesics to unparametrized  $(D', D')$  quasigeodesics, then  $G$  is coarse median and the QT embedding of  $G$  is quasimedian.*

*Sketch of proof.* We roughly explain how to prove Theorem 1.4 in the case that  $(G, \mathfrak{S})$  is an HHG with only type I domains, excluding the “moreover” part. This case contains most of the key ideas.

First, we introduce a class of *thick distances* on  $G$  each of which is defined using a class of *thick segments* of hierarchy paths on  $G$ . We prove in Section 3 a thick distance formula saying that the word metric of  $G$  can be recovered by summing up these thick distances. This is an analogue of the distance formula for HHGs.

Then we show that any class of thick segments is cofinite up to the group action. Furthermore, these thick segments can be extended to a cofinite collection of quasi-axes. Using projections to these quasi-axes, we can estimate the thick distance. This is done in Section 4.

Finally, we take a finite-index subgroup of  $G$ , say  $H$ , such that a collection of quasi-axes as above is divided into finitely many  $H$ -orbits. Each  $H$ -orbit satisfies the Bestvina–Bromberg–Fujiwara projection axioms, so it gives us a quasitree with an  $H$ -action. We prove that  $H$  equivariantly quasi-isometrically embeds in the product of these finitely many quasitrees in Section 5. Since property QT is commensurably invariant,  $G$  has property QT.  $\square$

As stated in the proof, property QT is commensurably invariant [BBF21, §2.2]. It follows that the conclusion of Theorem 1.4 also holds for any group that is virtually a relative HHG that satisfies our condition, even though such a group may not be a relative HHG itself [PS23].

For most examples of HHGs that emerged from the study, every domain satisfies the first three conditions of type I. Virtual colorability is also common in practice. Therefore, the biggest restriction of our theorem comes from the assumption of separable quasi-axes. We further discuss it in Section 6 and show how residual finiteness helps to give an easy-to-use criterion for having separable quasi-axes.

## 2. Background

**2.1. Quasi-isometric embeddings and acylindricity.** Given constants  $\lambda \geq 1$ ,  $c \geq 0$ , we say that a coarse map  $f : X \rightarrow Y$  between metric spaces  $(X, d_X)$  and  $(Y, d_Y)$  is a  $(\lambda, c)$ -quasi-isometric embedding if

$$\frac{1}{\lambda}d_X(x_1, x_2) - c \leq d_Y(f(x_1), f(x_2)) \leq \lambda d_X(x_1, x_2) + c$$

for all  $x_1, x_2 \in X$ . A  $(\lambda, c)$ -quasi-isometric embedding  $\gamma : [0, l] \rightarrow X$  is called a (parametrized)  $(\lambda, c)$ -quasigeodesic in  $X$ . A coarse map  $\gamma : [0, l] \rightarrow X$  is an *unparametrized*  $(\lambda, c)$ -quasigeodesic if there exists a strictly increasing function  $f : [0, l] \rightarrow [0, l]$  with  $f(0) = 0$ ,  $f(l) = l$  such that  $\gamma \circ f$  is a  $(\lambda, c)$ -quasigeodesic. We also use the term “quasigeodesic” to mean a quasi-isometric embedding of  $\mathbb{R}$ . We will not distinguish between a quasigeodesic and its image in  $X$ .

A geodesic metric space is called  $\delta$ -hyperbolic (or simply hyperbolic) for  $\delta \geq 0$  if for any geodesics  $\alpha, \beta, \gamma$  that form a triangle,  $\alpha$  is contained in the  $\delta$ -neighborhood of  $\beta \cup \gamma$  [Gro87]. For a  $\delta$ -hyperbolic space  $X$ , an isometry  $g : X \rightarrow X$  is called *loxodromic* if the  $g$ -orbit  $n \mapsto g^n x$  is a quasigeodesic for some (equivalently, any)  $x \in X$ .

Let  $X$  be a hyperbolic space and  $G$  be a group acting by isometries on  $X$  with a loxodromic element  $g$ . Given constants  $\lambda \geq 1$  and  $c \geq 0$ , a  $(\lambda, c)$ -quasigeodesic  $\gamma \subset X$  is called a  $(\lambda, c)$ -quasi-axis for  $g$  if  $\gamma$  is  $g$ -invariant. The *elementary closure* of  $g$  in  $G$ ,  $EC_G(g)$ , is the subgroup of  $G$  that stabilizes  $\gamma$  up to bounded Hausdorff distance. If there is no ambiguity in  $G$ , we often simplify the notation as  $EC(g)$ . Equivalently, it is the stabilizer of the set  $\gamma(\pm\infty)$ , the points at infinity of  $\gamma$ . Thus, the elementary closure does not depend on the choice of  $\gamma$ . Everything that commutes with  $g$  is contained in  $EC(g)$  (including powers and roots), but there may be other elements.

A group action  $G \curvearrowright X$  by isometries is called *acylindrical* [Bow08] if for any  $r \geq 0$  there exist constants  $R, N \geq 0$  such that, for any pair  $a, b \in X$  with  $d(a, b) \geq R$ ,

$$\#\{g \in G \mid d(ga, a) \leq r \text{ and } d(gb, b) \leq r\} \leq N.$$

Let  $X$  be a hyperbolic space and  $G$  be a group acting acylindrically on  $X$  with a loxodromic element  $g$ . Some basic properties of this kind of action can be found in [Osi16]. In this case, the elementary closure  $EC(g)$  is the unique maximal virtually cyclic subgroup of  $G$  that contains  $g$  [DGO17, Lemma 6.5]. Moreover,  $EC(g)$  has a subgroup of index at most 2 that is a centralizer of a large power of  $g$  in  $G$  [DGO17, Corollary 6.6].

In this paper, we will consider group actions with a large kernel, in which case the action cannot be acylindrical. As in [BBF21], an action  $G \curvearrowright X$  is said to have *acylindrical image* if the image of  $G$  in the isometry group of  $X$  is acylindrical.

**2.2. Projection axioms.** In this section, we review the construction of a quasitree of spaces in [BBF15] with improvements from [BBFS19].

Let  $\mathbf{Y}$  be a collection of geodesic metric spaces, and  $\pi_Y(X) \subset Y$  be specified subsets whenever  $X \neq Y$  are elements of  $\mathbf{Y}$ . Set  $d_Y^\pi(X, Z) := \text{diam}(\pi_Y(X) \cup \pi_Y(Z))$  for  $X \neq Y \neq Z$ . We say that  $(\mathbf{Y}, \{\pi_Y\})$  is a *projection system* with *projection constant*  $\xi \geq 0$  if it satisfies the following *projection axioms*:

- (P0) (*bounded projection*)  $\text{diam}(\pi_Y(X)) \leq \xi$  when  $X \neq Y$ .
- (P1) (*Behrstock inequality*) If  $X, Y, Z$  are distinct and  $d_Y^\pi(X, Z) > \xi$ , then  $d_X^\pi(Y, Z) \leq \xi$ .
- (P2) (*finiteness*) For  $X \neq Z$  the set

$$\{Y \in \mathbf{Y} \mid d_Y^\pi(X, Z) > \xi\}$$

is finite.

We say that  $(Y, \{\pi_Y\})$  is a  $G$ -projection system if a group  $G$  acts on the set  $Y$  in such a way that every  $g \in G$  acts as an isometry from  $Y$  to  $gY$  and the projections  $\pi_Y$  are  $G$ -equivariant, that is,  $\pi_{gY}(gX) = g\pi_Y(X)$ .

We say that  $(Y, \{\pi_Y\})$  satisfies the *strong projection axioms* if (P0), (P2) and the following replacement for (P1) are satisfied:

(P1') If  $X, Y, Z$  are distinct and  $d_Y^x(X, Z) > \xi$ , then  $\pi_X(Y) = \pi_X(Z)$ .

While there are many natural situations where the projection axioms hold, the strong projection axioms are not as natural. However, we can modify the projections so that they do hold.

**Theorem 2.1** [BBFS19, Theorem 4.1]. *If  $(Y, \{\pi_Y\})$  is a projection system with constant  $\xi$ , then there are projections  $\{\pi'_Y\}$  such that  $(Y, \{\pi'_Y\})$  satisfies the strong projections axioms with constant  $\xi'$ , where  $\pi'_Y(X)$  and  $\pi_Y(X)$  are apart from each other within a uniform Hausdorff distance  $\epsilon$ , and  $\epsilon$  and  $\xi'$  only depend on  $\xi$ .*

*If  $(Y, \{\pi_Y\})$  is a  $G$ -projection system, then  $(Y, \{\pi'_Y\})$  is still a  $G$ -projection system.*

Let  $\mathcal{C}_K Y$  denote the space obtained from the disjoint union

$$\bigsqcup_{Y \in \mathcal{Y}} Y$$

by joining points in  $\pi_X(Z)$  with points in  $\pi_Z(X)$  by an edge of length one whenever  $d_Y(X, Z) < K$  for all  $Y \in \mathcal{Y} - \{X, Z\}$ . When the spaces are graphs and projections are subgraphs, we can join just the vertices in these projections. If  $\mathcal{Y}$  is a  $G$ -projection system, then  $G$  acts isometrically on  $\mathcal{C}_K Y$ .

**Theorem 2.2** [BBF15, §4]. *Let  $(Y, \{\pi_Y\})$  satisfy the strong projection axioms with constant  $\xi$ , and take  $K > 2\xi$ . Then:*

- $\mathcal{C}_K Y$  is hyperbolic if all  $Y \in \mathcal{Y}$  are  $\delta$ -hyperbolic.
- $\mathcal{C}_K Y$  is a quasitree if all  $Y \in \mathcal{Y}$  are quasitrees with uniform QI constants.

There is a very useful distance formula in  $\mathcal{C}_K Y$ . Let  $X, Z \in \mathcal{Y}$  and  $x \in X, z \in Z$ . We define  $\pi_Y(x) = \pi_Y(X)$  if  $Y \neq X$  and define  $\pi_X(x) = x$ . Then define  $d_Y(x, z) = \text{diam}(\pi_Y(x) \cup \pi_Y(z))$ .

**Notation.** Given  $A, B \geq 0$ , we define a threshold function by

$$\{\!\!\{A\}\!\!\}_B = \begin{cases} A & \text{if } A \geq B, \\ 0 & \text{otherwise.} \end{cases}$$

**Theorem 2.3** [BBFS19, Theorem 6.3]. *Let  $(Y, \{\pi_Y\})$  satisfy the strong projection axioms with constant  $\xi$ . Let  $x \in X$  and  $z \in Z$  be two points of  $\mathcal{C}_K(Y)$  with  $X, Z \in Y$ . Then*

$$\frac{1}{4} \sum_{Y \in Y} \{\{d_Y(x, z)\}\}_K \leq d_{\mathcal{C}_K Y}(x, z) \leq 2 \sum_{Y \in Y} \{\{d_Y(x, z)\}\}_K + 3K$$

for all  $K \geq 4\xi$ .

Next we recall a theorem that allows us to pass projection axioms from a projection system to a collection of certain subspaces. Let  $Y$  be a collection of  $\delta$ -hyperbolic spaces and  $(Y, \{\pi_Y\})$  be a projection system with constant  $\xi$ . For each  $Y \in Y$ , let  $\mathcal{A}_Y$  be a collection of quasigeodesics in  $Y$ . Let  $\mathcal{A}$  be the disjoint union of all  $\mathcal{A}_Y$ 's. We also make the following assumptions.

- As a collection of quasigeodesics,  $\mathcal{A}$  has uniform coarse constants.
- For  $\alpha, \beta \in \mathcal{A}_Y$ , we define  $\pi_\alpha(\beta)$  to be the closest-point projection of  $\beta$  to  $\alpha$ .
- For  $\alpha \in \mathcal{A}_X$  and  $\beta \in \mathcal{A}_Y$  where  $X \neq Y$ , we define  $\pi_\alpha(\beta)$  to be the closest point projection of  $\pi_X(Y)$  to  $\alpha$ .

**Theorem 2.4** [BBF21, Theorem 4.17]. *For any  $\theta > 0$ , there exists  $\xi' > 0$ , depending only on  $\theta, \delta, \xi$  and coarse constants of  $\mathcal{A}$ , such that, if  $\text{diam}(\pi_\alpha(\beta)) \leq \theta$  whenever  $\alpha$  and  $\beta$  are distinct elements in the same  $\mathcal{A}_Y$ , then  $(\mathcal{A}, \{\pi_\alpha\})$  is a projection system with constant  $\xi'$ .*

**2.3. (Relatively) hierarchically hyperbolic spaces.** In this paper, we deal with (relatively) hierarchically hyperbolic spaces and (relatively) hierarchically hyperbolic groups. Coarsely speaking, a (relative) HHS is a pair  $(\mathcal{X}, \mathfrak{S})$ , where  $\mathcal{X}$  is a quasigeodesic space and  $\mathfrak{S}$  is an index set, with some extra structure. A full definition can be found in [BHS19, Definition 1.1,1.21]. Some important information from the definition is collected below.

- An element  $U \in \mathfrak{S}$  is called a *domain* of  $\mathcal{X}$ .  $\mathfrak{S}$  has a partial order  $\sqsubseteq$ , called *nesting*, and a symmetric relation  $\perp$ , called *orthogonality*. These two relations are required to be mutually exclusive. Any two elements that are neither comparable under the partial order nor mutually orthogonal are by definition mutually *transversal*; we denote this relation by  $\pitchfork$ . We denote by  $\mathfrak{S}_U$  the set of domains nested in  $U$ , and by  $\mathfrak{S}_U^\circ$  the set of domains properly nested in  $U$ .
- There is a unique  $\sqsubseteq$ -maximal element  $S$  in  $\mathfrak{S}$  and a uniform bound on the length of  $\sqsubseteq$ -chains in  $\mathfrak{S}$ , called the *complexity* of  $(\mathcal{X}, \mathfrak{S})$ . The *level*  $\ell(V)$  of  $V \in \mathfrak{S}$  is defined inductively as follows. If  $V$  is  $\sqsubseteq$ -minimal then  $\ell(V) = 1$ . The element  $V$  has level  $k+1$  if  $k$  is the maximal integer such that there exists  $U \sqsubset V$  with  $\ell(U) = k$ .
- For HHSes, there is a set  $\{\{\mathcal{C}U, d_U\} : U \in \mathfrak{S}\}$  of uniformly hyperbolic spaces and a set of uniformly coarsely Lipschitz and coarsely surjective maps  $\pi_U : \mathcal{X} \rightarrow \mathcal{C}U$

for all  $U \in \mathfrak{S}$ . For relative HHSes, the complexity is at least 2. If  $U$  is  $\sqsubseteq$ -minimal,  $\mathcal{C}U$  is not required to be hyperbolic. This is the only difference between HHSes and relative HHSes in definition.

- For  $U \sqsubset V$  or  $U \pitchfork V$ , there is a uniformly bounded set  $\rho_V^U \subset \mathcal{C}V$ .
- For  $U \sqsubseteq V$ , there is a coarse map  $\rho_U^V : \mathcal{C}V \rightarrow \mathcal{C}U$ .
- Whenever  $V \sqsubseteq W$  and  $W \perp U$ , we require that  $V \perp U$ .
- (*orthogonal containers*) For each  $T \in \mathfrak{S}$  and each  $U \in \mathfrak{S}_T$  for which the set  $\{V \in \mathfrak{S}_T \mid V \perp U\} \neq \emptyset$ , there exists  $W \in \mathfrak{S}_T^\circ$ , so that whenever  $V \perp U$  and  $V \sqsubseteq T$ , we have  $V \sqsubseteq W$ . We say that  $W$  is an *orthogonal container* of  $U$  in  $T$  if  $W$  is a  $\sqsubseteq$ -minimal element satisfying the above property. Let  $\text{cont}_T^\perp U$  denote the set of all orthogonal containers of  $U$  in  $T$ . If  $T$  is the maximal element of  $\mathfrak{S}$ , then we suppress it from the notation and write  $\text{cont}^\perp U$ . We set  $\mathfrak{S}_U^\perp = \{V \in \mathfrak{S} \mid V \perp U\} \cup \{A\}$ , where  $A$  is an arbitrary element of  $\text{cont}^\perp U$ .
- (*consistency*) For every  $x \in X$ , the tuple  $(\pi_U(x))_{U \in \mathfrak{S}}$  is  $\kappa_0$ -consistent (defined below). If  $U \sqsubseteq V$ , then  $d_W(\rho_W^U, \rho_W^V) \leq \kappa_0$  whenever  $W \in \mathfrak{S}$  satisfies either  $V \sqsubset W$  or  $V \pitchfork W$  and  $U \not\sqsubseteq W$ .
- (*bounded geodesic image*) There exists  $E > 0$  such that for all  $W \in \mathfrak{S}$ , all  $V \in \mathfrak{S}_W^\circ$ , and all  $x, y \in \mathcal{X}$  such that some geodesic from  $\pi_W(x)$  to  $\pi_W(y)$  stays  $E$ -far from  $\rho_W^V$ , we have  $d_V(\pi_V(x), \pi_V(y)) \leq E$ . We will refer this property as BGI in this paper.

**Definition 2.5** ( $\kappa$ -consistent tuple). For  $\kappa \geq 0$ , let  $\vec{b} = (b_U)_{U \in \mathfrak{S}} \in \prod_{U \in \mathfrak{S}} 2^{\mathcal{C}U}$  be a tuple such that every set  $b_U$  has diameter at most  $\kappa$ . We say that  $\vec{b}$  is  $\kappa$ -consistent if

$$\begin{aligned} \min \{d_U(b_U, \rho_U^V), d_V(b_V, \rho_V^U)\} &\leq \kappa \quad \text{whenever } U \pitchfork V, \text{ and} \\ \min \{d_V(b_V, \rho_V^U), \text{diam}_U(b_U \cup \rho_U^V(b_V))\} &\leq \kappa \quad \text{whenever } U \sqsubset V. \end{aligned}$$

For convenience, we always take  $E$  to be the greatest constant in all coarseness from the above list (see [BHS19, Remark 1.6] for discussion on these constants). For the rest of this subsection, let  $(\mathcal{X}, \mathfrak{S})$  be a relative HHS.

**Notation.** Given  $x, y \in \mathcal{X}$ , we write  $d_U(x, y)$  to mean  $d_U(\pi_U(x), \pi_U(y))$ . If  $U \pitchfork V$  or  $U \sqsubset V$ , we write  $d_V(x, \rho_V^U)$  to mean  $d_V(\pi_V(x), \rho_V^U)$ .

**Notation.** Given two functions  $f, g : X \rightarrow \mathbb{R}$  and  $A, B > 0$ , we write  $f \preceq_{(A,B)} g$  to mean  $f(x) \leq Ag(x) + B$  for any  $x \in X$ . We write  $f \succ_{(A,B)} g$  to mean

$$\frac{1}{A}f(x) - B \leq g(x) \leq Af(x) + B$$

for any  $x \in X$ . Sometimes we omit the constants, meaning that the inequality holds for some constants.

The powerful Masur–Minsky distance formula [MM00] shows that the distance between points in a mapping class group is coarsely the sum of the distances

between the projections of these points to the curve graphs of all subsurfaces. Like mapping class groups, relative HHSes also satisfy a Masur–Minsky-style distance formula.

**Theorem 2.6** (distance formula [BHS19, Theorem 6.10]). *There exists  $s_0$  such that for all  $s \geq s_0$  there exists a constant  $C > 0$  such that, for all  $x, y \in \mathcal{X}$ ,*

$$d_{\mathcal{X}}(x, y) \asymp_{(C,C)} \sum_{W \in \mathfrak{S}} \{d_W(x, y)\}_s.$$

Closely related to the distance formula is the existence of *hierarchy paths*.

**Definition 2.7** (hierarchy path). A  $(D, D)$ -quasigeodesic  $\gamma \subset \mathcal{X}$  is a  $D$ -*hierarchy path* if  $\pi_U(\gamma)$  is an unparametrized  $(D, D)$ -quasigeodesic for each  $U \in \mathfrak{S}$ .

**Theorem 2.8** (existence of hierarchy paths [BHS19, Theorem 6.11]). *There exists  $D_0$  such that any two points in  $\mathcal{X}$  are joined by a  $D_0$ -hierarchy path.*

**Remark 2.9.** Let  $\gamma$  be a  $D$ -hierarchy path connecting  $x$  and  $y$ . By the construction of hierarchy paths in [BHS19],  $\pi_U(\gamma)$  is contained in the  $D$ -neighborhood of a geodesic connecting  $\pi_U(x)$  and  $\pi_U(y)$ . If  $CU$  is hyperbolic, this is easy to see from the Morse lemma. In the general case, it deserves its own mention.

There is an important class of subspaces in relative HHSes. We will consider them in Section 4.

**Definition 2.10** (standard product region, standard nested region, standard orthogonal region). Fix  $U \in \mathfrak{S}$  and  $\kappa \geq \kappa_0$ . Let  $F_U$  be the set of  $\kappa$ -consistent tuples in  $\prod_{V \in \mathfrak{S}_U} 2^{C^V}$ . Let  $E_U$  be the set of  $\kappa$ -consistent tuples in  $\prod_{V \in \mathfrak{S}_U^{\perp} - \{A\}} 2^{C^V}$ . Let  $P_U = F_U \times E_U$ . We can define a coarse map  $\phi_U : P_U \rightarrow \mathcal{X}$  as follows.

For each  $(\vec{a}, \vec{b}) \in F_U \times E_U$  and each  $V \in \mathfrak{S}$ , define the coordinate  $(\phi_U(\vec{a}, \vec{b}))_V$  as follows. If  $V \sqsubseteq U$ , then  $(\phi_U(\vec{a}, \vec{b}))_V = a_V$ ; if  $V \perp U$ , then  $(\phi_U(\vec{a}, \vec{b}))_V = b_V$ ; if  $V \pitchfork U$ , then  $(\phi_U(\vec{a}, \vec{b}))_V = \rho_V^U$ ; and if  $U \sqsubset V$ , then  $(\phi_U(\vec{a}, \vec{b}))_V = \rho_V^U$ . We can check that the tuple  $\phi_U(\vec{a}, \vec{b})$  is  $\kappa$ -consistent, and thus the realization theorem [BHS19, Theorem 3.1] supplies the map  $\phi_U : P_U \rightarrow \mathcal{X}$ . (See [BHS19, §5B] for details.)

For convenience, we do not distinguish between  $P_U$  and its image in  $\mathcal{X}$ . We call  $P_U$  the *standard product region*. By choosing any copy of  $F_U$  in the direct product,  $\phi_U$  restricts to a coarse map  $\phi^{\square} : F_U \rightarrow \mathcal{X}$ . We also define  $\phi^{\perp} : E_U \rightarrow \mathcal{X}$  in the same way. We call  $F_U$  and  $E_U$  the *standard nested region* and the *standard orthogonal region*, respectively.

**Remark 2.11.** By definition,  $F_U, E_U$  and  $P_U$  depend on the constant  $\kappa$ . *From now on, we fix some  $\kappa \geq \kappa_0$  and do not mention it again.*

It is known that  $(F_U, \mathfrak{S}_U)$ ,  $(E_U, \mathfrak{S}_U^\perp)$  are both relatively hierarchically hyperbolic. By definition of  $F_U$ ,  $E_U$  and  $P_U$ , there are natural retractions from  $\mathcal{X}$  to these subspaces. We call such a map a *gate map*. Take  $F_U$  for example. We denote the gate map to  $F_U$  by  $\mathfrak{g}_{F_U}$ . For all  $x \in \mathcal{X}$  and all  $V \in \mathfrak{S}$  such that  $\mathcal{C}V$  is hyperbolic,  $\pi_V(\mathfrak{g}_{F_U}(x))$  uniformly coarsely coincides with the closest point projection of  $\pi_V(x)$  to the quasiconvex subset  $\pi_V(F_U)$ . In fact, gate maps can be defined for all “hierarchically quasiconvex” subspaces, which form a larger class of subspaces of relative HHSes (see [BHS19, §5] for HHSes).

For any (relative) HHS  $(\mathcal{X}, \mathfrak{S})$ , an automorphism is roughly speaking a bijection from  $(\mathcal{X}, \mathfrak{S})$  to itself that preserves its (relative) HHS structure [BHS19, §1G]. The automorphisms of  $(\mathcal{X}, \mathfrak{S})$  form a group  $\text{Aut}(\mathfrak{S})$ , which we call the *automorphism group* of  $(\mathcal{X}, \mathfrak{S})$ .

**Definition 2.12** (hierarchically hyperbolic and relatively hierarchically hyperbolic groups). A finitely generated group  $G$  is (relatively) *hierarchically hyperbolic* if there exists a (relatively) hierarchically hyperbolic space  $(\mathcal{X}, \mathfrak{S})$  and an action  $G \rightarrow \text{Aut}(\mathfrak{S})$  such that the action  $G \curvearrowright \mathcal{X}$  is metrically proper and cobounded, and such that the induced action on  $\mathfrak{S}$  is cofinite.

Note that if  $G$  is (relatively) hierarchically hyperbolic by virtue of its action on the (relatively) hierarchically hyperbolic space  $(\mathcal{X}, \mathfrak{S})$ , then  $(G, \mathfrak{S})$  is a (relatively) hierarchically hyperbolic structure with respect to any word metric on  $G$ .

Let  $\text{Aut}(\mathfrak{S}; V)$  be the group of automorphisms  $g \in \text{Aut}(\mathfrak{S})$  such that  $g \cdot V = V$ . Then there is a *restriction homomorphism*  $\theta_V : \text{Aut}(\mathfrak{S}; V) \rightarrow \text{Aut}(\mathfrak{S}_V)$  defined as follows. Given  $g \in \text{Aut}(\mathfrak{S}; V)$ , let  $\theta_V(g)$  act like  $g$  on the substructure  $\mathfrak{S}_V$ . For a group  $G < \text{Aut}(\mathfrak{S})$ , we write  $\text{Stab}_G(V)$  to mean  $G \cap \text{Aut}(\mathfrak{S}; V)$  and write  $G_V$  to mean the image of  $\text{Stab}_G(V)$  under  $\theta_V$ .

For many HHGs (for example, the case of mapping class groups), every  $G_V$  acts acylindrically on  $\mathcal{C}V$ . However, not all HHGs have this property [DHS20].

**Definition 2.13** (colorability). Let  $(G, \mathfrak{S})$  be a relative HHG. Let  $\mathfrak{S}' \subset \mathfrak{S}$  be a  $G$ -invariant subset. We say  $\mathfrak{S}'$  is *colorable* if  $\mathfrak{S}'$  admits a decomposition  $\mathfrak{S}' = \bigsqcup_{i=1}^k \mathfrak{S}'_i$  into finitely many  $G$ -invariant families  $\mathfrak{S}'_i$  such that any two domains in the same family are transverse. Such a decomposition is called a *coloring* of  $\mathfrak{S}'$ . We say a relative HHG  $(G, \mathfrak{S})$  is *colorable* if  $\mathfrak{S}$  is colorable.

The notion of colorability is formalized in [DMS23; HP22]. There are many classes of (virtually) colorable HHGs, as listed in those papers. In particular, a coloring is constructed for (a finite-index subgroup of) a mapping class group in [BBF15, §5], from which the notion comes. However, one cannot expect that all HHGs are virtually colorable [Hag23]. Nevertheless, Proposition 3.2 of [HP22] provides a sufficient condition for an HHG to be virtually colorable.

**Remark 2.14.** In this paper, we only treat unbounded domains, i.e., domains with unbounded associated hyperbolic spaces. Abusing terminology, we say a relative HHG  $(G, \mathfrak{S})$  is colorable if the collection of unbounded domains is colorable.

### 3. Thick distance formula

In this section, we will prove a *thick distance formula* that is similar to [BBF21, Theorem 4.13]. This allows us to estimate the distance in a relative HHS by counting only “thick” segments of a hierarchy path instead of the whole hierarchy path. The reader should be aware that the definitions in this section are different from those in [BBF21]. In particular, we do not have tight geodesics in a general HHG.

Let  $(\mathcal{X}, \mathfrak{S})$  be a relative HHS and fix  $T > 100E + 10D_0$  (see Section 2.3 for constants associated with a relative HHS). As in [BHS19, §2B], we say a domain  $U \in \mathfrak{S}$  is  $T$ -relevant for  $x, y \in \mathcal{X}$  if  $d_U(x, y) > T$ . We write  $\text{Rel}_T(x, y)$  for the set of  $T$ -relevant domains for  $x, y$ , and define  $\text{Rel}_T(V; x, y) := \text{Rel}_T(x, y) \cap \mathfrak{S}_V^\circ$ . We write  $\text{Rel}_T^m(V; x, y)$  for the set of  $\sqsubseteq$ -maximal elements in  $\text{Rel}_T(V; x, y)$ . When  $x$  and  $y$  are fixed, we often omit them from the notation.

**Lemma 3.1.** *Given  $x, y \in \mathcal{X}$  and  $U \in \text{Rel}_T(x, y)$ , there exist at most two domains  $V_1, V_2 \in \text{Rel}_T(x, y)$  such that  $U \in \text{Rel}_T^m(V_i; x, y)$  for  $i = 1, 2$ .*

*Proof.* Suppose there exist three such domains  $V_1, V_2, V_3$ . Since  $U$  is maximal in each  $\text{Rel}_T(V_i; x, y)$ , we know that the  $V_i$  are not  $\sqsubseteq$ -comparable. No two of them can be orthogonal since  $U \sqsubset V_i$ . Thus, the  $V_i$  must be pairwise transverse.

By [BHS19, Proposition 2.8], any set of pairwise transverse elements in  $\text{Rel}_T(x, y)$  has a total order  $<$ , obtained by setting  $U < V$  whenever  $d_U(y, \rho_U^V) \leq E$ . We assume that  $V_1 < V_2 < V_3$ .

On the one hand,  $d_{V_2}(\rho_{V_2}^{V_1}, \rho_{V_2}^{V_3}) \geq d_{V_2}(x, y) - 2E > T - 2E$  by the triangle inequality. On the other hand,  $d_{V_2}(\rho_{V_2}^{V_1}, \rho_{V_2}^U) \leq \kappa_0$  and  $d_{V_2}(\rho_{V_2}^{V_3}, \rho_{V_2}^U) \leq \kappa_0$  by consistency, which gives  $d_{V_2}(\rho_{V_2}^{V_1}, \rho_{V_2}^{V_3}) \leq 2\kappa_0 < T - 2E$ . This gives a contradiction.  $\square$

**Definition 3.2** ( $T$ -thickness). Given  $\mathfrak{S}' \subset \mathfrak{S}$ , we say a pair of points  $(x, y) \in \mathcal{X} \times \mathcal{X}$  is  $T$ -thick for  $\mathfrak{S}'$  if  $\text{diam}(\pi_U(x) \cup \pi_U(y)) \leq T$  for all  $U \in \mathfrak{S}'$ . We define  $\mathcal{P}_T(\mathfrak{S}')$  to be the set of all  $T$ -thick pairs of points for  $\mathfrak{S}'$ . If  $\mathfrak{S}' = \{U\}$ , we also say  $(x, y)$  is  $T$ -thick for  $U$  and write  $(x, y) \in \mathcal{P}_T(U)$ .

Note that  $(x, y) \in \mathcal{P}_T(U)$  if and only if  $U \notin \text{Rel}_T(x, y)$ , and that  $(x, y) \in \mathcal{P}_T(\mathfrak{S}_V^\circ)$  if and only if  $\text{Rel}_T(V; x, y) = \emptyset$ .

**Lemma 3.3.** *Let  $D_0$  be the constant provided by Theorem 2.8. For any  $x, y \in \mathcal{X}$ , let  $\gamma$  be a  $D_0$ -hierarchy path between  $x, y$ . Given any  $U \in \mathfrak{S}$  and any  $x', y' \in \gamma$ , then*

$$d_U(x', y') \leq d_U(x, y) + 2D_0.$$

*In particular, if  $(x, y) \in \mathcal{P}_T(U)$ , then  $(x', y') \in \mathcal{P}_{T+2D_0}(U)$ .*

*Proof.* By Remark 2.9,  $\pi_U(\gamma)$  lies in the  $D_0$ -neighborhood of a geodesic connecting  $\pi_U(x)$  and  $\pi_U(y)$ . The conclusion then follows from the triangle inequality.  $\square$

**Notation.** Throughout this paper, let  $\hat{T} = T + 2D_0$  and  $\check{T} = T - 2D_0$  for every constant  $T > 2D_0$ .

**Notation.** Given two points  $x, y$  in a hyperbolic space, we write  $[x, y]$  to mean a geodesic segment between  $x, y$ , which is coarsely unique. For an interval  $I$  or a path  $\gamma$ , we write  $I^-, I^+$  or  $\gamma^-, \gamma^+$  to mean their endpoints.

**Definition 3.4** ( $(T, R)$ -thick distance). Fix sufficiently large constants  $T, R$ . Let  $\gamma$  be a  $D_0$ -hierarchy path between  $x$  and  $y$ . Let  $\gamma_1, \dots, \gamma_n \subset \gamma$  be disjoint subpaths occurring in this order such that  $(\gamma_i^-, \gamma_i^+) \in \mathcal{P}_T(\mathfrak{S}_V^\circ)$  for each  $i$ .

The  $(T, R)$ -thick distance in  $V$  is denoted by  $d_V^{T,R}(x, y)$  and is defined to be the supremum of  $\sum_{i=1}^n \{\{d_V(\gamma_i^-, \gamma_i^+)\}_R\}$  over all such choices for the  $\gamma_i$ , and for all  $D_0$ -hierarchy paths from  $x$  to  $y$ .

It is always true that  $d_V^{T,R}(x, y) \leq \{\{d_V(x, y)\}_R\}$ . This becomes an equality if  $V$  is  $\sqsubseteq$ -minimal. For the opposite direction, we have the following estimate.

**Lemma 3.5.** Fix constants  $T, R > 100E$ . For any  $x, y \in \mathcal{X}$  and  $W \in \mathfrak{S}$ , we have

$$\{\{d_W(x, y)\}_R\} \leq d_W^{T,R}(x, y) + (6E + 2R) |\text{Rel}_T^m(W; x, y)|.$$

*Proof.* If  $\text{Rel}_T^m(W) = \emptyset$ , then  $(x, y) \in \mathcal{P}_T(\mathfrak{S}_W^\circ)$ . Thus, both sides of the above inequality are equal.

Now assume that  $\text{Rel}_T^m(W) \neq \emptyset$  and  $d_W(x, y) \geq R$ . Then  $CW$  is not  $\sqsubseteq$ -minimal, so it is hyperbolic. Let  $\gamma : I \rightarrow \mathcal{X}$  be a  $D_0$ -hierarchy path realizing  $d_W^{T,R}(x, y)$ , where  $I$  is an interval of  $\mathbb{R}$ . For any  $V \sqsubset W$ , we define

$$\begin{aligned} s_V^- &:= \inf\{s \in I \mid \exists U \sqsubseteq V \text{ such that } d_W(\gamma(s), \rho_W^U) \leq 2E\}, \\ s_V^+ &:= \sup\{s \in I \mid \exists U \sqsubseteq V \text{ such that } d_W(\gamma(s), \rho_W^U) \leq 2E\}. \end{aligned}$$

For any  $U \sqsubseteq V \sqsubset W$ , we know that  $d_W(\rho_W^U, \rho_W^V) \leq \kappa_0$  by consistency. Thus,

$$d_W(\gamma(s_V^-), \rho_W^V) \leq 2E + \kappa_0 \leq 3E \quad \text{and} \quad d_W(\gamma(s_V^+), \rho_W^V) \leq 2E + \kappa_0 \leq 3E.$$

Therefore,

$$d_W(\gamma(s_V^-), \gamma(s_V^+)) \leq d_W(\gamma(s_V^-), \rho_W^V) + d_W(\gamma(s_V^+), \rho_W^V) \leq 6E.$$

Let  $J_0, \dots, J_n$  be the collection of maximal intervals in  $I - \bigcup_{V \in \text{Rel}_T^m(W)} (s_V^-, s_V^+)$ . Note that  $n \leq |\text{Rel}_T^m(W)|$ . We now prove that  $(\gamma(J_i^-), \gamma(J_i^+)) \in \mathcal{P}_T(\mathfrak{S}_W^\circ)$ .

On the one hand,  $d_W(\rho_W^U, \gamma(J_i)) \geq 2E$  for any  $U \in \text{Rel}_T^m(W)$  by the definition of  $J_i$ . By the Morse lemma,

$$d_W(\rho_W^U, [\gamma(J_i^-), \gamma(J_i^+)]) \geq d_W(\rho_W^U, \gamma(J_i)) - E \geq E.$$

Therefore,  $d_U(\gamma(J_i^-), \gamma(J_i^+)) \leq E < T$  by BGI. On the other hand,  $(x, y)$  is  $\check{T}$ -thick for  $\mathfrak{S}_W^\circ - \text{Rel}_{\check{T}}(W)$  by definition. It follows from Lemma 3.3 that  $(\gamma(J_i^-), \gamma(J_i^+))$  is  $T$ -thick for  $\mathfrak{S}_W^\circ - \text{Rel}_{\check{T}}(W)$ . In sum,  $(\gamma(J_i^-), \gamma(J_i^+)) \in \mathcal{P}_T(\mathfrak{S}_W^\circ)$ .

Finally, we estimate

$$\begin{aligned} d_W(x, y) &\leq \sum_{i=0}^n d_W(\gamma(J_i^+), \gamma(J_i^-)) + 6E |\text{Rel}_{\check{T}}^m(W)| \\ &\leq d_W^{T,R}(x, y) + R(n+1) + 6E |\text{Rel}_{\check{T}}^m(W)| \\ &\leq d_W^{T,R}(x, y) + (6E + 2R) |\text{Rel}_{\check{T}}^m(W)|. \quad \square \end{aligned}$$

Let  $S$  denote the unique maximal domain in  $\mathfrak{S}$ . Recall that the level  $\ell(S)$  of  $S$  is equal to the complexity of  $(\mathcal{X}, \mathfrak{S})$ .

**Theorem 3.6.** *Fix constants  $T, R$  with  $\check{T} \geq R > 100E$ . Let  $x, y \in \mathcal{X}$ . Then, for each  $n$ ,*

$$\sum_{\ell(W) \leq n} \{\{d_W(x, y)\}\}_R \leq \sum_{\ell(W)=n} d_W^{T,R}(x, y) + 7 \sum_{\ell(W) < n} \{\{d_W(x, y)\}\}_R.$$

Note that each sum is has finitely many terms since there are only finitely many  $W$  such that  $d_W(x, y) \geq R$  for given  $x, y$ , by the distance formula (Theorem 2.6).

*Proof.* If  $\ell(W) = n$  then by Lemma 3.5,

$$\begin{aligned} \{\{d_W(x, y)\}\}_R &\leq d_W^{T,R}(x, y) + (6E + 2R) |\text{Rel}_{\check{T}}^m(W)| \\ &\leq d_W^{T,R}(x, y) + (6E + 2R) \sum_{V \in \text{Rel}_{\check{T}}^m(W)} \frac{\{\{d_V(x, y)\}\}_{\check{T}}}{\check{T}} \\ &\leq d_W^{T,R}(x, y) + 3 \sum_{V \in \text{Rel}_{\check{T}}^m(W)} \{\{d_V(x, y)\}\}_{\check{T}} \\ &\leq d_W^{T,R}(x, y) + 3 \sum_{V \in \text{Rel}_{\check{T}}^m(W)} \{\{d_V(x, y)\}\}_R. \end{aligned}$$

By Lemma 3.1, any  $V$  appears in at most two  $\text{Rel}_{\check{T}}^m(W)$ . Therefore, if we sum up the left side over all  $W$  with  $\ell(W) = n$ , we have

$$\sum_{\ell(W)=n} \{\{d_W(x, y)\}\}_R \leq \sum_{\ell(W)=n} d_W^{T,R}(x, y) + 6 \sum_{\ell(W) < n} \{\{d_W(x, y)\}\}_R.$$

Adding  $\sum_{\ell(W) < n} \{\{d_W(x, y)\}\}_R$  to both sides gives the desired inequality. □

**Corollary 3.7.** *Fix constants  $T, R$  with  $\check{T} \geq R > 100E$ . Let  $x, y \in \mathcal{X}$ . Then*

$$\frac{1}{D_0} \sum_{W \in \mathfrak{S}} d_W^{T,R}(x, y) - D_0 \leq \sum_{W \in \mathfrak{S}} \{\{d_W(x, y)\}\}_R \leq 7^{\ell(S)-1} \sum_{W \in \mathfrak{S}} d_W^{T,R}(x, y).$$

*Proof.* The first inequality is trivial since  $(1/D_0)d_W^{T,R}(x, y) - D_0 \leq \{\!\!\{d_W(x, y)\}\!\!\}_R$  for all  $W$ . By inductively applying Theorem 3.6, with base case  $n = \ell(S)$ , we have

$$\sum_{W \in \mathfrak{S}} \{\!\!\{d_W(x, y)\}\!\!\}_R \leq 7^{\ell(S)-n} \left( \sum_{n \leq \ell(W) \leq \ell(S)} d_W^{T,R}(x, y) + 7 \sum_{\ell(W) < n} \{\!\!\{d_W(x, y)\}\!\!\}_R \right).$$

When  $n = 1$ , the last term on the right is zero, and the result follows. □

Combining the distance formula (Theorem 2.6) with Corollary 3.7, we obtain:

**Theorem 3.8** (thick distance formula). *There exists  $R_0$  such that, for all  $T, R$  with  $\check{T} \geq R > R_0$ , there exists a constant  $L > 0$  such that, for all  $x, y \in \mathcal{X}$ ,*

$$d_{\mathcal{X}}(x, y) \asymp_{(L,L)} \sum_{W \in \mathfrak{S}} d_W^{T,R}(x, y).$$

#### 4. Estimation of thick distance via quasi-axes

Our proof in this section is inspired by [NY23]. The main technique in the proof of [NY23, Lemma 5.5] that is different from [BBF21] is the use of the extension lemma [Yan19, Lemma 2.14]. Lemma 4.1 below is a trimmed version for acylindrical actions on hyperbolic spaces. In the original statement of Lemma 4.1 in [NY23], the group action is required to be cobounded, but it is easy to see from the proof that this condition can be removed.

**Lemma 4.1** (extension lemma [NY23, Lemma 4.13]). *Let  $H$  be a group acting nonelementarily and acylindrically on a  $\delta$ -hyperbolic space  $Y$ . Fix a base point  $o \in Y$ . There exists a set  $F \subset H$  of three loxodromic elements and constants  $\lambda \geq 1$ , and  $c \geq 0$  with the following property.*

*For any  $h \in H$  there exists  $f \in F$  such that  $hf$  is a loxodromic element and the bi-infinite path*

$$\gamma_h = \bigcup_{i \in \mathbb{Z}} (hf)^i ([o, ho][ho, hfo])$$

*is a  $(\lambda, c)$ -quasigeodesic.*

Let  $(\mathcal{X}, \mathfrak{S})$  be a relative HHS with a coarse constant  $E$ , and let  $G$  be a relative HHG by virtue of its action on  $\mathcal{X}$ . Corresponding to Definition 1.3, we say that a domain  $U \in \mathfrak{S}$  has

- (1) *hyperbolicity* if  $\mathcal{C}U$  is hyperbolic.
- (2) *acylindrical image* if  $G_U$  acts on  $\mathcal{C}U$  acylindrically.
- (3) *cobounded nested region* if  $G_U$  acts on  $F_U$  coboundedly.
- (4) *separable quasi-axes* if for any element  $g \in \text{Stab}_G(U)$  that acts loxodromically on  $\mathcal{C}U$ , the elementary closure  $EC(g)$  is *separable* in  $G$ , i.e.,  $EC(g)$  equals the intersection of all finite-index subgroups of  $G$  that contain  $EC(g)$ .

Till the end of this section, let  $V \in \mathfrak{S}$  be an unbounded domain that has hyperbolicity, cobounded nested region and acylindrical image.

The next statement could be compared with Theorem 4.19 of [BBF21] for mapping class groups.

**Lemma 4.2** (extension of thick segments). *There exist constants  $\lambda \geq 1$ ,  $c \geq 0$ ,  $B \geq 0$  such that the following holds. For any  $T, R > 0$ , there exists a  $G_V$ -finite collection  $\mathcal{A}_V = \mathcal{A}_V^{T,R}$  of  $(\lambda, c)$ -quasi-axes in  $\mathcal{CV}$  such that for any pair of points  $(x, y) \in \mathcal{P}_{\max\{\hat{T}, R\}}(\mathfrak{S}_V)$  there exists  $\gamma \in \mathcal{A}_V$  such that  $[\pi_V(x), \pi_V(y)] \subset \mathcal{N}_B(\gamma)$ .*

*Proof.* Fix a base point  $o \in F_V$  and project it to a base point in  $\mathcal{CV}$ . If the action  $G_V \curvearrowright \mathcal{CV}$  is elementary,  $\mathcal{CV}$  itself is a quasi-axis of a loxodromic element, and thus satisfies the requirement. Now assume that the action  $G_V \curvearrowright \mathcal{CV}$  is nonelementary. Lemma 4.1 provides a finite set  $F \subset G_V$  and constants  $\lambda \geq 1$ ,  $c \geq 0$ . Recall that  $(F_V, \mathfrak{S}_V)$  is a relative HHS.  $F_V$  is proper because  $\mathcal{X}$  is proper.

Since  $G_V$  acts coboundedly on  $F_V$ , there exists  $\epsilon > 0$  such that  $F_V$  is covered by the  $G_V$ -translates of any  $\epsilon$ -ball. Let  $T' = \max\{\hat{T}, R\}$ . By Theorem 2.6, there exists  $r > 0$ , depending only on  $E, T', \epsilon$ , such that the distance between any pair of points in  $F_V$  that is  $(T' + \epsilon)$ -thick for  $\mathfrak{S}_V$  is bounded above by  $r$ . Fix any base point  $o \in F_V$ . Since  $F_V$  is proper, there exists a finite subset  $S \subset G_V$  such that  $\mathcal{N}_{r+\epsilon}(o)$  is covered by  $\bigcup_{s \in S} s \cdot \mathcal{N}_\epsilon(o)$ .

Lemma 4.1 tells us that for each  $s \in S$ , there exists  $f \in F$  such that  $sf$  is a loxodromic element acting on  $\mathcal{CV}$ . Let  $\mathcal{A}_V = \mathcal{A}_V^{T,R}$  be the collection of  $G_V$ -translates of the  $(\lambda, c)$ -quasi-axes provided by Lemma 4.1 for all loxodromic elements of the form  $sf$ .

Now we verify that  $\mathcal{A}_V$  meets our requirements. Let  $(x, y) \in \mathcal{P}_{T'}(\mathfrak{S}_V)$ . We can choose  $g \in G_V$  such that  $d_{F_V}(x, go) < \epsilon$ . Then

$$d_{F_V}(o, g^{-1}y) \leq d_{F_V}(x, y) + d_{F_V}(x, go) < r + \epsilon.$$

By our choice of  $S$ , there exists  $s \in S$  such that  $d_{F_V}(g^{-1}y, so) < \epsilon$ . Thus,

$$d_V(x, go) < E\epsilon + E, \quad d_V(y, gso) < E\epsilon + E,$$

because  $\pi_V$  is  $E$ -coarsely Lipschitz. Since  $\mathcal{CV}$  is  $E$ -hyperbolic, we can find  $B > 0$  by the fellow-traveler property such that  $[\pi_V(x), \pi_V(y)] \subset \mathcal{N}_B([g \cdot \pi_V(o), gs \cdot \pi_V(o)])$ . By construction,  $[g \cdot \pi_V(o), gs \cdot \pi_V(o)]$  is contained in some  $\gamma \in \mathcal{A}_V$ , so we are done. □

**Notation.** Assume that  $\gamma$  is a quasigeodesic in a hyperbolic space  $Y$ . We write  $\pi_\gamma : Y \rightarrow \gamma$  to mean the closest point projection. For  $x, y \in Y$ , we write  $d_\gamma(x, y)$  to mean  $\text{diam}(\pi_\gamma(x) \cup \pi_\gamma(y))$ .

**Notation.** Assume that  $\gamma$  is a quasigeodesic in  $\mathcal{CV}$ . We write  $\pi_\gamma^\mathcal{X}$  to mean  $\pi_\gamma \circ \pi_V$ . For  $x, y \in \mathcal{X}$ , we write  $d_\gamma^\mathcal{X}(x, y)$  to mean  $\text{diam}(\pi_\gamma^\mathcal{X}(x) \cup \pi_\gamma^\mathcal{X}(y))$ .

The following lemma is a well-known corollary of the Morse lemma for quasi-geodesics in  $\delta$ -hyperbolic spaces so we omit the proof.

**Lemma 4.3.** *Let  $\gamma$  and  $\alpha$  be two  $(\lambda, c)$ -quasigeodesics in a  $\delta$ -hyperbolic space. Then for any  $B > 0$ , there exists a constant  $C = C(\lambda, c, B, \delta) > 0$  such that*

$$d_\gamma(\alpha^-, \alpha^+) \geq \text{diam}(\alpha \cap \mathcal{N}_B(\gamma)) - C.$$

The main result of this section is the following estimate, which generalizes Proposition 4.18 of [BBF21].

**Proposition 4.4.** *For any  $K > 0$ , there exists  $R > 0$  such that the following holds. Given any  $T > 0$ , let  $\mathcal{A}_V = \mathcal{A}_V^{T,R}$  be the collection of  $(\lambda, c)$ -quasi-axes provided by Lemma 4.2. Then for any two points  $x, y \in \mathcal{X}$ ,*

$$d_V^{T,R}(x, y) \leq 2(D_0 + 1) \sum_{\gamma \in \mathcal{A}_V} \{\{d_\gamma^{\mathcal{X}}(x, y)\}\}_K,$$

where  $D_0$  is the constant provided by Theorem 2.8.

*Proof.* Let  $C = C(\lambda + D_0, c + D_0, B, E)$  be the constant provided by Lemma 4.3. Let  $R > 2D_0(C + 1) + K$ . We will show that projections to quasi-axes  $\mathcal{A}_V = \mathcal{A}_V^{T,R}$  bound the  $(T, R)$ -thick distance in  $V$  from above.

For any two points  $x, y \in \mathcal{X}$ , let  $\beta$  be a  $(D_0, D_0)$ -hierarchy path connecting  $x$  and  $y$  realizing  $d_V^{T,R}(x, y)$ . Let  $\{\alpha_1, \dots, \alpha_n\}$  be the collection of disjoint subpaths of  $\beta$  with  $d_V(\alpha_i^-, \alpha_i^+) \geq R$  and  $(\alpha_i^-, \alpha_i^+) \in \mathcal{P}_T(\mathfrak{S}_V^\circ)$  such that

$$d_V^{T,R}(x, y) = \sum_{i=1}^n d_V(\alpha_i^-, \alpha_i^+).$$

By the definition of gate maps,  $\pi_V(z)$  is coarsely  $\pi_V(\mathfrak{g}_{F_V}(z))$  for any  $z \in \mathcal{X}$ . Thus, the difference between  $d_V(\alpha_i^-, \alpha_i^+)$  and  $d_V(\mathfrak{g}_{F_V}(\alpha_i^-), \mathfrak{g}_{F_V}(\alpha_i^+))$  is uniformly bounded. This enables us to replace  $\alpha_i$  with  $\mathfrak{g}_{F_V}(\alpha_i)$  from now on. We divide each  $\alpha_i$  into several consecutive subpaths  $\{\tilde{\alpha}_{i,j} \mid 1 \leq j \leq m_i\}$  with  $d_V(\tilde{\alpha}_{i,j}^-, \tilde{\alpha}_{i,j}^+) = R$  for  $j = 1, \dots, m_i - 1$  and  $d_V(\tilde{\alpha}_{i,m_i}^-, \tilde{\alpha}_{i,m_i}^+) \leq R$ . By Lemma 3.3, we already know that  $(\tilde{\alpha}_{i,j}^-, \tilde{\alpha}_{i,j}^+) \in \mathcal{P}_{\hat{T}}(\mathfrak{S}_V^\circ)$  for every pair  $(i, j)$ . Thus,

$$d_U(\tilde{\alpha}_{i,j}^-, \tilde{\alpha}_{i,j}^+) \leq \max\{\hat{T}, R\}$$

for all  $U \in \mathfrak{S}_V$ . By Lemma 4.2, there exists  $\gamma_{i,j} \in \mathcal{A}_V$  such that  $\pi_U(\tilde{\alpha}_{i,j}) \subset \mathcal{N}_B(\gamma_{i,j})$  (with an increased  $B$  by a uniform constant), which yields

$$\text{diam}(\pi_V(\alpha_i) \cap \mathcal{N}_B(\gamma_{i,j})) \geq R.$$

Let  $\mathcal{A}'_V$  be the collection of all distinct  $\gamma_{i,j}$ . We see that

$$\pi_V(\alpha_i \setminus \tilde{\alpha}_{i,m_i}) \subset \bigcup_{\gamma \in \mathcal{A}'_V} \pi_V(\alpha_i) \cap \mathcal{N}_B(\gamma).$$

Thus, we have

$$d_V(\alpha_i^-, \alpha_i^+) \leq \sum_{\gamma \in \mathcal{A}'_V} \text{diam}(\pi_V(\alpha_i) \cap \mathcal{N}_B(\gamma)) + R \leq 2 \sum_{\gamma \in \mathcal{A}'_V} \text{diam}(\pi_V(\alpha_i) \cap \mathcal{N}_B(\gamma)).$$

Summing up from  $i = 1$  to  $n$  yields

$$(1) \quad d_V^{T,R}(x, y) \leq 2 \sum_{\gamma \in \mathcal{A}'_V} (D_0 \text{diam}(\pi_V(\beta) \cap \mathcal{N}_B(\gamma)) + D_0).$$

Note that  $R > 2D_0(C + 1) + K$ . Thus, Lemma 4.3 tells us that

$$d_\gamma^{\mathcal{X}}(x, y) \geq \text{diam}(\pi_V(\beta) \cap \mathcal{N}_B(\gamma)) - C \geq R - C > D_0(C + 1) + K$$

for each  $\gamma \in \mathcal{A}'_V$ . We now estimate by Lemma 4.3 that

$$D_0 \text{diam}(\pi_V(\beta) \cap \mathcal{N}_B(\gamma)) + D_0 \leq D_0(d_\gamma^{\mathcal{X}}(x, y) + C) + D_0 < (D_0 + 1)d_\gamma^{\mathcal{X}}(x, y).$$

Combining this with (1), we obtain

$$\begin{aligned} d_V^{T,R}(x, y) &\leq 2(D_0 + 1) \sum_{\gamma \in \mathcal{A}'_V} d_\gamma^{\mathcal{X}}(x, y) \\ &= 2(D_0 + 1) \sum_{\gamma \in \mathcal{A}'_V} \llbracket d_\gamma^{\mathcal{X}}(x, y) \rrbracket_K \leq 2(D_0 + 1) \sum_{\gamma \in \mathcal{A}_V} \llbracket d_\gamma^{\mathcal{X}}(x, y) \rrbracket_K. \quad \square \end{aligned}$$

### 5. Construction of quasitrees

This section is devoted to the proof of Theorem 1.4. Let  $(G, \mathfrak{S})$  be a relative HHG that is virtually colorable and assume that every domain in  $\mathfrak{S}$  is of type I or II. The index set  $\mathfrak{S}$  admits a  $G$ -invariant decomposition  $\mathfrak{S} = \mathfrak{S}^I \sqcup \mathfrak{S}^{II}$ , where  $\mathfrak{S}^I$  (respectively,  $\mathfrak{S}^{II}$ ) only contains domains of type I (respectively, type II). Note that types I and II are not mutually exclusive, but for those domains of both types, we can simply put them in  $\mathfrak{S}^{II}$ .

Before starting the proof, we summarize the dependencies of some important constants that will be used in the proof:

$$(E, D_0, A) \xrightarrow{\text{Corollary 5.3}} \theta \xrightarrow{\text{Theorem 2.4}} \xi \xrightarrow{\text{Lemma 5.4}} K \xrightarrow{\text{Proposition 4.4}} R \xrightarrow{\text{Theorem 3.8}} T.$$

Here  $A$  stands for the acylindrical constants. We draw an arrow from a constant  $M$  to  $N$  if  $N$  depends on  $M$ . Remember that the dependency graph shown above is incomplete, but we hope it is helpful to the reader.

#### 5.1. Quasitrees from domains of type I.

**Proposition 5.1.** *There exists a finite-index subgroup  $H < G$  satisfying the following. For any sufficiently large constant  $R$  and any  $T > 0$ , there exist quasitrees  $\mathcal{T}_1, \dots, \mathcal{T}_n$*

such that  $H$  acts on  $\prod_{j=1}^n \mathcal{T}_j$  diagonally and for any choice of base points  $o_j \in \mathcal{T}_j$  we have

$$\sum_{V \in \mathfrak{S}^I} d_V^{T,R}(1, h) \leq \sum_{j=1}^n d_{\mathcal{T}_j}(o_j, ho_j)$$

for any  $h \in H$ .

Before proving this, we recall an auxiliary result.

**Proposition 5.2** [BBF21, Proposition 3.4]. *Let a group  $H$  act on a  $\delta$ -hyperbolic space  $Y$ . Assume that the image of  $H$  in  $\text{Isom}(Y)$  is acylindrical. Consider a loxodromic element  $g \in H$  and the collection  $\mathbb{A}$  of all  $H$ -translates of a fixed  $(\lambda, c)$ -quasi-axis of  $g$ . Then there exists a constant  $\theta > 0$  depending only on  $\lambda, c, \delta$  and the acylindrical constants such that for any  $\gamma \in \mathbb{A}$ , the set*

$$\{h \in H \mid \text{diam}(\pi_\gamma(h\gamma)) \geq \theta\}$$

is a finite union of double  $EC(g)$ -cosets.

**Corollary 5.3.** *Let  $U \in \mathfrak{S}^I$ . Consider a  $(\lambda, c)$ -quasi-axis  $\gamma \subset \mathcal{C}U$  for some loxodromic element of the acylindrical action of  $G_U$ . Then there exists  $\theta > 0$ , only depending on  $\lambda, c, E$  and the acylindrical constants, and a finite-index subgroup  $G_\gamma < G$  such that every translate of  $\gamma$  by an element of  $G_\gamma \cap \text{Stab}_G(U)$  either has finite Hausdorff distance with  $\gamma$  or has  $\theta$ -bounded projection to  $\gamma$ .*

*Proof.* This is clear by Proposition 5.2 and separability of quasi-axes. □

By definition of relative HHGs,  $\mathfrak{S}^I$  consists of finitely many  $G$ -orbits, so acylindrical constants for  $U \in \mathfrak{S}^I$  can be chosen uniformly. Thus, Lemma 4.2 provides uniform constants  $\lambda \geq 1, c \geq 0$  for every  $U \in \mathfrak{S}^I$ . This further gives a uniform constant  $\theta > 0$  by Corollary 5.3.

Let  $\mathcal{U}$  be a  $G$ -representative set of  $\mathfrak{S}^I$  such that  $1 \in P_U$  for any  $U \in \mathcal{U}$ . Let  $U \in \mathcal{U}$ . Let  $T > 0$  and let  $K > 0$  be a sufficiently large constant that will be decided by Lemma 5.4. Lemma 4.2 provides a  $G_U$ -finite collection  $\mathcal{A}_U = \mathcal{A}_U^{T,R}$  of  $(\lambda, c)$ -quasi-axes, where  $R$  is provided by Proposition 4.4. By Corollary 5.3, we can find a finite-index subgroup  $H_U < G$  such that for any  $\gamma \in \mathcal{A}_U$  and  $h \in H_U \cap \text{Stab}_G(U)$ , either  $d_{\text{Haus}}(h\gamma, \gamma) < \infty$  or  $\text{diam } \pi_\gamma(h\gamma) < \theta$ . For any  $g \in G$ , we define  $\mathcal{A}_{gU} := \{g\gamma \mid \gamma \in \mathcal{A}_U\}$ .

Let  $\mathcal{A} := \bigsqcup_{U \in \mathfrak{S}} \mathcal{A}_U$  and  $H := \bigcap_{U \in \mathcal{U}} H_U$ . Since  $\mathcal{U}$  is finite,  $H$  is of finite index in  $G$ . By adding finitely many domains to  $\mathcal{U}$  so that there is one representative for each  $H$ -orbit on  $\mathfrak{S}^I$ , we obtain an  $H$ -representative set  $\tilde{\mathcal{U}}$  of  $\mathfrak{S}^I$ . We still assume that  $1 \in P_U$  for any  $U \in \tilde{\mathcal{U}}$ . Let  $\{\gamma_1, \dots, \gamma_n\}$  be an  $H$ -representative set of  $\mathcal{A}$ . We assume that every representative  $\gamma_j$  is contained in  $\mathcal{C}U$  for some  $U \in \tilde{\mathcal{U}}$ . Let  $\mathcal{A}_j \subset \mathcal{A}$  be the  $H$ -orbit of  $\gamma_j$ .

Without loss of generality, we assume that  $H$  is colorable instead of virtually colorable. Thus, the  $H$ -orbit of any domain is pairwise transverse. By [HP22, Lemma 3.4], every  $H$ -orbit of  $\mathfrak{S}^I$  is an  $H$ -projection system with constant  $s_0 + 4E$ , where  $s_0$  is the constant provided by Theorem 2.6. Thus, every  $\mathcal{A}_j$  is an  $H$ -projection system with a uniform projection constant  $\xi = \xi(\theta, s_0, \lambda, c, E)$  by Theorem 2.4. The projections defined there will be denoted by  $\Pi_\gamma$ .

Using Theorem 2.1, we obtain modified projections  $\Pi'_\gamma$  such that  $(\mathcal{A}_j, \{\Pi'_\gamma\})$  satisfies the strong projection axioms with constant  $\xi' = \xi'(\xi)$  and that  $\Pi_\gamma(\alpha)$  and  $\Pi'_\gamma(\alpha)$  are apart from each other within a uniform Hausdorff distance  $\epsilon = \epsilon(\xi)$ . For any  $K' \geq 4\xi'$ ,  $\mathcal{C}_{K'}\mathcal{A}_j$  is a quasitree by Theorem 2.2. The following lemma is an estimate via the orbit map between the projections  $\pi_\gamma^H = \pi_\gamma \circ \pi_V$  in the relative HHG structure and the projections  $\Pi'_\gamma$  in the quasitree  $\mathcal{C}_{K'}\mathcal{A}_j$ . Recall from Section 4 that  $d_\gamma^H(g, h)$  is defined as  $\text{diam}(\pi_\gamma^H(g) \cup \pi_\gamma^H(h))$  for any  $g, h \in H$ .

**Lemma 5.4.** *Fix a base point  $o_j \in \gamma_j$  for each  $j = 1, \dots, n$ . There exists a sufficiently large constant  $K' = K'(\xi, \lambda, c, E)$  and a constant  $\Delta > 0$  such that if  $K \geq 2K'$  then*

$$\sum_{\gamma \in \mathcal{A}_j} \{\{d_\gamma^H(1, h)\}\}_K \leq 8d_{\mathcal{C}_{K'}\mathcal{A}_j}(o_j, ho_j) + \Delta$$

for any  $h \in H$  and any  $j = 1, \dots, n$ .

The proof depends on another lemma.

**Lemma 5.5.** *For any constants  $A, B \geq 0$  and constants  $L, M > 0$ ,*

$$\frac{\{\{A + B\}\}_{L+M}}{L + M} \leq \frac{\{\{A\}\}_L}{L} + \frac{\{\{B\}\}_M}{M}.$$

*Proof.* Assume that  $A + B \geq L + M$ . First, if  $A < L$  then  $B > M$ . Thus,  $\{\{B\}\}_M = B \geq \frac{M}{L+M}(A + B)$ . Next, if  $B < M$  then  $A > L$  and the same argument holds. Finally, if  $A \geq L$  and  $B \geq M$ , then all thresholds are reached and the inequality is obviously true.  $\square$

*Proof of Lemma 5.4.* For simplicity, we use  $|p - q|$  to mean the distance between two points  $p, q$  in the same space. Assume that  $\gamma_j \subset \mathcal{C}U$  for some  $U \in \tilde{\mathcal{U}}$ . For any  $g \in H - \text{Stab}_H(U)$ , since  $1 \in \mathbf{P}_U$  and  $U \cap gU$ , we have  $|\pi_{gU}(1) - \rho_{gU}^U| \leq E$  by the definition of standard product regions.

Assume  $\gamma = g\gamma_j$ . By hyperbolicity, there exists a constant  $F = F(E, \lambda, c)$  such that if  $\gamma \subset \mathcal{C}gU \neq \mathcal{C}U$  then  $|\pi_\gamma^H(1) - \Pi'_\gamma(o_j)| \leq |\pi_{gU}(1) - \rho_{gU}^U| + F + \epsilon \leq E + F + \epsilon$ . Let  $M > \xi' + E + F + \epsilon$  and define  $\delta_\gamma(h) = |\pi_\gamma^H(h) - \Pi'_\gamma(ho_j)|$ . We see that if  $\delta_\gamma(h) = \delta_{h^{-1}\gamma}(1) \geq M$  then  $\gamma \subset ChU$ . Thus, for a fixed  $h \in H$ , there are only finitely many  $\gamma \in \mathcal{A}_j$  such that  $\delta_\gamma(h) \geq M$  by projection axiom (P2).

Let  $K' > 2M + 4\xi'$ . Define  $D_\gamma(1, h) = |\Pi'_\gamma(o_j) - \Pi'_\gamma(ho_j)|$ . By the triangle inequality and Lemma 5.5, we obtain

$$\{\{d_\gamma^H(1, h)\}\}_{K'+2M} \leq \frac{K'+2M}{K'} \{\{D_\gamma(1, h)\}\}_{K'} + \frac{K'+2M}{2M} \{\{\delta_\gamma(1) + \delta_\gamma(h)\}\}_{2M}.$$

Therefore,

$$\{\{d_\gamma^H(1, h)\}\}_K \leq 2\{\{D_\gamma(1, h)\}\}_{K'} + \frac{K'+2M}{M} \{\{\delta_\gamma(1)\}\}_M.$$

Summing over  $\gamma \in \mathcal{A}_j$ , we obtain

$$\sum_{\gamma \in \mathcal{A}_j} \{\{d_\gamma^H(1, h)\}\}_K \leq 2 \sum_{\gamma \in \mathcal{A}_j} \{\{D_\gamma(1, h)\}\}_{K'} + \frac{K'+2M}{M} \sum_{\gamma \in \mathcal{A}_j} \{\{\delta_\gamma(1)\}\}_M.$$

By the preceding discussion,

$$\Delta_j = \frac{K'+2M}{M} \sum_{\gamma \in \mathcal{A}_j} \{\{\delta_\gamma(1)\}\}_M$$

is a finite constant. Let  $\Delta = \max_{1 \leq j \leq n} \Delta_j$ . We conclude using Theorem 2.3.  $\square$

*Proof of Proposition 5.1.* Let  $K$  and  $K'$  be the constants provided by Lemma 5.4. Let  $T > 0$ . From the discussion before Lemma 5.4, we know that the collection  $\mathcal{A}$  of quasi-axes provided by Proposition 4.4 forms  $n$  quasitrees  $\mathcal{C}_{K'}\mathcal{A}_j$ ,  $j = 1, \dots, n$ . Moreover, Proposition 4.4 tells us that there exists  $R > 0$  such that

$$\sum_{V \in \mathfrak{S}'} d_V^{T,R}(1, h) \leq \sum_{\gamma \in \mathcal{A}} \{\{d_\gamma^H(1, h)\}\}_K.$$

Finally, we conclude by Lemma 5.4.  $\square$

**5.2. Quasitrees from domains of type II.** We now prove an analogue of Proposition 5.1 for domains of type II.

Let  $H < G$  be the subgroup provided by Proposition 5.1. Fix an  $H$ -representative set  $\mathcal{V}$  of  $\mathfrak{S}^H$  such that  $1 \in \mathbf{P}_V$  for any  $V \in \mathcal{V}$ . For  $V \in \mathcal{V}$  we write  $[V]$  to mean its  $H$ -orbit. By colorability and [HP22, Lemma 3.4], every  $[V]$  is an  $H$ -projection system with constant  $s_0 + 4E$ .

Fix any  $V \in \mathcal{V}$ . By property QT<sub>0</sub> of the action  $\text{Stab}_G(V) \curvearrowright \mathcal{C}V$ , there exist quasitrees  $T_V^i$  along with  $\text{Stab}_H(V)$ -equivariant maps  $\iota_V^i : \mathcal{C}V \rightarrow T_V^i$  for  $i = 1, \dots, n_V$  such that

$$\prod_{i=1}^{n_V} \iota_V^i : \mathcal{C}V \rightarrow \prod_{i=1}^{n_V} T_V^i$$

is a  $(\lambda', c')$ -quasi-isometric embedding. Thus  $\iota_V^i$  is  $(\lambda', c')$ -coarsely Lipschitz. Fix  $i \in \{1, \dots, n_V\}$ . It is conventional to extend the map  $\iota_V^i$  on  $[V]$  in an  $H$ -equivariant way. This means that we can construct a collection of quasitrees  $\mathbf{T}_{[V]}^i = \{T_U^i \mid U \in [V]\}$

with an  $H$ -action and a collection of coarsely Lipschitz maps  $\{t_U^i : \mathcal{C}U \rightarrow T_U^i \mid U \in [V]\}$  such that the following diagram commutes for any  $h \in H$  and  $U \in [V]$ :

$$\begin{array}{ccc} \mathcal{C}U & \xrightarrow{h} & ChU \\ t_U^i \downarrow & & \downarrow t_{hU}^i \\ T_U^i & \xrightarrow{h} & T_{hU}^i \end{array}$$

Define  $\Pi_{T_{hU}^i}(T_U^i) := t_{hU}^i(\rho_{hU}^U)$  for any  $U \in [V]$  and  $hU \neq U$ . Clearly, these projections are  $H$ -equivariant and the projection axioms pass to the collection  $(T_{[V]}^i, \{\Pi_{T_U^i}\})$  under coarsely Lipschitz maps  $\{t_U^i\}$ . We modify the projections within an error  $\epsilon$  such that  $(T_{[V]}^i, \{\Pi'_{T_U^i}\})$  satisfies the strong projection axioms with constant  $\zeta = \zeta(s_0, \lambda', c', E)$ . For any  $K'' \geq 4\zeta$ ,  $\mathcal{C}_{K''} T_{[V]}^i$  is a quasitree by Theorem 2.2. Define  $d_{T_U^i}(1, h) := |t_U^i(\pi_U(1)) - t_U^i(\pi_U(h))|$  for any  $U \in [V]$ . For any  $V \in \mathcal{V}$  and  $i \in \{1, \dots, n_V\}$ , fix a base point  $o_V^i \in T_V^i$ . The following proposition is an analogue of Proposition 5.1.

**Proposition 5.6.** *There is a constant  $K'' = K''(\zeta, \lambda', c', E)$  such that  $R \geq 2K''$  implies*

$$\sum_{U \in \mathfrak{S}^H} \{d_U(1, h)\}_R \leq \sum_{V \in \mathcal{V}} \sum_{i=1}^{n_V} d_{\mathcal{C}_{K''} T_{[V]}^i}(o_V^i, ho_V^i),$$

for any  $h \in H$ .

*Proof.* Fix  $V \in \mathcal{V}$  and  $i \in \{1, \dots, n_V\}$ . For any  $U \in [V] - \{V\}$ , we have

$$|t_U^i(\pi_U(1)) - \Pi'_{T_U^i}(o_V^i)| \leq \lambda' |\pi_U(1) - \rho_U^V| + c' + \epsilon \leq \lambda' E + c' + \epsilon.$$

Let  $K'' > 4\zeta + 2(\lambda' E + c' + \epsilon)$ . As in the proof of Lemma 5.4, we estimate that

$$\sum_{U \in [V]} \{d_{T_U^i}(1, h)\}_R \leq \{|\Pi'_{T_U^i}(o_i) - \Pi'_{T_U^i}(ho_i)|\}_{K''} \leq d_{\mathcal{C}_{K''} T_{[V]}^i}(o_V^i, ho_V^i).$$

Here the first inequality follows from the triangle inequality and Lemma 5.5, and the second holds by Theorem 2.3. Since the map  $\prod_{i=1}^{n_U} t_U^i : \mathcal{C}U \rightarrow \prod_{i=1}^{n_U} T_U^i$  is a quasi-isometric embedding for any  $U \in \mathfrak{S}^H$ , we conclude by summing the inequality over  $1 \leq i \leq n_V$  for all  $V \in \mathcal{V}$ .  $\square$

**5.3. Proof of Theorem 1.4.**

*Proof.* Let  $R > 0$  be large enough to satisfy Propositions 5.1 and 5.6 and Theorem 3.8. Let  $T \geq R + 2D_0$ . By Proposition 5.1 and Proposition 5.6, there exists quasitrees  $\mathcal{T}_1, \dots, \mathcal{T}_m$  such that  $H$  acts on  $\prod_{k=1}^m \mathcal{T}_k$  diagonally and for any choice of base points  $o_k \in \mathcal{T}_k$  and any  $h \in H$ ,

$$\sum_{V \in \mathfrak{S}^T} d_V^{T,R}(1, h) + \sum_{U \in \mathfrak{S}^H} \{d_U(1, h)\}_R \leq \sum_{k=1}^m d_{\mathcal{T}_k}(o_k, ho_k).$$

By definition of thick distance,  $d_U^{T,R}(1, h) \leq \llbracket d_U(1, h) \rrbracket_R$ . Thus, by Theorem 3.8,

$$\sum_{V \in \mathfrak{S}} d_V^{T,R}(1, h) \leq \sum_{k=1}^m d_{\mathcal{T}_k}(o_k, ho_k)$$

for any  $h \in H$ .

On the other hand, the orbit map from  $H$  to  $\prod_{k=1}^m \mathcal{T}_k$  is coarsely Lipschitz since  $H$  is finitely generated. Therefore,  $H$  embeds quasi-isometrically into  $\prod_{k=1}^m \mathcal{T}_k$ , which means that  $H$  has property  $QT_0$ . Finally, we know that  $G$  has property  $QT$  since property  $QT$  passes to any finite-index supergroup.

For the “moreover” part, first note that  $G$  is coarse median for the same reason as [BHS19, Theorem 7.3]. The rest of the proof is just a combination of [HP22; Pet21]. The proof of [HP22, Proposition 3.9] can be naturally generalized to quasitrees from domains of type II. The proof in [Pet21, §3] for mapping class groups can be repeated verbatim to deal with quasitrees from domains of type I.  $\square$

For relative HHGs with only type II domains, we obtain the following stronger theorem.

**Theorem 5.7.** *Let  $(G, \mathfrak{S})$  be a relative HHG that is colorable. If every  $U \in \mathfrak{S}$  is of type II, then  $G$  has property  $QT_0$ .*

*Proof.* The proof is a simplified version of that of Theorem 1.4. Since  $G$  is colorable and every domain is of type II, the finite-index subgroup  $H$  in the above proof can be replaced with  $G$  itself. This gives  $QT_0$  rather than just  $QT$ .  $\square$

### 6. A criterion for having separable quasi-axes

In this section, we provide a criterion for a relative HHG to have separable quasi-axes that is easy to use in application. For an acylindrical action on a hyperbolic space, we have seen in Section 2.1 that the elementary closure of any loxodromic element is a *virtual centralizer*. Inspired by the discussion in [BBF21, §4.3], the following lemma generalizes this fact. In general, we cannot expect the elementary closure to be a centralizer as in [BBF21, §4.3], because it may contain a flip.

**Lemma 6.1.** *Let  $G$  be a group acting on a  $\delta$ -hyperbolic space  $X$  with an acylindrical image. Let  $K$  be the kernel of this action. Assume that there is a subgroup  $H < G$  such that  $[H, K] = 1$  and  $H$  is mapped to a finite-index subgroup of  $G/K$ . Then for any loxodromic element  $g \in G$ , the elementary closure  $EC_G(g)$  is a virtual centralizer in  $G$  of some loxodromic element in  $H$ .*

*Proof.* Let  $\bar{G} = G/K$ , and let  $\theta : G \rightarrow \bar{G}$  be the quotient map. For any  $g \in G$ , denote the image  $\theta(g) \in \bar{G}$  by  $\bar{g}$ .

Let  $g \in G$  be a loxodromic element. Since  $\theta(H)$  is a finite-index subgroup of  $\bar{G}$ , we can choose  $h \in H$  and  $n > 0$  such that  $\bar{h} = \bar{g}^n$ . Since  $\bar{G}$  acts acylindrically,

$EC_{\bar{G}}(\bar{g}) = EC_{\bar{G}}(\bar{h})$ , which contains the cyclic subgroup  $\langle \bar{h} \rangle$  as a finite-index subgroup. By definition,  $EC_G(g) = \theta^{-1}(EC_{\bar{G}}(\bar{g}))$ . Thus, the preimage  $\theta^{-1}(\langle \bar{h} \rangle) = K \cdot \langle h \rangle$  is a finite-index subgroup of  $EC_G(g)$ .

On the other hand, any element in  $G$  that commutes with  $h$  has an image in  $\bar{G}$  that commutes with  $\bar{h}$ . Thus,  $C_G(h) < EC_G(g)$ . Note that  $K \cdot \langle h \rangle < C_G(h)$  since  $[H, K] = 1$ . In conclusion,  $C_G(h)$  is a finite-index subgroup of  $EC_G(g)$ .  $\square$

**Definition 6.2.** For a relative HHG  $(G, \mathfrak{S})$ , we say a domain  $V \in \mathfrak{S}$  has *neat kernel* if there exists a subgroup  $H_V < \text{Stab}_G(V)$  such that  $[H_V, \ker(\theta_V)] = 1$  and  $\theta_V$  maps  $H_V$  to a finite-index subgroup of  $G_V$ .

**Proposition 6.3.** *Let  $(G, \mathfrak{S})$  be a relative HHG that is residually finite. Let  $V \in \mathfrak{S}$ . If  $V$  has hyperbolicity, acylindrical image and neat kernel, then  $V$  has separable quasi-axes.*

*Proof.* Let  $g \in \text{Stab}_G(V)$  be a loxodromic element. Since  $V$  has neat kernel,  $EC(g)$  is a virtual centralizer in  $\text{Stab}_G(V)$  of some loxodromic element  $h \in H_V$  by Lemma 6.1. Every element that commutes with  $h$  stabilizes  $V$ . Therefore,  $EC(g)$  is a virtual centralizer of  $h$  in  $G$ . Moreover, a centralizer in a residually finite group is separable (see [BBF21, Lemma 2.1] or the proof of [Lon87, Proposition]). It is also known that a finite-index supergroup of a separable subgroup is still separable (easy to see from the profinite topology). Therefore,  $V$  has separable quasi-axes.  $\square$

**Corollary 6.4.** *Let  $(G, \mathfrak{S})$  be a relative HHG that is residually finite. Let  $S \in \mathfrak{S}$  be the unique maximal domain. Then  $S$  is of type I.*

*Proof.* By [BHS17b, Theorem 14.3],  $G$  acts on  $\mathcal{CS}$  acylindrically. Now  $\ker(\theta_V)$  is trivial, so  $S$  has neat kernel. Thus,  $S$  has separable quasi-axes by Proposition 6.3. Moreover,  $S$  has cobounded nested region because  $G$  acts on  $\mathcal{X}$  coboundedly. In conclusion,  $S$  is of type I.  $\square$

## 7. Applications

**7.1. Mapping class groups.** We now explain how Theorem 1.4 applies to mapping class groups to recover the following theorem.

**Theorem 7.1** [BBF21, Theorem 1.2]. *Mapping class groups of finite-type surfaces have property QT.*

*Proof.* Let  $\Sigma$  be a finite-type surface, i.e., a closed oriented surface with finitely many marked points. Let  $\mathcal{M}(\Sigma)$  be its marking complex [MM00]. Let  $\mathfrak{S}$  be the collection of isotopy classes of essential non-pants-closed subsurfaces of  $\Sigma$ , where disconnected subsurfaces are also allowed. Given any  $V \in \mathfrak{S}$ , let  $\hat{V}$  be the surface obtained by gluing a once-punctured disk to each boundary component of  $V$ . Let  $\mathcal{CV}$  be the curve graph of  $\hat{V}$ . Here the curve graph of a disconnected surface is

defined as the join of the curve graphs of its components, and thus is bounded. It is known that  $(\mathcal{M}(\Sigma), \mathfrak{S})$  is an HHS. Moreover, the mapping class group  $\text{MCG}(\Sigma)$  is an HHG by virtue of its action on  $(\mathcal{M}(\Sigma), \mathfrak{S})$  [BHS19, §11].

For any  $V \in \mathfrak{S}^\circ$ , let  $V^\perp$  be the closure of  $\Sigma - V$  in  $\Sigma$ , and let

$$\text{MCG}(\Sigma; V) < \text{MCG}(\Sigma)$$

be the stabilizer of  $V$ . Let  $\eta_V : \text{MCG}(V) \rightarrow \text{MCG}(\Sigma)$  be the homomorphism induced by the inclusion  $V \hookrightarrow \Sigma$ . Denote the image of  $\eta_V$  by  $\overline{\text{MCG}}(V)$ . By Theorem 3.18 of [FM12],  $\overline{\text{MCG}}(V^\perp)$  is exactly the kernel of the restriction homomorphism  $\theta_V : \text{MCG}(\Sigma; V) \rightarrow G_V$ , where  $G_V$  is a finite-index subgroup of  $\text{MCG}(\hat{V})$ . It is clear that  $\overline{\text{MCG}}(V)$  commutes with  $\overline{\text{MCG}}(V^\perp)$  and  $\theta_V$  maps  $\overline{\text{MCG}}(V)$  to a finite-index subgroup of  $G_V$ . Therefore,  $V$  has neat kernel.

It is known that  $\text{MCG}(\Sigma)$  is virtually colorable [BBF15, §5] and residually finite [Gro74]. We only need to prove that every unbounded domain  $V \in \mathfrak{S}$  is of type I. First, it is clear that  $F_V$  is coarsely  $\mathcal{M}(V)$ . Since  $\text{MCG}(\hat{V})$  acts coboundedly on  $\mathcal{M}(V)$  and acts acylindrically on  $\mathcal{C}\hat{V}$  [Bow08],  $V$  has cobounded nested region and acylindrical image. Furthermore,  $V$  has separable quasi-axes by the above discussion and Proposition 6.3. Therefore,  $V$  is of type I.

In conclusion, mapping class groups of finite-type surfaces have property QT by Theorem 1.4. □

Certain quotients of a mapping class group are again HHGs, as proved in [BHS17a] and [BHMS24]. In particular, the quotient by the normal closure of a suitable power of a pseudo-Anosov element or by the normal closure of suitable powers of all Dehn twists is again an HHG. It would be interesting to determine whether these quotient groups are still residually finite, thereby satisfying the assumption of Theorem 1.4.

**7.2. Admissible graphs of groups.** *Admissible groups* were introduced by Croke and Kleiner in [CK02]; they generalize the fundamental groups of nongeometric 3-dimensional graph manifolds.

**Definition 7.2.** Let  $\mathcal{G} = (\Gamma, \{G_v\}, \{G_e\})$  be a graph of groups. We say  $\mathcal{G}$  is *admissible* if the following hold:

- (1)  $\Gamma$  is a finite graph with at least one edge.
- (2) Each vertex group  $G_v$  has center  $Z_v \cong \mathbb{Z}$ , and  $H_v := G_v/Z_v$  is a nonelementary hyperbolic group.
- (3) Every edge group  $G_e$  is isomorphic to  $\mathbb{Z}^2$ .
- (4) If  $e$  is an edge with  $v = e^+$  and  $w = e^-$ , and  $\tau_e, \tau_{\bar{e}}$  are the edge monomorphisms, then the subgroup  $\langle \tau_e^{-1}(Z_v), \tau_{\bar{e}}^{-1}(Z_w) \rangle$  has finite index in  $G_e$ .

- (5) Let  $e_1$  and  $e_2$  be distinct edges entering a vertex  $v$ , and let  $K_i \subset G_v$  be the image of the edge homomorphism  $\tau_{e_i}$  for  $i = 1, 2$ . Then
- for every  $g \in G_v$ ,  $gK_1g^{-1}$  is not commensurable with  $K_2$ ;
  - for every  $g \in G_v - K_i$ ,  $gK_ig^{-1}$  is not commensurable with  $K_j$ .

A group  $G$  is *admissible* if it is the fundamental group of an admissible graph of groups.

Every admissible group  $G$  has a (combinatorial) HHG structure by [HRSS24, Theorem 1.4]. According to the classification of simplices by [HRSS24, Lemma 6.2], if  $\Delta \perp g\Delta$ , where  $\Delta$  corresponds to an unbounded hyperbolic space, then  $\Delta$  is of type 8 and  $g$  exchanges two adjacent vertices in the Bass–Serre tree (see [BHMS24, Definition 1.11] for definition of orthogonality in a combinatorial HHS). Therefore, it is easy to see that  $G$  has a subgroup of index at most 2 that is colorable (see [NY23, Lemma 4.6] for example). This shows the virtual colorability of  $G$ . Thus every nongeometric graph manifold group has a virtually colorable HHG structure with all associated hyperbolic spaces being quasitrees. Thus, nongeometric graph manifold groups have property QT by [HP22, Theorem 3.1] or Theorem 1.4. However, in the HHG structure of an admissible group, associated hyperbolic spaces are not necessarily quasitrees. As an application of Theorem 1.4, we show that property QT still holds true in this case if we assume  $G$  to be residually finite.

**Theorem 7.3.** *Let  $\mathcal{G} = (\Gamma, \{G_v\}, \{G_e\})$  be an admissible graph of groups, and let  $G = \pi_1\mathcal{G}$ . If  $G$  is residually finite, then  $G$  has property QT.*

*Proof.* According to the classification of simplices [HRSS24, Lemma 6.2], any simplex that is not of type 7 corresponds to a quasitree so it is a domain of type II. Thus, we only need to check that simplices of type 7 are of type I. The stabilizer of such a simplex  $\Delta$  is exactly a vertex group  $G_v$  that acts on  $\mathcal{C}(\Delta)$  with image  $H_v = G_v/Z_v$ . Now  $\mathcal{C}(\Delta)$  is coarsely the hyperbolic space obtained by coning off  $H_v$  as a relatively hyperbolic group, and  $F_\Delta$  is coarsely  $H_v$  itself. Therefore, acylindrical image and cobounded nested region hold true (see [Osi16, Proposition 5.2] for acylindricity). Since  $G_v$  is a central extension of  $H_v$  by  $Z_v$ ,  $\Delta$  has neat kernel. Therefore, we conclude by Proposition 6.3 and Theorem 1.4.  $\square$

There is another approach to property QT of nongeometric graph manifold groups in [HNY25]. For graph manifolds with nonempty boundary, they actually prove in a more general setting. A *Croke–Kleiner admissible group* (abbreviated as CKA group) is an admissible group that admits a geometric action on a complete proper CAT(0) space. As a corollary of Theorem 7.3, we recover the following theorem.

**Corollary 7.4** ([HNY25, Theorem 1.3]). *Let  $G$  be a CKA group where for every vertex  $v$  the central extension  $1 \rightarrow Z_v \rightarrow G_v \rightarrow H_v \rightarrow 1$  has an omnipotent hyperbolic quotient group  $H_v$ . Then  $G$  has property QT.*

For definition of omnipotence, we refer the reader to [Wis00]. Note that if every hyperbolic group is residually finite, then every hyperbolic group is omnipotent by [Wis00, Remark 3.4]. Under the assumption of Corollary 7.4, the central extension associated with any vertex virtually splits by [BH99, Theorem II.7.1]. Therefore, Theorem 7.3 implies Corollary 7.4 due to the following lemma.

**Lemma 7.5.** *Let  $G$  be an admissible group where for every vertex  $v$  the central extension  $1 \rightarrow Z_v \rightarrow G_v \rightarrow H_v \rightarrow 1$  virtually splits and the hyperbolic quotient group  $H_v$  is omnipotent. Then  $G$  is residually finite.*

We omit the proof of Lemma 7.5 since it is almost the same as the proof of residual finiteness for graph manifold groups by Hempel [Hem87]. The reader can also see [Ngu26] for an improved result.

**7.3. Hyperbolic-2-decomposable groups.** A group  $G$  is hyperbolic-2-decomposable if  $G$  splits as a graph of hyperbolic groups with 2-ended edge groups.

**Theorem 7.6.** *Let  $G$  be a residually finite hyperbolic-2-decomposable group. Then  $G$  has property QT if and only if  $G$  does not contain any distorted element.*

*Proof.* If  $G$  has property QT, then  $G$  does not contain any distorted element by [HNY25, Lemma 2.5]. Now assume that  $G$  does not contain any distorted element. Let  $\mathfrak{S}$  be the HHG structure of  $G$  given by [RS20]. By construction, there is no orthogonality in  $\mathfrak{S}$ . Thus,  $G$  is colorable. Let  $U \in \mathfrak{S}$ . Then  $\mathcal{C}U$  is either a quasitree or a hyperbolic space obtained by coning off a vertex group  $G_v$  as a relatively hyperbolic group. In the former case,  $U$  is of type II. Now we only need to consider the latter case. Similarly to Theorem 7.3, we have  $\text{Stab}_G(U) = G_v$  and  $F_U$  is coarsely  $G_v$  itself. It is easy to see that acylindrical image, cobounded nested region and neat kernel hold true. Therefore, we conclude by Proposition 6.3 and Theorem 1.4.  $\square$

Similarly to Lemma 7.5, if  $G$  is a hyperbolic-2-decomposable group without any distorted element such that every vertex group is omnipotent, then  $G$  is residually finite (see [Wis00, §4]).

**7.4. Artin groups and extensions of lattice Veech groups.** Let  $G$  be either

- an Artin group of large and hyperbolic type, or
- the  $\pi_1(\Sigma)$ -extension group of a lattice Veech group in the mapping class group  $\text{MCG}(\Sigma)$  of a closed surface  $\Sigma$ .

As shown in [HMS24] and [DDLS24] respectively,  $G$  is a virtually colorable HHG. Moreover, the associated hyperbolic spaces of  $G$  are all quasitrees except the maximal one. By Corollary 6.4 and Theorem 1.4,  $G$  has property QT if  $G$  is residually finite. Hence:

**Theorem 7.7.** *Every residually finite Artin group of large and hyperbolic type has property QT.*

It is proved in [Jan22] that any 3-generator Artin groups with labels  $\geq 4$  except for  $(2m + 1, 4, 4)$  for any  $m \geq 2$  is residually finite. Therefore, any 3-generator Artin group with labels  $\geq 4$  except for  $(2m + 1, 4, 4)$  for any  $m \geq 2$  has property QT. On the other hand, we ask

**Question 7.8.** *When is the  $\pi_1(\Sigma)$ -extension group of a lattice Veech group residually finite?*

### 7.5. Graph products.

**Definition 7.9** (graph product). Let  $\Gamma$  be a finite simplicial graph with the vertex set  $V(\Gamma)$  and the edge set  $E(\Gamma)$ . Each vertex  $v \in V(\Gamma)$  is labeled by a group  $G_v$ . The graph product  $G_\Gamma$  is the group

$$G_\Gamma = \left( \bigstar_{v \in V(\Gamma)} G_v \right) / \langle\langle [g_v, g_w] \mid g_v \in G_v, g_w \in G_w, \{v, w\} \in E(\Gamma) \rangle\rangle.$$

We call the  $G_v$  the *vertex groups* of the graph product  $G_\Gamma$ .

**Theorem 7.10.** *Any graph product of groups whose every vertex group has property  $QT_0$  still has property  $QT_0$ .*

*Proof.* Any graph product  $G_\Gamma$  has a relative HHG structure  $\mathfrak{S}_\Gamma$  by [BR22]. By definition of  $\mathfrak{S}_\Gamma$ , any  $G_\Gamma$ -orbit on  $\mathfrak{S}_\Gamma$  corresponds to a unique subgraph of  $\Gamma$  and is pairwise transversal. Thus,  $G_\Gamma$  is colorable. By [BR22, Theorem 4.4], for each domain  $[g\Lambda] \in \mathfrak{S}_\Gamma$ , either  $[g\Lambda]$  is  $\sqsubseteq$ -minimal or  $Cg\Lambda$  is a quasitree. Since each  $\sqsubseteq$ -minimal domain corresponds to a vertex group, this means that every domain is of type II. By Theorem 5.7,  $G_\Gamma$  has property QT.  $\square$

**Acknowledgements.** The author is grateful to his PhD supervisor, Koji Fujiwara, for many helpful discussions on this paper. He thanks Wenyan Yang for discussing with him the paper [HNY25] that became the motivation for this work. He thanks Shengkui Ye for his comments on Artin groups. He also thanks the anonymous referee for many helpful comments. This work was supported by JST SPRING, Grant Number JPMJSP2110.

## References

- [BBF15] M. Bestvina, K. Bromberg, and K. Fujiwara, “Constructing group actions on quasi-trees and applications to mapping class groups”, *Publ. Math. Inst. Hautes Études Sci.* **122** (2015), 1–64. MR
- [BBF21] M. Bestvina, K. Bromberg, and K. Fujiwara, “Proper actions on finite products of quasi-trees”, *Ann. H. Lebesgue* **4** (2021), 685–709. MR
- [BBFS19] M. Bestvina, K. Bromberg, K. Fujiwara, and A. Sisto, “Acylindrical actions on projection complexes”, *Enseign. Math.* **65**:1-2 (2019), 1–32. MR

- [BFG24] S. H. Balasubramanya, F. Fournier-Facio, and A. Genevois, “Property (NL) for group actions on hyperbolic spaces”, *Groups Geom. Dyn.* **19**:4 (2025), 1319–1371. MR
- [BH99] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, Grundle. Math. Wissen. **319**, Springer, 1999. MR
- [BHMS24] J. Behrstock, M. Hagen, A. Martin, and A. Sisto, “A combinatorial take on hierarchical hyperbolicity and applications to quotients of mapping class groups”, *J. Topol.* **17**:3 (2024), art. id. e12351, 94 pp. MR
- [BHS17a] J. Behrstock, M. F. Hagen, and A. Sisto, “Asymptotic dimension and small-cancellation for hierarchically hyperbolic spaces and groups”, *Proc. Lond. Math. Soc.* (3) **114**:5 (2017), 890–926. MR
- [BHS17b] J. Behrstock, M. F. Hagen, and A. Sisto, “Hierarchically hyperbolic spaces, I: Curve complexes for cubical groups”, *Geom. Topol.* **21**:3 (2017), 1731–1804. MR
- [BHS19] J. Behrstock, M. Hagen, and A. Sisto, “Hierarchically hyperbolic spaces II: Combination theorems and the distance formula”, *Pacific J. Math.* **299**:2 (2019), 257–338. MR
- [Bow08] B. H. Bowditch, “Tight geodesics in the curve complex”, *Invent. Math.* **171**:2 (2008), 281–300. MR
- [BR22] D. Berlyne and J. Russell, “Hierarchical hyperbolicity of graph products”, *Groups Geom. Dyn.* **16**:2 (2022), 523–580. MR
- [But25] J. O. Button, “Generalised Baumslag–Solitar groups and hierarchically hyperbolic groups”, *Algebr. Geom. Topol.* **25**:4 (2025), 2253–2279. MR
- [Che22] M. Chesser, “Stable subgroups of the genus 2 handlebody group”, *Algebr. Geom. Topol.* **22**:2 (2022), 919–971. MR
- [CK02] C. B. Croke and B. Kleiner, “The geodesic flow of a nonpositively curved graph manifold”, *Geom. Funct. Anal.* **12**:3 (2002), 479–545. MR
- [DDLS24] S. Dowdall, M. G. Durham, C. J. Leininger, and A. Sisto, “Extensions of Veech groups II: Hierarchical hyperbolicity and quasi-isometric rigidity”, *Comment. Math. Helv.* **99**:1 (2024), 149–228. MR
- [DGO17] F. Dahmani, V. Guirardel, and D. Osin, *Hyperbolically embedded subgroups and rotating families in groups acting on hyperbolic spaces*, vol. 245, Mem. Amer. Math. Soc. **1156**, Amer. Math. Soc., 2017. MR
- [DHS20] M. G. Durham, M. F. Hagen, and A. Sisto, “Correction to the article Boundaries and automorphisms of hierarchically hyperbolic spaces”, *Geom. Topol.* **24**:2 (2020), 1051–1073. MR
- [DJ99] A. Dranishnikov and T. Januszkiewicz, “Every Coxeter group acts amenably on a compact space”, *Topology Proc.* **24** (1999), 135–141. MR
- [DMS23] M. G. Durham, Y. N. Minsky, and A. Sisto, “Stable cubulations, bicombings, and barycenters”, *Geom. Topol.* **27**:6 (2023), 2383–2478. MR
- [FM12] B. Farb and D. Margalit, *A primer on mapping class groups*, Princeton Mathematical Series **49**, Princeton University Press, 2012. MR
- [Gro74] E. K. Grossman, “On the residual finiteness of certain mapping class groups”, *J. London Math. Soc.* (2) **9** (1974/75), 160–164. MR
- [Gro87] M. Gromov, “Hyperbolic groups”, pp. 75–263 in *Essays in group theory*, Math. Sci. Res. Inst. Publ. **8**, Springer, 1987. MR
- [Hag23] M. Hagen, “Non-colorable hierarchically hyperbolic groups”, *Internat. J. Algebra Comput.* **33**:2 (2023), 337–350. MR

- [Hem87] J. Hempel, “Residual finiteness for 3-manifolds”, pp. 379–396 in *Combinatorial group theory and topology* (Alta, UT, 1984), Ann. of Math. Stud. **111**, Princeton Univ. Press, Princeton, NJ, 1987. MR
- [HMS24] M. Hagen, A. Martin, and A. Sisto, “Extra-large type Artin groups are hierarchically hyperbolic”, *Math. Ann.* **388**:1 (2024), 867–938. MR
- [HNY25] S. Han, H. T. Nguyen, and W. Yang, “Property (QT) for 3-manifold groups”, *Algebr. Geom. Topol.* **25**:1 (2025), 107–159. MR
- [HP22] M. F. Hagen and H. Petyt, “Projection complexes and quasimedians maps”, *Algebr. Geom. Topol.* **22**:7 (2022), 3277–3304. MR
- [HRSS24] M. Hagen, J. Russell, A. Sisto, and D. Spriano, “Equivariant hierarchically hyperbolic structures for 3-manifold groups via quasimorphisms”, *Ann. Inst. Fourier (Grenoble)* **75**:2 (2025), 769–828. MR
- [Jan22] K. Jankiewicz, “Residual finiteness of certain 2-dimensional Artin groups”, *Adv. Math.* **405** (2022), art. id. 108487, 37 pp. MR
- [Lon87] D. D. Long, “Immersions and embeddings of totally geodesic surfaces”, *Bull. London Math. Soc.* **19**:5 (1987), 481–484. MR
- [Man06] J. F. Manning, “Quasi-actions on trees and property (QFA)”, *J. London Math. Soc. (2)* **73**:1 (2006), 84–108. MR
- [Man08] J. F. Manning, “Actions of certain arithmetic groups on Gromov hyperbolic spaces”, *Algebr. Geom. Topol.* **8**:3 (2008), 1371–1402. MR
- [MM00] H. A. Masur and Y. N. Minsky, “Geometry of the complex of curves, II: Hierarchical structure”, *Geom. Funct. Anal.* **10**:4 (2000), 902–974. MR
- [MM99] H. A. Masur and Y. N. Minsky, “Geometry of the complex of curves, I: Hyperbolicity”, *Invent. Math.* **138**:1 (1999), 103–149. MR
- [Ngu26] H. T. Nguyen, “Separability properties of extended admissible groups”, *Forum Math.* **38**:2 (2026), 519–537. MR
- [NY23] H. T. Nguyen and W. Yang, “Croke–Kleiner admissible groups: property (QT) and quasiconvexity”, *Michigan Math. J.* **73**:5 (2023), 971–1019. MR
- [Osi16] D. Osin, “Acylically hyperbolic groups”, *Trans. Amer. Math. Soc.* **368**:2 (2016), 851–888. MR
- [Pet21] H. Petyt, “Mapping class groups are quasicubical”, preprint, 2021. arXiv 2112.10681
- [PS23] H. Petyt and D. Spriano, “Unbounded domains in hierarchically hyperbolic groups”, *Groups Geom. Dyn.* **17**:2 (2023), 479–500. MR
- [PSZ25] H. Petyt, D. Spriano, and A. Zalloum, “Stable cylinders and fine structures for hyperbolic groups and curve graphs”, preprint, 2025. arXiv 2501.13600
- [RS20] B. Robbio and D. Spriano, “Hierarchical hyperbolicity of hyperbolic-2-decomposable groups”, preprint, 2020. arXiv 2007.13383
- [RS95] E. Rips and Z. Sela, “Canonical representatives and equations in hyperbolic groups”, *Invent. Math.* **120**:3 (1995), 489–512. MR
- [Ver24] I. Vergara, “Quasi-trees, Lipschitz free spaces, and actions on  $\ell^1$ ”, preprint, 2024. arXiv 2409.14186
- [Wis00] D. T. Wise, “Subgroup separability of graphs of free groups with cyclic edge groups”, *Q. J. Math.* **51**:1 (2000), 107–129. MR
- [Yan19] W.-y. Yang, “Statistically convex-cocompact actions of groups with contracting elements”, *Int. Math. Res. Not.* **2019**:23 (2019), 7259–7323. MR

Received January 1, 2025. Revised December 12, 2025.

BINGXUE TAO  
DEPARTMENT OF MATHEMATICS  
KYOTO UNIVERSITY  
KYOTO  
JAPAN  
[tao.bingxue.s66@kyoto-u.jp](mailto:tao.bingxue.s66@kyoto-u.jp)

## Guidelines for Authors

Authors may submit articles at [msp.org/pjm/about/journal/submissions.html](http://msp.org/pjm/about/journal/submissions.html) and choose an editor at that time. Exceptionally, a paper may be submitted in hard copy to one of the editors; authors should keep a copy.

By submitting a manuscript you assert that it is original and is not under consideration for publication elsewhere. Instructions on manuscript preparation are provided below. For further information, visit the web address above or write to [journals+pjm@msp.org](mailto:journals+pjm@msp.org) or to Pacific Journal of Mathematics, University of California, Los Angeles, CA 90095–1555. Correspondence by email is requested for convenience and speed.

Manuscripts must be in English, French or German. A brief abstract of about 150 words or less in English must be included. The abstract should be self-contained and not make any reference to the bibliography. Also required are keywords and subject classification for the article, and, for each author, postal address, affiliation (if appropriate) and email address if available. A home-page URL is optional.

At submission time a PDF file is required, and it must be derived from a  $\LaTeX$  source file in which the command `\documentclass[12pt]{amsart}` is active. No changes to horizontal or vertical margins, or line spacing, should be made. Please include in the preamble the command `\usepackage[pagewise]{lineno}\linenumbers` — this will spare referees tedious effort in line counting.

Carefully preserve all related files, such as  $\LaTeX$  sources and individual files for each figure; you will be asked to submit them upon acceptance of the paper.

Bibliographical references should be listed alphabetically at the end of the paper. All references in the bibliography should be cited in the text. Use of  $\BibTeX$  is preferred but not required. Any bibliographical citation style may be used but tags will be converted to the house format (see a current issue for examples).

Figures, whether prepared electronically or hand-drawn, must be of publication quality. Figures prepared electronically should be submitted in Encapsulated PostScript (EPS) or in a form that can be converted to EPS, such as GnuPlot, Maple or Mathematica. Many drawing tools such as Adobe Illustrator and Aldus FreeHand can produce EPS output. Figures containing bitmaps should be generated at the highest possible resolution. If there is doubt whether a particular figure is in an acceptable format, the authors should check with production by sending an email to [journals+pjm@msp.org](mailto:journals+pjm@msp.org).

Each figure should be captioned and numbered, so that it can float. Small figures occupying no more than three lines of vertical space can be kept in the text (“the curve looks like this:”). It is acceptable to submit a manuscript with all figures at the end, if their placement is specified in the text by means of comments such as “Place Figure 1 here”. The same considerations apply to tables, which should be used sparingly.

Forced line breaks or page breaks should not be inserted in the document. There is no point in your trying to optimize line and page breaks in the original manuscript. The manuscript will be reformatted to use the journal’s preferred fonts and layout.

Page proofs will be made available to authors (or to the designated corresponding author) at a website in PDF format. Failure to acknowledge the receipt of proofs or to return corrections within the requested deadline may cause publication to be postponed.

# PACIFIC JOURNAL OF MATHEMATICS

Volume 343    No. 1    July 2026

---

The generic extension map and modular standard modules	1
JOHANNES DROSCHL	
Transverse minimal foliations on unit tangent bundles and applications	39
SÉRGIO R. FENLEY and RAFAEL POTRIE	
Refined bounds for the eigenvalues of the Stokes operator	119
ZHENGCHAO JI and TÜRKAY YOLCU	
The class $\mathcal{Q}$ and mixture distributions with dominated continuous singular parts	139
ALEXEY A. KHARTOV	
Data for Shimura varieties intersecting the Torelli locus	179
WANLIN LI, ELENA MANTOVAN and RACHEL PRIES	
The derived series of GGS groups	211
J. MORITZ PETSCHICK	
Property QT of relatively hierarchically hyperbolic groups	231
BINGXUE TAO	