

*Pacific
Journal of
Mathematics*

**REFINED BOUNDS FOR THE EIGENVALUES
OF THE STOKES OPERATOR**

ZHENGCHAO JI AND TÜRKAY YOLCU

Volume 343 No. 1

July 2026

REFINED BOUNDS FOR THE EIGENVALUES OF THE STOKES OPERATOR

ZHENGCHAO JI AND TÜRKAY YOLCU

We analyze bounds for the sums of eigenvalues of the Stokes operator restricted to a bounded domain $\Omega \subset \mathbb{R}^d$ with $d \geq 2$. We improve upon existing lower bound estimates due to Ilyin as well as those by the second author and S. Yıldırım Yolcu, while preserving sharpness in the sense of Weyl asymptotics.

1. Introduction

Let Ω be an open bounded set in \mathbb{R}^d , $d \geq 2$. We derive sharper estimates for the eigenvalues $\{\lambda_k\}_{k=1}^\infty$ of Stokes problem defined by

$$(1) \quad \begin{cases} -\Delta \mathbf{u}_k + \nabla p_k = \lambda_k \mathbf{u}_k & \text{in } \Omega, \\ \operatorname{div} \mathbf{u}_k = 0 & \text{in } \Omega, \\ \mathbf{u}_k = 0 & \text{on } \partial\Omega. \end{cases}$$

The eigenvalues of the Stokes problem in (1) are important due to their numerous applications in fluid mechanics. They can be interpreted as the eigenvalues associated with linear (small or infinitesimal) self-oscillations of the fluid within the domain Ω [4]. When the eigenfunctions are required to meet specific conditions that are crucial from a physical perspective in fluid dynamics, exact eigenvalues for the Stokes problem are not obtainable. Consequently, both theoretical and practical aspects necessitate the close identification of the eigenvalues. The literature concerning the eigenvalues of the Stokes problem is vast, and recent studies aimed at deriving estimates for these eigenvalues are documented in [6; 9; 10; 12; 24].

Before presenting the results, we first review some basic facts about the theory of the Stokes operator. Let \mathcal{U} denote the set of smooth, divergence-free vector functions with compact supports. Specifically,

$$\mathcal{U} = \{\mathbf{u} : \Omega \rightarrow \mathbb{R}^d, \mathbf{u} \in C_0^\infty(\Omega), \operatorname{div} \mathbf{u} = 0\}$$

MSC2020: primary 35P15; secondary 35P30.

Keywords: Stokes operator, Berezin–Li–Yau inequality, eigenvalue, inequality, Weyl asymptotics.

and let the closure of \mathcal{U} in $L^2(\Omega)$ and $\mathbf{H}_0^1(\Omega)$ be denoted by L and U , respectively. In particular,

$$U \subseteq \{\mathbf{u} \in \mathbf{H}_0^1(\Omega), \operatorname{div} \mathbf{u} = 0\}$$

and if Ω is open, bounded and locally Lipschitz, we have the equality [3]:

$$U = \{\mathbf{u} \in \mathbf{H}_0^1(\Omega), \operatorname{div} \mathbf{u} = 0\}.$$

We note that $L^2(\Omega)$ can be written as $L^2(\Omega) = L \oplus L^\perp$, where

$$L^\perp = \{\mathbf{u} \in L^2(\Omega), \mathbf{u} = \nabla p, p \in L_{\text{loc}}^2(\Omega)\}$$

(see [3; 4; 19]). The Stokes operator is formally expressed as

$$A\mathbf{u} = -P_L(\Delta\mathbf{u}),$$

where the linear operator $P_L : L^2(\Omega) \rightarrow L$, defined by

$$P_L(\mathbf{v}) = \mathbf{v} - \nabla \Delta^{-1}(\operatorname{div} \mathbf{v}),$$

is called the Leray projection [4], i.e., $P_L(P_L(\mathbf{v})) = P_L(\mathbf{v})$. Notice that P_L becomes the identity operator for the divergence-free vector fields because $P_L(\mathbf{v}) = \mathbf{v}$ for $\operatorname{div} \mathbf{v} = 0$. Furthermore, P_L can be understood as the projection onto divergence-free vector fields. This projection is particularly employed to remove some terms and components in the Stokes and Navier–Stokes equations. Going back to the Stokes problem in (1), one can also see that

$$A\mathbf{u} = -\Delta\mathbf{u} + \nabla p, \quad p = \Delta^{-1}(\operatorname{div} \Delta\mathbf{u}),$$

and that for all \mathbf{u}, \mathbf{w} in U , A is defined by

$$(A\mathbf{u}, \mathbf{w}) = (\nabla\mathbf{u}, \nabla\mathbf{w}),$$

i.e.,

$$\int_{\Omega} A\mathbf{u}(\mathbf{x}) \cdot \mathbf{w}(\mathbf{x}) \, d\mathbf{x} = \int_{\Omega} \sum_{i=1}^d \frac{\partial \mathbf{u}}{\partial x_i} \frac{\partial \mathbf{w}}{\partial x_i} \, d\mathbf{x}.$$

The Stokes operator A is an unbounded, linear, self-adjoint, positive definite operator in L . In addition to these nice properties, its inverse A^{-1} is a compact, self-adjoint operator [3; 4]. Thus, there exists an orthonormal basis $\{\mathbf{u}_k\}_{k=1}^{\infty}$ in L and a set of positive eigenvalues $\{\delta_k\}_{k=1}^{\infty}$ accumulating at zero such that

$$A^{-1} \mathbf{u}_k = \delta_k \mathbf{u}_k,$$

for $k = 1, 2, 3, \dots$. Therefore, $\{\mathbf{u}_k\}_{k=1}^{\infty} \in U$ with corresponding eigenvalues $\{\lambda_k\}_{k=1}^{\infty}$ with $\lambda_k = 1/\delta_k$ are such that

$$(2) \quad A\mathbf{u}_k = \lambda_k \mathbf{u}_k.$$

The eigenvalues (including multiplicities) satisfy

$$(3) \quad 0 < \lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_k \leq \dots \rightarrow \infty \quad \text{as } k \rightarrow \infty.$$

Observe that by taking the scalar product with orthonormal \mathbf{u}_k we have

$$(4) \quad \lambda_k = \|\nabla \mathbf{u}_k\|^2 = \int_{\Omega} \sum_{i=1}^d \frac{\partial \mathbf{u}_k}{\partial x_i} \cdot \frac{\partial \mathbf{u}_k}{\partial x_i} dx.$$

The eigenvalues also satisfy the Weyl asymptotic formulas [4; 8; 17] for $d \geq 2$:

$$(5) \quad \lambda_k \sim 4\pi \left(\frac{\Gamma(1 + \frac{d}{2})}{(d-1)|\Omega|} \right)^{\frac{2}{d}} k^{\frac{2}{d}},$$

where $\Gamma(x)$ denotes the gamma function $\Gamma(x) = \int_0^\infty t^{x-1} e^{-t} dt$ for $x > 0$ and $|\Omega|$ denotes the Lebesgue measure of Ω .

The eigenvalue bounds that will be established in this paper closely resemble those associated with the Dirichlet Laplacian. A substantial body of literature exists regarding inequalities for the eigenvalues of the Dirichlet Laplacian; notable references include the survey articles cited in [1; 2; 7]. In his renowned paper [18], G. Pólya demonstrated that for small values of k , the ratio $4\pi k/|\Omega|$ serves as a lower bound for the eigenvalues μ_k of the Dirichlet Laplacian in tiling domains in \mathbb{R}^2 . Pólya conjectured that this result could be extended to any bounded domain in \mathbb{R}^d . This conjecture remains unresolved. A significant advance in establishing a lower bound was achieved by P. Li and S.-T. Yau [14], who demonstrated the following inequality related to the sums of the eigenvalues μ_j of the Dirichlet Laplacian on the domain Ω :

$$(6) \quad \sum_{j=1}^k \mu_j \geq 4\pi \frac{d}{d+2} \left(\frac{\Gamma(1 + \frac{d}{2})}{|\Omega|} \right)^{\frac{2}{d}} k^{1+\frac{2}{d}}.$$

Recent research has focused significantly on these types of bounds, along with their extensions and enhancements, particularly in relation to other operators. For instance, A. Melas [16] provided an improvement on (6) that incorporates moments of inertia. In two dimensions, H. Kovařík, S. Vugalter, and T. Weidl [13] further advanced Melas’s findings by introducing a positive correction term that involves the size of the boundary. T. Weidl [21] achieved an enhancement of the sharp Berezin-type bounds concerning the Riesz means $\sum_k (z - \mu_k)_+^\sigma$ of the eigenvalues associated with the Dirichlet Laplacian operator within a specified domain for $\sigma \geq 3/2$. Regarding other operators, Harrell and Yıldırım Yolcu [5] along with Yolcu [29] proved inequalities of Berezin–Li–Yau type applicable to the eigenvalues of Klein–Gordon operators. Yıldırım Yolcu and Yolcu [25; 28; 27] established

Melas-type improvements for the eigenvalues of fractional Laplacian operators $(-\Delta)^{\alpha/2}$, $\alpha \in (0, 2]$.

It is noteworthy that the coefficient of $k^{1+\frac{2}{d}}$ in (6) is slightly larger for the tiling domains [18]. See [23] for more extensive research and a generalized conjecture.

Lower bounds on the sums of eigenvalues are important for the theoretical framework of attractors associated with the Navier–Stokes equations [9; 10]. The eigenvalue bounds of the Stokes operator have garnered attention following the work by A. A. Ilyin [9], who proved an inequality of Berezin–Li–Yau type for the Stokes operator:

$$(7) \quad \sum_{j=1}^k \lambda_j \geq 4\pi \frac{d}{d+2} \left(\frac{\Gamma(1+\frac{d}{2})}{(d-1)|\Omega|} \right)^{\frac{2}{d}} k^{1+\frac{2}{d}},$$

where $k \geq 1$. In this article, we first build upon (7) by establishing upper bounds for the sums of negative powers and lower bounds for the sums of positive powers of the eigenvalues of the Stokes operator as follows:

Theorem 1. *For $k \geq 1$, $0 < b < d/2$, and $d \geq 2$, the sums of negative powers of eigenvalues $\{\lambda_i\}_{i=1}^{\infty}$ of the Stokes operator on Ω in (1) satisfy*

$$(8) \quad \sum_{j=1}^k \lambda_j^{-b} \leq (4\pi)^{-b} \frac{d}{d-2b} \left(\frac{(d-1)|\Omega|}{\Gamma(1+\frac{d}{2})} \right)^{\frac{2b}{d}} k^{1-\frac{2b}{d}}.$$

Theorem 2. *For $k \geq 1$, $0 < a \leq 1$, and $d \geq 2$, the sums of positive powers of eigenvalues $\{\lambda_i\}_{i=1}^{\infty}$ of the Stokes operator on Ω in (1) satisfy*

$$(9) \quad \sum_{j=1}^k \lambda_j^a \geq (4\pi)^a \frac{d}{d+2a} \left(\frac{\Gamma(1+\frac{d}{2})}{(d-1)|\Omega|} \right)^{\frac{2a}{d}} k^{1+\frac{2a}{d}}.$$

By appropriately translating the open set Ω if needed, we can assume that the second moment of inertia $I(\Omega)$ is defined as

$$I(\Omega) = \int_{\Omega} |\mathbf{x}|^2 d\mathbf{x}.$$

Ilyin [10] was able to improve the Berezin–Li–Yau inequality in (7) for the Stokes operator by adding a lower-order term involving the moment of inertia $I(\Omega)$ in dimensions 2, 3, and 4 as follows:

$$(10) \quad \sum_{j=1}^k \lambda_j \geq 4\pi \frac{d}{d+2} \left(\frac{\Gamma(1+\frac{d}{2})}{(d-1)|\Omega|} \right)^{\frac{2}{d}} k^{1+\frac{2}{d}} + \frac{d-1}{48} \beta_d^S \frac{|\Omega|}{I(\Omega)} k,$$

where in the two-dimensional case $\beta_2^S = \frac{239}{240}$, while for $d = 3, 4$ it suffices to take $\beta_3^S = 0.986$ and $\beta_4^S = 0.978$.

Let

$$B_R(\mathbf{x}) := \{\mathbf{y} \in \mathbb{R}^d : |\mathbf{y} - \mathbf{x}| \leq R\}$$

be the ball of radius R centered at \mathbf{x} in \mathbb{R}^d . Let

$$(11) \quad \omega_d = \frac{\pi^{\frac{d}{2}}}{\Gamma(1 + \frac{d}{2})}$$

be the volume of the unit ball $B_1(\mathbf{x})$ in \mathbb{R}^d .

Building on the improvements in (10), Yıldırım Yolcu and Yolcu [24] introduced additional correction terms beyond those dependent on $k^{1+\frac{2}{d}}$ and $k \geq 1$, along with their explicit coefficients, applicable for any $d \geq 2$:

$$(12) \quad \sum_{j=1}^k \lambda_j \geq 4\pi \frac{d}{d+2} \left(\frac{\Gamma(1 + \frac{d}{2})}{(d-1)|\Omega|} \right)^{\frac{2}{d}} k^{1+\frac{2}{d}} + \frac{d-1}{24(d^2+2d)} \frac{|\Omega|}{I(\Omega)} k \\ + \frac{((d-1)|\Omega|)^{\frac{3d+2}{2d}}}{144\sqrt{\pi} d^{\frac{3}{2}} (d+2) I(\Omega)^{\frac{3}{2}} \Gamma(1 + \frac{d}{2})^{\frac{1}{d}}} k^{1-\frac{1}{d}}.$$

In this paper we establish several effective lower bounds for the sums of the eigenvalues of the Stokes problem, including the following improved estimates:

Theorem 3 (refinement of the Berezin–Li–Yau inequality for the Stokes operator). *Let $\Omega \subset \mathbb{R}^d$. For any $d \geq m+1 \geq 2$, the eigenvalues $\{\lambda_i\}_{i=1}^{\infty}$ of the Stokes operator in (1) satisfy*

$$\sum_{i=1}^k \lambda_i \geq 4\pi \left(\frac{\Gamma(1 + \frac{d}{2})}{(d-1)|\Omega|} \right)^{\frac{2}{d}} k^{1+\frac{2}{d}} - \frac{2\omega_d^{\frac{m-1}{d}} A_{m+2} \varrho^{\frac{(m+1)d+m-1}{d}}}{(d+2)M^{m+1}} k^{1-\frac{m-1}{d}} \\ + c \frac{2\omega_d^{\frac{m}{d}} (m+1) A_{m+3} \varrho^{\frac{(m+2)d+m}{d}}}{(d+2)(m+3)M^{m+2}} k^{1-\frac{m}{d}},$$

where $k \geq 1$, a is defined by the equality in (39),

$$(13) \quad \varrho = \frac{d-1}{(2\pi)^d} |\Omega|, \quad m_s = \frac{2\sqrt{(d^2-d)|\Omega|I(\Omega)}}{(2\pi)^d}, \quad M = \frac{\sqrt{d^2-d}}{(2\pi)^d} |\Omega|^{\frac{d+1}{d}} \omega_d^{-\frac{1}{d}},$$

and the remaining notation is defined by

$$A_r = (a+1)^r - a^r, \quad c = \min\{1, \max\{a_1, a_2\}\}, \\ a_1 = \frac{(d+2)(m+3)}{(m+1)((m+2)d+m)} \frac{(2\pi)^{m+2}}{A_{m+3}} d^{\frac{m+2}{2}} (d-1)^{-\frac{m(d+2)+2m}{2d}},$$

and

$$a_2 = \frac{\sqrt{2}d^{\frac{1}{d}}}{2(d-1)^{\frac{d+2}{2d}}} \frac{[(m+1)d+m-1](m+3)}{[(m+2)d+m](m+1)} \frac{A_{m+2}}{A_{m+3}}.$$

Theorem 3 yields no improvement when $d = 2$, but it quickly yields the following enhanced lower bound for any dimension $d \geq 3$:

Corollary 1. *For any bounded domain $\Omega \subseteq \mathbb{R}^d$, with $d \geq 3$, and any $k \geq 1$, the eigenvalues $\{\lambda_i\}_{i=1}^\infty$ of the Stokes operator in (1) satisfy*

$$\begin{aligned} \sum_{i=1}^k \lambda_i \geq \frac{4\pi d}{d+2} \left(\frac{\Gamma(1+\frac{d}{2})}{(d-1)|\Omega|} \right)^{\frac{2}{d}} k^{1+\frac{2}{d}} + \frac{\varrho^2}{6(d+2)m_s^2} k + \frac{\omega_d^{-\frac{1}{d}} \varrho^{\frac{3d+1}{d}}}{9(d+2)m_s^3} k^{\frac{d-1}{d}} \\ + \frac{3\omega_d^{-\frac{1}{d}} \varrho^{\frac{4d+2}{d}}}{80(n+2)m_s^4} k^{\frac{d-2}{d}}, \end{aligned}$$

where ϱ and m_s are as in (13).

When $d = 3$, we have an improvement:

Theorem 4. *Let $\Omega \subset \mathbb{R}^d$ with $d = 3$. Then the eigenvalues $\{\lambda_i\}_{i=1}^\infty$ of the Stokes operator in (1) satisfy*

$$(14) \quad \sum_{i=1}^k \lambda_i \geq \frac{2\pi\varrho^6}{15m_s^5} ((t+1)^6 - t^6),$$

where m_s and ϱ are defined in (13) and

$$t = \frac{1}{2} \left(\left(T + \sqrt{T^2 + \frac{1}{27}} \right)^{\frac{1}{3}} - \left(-T + \sqrt{T^2 + \frac{1}{27}} \right)^{\frac{1}{3}} - 1 \right), \quad T = \frac{(d+1)m_s^d}{\omega_d \varrho^{d+1}} k.$$

Furthermore,

$$\sum_{i=1}^k \lambda_i \geq \frac{3}{5} \left(\frac{2\pi^3}{2\omega_3|\Omega|} \right)^{\frac{2}{3}} k^{\frac{5}{3}} + \frac{1}{24} \frac{|\Omega|}{I(\Omega)} k - \frac{11\pi\omega_3^{-\frac{1}{3}} \varrho^{\frac{14}{3}}}{180 m_s^4} k^{\frac{1}{3}} + \frac{659\pi\omega_3^{\frac{1}{3}} \varrho^{\frac{22}{3}}}{75000 \cdot 4^{\frac{1}{3}} m_s^6} k^{-\frac{1}{3}}.$$

Inspired by Melas's research [16], we adopted a similar methodology using fundamental techniques from prior studies [10; 22; 24; 27; 28; 29; 26]. While preserving the core strategy of these works, we introduced significant modifications to achieve stronger lower bounds.

Outline. Section 2 presents results needed for proving the theorems discussed in this work. The main content is found in Sections 3–6, which respectively establish Theorems 1, 2, 3 and 4, proving intermediate results as needed.

2. Preliminaries

The set of eigenfunctions $\{\mathbf{u}_j\}_{j=1}^\infty$ of the Stokes operator is orthonormal in $L^2(\Omega)$, and so the set of Fourier transforms $\{\hat{\mathbf{u}}_j\}_{j=1}^\infty$ also forms an orthonormal set in $L^2(\mathbb{R}^d)$, by Plancherel’s theorem. Let us set

$$(15) \quad U_k(\xi) = \sum_{j=1}^k |\hat{\mathbf{u}}_j(\xi)|^2 = \sum_{j=1}^k \left| \frac{1}{(2\pi)^{\frac{d}{2}}} \int_{\Omega} e^{-i\mathbf{x}\cdot\xi} \mathbf{u}_j(\mathbf{x}) \, d\mathbf{x} \right|^2 \geq 0.$$

The integral is taken over Ω instead of \mathbb{R}^d because the support of \mathbf{u}_j is Ω .

The following result, proven by A. A. Ilyin [10, Section 2], is essential to our calculations.

Lemma 1 [10]. *The functions U_k defined by (15) satisfy*

$$(16) \quad \int_{\mathbb{R}^d} U_k(\xi) \, d\xi = k,$$

$$(17) \quad U_k(\xi) \leq \frac{d-1}{(2\pi)^d} |\Omega|, \quad \xi \in \mathbb{R}^d,$$

$$(18) \quad \int_{\mathbb{R}^d} |\xi|^2 U_k(\xi) \, d\xi = \sum_{j=1}^k \lambda_j.$$

Note that equations (16)–(18) bear a striking resemblance to their Laplacian counterparts. However, the proof of Lemma 1 employs suborthonormal functions [10].

We highlight some important properties of the function U . Let $U^*(\xi)$ designate the radial decreasing rearrangement of $U(\xi)$. By approximating U if needed, we may assume that there exists a real-valued absolutely continuous function

$$\zeta : [0, \infty) \rightarrow \left[0, \frac{d-1}{(2\pi)^d} |\Omega|\right]$$

such that $U^*(\xi) = \zeta(|\xi|)$. Then

$$(19) \quad 0 \leq -\zeta' \leq m_s,$$

where m_s depends only on d and Ω and is explicitly defined in (13). To see (19), let μ be the distribution function defined by

$$\mu(s) = |\{U(\xi) > s\}| = |\{U^*(\xi) > s\}|.$$

Then, we observe that

$$\mu(\zeta(t)) = |\{U^*(\xi) > \zeta(t)\}| = |\{\xi : |\xi| < t\}| = |B_t(0)| = w_d t^d.$$

Invoking the coarea formula in view of (17), we have

$$\mu(s) = \int_s^\infty \int_{\{U^{-1}(t)\}} \frac{1}{|\nabla U|} d\mathcal{H} dt = \int_s^{\frac{d-1}{(2\pi)^d} |\Omega|} \int_{\{U=t\}} \frac{1}{|\nabla U|} d\mathcal{H} dt,$$

where \mathcal{H} is the $(d-1)$ -dimensional Hausdorff measure. Let us consider $t > 0$ values such that $\zeta'(t) < 0$.

Let \bar{K} denote the closure of $K \subset \mathbb{R}^d$ and ∂K the boundary of K . The isoperimetric inequality,

$$\mathcal{H}(\partial K) \geq dw_d^{\frac{1}{d}} |\bar{K}|^{\frac{d-1}{d}},$$

enables us to deduce

$$\begin{aligned} \frac{dw_d t^{d-1}}{\zeta'(t)} &= \mu'(\zeta(t)) = - \int_{\{U=\zeta(t)\}} \frac{1}{|\nabla U|} d\mathcal{H} \leq -\frac{1}{m_S} \mathcal{H}(\{U = \zeta(t)\}) \\ &\leq -\frac{1}{m_S} dw_d^{\frac{1}{d}} \mu(\zeta(t))^{\frac{d-1}{d}} = -\frac{1}{m_S} dw_d t^{d-1}. \end{aligned}$$

This inequality together with $\zeta' \leq 0$ easily yields (19). Note that (19) essentially states that if the gradient vector of the original function is bounded, then the gradient of the rearrangement retains the same bound. (An alternative derivation is possible, using the Pólya–Szegő inequality.)

Now, let r represent the real number such that $|\Omega| = w_d r^d$. We can get a lower bound for $I(\Omega)$:

$$I(\Omega) \geq \int_{B_r(0)} |\mathbf{x}|^2 d\mathbf{x} = \frac{dw_d}{d+2} r^{d+2} = \frac{d}{d+2} w_d^{-\frac{2}{d}} |\Omega|^{\frac{d+2}{d}},$$

which quickly results in a lower bound for m_S as follows:

$$(20) \quad m_S = \frac{2\sqrt{(d^2-d)|\Omega|I(\Omega)}}{(2\pi)^d} \geq \frac{\sqrt{d^2-d}}{(2\pi)^d} |\Omega|^{\frac{d+1}{d}} w_d^{-\frac{1}{d}} =: M.$$

3. Proof of Theorem 1

The following proof is inspired from the proof of the Berezin–Li–Yau inequality in [5; 14; 28; 29]. An analogous proof is also given in [20] by means of the bathtub principle [15].

Proof of Theorem 1. Assume the properties (16)–(18). Since $|\hat{\mathbf{u}}_j(\xi)|^2 d\xi$ is a probability measure on \mathbb{R}^d and $t \mapsto t^{-b}$ is convex for $t > 0$ and $b > 0$, employing Jensen's inequality and (15), we obtain

$$(21) \quad \sum_{j=1}^k \lambda_j^{-b} = \sum_{j=1}^k \left(\int_{\mathbb{R}^d} |\xi|^2 |\hat{\mathbf{u}}_j(\xi)|^2 d\xi \right)^{-b} \leq \int_{\mathbb{R}^d} \frac{U_k(\xi)}{|\xi|^{2b}} d\xi.$$

Let $\mathbf{1}_\Omega(t)$ denote the characteristic function of the set Ω . Define

$$(22) \quad \begin{aligned} \Psi_k(\xi) &= \frac{d-1}{(2\pi)^d} |\Omega| \mathbf{1}_{B_{S_k}(0)}(\xi) \quad \text{and} \\ S_k &= \left(\frac{(d-2b)(2\pi)^d (\sum_{j=1}^k \lambda_j^{-b})}{d(d-1)\omega_d |\Omega|} \right)^{\frac{1}{d-2b}}, \end{aligned}$$

so that

$$(23) \quad \begin{aligned} \int_{\mathbb{R}^d} \frac{\Psi_k(\xi)}{|\xi|^{2b}} d\xi &= \frac{d-1}{(2\pi)^d} |\Omega| d\omega_d \int_0^{S_k} \frac{1}{t^{2b}} t^{d-1} dt \\ &= \frac{d-1}{(2\pi)^d} |\Omega| \frac{d\omega_d}{d-2b} S_k^{d-2b} = \sum_{j=1}^k \lambda_j^{-b}. \end{aligned}$$

Notice that

$$(24) \quad \left(\frac{1}{|\xi|^{2b}} - \frac{1}{S_k^{2b}} \right) (U_k(\xi) - \Psi_k(\xi)) \leq 0.$$

Integrating (24) on \mathbb{R}^d and using (23) we arrive at

$$(25) \quad \frac{1}{S_k^{2b}} \int_{\mathbb{R}^d} (U_k(\xi) - \Psi_k(\xi)) d\xi \geq \int_{\mathbb{R}^d} \frac{U_k(\xi) - \Psi_k(\xi)}{|\xi|^{2b}} d\xi \geq 0$$

from which it follows that

$$\int_{\mathbb{R}^d} U_k(\mu) d\mu \geq \int_{\mathbb{R}^d} \Psi_k(\mu) d\mu.$$

Thus, by (16), we obtain

$$(26) \quad k \geq \int_{\mathbb{R}^d} \Psi_k(\mu) d\mu = \frac{d-1}{(2\pi)^d} |\Omega| d\omega_d \left(\frac{S_k^d}{d} \right).$$

Substituting ω_d given by (11) and S_k given by (22) into (26) and rearranging the terms, we obtain (8). \square

4. Proof of Theorem 2

We sketch a direct proof of (9) for the sake of completeness.

Proof of Theorem 2. Recall that U_k satisfies (16)–(18). Since $|\hat{\mathbf{u}}_j(\xi)|^2 d\xi$ is a probability measure on \mathbb{R}^d and $r \mapsto r^a$ is concave for $r > 0$ and $0 < a \leq 1$, we can use Jensen's inequality to derive that

$$(27) \quad \sum_{j=1}^k \lambda_j^a = \sum_{j=1}^k \left(\int_{\mathbb{R}^d} |\xi|^{2a} |\hat{\mathbf{u}}_j(\xi)|^2 d\xi \right)^a \geq \int_{\mathbb{R}^d} |\xi|^{2a} U_k(\xi) d\xi.$$

Setting

$$(28) \quad \Theta_k(\xi) = \frac{d-1}{(2\pi)^d} |\Omega| \mathbf{1}_{B_{T_k}(0)}(\xi), \quad T_k = \left(\frac{(d+2a)(2\pi)^d (\sum_{j=1}^k \lambda_j^a)}{d(d-1)\omega_d |\Omega|} \right)^{\frac{1}{d+2a}},$$

we have

$$(29) \quad \int_{\mathbb{R}^d} |\xi|^{2a} \Theta_k(\xi) d\xi = \frac{d-1}{(2\pi)^d} |\Omega| d \omega_d \int_0^{T_k} r^{2a} r^{d-1} dr \\ = \frac{d-1}{(2\pi)^d} |\Omega| \frac{d \omega_d}{d+2a} T_k^{d+2a} = \sum_{j=1}^k \lambda_j^a.$$

Now observe that

$$(30) \quad (|\xi|^{2a} - T_k^{2a})(U_k(\xi) - \Theta_k(\xi)) \geq 0.$$

Integrating (30) on \mathbb{R}^d and using (27) and (29) we conclude that

$$(31) \quad T_k^{2a} \int_{\mathbb{R}^d} (U_k(\xi) - \Theta_k(\xi)) d\xi \leq \int_{\mathbb{R}^d} |\xi|^{2a} (U_k(\xi) - \Theta_k(\xi)) d\xi \leq 0,$$

which yields

$$\int_{\mathbb{R}^d} U_k(\xi) d\xi \leq \int_{\mathbb{R}^d} \Theta_k(\xi) d\xi.$$

Now we use (16) to obtain

$$(32) \quad k \leq \int_{\mathbb{R}^d} \Theta_k(\xi) d\xi = \frac{d-1}{(2\pi)^d} |\Omega| d \omega_d \left(\frac{T_k^d}{d} \right).$$

Substituting the values of ω_d and T_k given in (11) and (28) into (32) and simplifying, we deduce (9). \square

5. Proof of Theorem 3

Before the proof of Theorem 3, we give some elementary results from [11; 24]. To connect two key integrals in our proof, we introduce an important quantity, a , using the idea in [24]:

Lemma 2. For any function $\alpha : [0, \infty) \rightarrow [0, 1]$ and integer $d \geq 1$ with

$$\int_0^\infty \phi(t) dt = 1, \quad \int_0^\infty t^d \phi(t) dt < \infty, \quad \int_0^\infty t^{d+2} \phi(t) dt \leq \infty,$$

there exists an $a \geq 0$ such that

$$\int_a^{a+1} t^d dt = \int_0^\infty t^d \phi(t) dt \quad \text{and} \quad \int_a^{a+1} t^{d+2} dt \leq \int_0^\infty t^{d+2} \phi(t) dt.$$

The following sharp inequality is key in estimating the lower bounds for $\sum_{i=1}^k \lambda_i$.

Lemma 3 [11]. *For an integer $d \geq m + 1 \geq 2$ and positive real numbers t and τ we have*

$$(33) \quad dt^{d+2} - (d+2)\tau^2 t^d + 2\tau^{d+2} - \sum_{k=1}^{m+1} 2k t^{k-1} \tau^{n-k+1} (\tau - t)^2 \geq 0.$$

Considering all of these findings, we arrive at the following intermediate estimate.

Proposition 1. *If $d \geq m + 1 \geq 2$, we have*

$$(34) \quad \sum_{i=1}^k \lambda_i \geq \omega_d (dA)^{\frac{d+2}{d}} (\zeta(0))^{-\frac{2}{d}} - \frac{2\omega_d A_{m+2} (dA)^{\frac{d-m+1}{d}} (\zeta(0))^{\frac{(m+1)d+m-1}{d}}}{(d+2)m_s^{m+1}} + \frac{2\omega_d (m+1) A_{m+3} (dA)^{\frac{d-m}{d}} (\zeta(0))^{\frac{(m+2)d+m}{d}}}{(d+2)(m+3)m_s^{m+2}},$$

where m_s is defined in (13) and

$$A_r := (a+1)^r - a^r \geq 1, \quad A := \frac{1}{d} \int_a^{a+1} t^d dt, \quad 0 \leq \zeta(0) \leq \frac{d-1}{(2\pi)^d} |\Omega|,$$

the quantity a being defined by

$$\int_a^{a+1} t^d dt = \frac{k}{\omega_d}.$$

Proof. We use a strategy similar to that of [11; 24]. Since the map $\xi \mapsto |\xi|^2$ is radial and increasing, Lemma 1 implies that

$$(35) \quad k = \int_{\mathbb{R}^d} U(\xi) d\xi = \int_{\mathbb{R}^d} U^*(\xi) d\xi = d\omega_d \int_0^\infty t^{d-1} \zeta(t) dt,$$

$$(36) \quad \sum_{i=1}^k \lambda_i = \int_{\mathbb{R}^d} |\xi|^2 U(\xi) d\xi \geq \int_{\mathbb{R}^d} |\xi|^2 U^*(\xi) d\xi = d\omega_d \int_0^\infty t^{d+1} \zeta(t) dt.$$

For convenience, we rescale ζ as

$$S(t) = \frac{1}{\zeta(0)} \zeta\left(\frac{\zeta(0)}{m_s} t\right).$$

Obviously, $S(t)$ is a positive function with $S(0) = 1$ and $0 \leq -S'(t) \leq 1$. Let $\phi(t) := -S'(t)$ and define

$$(37) \quad A = \int_0^\infty t^{d-1} S(t) dt, \quad B = \int_0^\infty t^{d+1} S(t) dt.$$

Integrating by parts, we get

$$(38) \quad \int_0^\infty t^d \phi(t) dt = dA, \quad \int_0^\infty t^{d+2} \phi(t) dt = (d+2)B.$$

According to (38) and Lemma 2, there exists $a \geq 0$ such that

$$(39) \quad \int_a^{a+1} t^d dt = dA \quad \text{and} \quad \int_a^{a+1} t^{d+2} dt \leq (d+2)B.$$

For $i \geq 0$, $\tau \geq \frac{1}{2}$ and $a \geq 0$, we have

$$(40) \quad \int_a^{a+1} t^i (\tau - t)^2 dt = \frac{A_{i+3}}{i+3} - \frac{2A_{i+2}}{i+2} \tau + \frac{A_{i+1}}{i+1} \tau^2,$$

where

$$A_r = (a+1)^r - a^r \geq 1.$$

Taking into account Lemma 3 and (40), we may integrate (33) in t from a to $a+1$ to get

$$(41) \quad d(d+2)B \geq d(d+2)A\tau^2 - 2\tau^{d+2} + \sum_{i=1}^{m+1} 2i\tau^{d-i+1} \left(\frac{A_i}{i} \tau^2 - \frac{2A_{i+1}}{i+1} \tau + \frac{A_{i+2}}{i+2} \right).$$

The summation on the second row of (41) can be rewritten as

$$\begin{aligned} &= 2\tau^{d+2} + 2 \sum_{i=1}^m A_{i+1} \tau^{d-i+2} - 2 \sum_{i=1}^{m+1} \frac{2iA_{i+1}}{i+1} \tau^{d-i+2} + 2 \sum_{i=1}^{m+1} \frac{iA_{i+2}}{i+2} \tau^{d-i+1} \\ &= 2\tau^{d+2} + 2A_2\tau^{d+1} + \frac{2mA_{m+2}}{m+2} \tau^{d-m+1} + \frac{2(m+1)A_{m+3}}{m+3} \tau^{d-m} \\ &\quad - 2A_2\tau^{d+1} - \frac{4(m+1)A_{m+2}}{m+2} \tau^{d-m+1} + 2 \sum_{i=2}^m \left(1 + \frac{i-1}{i+1} - \frac{2i}{i+1} \right) A_{i+1} \tau^{d-i+2}. \end{aligned}$$

The parenthetical factor in this last summation vanishes, so the right-hand side of the preceding display boils down to

$$\frac{2(m+1)A_{m+3}}{m+3} \tau^{d-m} + 2\tau^{d+2} - 2A_{m+2} \tau^{d-m+1},$$

and we deduce that

$$(42) \quad d(d+2)B - (d+2)\tau^2 dA \geq \frac{2(m+1)A_{m+3}}{m+3} \tau^{d-m} - 2A_{m+2} \tau^{d-m+1}.$$

Using Jensen's inequality, we obtain $(dA)^{\frac{1}{d}} \geq \int_\delta^{\delta+1} t dt \geq \frac{1}{2}$ for any $\delta \geq 0$. Hence,

putting $\tau = (dA)^{\frac{1}{d}}$ in (42), we arrive at

$$(43) \quad B \geq \frac{(dA)^{\frac{d+2}{d}}}{d} - \frac{2A_{m+2}(dA)^{\frac{d-m+1}{n}}}{d(d+2)} + \frac{2(m+1)A_{m+3}(nA)^{\frac{d-m}{d}}}{d(d+2)(m+3)}.$$

Taking the rescaling of $\zeta(t)$ into consideration, we obtain from (43) the integral inequality

$$\int_0^\infty t^{s+1} \zeta(t) dt \geq \frac{(dA)^{\frac{d+2}{d}} (\zeta(0))^{-\frac{2}{d}}}{d} - \frac{2S_{m+2}(dA)^{\frac{d-m+1}{d}} (\zeta(0))^{\frac{(m+1)d+m-1}{d}}}{d(d+2)m_s^{m+1}} + \frac{2(m+1)S_{m+3}(dA)^{\frac{d-m}{d}} (\zeta(0))^{\frac{(m+2)d+m}{d}}}{d(d+2)(m+3)m_s^{m+2}}.$$

Notice that $\inf_t \zeta(t) \leq \zeta(0) \leq \sup_t \zeta(t)$. Due to (36), the estimate of $\int_0^\infty t^{s+1} \zeta(t) dt$ provides a lower bound

$$\sum_{i=1}^k \lambda_i \geq \omega_d (dA)^{\frac{d+2}{d}} (\zeta(0))^{-\frac{2}{d}} - \frac{2\omega_d A_{m+2}(dA)^{\frac{d-m+1}{d}} (\zeta(0))^{\frac{(m+1)d+m-1}{d}}}{(d+2)m_s^{m+1}} + \frac{2\omega_d(m+1)A_{m+3}(dA)^{\frac{d-m}{d}} (\zeta(0))^{\frac{(m+2)d+m}{d}}}{(d+2)(m+3)m_s^{m+2}}.$$

This is what we required in Proposition 1. \square

The right-hand side of the intermediate lower bound (34) depends on $\zeta(0)$. Our goal is to derive a final eigenvalue inequality that depends solely on d and k . To achieve this, we adopt a strategy of minimizing (34) with respect to $\zeta(0)$ over the interval $[0, \varrho]$, where $\varrho = \frac{d-1}{(2\pi)^d} |\Omega|$ as defined in (13).

Proof of Theorem 3. Using (35) and (38) in combination with integration by parts, we obtain

$$\frac{k}{\omega_n} = d \int_0^\infty t^{d-1} \zeta(t) dt = \int_0^\infty -t^d \zeta'(t) dt = \int_0^\infty t^d \phi(t) dt = dA.$$

Using the fact that $dA = k/\omega_d$, we define $G_1(x)$, $G_2(x)$ and $G(x)$ for $x \in [0, \varrho]$ by

$$G_1(x) = \omega_d^{-\frac{2}{d}} x^{-\frac{2}{d}} k^{\frac{d+2}{d}} + c_2 \frac{2\omega_d^{\frac{m}{d}} (m+1)A_{m+3} x^{\frac{(m+2)d+m}{d}}}{(d+2)(m+3)m_s^{m+2}} k^{\frac{d-m}{d}},$$

$$G_2(x) = -\frac{2\omega_d^{\frac{m-1}{d}} A_{m+2} x^{\frac{(m+1)d+m-1}{d}}}{(d+2)m_s^{m+1}} k^{\frac{d-m+1}{d}},$$

$$G(x) = G_1(x) - G_2(x).$$

Here $0 < c_2 \leq 1$ is a positive number to be determined. Since $m \geq 1$, we conclude that $G_2(x)$ is decreasing on $[0, \varrho]$. By direct calculation, we have

$$\frac{d}{2} \frac{\omega_d^{\frac{2}{d}} G_1'(x)}{k^{\frac{d-m}{d}}} = -k^{\frac{2+d}{d}} x^{-\frac{d+2}{d}} + c_2 \omega_d^{\frac{m+2}{d}} \frac{(m+1)((m+2)d+m)}{(d+2)(m+3)} \frac{A_{m+3}}{m_s^{m+2}} x^{\frac{(m+1)d+m}{d}}.$$

Therefore, when

$$c_2 \leq \frac{k^{\frac{d+2}{d}}}{\omega_d^{\frac{m+2}{d}}} \frac{(d+2)(m+3)}{(m+1)[(m+2)d+m]} \frac{m_s^{m+2}}{A_{m+3}} \varrho^{-\frac{(m+2)(d+1)}{d}},$$

we conclude that $G_1'(x) \leq 0$ on $[0, \varrho]$. Since

$$(44) \quad m_s \geq M = \frac{\sqrt{d^2-d}}{(2\pi)^d \omega_d^{\frac{1}{d}}} |\Omega|^{\frac{d+1}{d}}, \quad \varrho = \frac{d-1}{(2\pi)^d} |\Omega|,$$

we can choose $c_1 = \min\{1, a_1\}$, where a_1 is defined by

$$(45) \quad a_1 = \frac{(d+2)(m+3)}{(m+1)[(m+2)d+m]} \frac{(2\pi)^{m+2}}{A_{m+3}} \frac{d^{\frac{m+2}{2}}}{(d-1)^{\frac{m(d+2)+2m}{2d}}},$$

to obtain

$$(46) \quad \sum_{i=1}^k \lambda_i \geq \omega_d^{-\frac{2}{d}} \varrho^{-\frac{2}{d}} k^{\frac{d+2}{d}} - \frac{2\omega_d^{\frac{m-1}{d}} A_{m+2} \varrho^{\frac{(m+1)d+m-1}{d}}}{(d+2)M^{m+1}} k^{\frac{d-m+1}{d}} + c_1 \frac{2\omega_d^{\frac{m}{d}} (m+1) A_{m+3} \varrho^{\frac{(m+2)d+m}{d}}}{(d+2)(m+3)M^{m+2}} k^{\frac{d-m}{d}},$$

where ϱ and M are defined in (13). Similarly, we can split $G(x)$ as a sum of two functions:

$$G(x) = g_1(x) + g_2(x), \quad x \in [0, \varrho],$$

where g_1 and g_2 are defined by $g_1(x) = \omega_d^{-\frac{2}{d}} x^{-\frac{2}{d}} k^{\frac{d+2}{d}}$ and

$$g_2(x) = c_2 \frac{2\omega_d^{\frac{m}{d}} (m+1) A_{m+3} x^{\frac{(m+2)d+m}{d}}}{(d+2)(m+3)m_s^{m+2}} k^{\frac{d-m}{d}} - \frac{2\omega_d^{\frac{m-1}{d}} A_{m+2} x^{\frac{(m+1)d+m-1}{d}}}{(d+2)m_s^{m+1}} k^{\frac{d-m+1}{d}}.$$

Obviously, $g_1(x)$ is decreasing on $[0, \varrho]$. Rewriting $g_2(x)$ as

$$\frac{d(d+2)}{2} \frac{m_s}{\omega_d^{\frac{m}{d}} k^{\frac{d-m}{d}}} g_2(x) = c_2 \frac{[(m+2)d+m](m+1)A_{m+3}}{(m+3)m_s} x^{\frac{(m+1)d+m}{d}} - ((m+1)d+m-1)\omega_d^{-\frac{1}{d}} A_{m+2} k^{\frac{1}{d}} x^{\frac{(d+1)m-1}{d}}$$

allows us to observe that if

$$c_2 \leq \frac{[(m+1)d+m-1](m+3)}{[(m+2)d+m](m+1)} \frac{A_{m+2}}{A_{m+3}} \frac{m_s}{\omega_d^{1/d}} \varrho^{-\frac{d+1}{d}},$$

then $g'_2(x) \leq 0$ on $[0, \varrho]$. Using (44) again, we choose $c_3 = \min\{1, a_2\}$, where a_2 is defined by

$$(47) \quad a_2 = \frac{\sqrt{2}d^{\frac{1}{d}}}{2(d-1)^{\frac{d+2}{2d}}} \frac{[(m+1)d+m-1](m+3)}{[(m+2)d+m](m+1)} \frac{A_{m+2}}{A_{m+3}},$$

to derive that

$$(48) \quad \sum_{i=1}^k \lambda_i \geq \omega_d^{-\frac{2}{d}} \varrho^{-\frac{2}{d}} k^{\frac{d+2}{d}} - \frac{2\omega_d^{\frac{m-1}{d}} A_{m+2} \varrho^{\frac{(m+1)d+m-1}{d}}}{(d+2)M^{m+1}} k^{\frac{d-m+1}{d}} + c_3 \frac{2\omega_d^{\frac{m}{d}} (m+1) A_{m+3} \varrho^{\frac{(m+2)d+m}{d}}}{(d+2)(m+3)M^{m+2}} k^{\frac{d-m}{d}}.$$

In light of equations (46) and (48), we can use $c = \min\{1, \max\{a_1, a_2\}\}$, where a_1 and a_2 are defined in (45) and (47), to get

$$(49) \quad \sum_{i=1}^k \lambda_i \geq \omega_d^{-\frac{2}{d}} \varrho^{-\frac{2}{d}} k^{\frac{d+2}{d}} - \frac{2\omega_d^{\frac{m-1}{d}} A_{m+2} \varrho^{\frac{(m+1)d+m-1}{d}}}{(d+2)M^{m+1}} k^{\frac{d-m+1}{d}} + c \frac{2\omega_d^{\frac{m}{d}} (m+1) A_{m+3} \varrho^{\frac{(m+2)d+m}{d}}}{(d+2)(m+3)M^{m+2}} k^{\frac{d-m}{d}}.$$

Finally, we use

$$(50) \quad \omega_d^{-\frac{2}{d}} \varrho^{-\frac{2}{d}} k^{\frac{d+2}{d}} = 4\pi \left(\frac{\Gamma(1+\frac{d}{2})}{(d-1)|\Omega|} \right)^{\frac{2}{d}} k^{1+\frac{2}{d}}$$

to complete the proof of Theorem 3. □

Proof of Corollary 1. An argument similar to that of [11] gives

$$d(d+2)B - d(d+2)A\tau^2 + 2\tau^{d+2} \geq \frac{1}{6}(\tau^d) + \frac{1}{9}(\tau^{d-1}) + \frac{3}{80}\tau^{d-2}.$$

Hence, we get

$$\sum_{i=1}^k \lambda_i \geq \frac{d\omega_d^{-\frac{2}{d}} \zeta(0)^{-\frac{2}{d}}}{d+2} k^{\frac{d+2}{d}} + \frac{\zeta(0)^2}{6(d+2)m_s^2} k + \frac{c_4 \omega_d^{-\frac{1}{d}} \zeta(0)^{\frac{3d+1}{d}}}{(d+2)m_s^3} k^{\frac{d-1}{d}} + \frac{c_5 \omega_d^{-\frac{1}{d}} \zeta(0)^{\frac{4d+2}{d}}}{(n+2)m_s^4} k^{\frac{d-2}{d}},$$

where $0 < c_4 \leq \frac{1}{9}$ and $0 < c_5 \leq \frac{3}{80}$ are two constants to be chosen. Following an approach analogous to the discussion in the proof of Theorem 3, we can choose $c_4 = \frac{1}{9}$ and $c_5 = \frac{3}{80}$. Using the equality in (50) completes the proof. \square

6. Proof of Theorem 4

The following lemma from [10] plays a pivotal role in establishing our eigenvalue inequality and makes use of the piecewise continuous function

$$\Phi_s(r) = \begin{cases} m_s & \text{if } 0 \leq r \leq s, \\ m_s - \varrho(r - s) & \text{if } s \leq r \leq s + m_s/\varrho, \\ 0 & \text{if } s + m_s/\varrho \leq r, \end{cases}$$

where m_s and ϱ are defined in (13).

Lemma 4 [10]. *Let $\alpha > 0$ be a real number and $\Psi(r)$ be a decreasing and absolutely continuous function such that*

$$\int_0^\infty r^\alpha \Psi(r) dr = \int_0^\infty r^\alpha \Phi_s(r) dr, \quad 0 \leq \Psi \leq m_s, \quad -L < \Psi' < 0.$$

Then, for any $\beta \geq \alpha$, the following integral inequality holds:

$$\int_0^\infty r^\beta \Psi(r) dr \geq \int_0^\infty r^\beta \Phi_s(r) dr.$$

Moreover, for any $\gamma \geq 0$, we have

$$\int_0^\infty r^\gamma \Phi_s(r) dr = \frac{\varrho^{\gamma+2}}{(\gamma+1)(\gamma+2)m_s^{\gamma+1}} ((t+1)^{\gamma+2} - t^{\gamma+2}), \quad s = \frac{tm_s}{\varrho}.$$

Proof of Theorem 4. Let m_s be as in (13). Applying Lemma 4 to $\Psi(x) = \zeta(x)$ and setting $\alpha = d - 1$, we obtain

$$\int_0^\infty t^{d-1} \zeta(t) dt = \frac{k}{d\omega_d} = \frac{\varrho^{d+1}}{d(d+1)m_s^d} ((t+1)^{d+1} - t^{d+1}),$$

which in turn implies that t is the unique root of the equation

$$(t+1)^{d+1} - t^{d+1} = T := \frac{(d+1)m_s^d}{\omega_d \varrho^{d+1}} k.$$

Then Lemma 4 implies

$$(51) \quad \int_0^\infty t^{d+1} \zeta(t) dt \geq \frac{\varrho^{d+3}}{(d+2)(d+3)m_s^{d+2}} ((t+1)^{d+3} - t^{d+3}).$$

One can easily check that T satisfies

$$T \geq 1, \quad T \geq \frac{d+1}{d-1} \frac{(4\pi)^d}{\omega_d^2} \left(\frac{d^2}{(d-1)(d+2)} \right)^{\frac{d}{2}}.$$

In particular, when $d = 3$, we consider the equation

$$(t+1)^4 - t^4 = T = \frac{4\rho^3}{\omega_3 m_s^4} k,$$

whose positive root is

$$t = \frac{1}{2} \left(\left(T + \sqrt{T^2 + \frac{1}{27}} \right)^{\frac{1}{3}} - \left(-T + \sqrt{T^2 + \frac{1}{27}} \right)^{\frac{1}{3}} - 1 \right).$$

From (36), we obtain our desired result as

$$(52) \quad \sum_{i=1}^k \lambda_i \geq \frac{2\pi\rho^6}{15m_s^5} ((t+1)^6 - t^6).$$

Considering the Taylor series of t , we arrive at

$$t \geq -\frac{1}{2} + 4^{-\frac{1}{3}} T^{\frac{1}{3}} - \frac{1}{6} 2^{-\frac{1}{3}} T^{-\frac{1}{3}} + \frac{1}{324} 4^{-\frac{1}{3}} T^{-\frac{5}{3}} + \frac{1}{1944} 2^{-\frac{1}{3}} T^{-\frac{7}{3}} - \frac{1}{26244} 4^{-\frac{1}{3}} T^{-\frac{11}{3}} - \frac{1}{629856} 2^{-\frac{1}{3}} T^{-\frac{11}{3}},$$

where we used the fact that $T \geq 1$. Hence, we obtain

$$(t+1)^6 - t^6 \geq \frac{3}{4} 2^{-\frac{1}{3}} T^{\frac{5}{3}} + \frac{5}{8} T - \frac{11}{24} 2^{\frac{2}{3}} T^{\frac{1}{3}} + \frac{659}{10000} T^{-\frac{1}{3}}.$$

Putting the lower bound of $(t+1)^6 - t^6$ into (52), we get

$$\sum_{i=1}^k \lambda_i \geq \frac{3}{5} \left(\frac{2\pi^3}{2\omega_3 |\Omega|} \right)^{\frac{2}{3}} k^{\frac{5}{3}} + \frac{1}{24} \frac{|\Omega|}{I(\Omega)} k - \frac{11\pi}{180} \frac{\rho^{\frac{14}{3}}}{\omega_3^{\frac{1}{3}} m_s^4} k^{\frac{1}{3}} + \frac{659\pi\omega_3^{\frac{1}{3}}}{75000 \cdot 4^{\frac{1}{3}}} \frac{\rho^{\frac{22}{3}}}{m_s^6} k^{-\frac{1}{3}}.$$

This concludes the proof of Theorem 4. □

Acknowledgements

The authors wish to thank the referees for carefully reading this paper and making numerous valuable suggestions and corrections. Z. Ji’s research was supported by the National Natural Science Foundation of China, Grant no. 12501068, and Zhejiang Provincial Natural Science Foundation of China, Grant no. LQN25A010001. T. Yolcu acknowledges support from the Caterpillar Fellowship (no. 2511045) during the preparation of this article. Ji expresses his deep gratitude to Prof. Hongwei Xu, whose guidance and mentorship in mathematics have profoundly shaped his understanding and growth as a researcher. It is with great respect and appreciation that we dedicate this paper to him.

References

- [1] M. S. Ashbaugh, “The universal eigenvalue bounds of Payne–Pólya–Weinberger, Hile–Protter, and H. C. Yang”, *Proc. Indian Acad. Sci. Math. Sci.* **112**:1 (2002), 3–30. MR
- [2] M. S. Ashbaugh and R. D. Benguria, “Isoperimetric inequalities for eigenvalues of the Laplacian”, pp. 105–139 in *Spectral theory and mathematical physics: a Festschrift in honor of Barry Simon’s 60th birthday*, vol. 1, Proc. Sympos. Pure Math. **76**:1, Amer. Math. Soc., 2007. MR
- [3] P. Constantin and C. Foias, *Navier–Stokes equations*, University of Chicago Press, 1988. MR
- [4] C. Foias, O. Manley, R. Rosa, and R. Temam, *Navier–Stokes equations and turbulence*, Encyclopedia of Mathematics and its Applications **83**, Cambridge University Press, 2001. MR
- [5] E. M. Harrell, II and S. Yıldırım Yolcu, “Eigenvalue inequalities for Klein–Gordon operators”, *J. Funct. Anal.* **256**:12 (2009), 3977–3995. MR
- [6] J. Hu and Y. Huang, “Lower bounds for eigenvalues of the Stokes operator”, *Adv. Appl. Math. Mech.* **5**:1 (2013), 1–18. MR
- [7] D. Hundertmark, “Some bound state problems in quantum mechanics”, pp. 463–496 in *Spectral theory and mathematical physics: a Festschrift in honor of Barry Simon’s 60th birthday*, vol. 1, Proc. Sympos. Pure Math. **76**:1, Amer. Math. Soc., 2007. MR
- [8] A. A. Ilyin, “Attractors for Navier–Stokes equations in domains with finite measure”, *Nonlinear Anal.* **27**:5 (1996), 605–616. MR
- [9] A. A. Ilyin, “On the spectrum of the Stokes operator”, *Funktsional. Anal. i Prilozhen.* **43**:4 (2009), 14–25. In Russian; translated in *Funct. Anal. Appl.* **43**:4, 254–263 (2009). MR
- [10] A. A. Ilyin, “Lower bounds for the spectrum of the Laplace and Stokes operators”, *Discrete Contin. Dyn. Syst.* **28**:1 (2010), 131–146. MR
- [11] Z. Ji and H. Xu, “Lower bounds for eigenvalues of Laplacian operator and the clamped plate problem”, *Calc. Var. Partial Differential Equations* **62**:6 (2023), art. id. 175, 27 pp. MR
- [12] J. P. Kelliher, “Eigenvalues of the Stokes operator versus the Dirichlet Laplacian in the plane”, *Pacific J. Math.* **244**:1 (2010), 99–132. MR
- [13] H. Kovařík, S. Vugalter, and T. Weidl, “Two-dimensional Berezin–Li–Yau inequalities with a correction term”, *Comm. Math. Phys.* **287**:3 (2009), 959–981. MR
- [14] P. Li and S. T. Yau, “On the Schrödinger equation and the eigenvalue problem”, *Comm. Math. Phys.* **88**:3 (1983), 309–318. MR
- [15] E. H. Lieb and M. Loss, *Analysis*, 2nd ed., Graduate Studies in Mathematics **14**, Amer. Math. Soc., 2001. MR
- [16] A. D. Melas, “A lower bound for sums of eigenvalues of the Laplacian”, *Proc. Amer. Math. Soc.* **131**:2 (2003), 631–636. MR
- [17] G. Métivier, “Valeurs propres d’opérateurs définis par la restriction de systèmes variationnels à des sous-espaces”, *J. Math. Pures Appl.* (9) **57**:2 (1978), 133–156. MR
- [18] G. Pólya, “On the eigenvalues of vibrating membranes”, *Proc. London Math. Soc.* (3) **11** (1961), 419–433. MR
- [19] R. Temam, *Navier–Stokes equations: Theory and numerical analysis*, 3rd ed., Studies in Mathematics and its Applications **2**, North-Holland, Amsterdam, 1984. MR
- [20] V. Vougalter, “Sharp semiclassical bounds for the moments of eigenvalues for some Schrödinger type operators with unbounded potentials”, *Math. Model. Nat. Phenom.* **8**:1 (2013), 237–245. MR

- [21] T. Weidl, “Improved Berezin–Li–Yau inequalities with a remainder term”, pp. 253–263 in *Spectral theory of differential operators*, Amer. Math. Soc. Transl. Ser. 2 **225**, Amer. Math. Soc., 2008. MR
- [22] S. Yıldırım Yolcu, “An improvement to a Berezin–Li–Yau type inequality”, *Proc. Amer. Math. Soc.* **138**:11 (2010), 4059–4066. MR
- [23] S. Yıldırım Yolcu and T. Yolcu, “Bounds for the eigenvalues of the fractional Laplacian”, *Rev. Math. Phys.* **24**:3 (2012), art. id. 1250003, 18 pp. MR
- [24] S. Yıldırım Yolcu and T. Yolcu, “Multidimensional lower bounds for the eigenvalues of Stokes and Dirichlet Laplacian operators”, *J. Math. Phys.* **53**:4 (2012), art. id. 043508, 17 pp. MR
- [25] S. Yıldırım Yolcu and T. Yolcu, “Estimates for the sums of eigenvalues of the fractional Laplacian on a bounded domain”, *Commun. Contemp. Math.* **15**:3 (2013), art. id. 1250048, 15 pp. MR
- [26] S. Yıldırım Yolcu and T. Yolcu, “Estimates on the eigenvalues of the clamped plate problem on domains in Euclidean spaces”, *J. Math. Phys.* **54**:8 (2013), art. id. 043515. MR
- [27] S. Yıldırım Yolcu and T. Yolcu, “Refined eigenvalue bounds on the Dirichlet fractional Laplacian”, *J. Math. Phys.* **56**:7 (2015), art. id. 073506, 12 pp. MR
- [28] S. Yıldırım Yolcu and T. Yolcu, “Sharper estimates on the eigenvalues of Dirichlet fractional Laplacian”, *Discrete Contin. Dyn. Syst.* **35**:5 (2015), 2209–2225. MR
- [29] T. Yolcu, “Refined bounds for the eigenvalues of the Klein–Gordon operator”, *Proc. Amer. Math. Soc.* **141**:12 (2013), 4305–4315. MR

Received July 12, 2025. Revised February 11, 2026.

ZHENGCHAO JI
DEPARTMENT OF MATHEMATICS
CHINA JILIANG UNIVERSITY
HANGZHOU
CHINA
jizhengchao@zju.edu.cn

TÜRKAY YOLCU
DEPARTMENT OF MATHEMATICS
BRADLEY UNIVERSITY
PEORIA, IL
UNITED STATES
tyolcu@bradley.edu

PACIFIC JOURNAL OF MATHEMATICS

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

msp.org/pjm

EDITORS

Don Blasius (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Matthias Aschenbrenner
Fakultät für Mathematik
Universität Wien
Vienna, Austria
matthias.aschenbrenner@univie.ac.at

Vyjayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Atsushi Ichino
Department of Mathematics
Kyoto University
Kyoto 606-8502, Japan
atsushi.ichino@gmail.com

Kefeng Liu
School of Sciences
Chongqing University of Technology
Chongqing 400054, China
liu@math.ucla.edu

Sucharit Sarkar
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
sucharit@math.ucla.edu

Dimitri Shlyakhtenko
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
shlyakht@ipam.ucla.edu

Ruixiang Zhang
Department of Mathematics
University of California
Berkeley, CA 94720-3840
ruixiang@berkeley.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org


See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2026 is US \$710/year for the electronic version, and \$965/year for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004, U.S.A. The Pacific Journal of Mathematics is indexed by Mathematical Reviews, Zentralblatt MATH, PASCAL CNRS Index, Referativnyi Zhurnal, Current Mathematical Publications and Web of Knowledge (Science Citation Index).

The Pacific Journal of Mathematics (ISSN 1945-5844 electronic, 0030-8730 printed) at Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Mathematical Sciences Publishers, 2000 Allston Way # 59, Berkeley, CA 94701-4004.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2026 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 343 No. 1 July 2026

The generic extension map and modular standard modules JOHANNES DROSCHL	1
Transverse minimal foliations on unit tangent bundles and applications SÉRGIO R. FENLEY and RAFAEL POTRIE	39
Refined bounds for the eigenvalues of the Stokes operator ZHENGCHAO JI and TÜRKAY YOLCU	119
The class \mathcal{Q} and mixture distributions with dominated continuous singular parts ALEXEY A. KHARTOV	139
Data for Shimura varieties intersecting the Torelli locus WANLIN LI, ELENA MANTOVAN and RACHEL PRIES	179
The derived series of GGS groups J. MORITZ PETSCHICK	211
Property QT of relatively hierarchically hyperbolic groups BINGXUE TAO	231