

PROBABILITY and MATHEMATICAL PHYSICS

**OBSERVABLES OF COLOURED STOCHASTIC VERTEX MODELS AND
THEIR POLYMER LIMITS**

ALEXEI BORODIN AND MICHAEL WHEELER

1:1
2020



OBSERVABLES OF COLOURED STOCHASTIC VERTEX MODELS AND THEIR POLYMER LIMITS

ALEXEI BORODIN AND MICHAEL WHEELER

In the context of the coloured stochastic vertex model in a quadrant, we identify a family of observables whose averages are given by explicit contour integrals. The observables are certain linear combinations of q -moments of the coloured height functions of the model. In a polymer limit, this yields integral representations for moments of partition functions of strict-weak, semidiscrete Brownian, and continuum Brownian polymers with varying beginning and ending points of the polymers.

1. Introduction	205
2. Preliminaries	212
3. Extensions	224
4. Cauchy identities	231
5. Integral representations for f_μ	236
6. Observables of stochastic lattice models	244
7. Limit to polymers	253
Acknowledgments	263
References	263

1. Introduction

At least since the work of Kardar [1987], moments of polymer partition functions and related quantities have been an indispensable tool in analyzing models from the so-called Kardar–Parisi–Zhang (KPZ) universality class in (1+1) dimensions [Kardar et al. 1986], such as various (integrable) models of directed polymers in a random environment, exclusion and zero-range processes, and random growth models. An advanced (although nonrigorous) replica analysis of moments allowed Calabrese, Le Doussal and Rosso [2010] and Dotsenko [2010] to derive the long-time asymptotics of the so-called narrow wedge solution to the KPZ equation. An alternative approach to the asymptotics of this solution based on the pioneering work of Tracy and Widom [2008a; 2008b; 2009] was developed rigorously by Amir, Corwin and Quastel [2011] and in the physics literature by Sasamoto and Spohn [2010]. (These two approaches were mostly reconciled by Borodin, Corwin and Sasamoto [2014].) The moment method became rigorous with the appearance of explicit integral representations for q -moments of suitable q -deformed models together with the asymptotic analysis of their generating functions in [Borodin and Corwin 2014]. (A simpler approach to the asymptotics through moments was later suggested in [Borodin 2018])

MSC2010: 60K35.

Keywords: observables, stochastic vertex models, polymer models.

and [Borodin and Olshanski 2017].) Many papers with analysis of q -moments of various integrable probabilistic systems have been written since then. The stochastic six vertex model, first introduced by Gwa and Spohn [1992], was also found to be accessible via such a route in work of Borodin, Corwin and Gorin [2016b]; see also works by Borodin, Corwin, Petrov and Sasamoto [2015a; 2015b], Corwin and Petrov [2016] and Borodin and Petrov [2017; 2018a] for various approaches to the q -moments of this model.

Kuniba, Mangazeev, Maruyama and Okado [2016] introduced Yang–Baxter integrable *coloured* stochastic vertex models; see also works by Bosnjak and Mangazeev [2016] and Aggarwal, Borodin and Bufetov [2019]. Colours correspond to the simple roots in the underlying quantum affine algebra $U_q(\widehat{\mathfrak{sl}}_{n+1})$, with $n = 1$ corresponding to the *colourless* or rank-1 case considered previously.

Our recent paper [Borodin and Wheeler 2018] offered an extensive algebraic analysis of these coloured models and uncovered certain distributional correspondences between coloured and (much more studied) colourless ones. Such correspondences were further extended by Borodin and Bufetov [2019] and Borodin, Gorin and Wheeler [2019], and they gave access to various unknown marginals of the coloured models. However, so far no explicit formulas for observables of the coloured models have been found, apart from those that arise through matching with colourless models. The primary goal of this paper is to remedy this fact.

We obtain explicit integral representations for certain linear combinations of q -moments of coloured height functions for the coloured stochastic vertex model in a quadrant. Further, following a path worked out in [Borodin et al. 2019], we degenerate the (fully fused) coloured vertex model to directed polymers, thus obtaining formulas for joint moments of polymer partition functions with different starting points in the same noise field. The limiting objects include the KPZ equation (equivalently, the continuum Brownian polymer), the O’Connell–Yor semi-discrete Brownian polymer [2001], the strict-weak or Gamma polymer of Corwin, Seppäläinen and Shen [2015] and O’Connell and Ortmann [2015], and the Beta polymer of Barraquand and Corwin [2017].

Let us describe our results in more detail.

The coloured stochastic vertex model in a quadrant can be viewed as a Markovian recipe of constructing random coloured up-right paths in $\mathbb{Z}_{\geq 1} \times \mathbb{Z}_{\geq 1}$ with the colours labelled by natural numbers. Let us also initially assume that no horizontal edge of the lattice can be occupied by more than a single path; this restriction will eventually be removed. The model depends on a quantization parameter $q \in \mathbb{C}$, a spin parameter $s \in \mathbb{C}$, and row *rapidities* denoted by $x_1, x_2, \dots \in \mathbb{C}$.

Along the boundary of the quadrant, we demand that no paths enter the quadrant from the bottom. On the other hand, a single coloured path enters the quadrant from the left in each row. We assume that the colours of the paths entering on the left are weakly increasing in the upward direction, and denote by $\lambda_1 \geq 0$ the number of paths of colour 1, by $\lambda_2 \geq 0$ the number of paths of colour 2, etc. Let us also denote by

$$\ell_k = \lambda_1 + \dots + \lambda_k, \quad k \geq 1, \tag{1-1}$$

the partial sums of this sequence and also set $\ell_0 = 0$.

Once the paths are specified along the boundary, they progress in the up-right direction within the quadrant using certain interaction probabilities, also known as *vertex weights*. For each vertex of the

$\begin{array}{c} \mathbf{I} \\ \uparrow \\ 0 \text{ --- } \mathbf{I} \\ \rightarrow 0 \\ \mathbf{I} \end{array}$ $\frac{1-sxq^{I_{\geq 1}}}{1-sx}$	$\begin{array}{c} \mathbf{I} \\ \uparrow \\ i \text{ --- } \mathbf{I} \\ \rightarrow i \\ \mathbf{I} \end{array}$ $\frac{s(sq^{I_i} - x)q^{I_{\geq i+1}}}{1-sx}$	$\begin{array}{c} \mathbf{I}_i^- \\ \uparrow \\ 0 \text{ --- } \mathbf{I} \\ \rightarrow i \\ \mathbf{I} \end{array}$ $\frac{sx(q^{I_i} - 1)q^{I_{\geq i+1}}}{1-sx}$
$\begin{array}{c} \mathbf{I}_i^+ \\ \uparrow \\ i \text{ --- } \mathbf{I} \\ \rightarrow 0 \\ \mathbf{I} \end{array}$ $\frac{1-s^2q^{I_{\geq 1}}}{1-sx}$	$\begin{array}{c} \mathbf{I}_{ij}^{+-} \\ \uparrow \\ i \text{ --- } \mathbf{I} \\ \rightarrow j \\ \mathbf{I} \end{array}$ $\frac{sx(q^{I_j} - 1)q^{I_{\geq j+1}}}{1-sx}$	$\begin{array}{c} \mathbf{I}_{ji}^{+-} \\ \uparrow \\ j \text{ --- } \mathbf{I} \\ \rightarrow i \\ \mathbf{I} \end{array}$ $\frac{s^2(q^{I_i} - 1)q^{I_{\geq i+1}}}{1-sx}$

Table 1. The vertex weights.

lattice, once we know the colours of the entering paths along the bottom and left adjacent edges, we decide on the colours of the exiting paths along the top and right edges according to those probabilities.¹ They are given by Table 1, where it is assumed that x is the rapidity of the row to which the vertex belongs, $1 \leq i < j$, $\mathbf{I} = (I_1, I_2, \dots)$ denotes a vector whose coordinates I_k are equal to the number of paths of colour k that enter the vertex from the bottom, and $\mathbf{I}_i^\pm = \mathbf{I} \pm \mathbf{e}_i$ with \mathbf{e}_i being the standard basis vector with 1 as its i -th coordinate and all other coordinates equal to 0. We also use the notation $\mathbf{I}_{ab}^{+-} = \mathbf{I} + \mathbf{e}_a - \mathbf{e}_b$ and $\mathbf{I}_{\geq a} = I_a + I_{a+1} + \dots$. The weight of any vertex that does not fall into one of the six categories in Table 1 is set to zero. See Section 2 for the origin of these weights.

The *stochasticity* of the weights encodes the fact that these weights add up to 1 when summed over all possible states of the outgoing edges with the states of incoming edges being fixed. When the parameters of the model are such that all the weights are nonnegative, we obtain bona fide transition probabilities. However, if we agree to deal with complex-valued discrete distributions, we will not actually need the positivity assumption for our main algebraic result.


Let us now fix $n \geq 1$ and focus on the state of the model between row n and row $n + 1$. That is, let us record the locations where the paths of colours 1, 2, \dots exit the n -th row upwards as an n -dimensional coloured vector (equivalently, a *coloured composition*) ν with coordinates in $\mathbb{Z}_{\geq 1}$. By colour conservation that our weights observe, the counts of different colours in such a vector are provided by a (finite) sequence $\lambda = (\lambda_1, \lambda_2, \dots)$ corresponding to the colours that enter via the left edges of the first n rows; we call λ the *labelling composition*.² We will write the coordinates of ν as

$$\nu = (\nu_1 \geq \dots \geq \nu_{\ell_1} \mid \nu_{\ell_1+1} \geq \dots \geq \nu_{\ell_2} \mid \nu_{\ell_2+1} \geq \dots \geq \nu_{\ell_3} \mid \dots), \quad (1-2)$$

where the groups separated by vertical bars list the coordinates of the paths between rows n and $n + 1$ of colour 1, colour 2, etc. See Figure 1 for an example.

¹Such decisions at different vertices are independent.

²For convenience of notation, we assume that the left-incoming colours in rows n and $n + 1$ are different, so that we do not need to split the last coordinate in this sequence.



$$\kappa = (1, 1, 2, 0)$$

$$\mu = (4 | 4 | 2, 2) = 1^0 2^{(0,0,2,0)} 3^0 4^{(1,1,0,0)} 5^0 6^0 \dots$$

$$\lambda = (2, 1, 2, 1)$$

$$\nu = (6, 2 | 7 | 4, 2 | 5)$$

$$H_1^\mu(4) = H_2^\mu(4) = 1$$

$$H_3^\mu(4) = H_4^\mu(4) = 0$$

$$H_1^\nu(4) = H_2^\nu(4) = H_3^\nu(4) = H_4^\nu(4) = 1$$

$$H_1^{\nu/\mu} = H_2^{\nu/\mu}(4) = 0$$

$$H_3^{\nu/\mu}(4) = H_4^{\nu/\mu}(4) = 1$$

Figure 1. A possible configuration of the coloured vertex model.

Our observables on random ν 's are also indexed by coloured compositions. Let us fix a composition $\kappa = (\kappa_1, \kappa_2, \dots)$ satisfying $\kappa \leq \lambda$ coordinate-wise (otherwise the observable vanishes identically), and a coloured composition μ with coordinates in $\mathbb{Z}_{\geq 1}$ whose colour counts are given by κ . It is also convenient to parametrize μ differently by writing $\mu = 1^{m^{(1)}} 2^{m^{(2)}} \dots$, where the vectors $m^{(j)} = (m_1^{(j)}, m_2^{(j)}, \dots)$ are such that their coordinates $m_i^{(j)}$ count the number of parts of μ of colour i equal to j ; see Figure 1.

Yet another way to describe coloured compositions is through their *coloured height functions* defined as follows; see Figure 1:

$$H_i^\rho(x) = \#\{j : \text{colour}(\rho_j) = i, \rho_j \geq x\};$$

$$H_{>i}^\rho(x) = \sum_{k>i} H_k^\rho(x), \quad H_{\geq i}^\rho(x) = \sum_{k \geq i} H_k^\rho(x); \quad H_*^{\nu/\mu} \equiv H_*^\nu - H_*^\mu.$$

Here $\text{colour}(\rho_j)$ refers to the number of the block in the splitting of the form (1-2) for the coloured composition ρ that ρ_j belongs to.

For κ -coloured μ as above, let us now define an observable \mathcal{O}_μ , whose values on λ -coloured ν 's are given by

$$\mathcal{O}_\mu(\nu) = \prod_{i,j \geq 1} q^{m_i^{(j)} H_{>i}^{\nu/\mu}(j+1)} \binom{H_i^{\nu/\mu}(j+1)}{m_i^{(j)}}_q$$

$$= \prod_{i,j \geq 1} \frac{(q^{H_{>i}^{\nu/\mu}(j+1)} - q^{H_{\geq i}^{\nu/\mu}(j+1)}) (q^{H_{>i}^{\nu/\mu}(j+1)} - q^{H_{\geq i}^{\nu/\mu}(j+1)-1}) \dots (q^{H_{>i}^{\nu/\mu}(j+1)} - q^{H_{\geq i}^{\nu/\mu}(j+1)-m_i^{(j)}+1})}{(q; q)_{m_i^{(j)}}}.$$

In the *rainbow case* $\lambda = (1, 1, \dots, 1)$ of all colours being different, \mathcal{O}_μ simplifies to

$$\mathcal{O}_\mu^{\text{rainbow}}(\nu) = \prod_{i,j : m_i^{(j)}=1} \mathbf{1}_{H_i^{\nu/\mu}(j+1)=1} q^{H_{>i}^{\nu/\mu}(j+1)} = \prod_{i,j : m_i^{(j)}=1} \frac{q^{H_{>i+1}^{\nu/\mu}(j+1)} - q^{H_{>i}^{\nu/\mu}(j+1)}}{q - 1},$$

and in the *colour-blind case* $\lambda = (n)$ it is given by

$$\mathcal{O}_\mu^{\text{colour-blind}}(v) = \frac{(1 - q^{H^v(\mu_1+1)})(1 - q^{H^v(\mu_2+1)-1}) \cdots (1 - q^{H^v(\mu_m+1)-m+1})}{\prod_{j \geq 1} (q; q)_{\text{mult}_j(\mu)}},$$

where $H^v = H_{\geq 1}^v$ is the colour-blind height function, $\text{mult}_j(\mu) = \#\{i : \mu_i = j\}$, and $m = \ell(\mu)$ is the number of parts in μ .

In order to write down an integral representation for the average of \mathcal{O}_μ , we need to introduce certain rational functions $f_\mu(\kappa; z_1, \dots, z_m)$ with the number of variables $m = \ell(\mu)$ equal to the number of parts of μ . In the rainbow case of pairwise distinct colours, coloured compositions are the same as uncoloured ones, and for anti-dominant compositions $\delta = (\delta_1 \leq \delta_2 \leq \dots \leq \delta_m)$ these functions are completely factorized:

$$f_\delta(z_1, \dots, z_m) = \frac{\prod_{j \geq 0} (s^{2j}; q)_{\text{mult}_j(\delta)}}{\prod_{i=1}^m (1 - sz_i)} \prod_{i=1}^m \left(\frac{z_i - s}{1 - sz_i} \right)^{\delta_i}.$$

Note that we dropped κ from the notation for f_δ as it plays no role in the rainbow situation. For non-anti-dominant μ 's, one way to define f_μ is by the following recursion that allows us to move step by step from anti-dominant compositions toward the dominant ones: If $\mu_i < \mu_{i+1}$ for some $1 \leq i \leq m - 1$, then

$$T_i \cdot f_\mu(z_1, \dots, z_m) = f_{(\mu_1, \dots, \mu_{i+1}, \mu_i, \dots, \mu_m)}(z_1, \dots, z_m),$$

where

$$T_i \equiv q - \frac{z_i - qz_{i+1}}{z_i - z_{i+1}}(1 - \mathfrak{s}_i), \quad 1 \leq i \leq m - 1,$$

with elementary transpositions \mathfrak{s}_i acting by

$$\mathfrak{s}_i \cdot h(z_1, \dots, z_m) := h(z_1, \dots, z_{i+1}, z_i, \dots, z_m),$$

are the Demazure–Lusztig operators of the polynomial representation of the Hecke algebra of type A_{m-1} . The rainbow functions f_μ were thoroughly studied in [Borodin and Wheeler 2018] under the name of *spin nonsymmetric Hall-Littlewood functions*; they also play a central role in the present work.

For generic, not necessarily rainbow κ and κ -coloured μ , f_μ is defined as a suitable sum of rainbow functions. Concretely, let $\theta : \{1, \dots, m\} \rightarrow \{1, 2, \dots\}$ be the unique monotone map such that $|\theta^{-1}(j)| = \kappa_j$ for all $j \geq 1$. Then for any composition κ with m parts, we can define a κ -coloured composition $\theta_*(\kappa)$ by colouring the coordinate κ_i by colour j if and only if $\theta(i) = j$, for all $i = 1, \dots, m$. With this, we set

$$f_\mu(\kappa, z_1, \dots, z_m) = \sum_{\kappa : \theta_*(\kappa) = \mu} f_\kappa(z_1, \dots, z_m).$$

Finally, denote by $c_1 < \dots < c_\alpha$ the colours of parts of μ , and denote by $\mathfrak{m}_1, \dots, \mathfrak{m}_\alpha \geq 1$ the number of parts of μ of colours c_1, \dots, c_α , respectively (\mathfrak{m}_j 's are simply renumbered nonzero coordinates of κ). Set $\mathfrak{m}[a, b] = \mathfrak{m}_a + \mathfrak{m}_{a+1} + \dots + \mathfrak{m}_b$ and recall that

$$\ell_k = \lambda_1 + \dots + \lambda_k \quad \text{for } k \geq 1, \ell_0 = 0.$$

We can now state the main result of this paper.

Theorem 1.1. *With the above notation, we have*

$$\begin{aligned} \mathbb{E} \mathcal{O}_\mu &= \frac{q^{\sum_{u \geq 1} \sum_{i > j} m_i^{(u)} m_j^{(u)}} (-s)^{\mu_1 + \mu_2 + \dots}}{\prod_{j \geq 1} (s^2; q)_{|m^{(j)}|} (2\pi \sqrt{-1})^m} \oint \dots \oint_{\text{around}\{x_j^{-1}\}} \prod_{1 \leq i < j \leq m} \frac{y_j - y_i}{y_j - q y_i} \\ &\times \prod_{k=1}^{\alpha} \left(\sum_{j=0}^{m_k} \frac{(-1)^j q^{\binom{m_k-j}{2}}}{(q; q)_j (q; q)_{m_k-j}} \prod_{p > m[1, k-1]}^{j+m[1, k-1]} \prod_{a > \ell_{c_k-1}}^n \frac{1 - q x_a y_p}{1 - x_a y_p} \prod_{r > j+m[1, k-1]}^{m[1, k]} \prod_{b > \ell_{c_k}}^n \frac{1 - q x_b y_r}{1 - x_b y_r} \right) \\ &\times f_\mu(\kappa; y_1^{-1}, \dots, y_m^{-1}) \prod_{i=1}^m \frac{(y_i - s) dy_i}{y_i^2}, \quad (1-3) \end{aligned}$$

where (positively oriented) integration contours are chosen to encircle all points $\{x_j^{-1}\}_{j=1}^n$ and no other singularities of the integrand, or as *q-nested* closed simple curves with y_i -contour containing $q^{-1} \cdot (y_j\text{-contour})$ for all $i < j$, and all of the contours encircling $\{x_j^{-1}\}_{j=1}^n$. The contours can also be chosen to either encircle or not encircle the point 0.

Our proof of [Theorem 1.1](#) is deeply rooted in the formalism of spin nonsymmetric Hall-Littlewood functions f_μ developed in [\[Borodin and Wheeler 2018\]](#). The key new ingredient is the idea to treat skew-Cauchy identities for these functions as averages of certain observables over the measure given by terms of the nonskew Cauchy identity. The latter can then be identified, under a certain specialization, with the sum over states of the stochastic vertex model along a horizontal line. Accessing observables via deformed Cauchy identities was previously used in [\[Borodin and Petrov 2017; 2018a\]](#) in the colour-blind case, but the mechanism of deformation was different, and the path to integral representations was more complex. It should be noted, however, that [\[Borodin and Petrov 2018a\]](#) was able to reach an integral representation for fully inhomogeneous vertex models (which was later exploited analytically in [\[Borodin and Petrov 2018b\]](#)). So far we have not been able to reach the same level of inhomogeneity in the coloured case, although some inhomogeneity can be added, and it is actually necessary for the limit to polymers.

Let us briefly mention some algebraic corollaries of [Theorem 1.1](#).

First, in the colour-blind case $\lambda = (n)$, [\(1-3\)](#) readily leads to a formula for q -moments of the height function with a completely factorized integrand that could be viewed as a source of all the major asymptotic advances in the area. This colour-blind reduction is discussed in [Section 6B](#).

Next, as the dependence on the values of the coordinates of μ is concentrated in the f_μ -factor and f_μ 's are eigenfunctions of a transfer-matrix of our vertex model, the expectations \mathcal{O}_μ satisfy certain difference equations. Those can be seen as evidence that $\mathcal{O}_\mu(v)$ is actually a *duality functional* for our model; see [Section 6C](#) for details. Duality has served as a major tool for analyzing (q -)moments since [\[Kardar 1987\]](#). It would be very interesting to see if our prospective duality functionals can be related to those obtained by Kuan [\[2018\]](#).

In the rainbow case $\lambda = (1, 1, \dots, 1)$, together with simplification of observables \mathcal{O}_μ to $\mathcal{O}_\mu^{\text{rainbow}}$ mentioned above, the integrand also simplifies due to the lack of j -summations. This case carries additional colour-position symmetry both in the left-hand side of [\(1-3\)](#) and in the right-hand side; see [Section 6D](#). When applied to \mathcal{O}_μ , this yields another set of potential duality functionals. Let us also note

that for anti-dominant μ , when f_μ completely factorizes, the result can alternatively be obtained from the colour-blind case by applying a (highly nontrivial) shift-invariance property of [Borodin et al. 2019].

Finally, [Theorem 1.1](#) allows for *stochastic fusion*: A cluster of neighbouring rows with same left-entering colours and rapidities forming a geometric progression can be collapsed to a single “fat” row whose edges are allowed to carry multiple paths. One can further analytically continue in the parameters $q^{\text{number of rows in a cluster}}$, obtaining a corresponding result in the fully fused model.³ Details of this procedure can be found in [Borodin and Wheeler 2018; Borodin et al. 2019], and the resulting version of [Theorem 1.1](#) is [Corollary 6.9](#) in [Section 6E](#).

The source of analytic corollaries of [Theorem 1.1](#) is the fact that the coloured stochastic vertex model and its fully fused version degenerate, in various limits, to a variety of other probabilistic systems; see [Borodin and Wheeler 2018, Chapter 12] and [Borodin et al. 2019] for some of those degenerations, and the chart in the introduction to [Borodin et al. 2019] for a “big picture”. In this text we only consider, in [Section 7](#), the limit into directed random polymers that was worked out in [Borodin et al. 2019]. We obtain versions of [Theorem 1.1](#) for random Beta-polymers (first considered in [Barraquand and Corwin 2017]), strict-weak or Gamma-polymers (first considered in [Corwin et al. 2015; O’Connell and Ortmann 2015]), O’Connell–Yor semidiscrete Brownian polymers [O’Connell and Yor 2001], and fully continuous (also known as *continuum*) Brownian polymers (equivalently, the stochastic heat equation with multiplicative noise or the KPZ equation).

Two simplifications happen in these limits. First, the presence of colours in our vertex models translates into varying starting points of the polymers, while the general definitions of the polymer models remain the same as in the colourless situation. Second, the observables simplify to pure moments of partition functions (no linear combinations necessary) in three of the four polymer models we consider. Let us illustrate what happens on the example of the continuum Brownian polymer; see [Section 7E](#).

Let $\mathcal{Z}^{(y)}(t, x)$ be the unique solution of the following stochastic partial differential equation with the initial condition

$$\mathcal{Z}_t^{(y)} = \frac{1}{2} \mathcal{Z}_{xx}^{(y)} + \eta(t, x) \mathcal{Z}^{(y)}, \quad t > 0, x \in \mathbb{R}; \quad \mathcal{Z}^{(y)}(0, x) = \delta(x - y),$$

where $\eta = \eta(t, x)$ is the two-dimensional white noise. See, e.g., [Quastel 2012] and references therein for an extensive literature on this equation and its close relation to continuum Brownian path integrals and the Kardar–Parisi–Zhang equation.

We are interested in evaluating the average of the product of several $\mathcal{Z}^{(y)}(t, x)$ with varying x ’s and y ’s. We will use the notation \varkappa_j for different values of x , and we will group those according to which of them correspond to the same value of y . In order to do that, we will talk about a coloured real-valued vector $\varkappa = (\varkappa_1, \dots, \varkappa_m)$, with coordinates of the same colour corresponding to the same y ’s. Further, we will simply use the term “colour” for those y ’s and denote them as $s_1 < \dots < s_\alpha$; they are also real-valued. Denote

$$m_i^{(x)} = \#\{j : \varkappa_j = x \text{ and has colour } s_i\}, \quad m_i = \sum_x m_i^{(x)}, \quad 1 \leq i \leq \alpha, x \in \mathbb{R}.$$

³Our original setup can be viewed as a partially fused model because vertical edges are allowed to carry multiple paths; it could have been obtained from the *fundamental* model with no more than one path on any edge by clustering columns.

Then the limiting version of the fused version of [Theorem 1.1](#) reads (see [Proposition 7.9](#))

$$\begin{aligned} \mathbb{E} \left[\prod_{(i,x): m_i^{(x)} > 0} \frac{(\mathcal{Z}^{(s_i)}(t, x))^{m_i^{(x)}}}{m_i^{(x)}!} \right] \\ = \frac{1}{(2\pi\sqrt{-1})^m} \int \cdots \int \prod_{1 \leq i < j \leq m} \frac{w_j - w_i}{w_j - w_i - 1} \\ \times \prod_{k=1}^{\alpha} \frac{\exp\left(-s_k \cdot \sum_{r > m[1,k-1]}^{m[1,k]} w_r\right)}{m_k!} \cdot \mathfrak{e}_{\mathcal{x}}(w_1, \dots, w_m) \prod_{i=1}^m e^{tw_i^2/2} dw_i, \quad (1-4) \end{aligned}$$

where the integration is over upwardly oriented lines $w_i = a_i + \sqrt{-1} \cdot \mathbb{R}$ with $\Re a_j > \Re a_i + 1$ for $j > i$. The functions $\mathfrak{e}_{\mathcal{x}}$ in the integrand are the limiting versions of the functions f_{μ} in (1-3), and they are defined as follows. In the rainbow case, in the dominant sector $\kappa_1 \geq \kappa_2 \geq \dots \geq \kappa_m$ one has

$$\mathfrak{e}_{\mathcal{x}}(w_1, \dots, w_m) = \exp(\mathcal{x}_1 w_1 + \dots + \mathcal{x}_m w_m),$$

and for $\mathcal{x}_i > \mathcal{x}_{i+1}$ for some $1 \leq i \leq m - 1$ one uses the exchange relations

$$\mathfrak{F}_i \cdot \mathfrak{e}_{\mathcal{x}} = \mathfrak{e}_{(\mathcal{x}_1, \dots, \mathcal{x}_{i+1}, \mathcal{x}_i, \dots, \mathcal{x}_m)}, \quad \mathfrak{F}_i \equiv 1 - \frac{w_i - w_{i+1} + 1}{w_i - w_{i+1}} (1 - \mathfrak{s}_i), \quad 1 \leq i \leq m - 1,$$

to extend the definition to all rainbow vectors \mathcal{x} . For a more general colouring, we use the (unique) colour-identifying monotone map θ from $\{1, \dots, m\}$ to the set of colours $\{s_i\}$, and define $\mathfrak{e}_{\mathcal{x}} = \sum_{\text{rainbow } \mathcal{x}': \theta_*(\mathcal{x}') = \mathcal{x}} \mathfrak{e}_{\mathcal{x}'}$.

The moment formula (1-4) has a certain shift-invariance, see [Remark 7.10](#) below, that is partially explained by the KPZ-level degeneration of the results of [\[Borodin et al. 2019\]](#). This shift-invariance is, furthermore, a corollary of the conjecture in the introduction to [\[Borodin et al. 2019\]](#). It does not imply that conjecture though, because the moments are well-known to not determine the distributions of \mathcal{Z} 's uniquely.⁴

2. Preliminaries

The goal of this section is to summarize previously proved results that we will need later on. The notation and exposition largely follow [\[Borodin and Wheeler 2018\]](#).

2A. The weights. The vertex models that we consider assign weights to finite collections of finite paths drawn on a square grid. Each vertex for which there exists a path that enters and exits it produces a weight that depends on the configuration of all the paths that go through this vertex. The total weight for a collection of paths is the product of weights of the vertices that the paths traverse. We tacitly assume the normalization in which the weight of an empty vertex is always equal to 1.

Our paths are going to be *coloured*, i.e., each path carries a colour that will typically be a (positive) natural number, with colour 0 reserved for the absence of a path. Our vertex weights will actually depend on the ordering of the (nonzero) paths' colours, rather than on their exact values. The paths will always

⁴That conjecture was very recently proved, by two different methods, in [\[Dauvergne 2020\]](#) and [\[Galashin 2020\]](#).

They can also be given by the following explicit formula originating from [Bosnjak and Mangazeev 2016]; see also [Borodin and Wheeler 2018, Theorem C.1.1]:

$$W_{L,M} \left(x; q; \begin{array}{c} C \\ \leftarrow B \rightarrow \\ A \end{array} D \right) = (\mathbf{1}_{A+B=C+D}) x^{|D|-|B|} q^{|A|L-|D|M} \\ \times \sum_P \Phi(C-P, C+D-P; q^{L-M}x, q^{-M}x) \Phi(P, B; q^{-L}/x, q^{-L}), \quad (2-6)$$

where the sum is over compositions $P = (P_1, \dots, P_n)$ such that $0 \leq P_i \leq \min(B_i, C_i)$ for all $1 \leq i \leq n$; and for any two compositions $\lambda, \mu \in \mathbb{N}^n$ such that $\lambda_i \leq \mu_i$ for all $1 \leq i \leq n$, we used the notation

$$\Phi(\lambda, \mu; x, y) := \frac{(x; q)_{|\lambda|} (y/x; q)_{|\mu-\lambda|}}{(y; q)_{|\mu|}} (y/x)^{|\lambda|} q^{\sum_{i < j} (\mu_i - \lambda_i) \lambda_j} \prod_{i=1}^n \binom{\mu_i}{\lambda_i}_q.$$

The substitution of $L = M = 1$ into $W_{L,M}$ returns the (stochastic version of) the fundamental R -matrix for $U_q(\widehat{\mathfrak{sl}}_{n+1})$:

$$W_{1,1} \left(\begin{array}{c} x \\ y \end{array}; q; \begin{array}{c} C \\ \leftarrow B \rightarrow \\ A \end{array} D \right) = \begin{cases} R_{y/x}(A^*, B^*; C^*, D^*), & |A|, |B|, |C|, |D| \leq 1, \\ 0, & \text{otherwise,} \end{cases} \quad (2-7)$$

where we have defined

$$I^* = \begin{cases} 0, & I = \mathbf{0}, \\ i, & I = \mathbf{e}_i \text{ (} i\text{-th standard basis vector),} \end{cases}$$

for any composition $I = (I_1, \dots, I_n)$ such that $|I| \leq 1$, and where $R_{y/x}$ denotes the fundamental R -matrix depicted as (with $z = y/x$)

$$R_z(i, j; k, \ell) = \begin{array}{c} k \\ \uparrow \\ j \text{ --- } \ell \\ \downarrow \\ i \end{array}, \quad i, j, k, \ell \in \{0, 1, \dots, n\}, \quad (2-8)$$

and with matrix elements summarized in Table 2, in which we assume that $0 \leq i < j \leq n$. Observe that these weights are manifestly stochastic.

The general weights $W_{L,M}$ can be reconstructed from the fundamental ones in Table 2 via the procedure of *stochastic fusion*. To state how it works we need a bit of notation.

Let $N \geq 1$, and consider a vector of nonnegative integers $(i_1, \dots, i_N) \in \{0, 1, \dots, n\}^M$. From this we define another vector,

$$C(i_1, \dots, i_N) := (I_1, \dots, I_n), \quad I_a = \#\{k : i_k = a\}, \quad 1 \leq a \leq n, \quad (2-9)$$

which keeps track of the multiplicity of each colour $1 \leq a \leq n$ within (i_1, \dots, i_N) . Set

$$\text{inv}(i_1, \dots, i_N) = \#\{1 \leq a < b \leq N : i_a > i_b\}, \quad \widetilde{\text{inv}}(i_1, \dots, i_N) = \#\{1 \leq a < b \leq N : i_a < i_b\},$$


$\begin{array}{c} i \\ \uparrow \\ i \text{ --- } i \\ \downarrow \\ i \\ 1 \end{array}$	$\begin{array}{c} j \\ \uparrow \\ i \text{ --- } i \\ \downarrow \\ j \\ \frac{q(1-z)}{1-qz} \end{array}$	$\begin{array}{c} i \\ \uparrow \\ i \text{ --- } j \\ \downarrow \\ j \\ \frac{1-q}{1-qz} \end{array}$
	$\begin{array}{c} i \\ \uparrow \\ j \text{ --- } j \\ \downarrow \\ i \\ \frac{1-z}{1-qz} \end{array}$	$\begin{array}{c} j \\ \uparrow \\ j \text{ --- } i \\ \downarrow \\ i \\ \frac{(1-q)z}{1-qz} \end{array}$

Table 2. The matrix elements of the fundamental R -matrix.

and, denoting $I_0 := N - \sum_{a=1}^n I_a$,

$$Z_q(\mathbf{N}; \mathbf{I}) = \sum_{\mathcal{C}(i_1, \dots, i_N) = \mathbf{I}} q^{\text{inv}(i_1, \dots, i_N)} = \sum_{\mathcal{C}(j_1, \dots, j_N) = \mathbf{I}} q^{\widetilde{\text{inv}}(j_1, \dots, j_N)} = \frac{(q; q)_N}{(q; q)_{I_0} (q; q)_{I_1} \dots (q; q)_{I_n}}.$$

Then (see [Borodin et al. 2019, Appendix]),

$$W_{L,M} \left(\frac{x}{y}; q; \begin{array}{c} C \\ B \text{ --- } D \\ A \end{array} \right) = \frac{1}{Z_q(M; \mathbf{A}) Z_q(L; \mathbf{B})} \sum_{\substack{\mathcal{C}(j_1, \dots, j_L) = \mathbf{B} \\ \mathcal{C}(\ell_1, \dots, \ell_L) = \mathbf{D}}} q^{\widetilde{\text{inv}}(j_1, \dots, j_L)} \sum_{\substack{\mathcal{C}(i_1, \dots, i_M) = \mathbf{A} \\ \mathcal{C}(k_1, \dots, k_M) = \mathbf{C}}} q^{\text{inv}(i_1, \dots, i_M)} \quad (2-10)$$


where the figure on the right denotes the corresponding partition function with R -weights given by (2-8) and Table 2 (thus, summation over all possible states of interior edges is assumed), and the colours on the boundary edges, as well as row and column rapidities, are explicitly indicated.⁵

A key feature of the fused R -vertices that allows stacking them together is their q -exchangeability: In equation (2-10), the sum over (k_1, \dots, k_M) can be omitted at the expense of adding the factor of $q^{-\text{inv}(k_1, \dots, k_M)} Z_q(M; \mathbf{C})$ to the summands, and, independently, the sum over (ℓ_1, \dots, ℓ_L) can be removed at the expense of adding the factor of $q^{-\widetilde{\text{inv}}(\ell_1, \dots, \ell_L)} Z_q(L; \mathbf{D})$; see [Borodin and Wheeler 2018, Proposition B.2.2] for a proof.

⁵As in (2-8), the spectral parameter of an R -vertex is assumed to be equal to the ratio of the column rapidity and the row rapidity.

In what follows we will also need partially fused weights defined as (see (2-4))

$$L_x^{\text{stoch}}(A, b; C, d) := W_{1,M} \left(\frac{x}{s}; q; e_b \begin{array}{c} \xrightarrow{C} \\ \xrightarrow{A} \\ \xrightarrow{e_d} \end{array} \right) \Big|_{q^M \rightarrow s^{-2}}, \quad (2-11)$$

$$M_x^{\text{stoch}}(A, b; C, d) := W_{1,M} \left(\frac{s}{x}; \frac{1}{q}; e_b \begin{array}{c} \xrightarrow{C} \\ \xrightarrow{A} \\ \xrightarrow{e_d} \end{array} \right) \Big|_{q^M \rightarrow s^{-2}}, \quad (2-12)$$

where the substitution of a generic complex parameter s^2 for q^{-M} is based on the fact that the right-hand sides are rational in q^M . The weights L_x^{stoch} are tabulated in Table 1.

In addition, we will use gauge transformed (nonstochastic) versions

$$(-s)^{-1_{\ell \geq 1}} L_x^{\text{stoch}}(I, j; K, \ell) =: L_x(I, j; K, \ell) = x \rightarrow j \begin{array}{c} \uparrow \\ \text{---} \\ \downarrow \end{array} \begin{array}{c} K \\ \uparrow \\ \text{---} \\ \downarrow \\ I \end{array} \ell, \quad (2-13)$$

$$(-s)^{1_{j \geq 1}} M_x^{\text{stoch}}(I, j; K, \ell) =: M_x(I, j; K, \ell) = y \leftarrow \ell \begin{array}{c} \uparrow \\ \text{---} \\ \downarrow \end{array} \begin{array}{c} K \\ \uparrow \\ \text{---} \\ \downarrow \\ I \end{array} j, \quad (2-14)$$

that, as a consequence of (2-3), satisfy the following version of the Yang–Baxter equation; see [Borodin and Wheeler 2018, (2.3.5)]:

$$\sum_{0 \leq k_1, k_3 \leq n} \sum_{K \in \mathbb{N}^n} y \leftarrow j_3 \begin{array}{c} \uparrow \\ \text{---} \\ \downarrow \end{array} \begin{array}{c} K \\ \uparrow \\ \text{---} \\ \downarrow \\ I \end{array} \begin{array}{c} k_3 \\ \text{---} \\ k_1 \end{array} \begin{array}{c} j_1 \\ \text{---} \\ i_3 \end{array} \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \begin{array}{c} i_1 \\ \text{---} \\ x \rightarrow \end{array} = \sum_{0 \leq k_1, k_3 \leq n} \sum_{K \in \mathbb{N}^n} y \leftarrow j_3 \begin{array}{c} \uparrow \\ \text{---} \\ \downarrow \end{array} \begin{array}{c} K \\ \uparrow \\ \text{---} \\ \downarrow \\ I \end{array} \begin{array}{c} k_1 \\ \text{---} \\ k_3 \end{array} \begin{array}{c} j_1 \\ \text{---} \\ i_3 \end{array} \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \end{array} \begin{array}{c} i_1 \\ \text{---} \\ x \rightarrow \end{array} \quad (2-15)$$

with the spectral parameter of the R -vertex equal to $(qxy)^{-1}$.

The explicit values of the weights (2-13) are summarized in Table 3 where we assume that $1 \leq i < j \leq n$, and the notation $I_{[k,n]}$ stands for $\sum_{a=k}^n I_a$. For $n = 1$, these weights correspond to the image of the universal R -matrix for the quantum affine group $U_q(\widehat{\mathfrak{sl}}_2)$ in the tensor product of its vector representation (horizontal edges) and a Verma module (vertical edges), with parameter s encoding its highest weight; we call s the *spin parameter*.

2B. The q -Hahn specialization and a limit relation. The complicated expression (2-6) can simplify at special values of parameters; we have already seen this in the case of L^{stoch} and M^{stoch} . Another such specialization that allows L and M to remain generic is the following (see [Kuniba et al. 2016, Proposition 7])

$\begin{array}{c} I \\ \uparrow \\ 0 \text{ --- } \uparrow \text{ --- } 0 \\ \downarrow \\ I \end{array}$ $\frac{1-sxq^{I_{[1,n]}}}{1-sx}$	$\begin{array}{c} I \\ \uparrow \\ i \text{ --- } \uparrow \text{ --- } i \\ \downarrow \\ I \end{array}$ $\frac{(x-sq^{I_i})q^{I_{[i+1,n]}}}{1-sx}$	$\begin{array}{c} I_i^- \\ \uparrow \\ 0 \text{ --- } \uparrow \text{ --- } i \\ \downarrow \\ I \end{array}$ $\frac{x(1-q^{I_i})q^{I_{[i+1,n]}}}{1-sx}$
$\begin{array}{c} I_i^+ \\ \uparrow \\ i \text{ --- } \uparrow \text{ --- } 0 \\ \downarrow \\ I \end{array}$ $\frac{1-s^2q^{I_{[1,n]}}}{1-sx}$	$\begin{array}{c} I_{ij}^{+-} \\ \uparrow \\ i \text{ --- } \uparrow \text{ --- } j \\ \downarrow \\ I \end{array}$ $\frac{x(1-q^{I_j})q^{I_{[j+1,n]}}}{1-sx}$	$\begin{array}{c} I_{ji}^{+-} \\ \uparrow \\ j \text{ --- } \uparrow \text{ --- } i \\ \downarrow \\ I \end{array}$ $\frac{s(1-q^{I_i})q^{I_{[i+1,n]}}}{1-sx}$

Table 3. The explicit values of the weights (2-13).

and also [Bosnjak and Mangazeev 2016, (7.13)]: Assuming that $L \leq M$, we have

$$\begin{aligned} W_{L,M} \left(1; q; \begin{array}{c} C \\ \uparrow \\ B \text{ --- } \uparrow \text{ --- } D \\ \downarrow \\ A \end{array} \right) &= q^{(L-M)|D|} \frac{(q^{L-M}; q)_{|A|-|D|} (q^{-L}; q)_{|D|}}{(q^{-M}; q)_{|A|}} q^{\sum_{i < j} D_i (A_j - D_j)} \prod_{i=1}^n \binom{A_i}{A_i - D_i}_q \\ &= \Phi(D, A; q^{-L}, q^{-M}). \end{aligned} \quad (2-16)$$

The proof follows from the fact that setting $x = 1$ restricts the sum in (2-6) to a single term with $\mathbf{P} = \mathbf{B}$. This specialization is often referred to as the q -Hahn point because for $n = 1$ it reproduces the orthogonality weights for the classical q -Hahn orthogonal polynomials.

Note that replacing q^{-M} by a generic complex parameter s^2 as was done for $W_{1,M}$ in (2-11) can also be performed for $W_{L,M}$ for $L > 1$ either in (2-6) or in (2-16), because those are weights are still manifestly rational in q^M . Alternatively, the result of such a replacement could be seen as a stochastic fusion of the weights L^{stoch} in the spirit of (2-10), where only the outer sum over j_* 's and ℓ_* 's is present.

We will also need the following limiting relation for the weights (2-6); see [Borodin et al. 2019, Lemma 6.8]:

Assume that $\mathbf{B} = (0, \dots, 0, L)$. Then

$$\lim_{q^{-M}=s^2 \rightarrow 0} W_{L,M} \left(zq^M; q; \begin{array}{c} C \\ \uparrow \\ B \text{ --- } \uparrow \text{ --- } D \\ \downarrow \\ A \end{array} \right) = \begin{cases} \frac{(zq^L; q)_\infty}{(z; q)_\infty} \frac{(q^{-L}; q)_d}{(q; q)_d} (zq^L)^d & \text{if } \mathbf{D} = (0, \dots, 0, d), \ d \geq 0, \\ 0 & \text{otherwise.} \end{cases} \quad (2-17)$$

2C. Row operators and rational functions. Let V be an infinite-dimensional vector space obtained by taking the linear span of all n -tuples of nonnegative integers:

$$V = \text{Span}\{\mathbf{I}\} = \text{Span}\{\{i_1, \dots, i_n\}_{i_1, \dots, i_n \in \mathbb{N}}\}. \quad (2-18)$$

For a composition μ and n complex parameters x_1, \dots, x_n , define a rational function $f_\mu(x_1, \dots, x_n)$ as a partition function depicted below (vertex weights (2-13) are being used):

$$f_\mu(x_1, \dots, x_n) = \begin{array}{cccccc} & A(0) & A(1) & A(2) & \dots & \dots & \dots \\ & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \\ x_n \rightarrow n & \text{---} & \text{---} & \text{---} & \text{---} & \text{---} & \text{---} & 0 \\ \vdots & & & & & & & \vdots \\ \vdots & & & & & & & \vdots \\ x_2 \rightarrow 2 & \text{---} & \text{---} & \text{---} & \text{---} & \text{---} & \text{---} & 0 \\ x_1 \rightarrow 1 & \text{---} & \text{---} & \text{---} & \text{---} & \text{---} & \text{---} & 0 \\ & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \\ & 0 & 0 & 0 & \dots & \dots & \dots & \end{array} \quad (2-23)$$

with $A(k) = \sum_{j=1}^n \mathbf{1}_{\mu_j=k} e_j$. These are certain matrix elements of the operator $\mathcal{C}_1(x_1) \cdots \mathcal{C}_n(x_n)$ that can be symbolically written in the form $\langle \emptyset | \mathcal{C}_1(x_1) \cdots \mathcal{C}_n(x_n) | \mu \rangle$. A more general definition of the f_μ 's, as described in [Borodin and Wheeler 2018, Section 3.4], involves some of the operators \mathcal{C}_i being repeated with different arguments (equivalently, some of the paths entering the partition function (2-23) from the left being of the same colour); we will meet such functions in Section 2H.

Similarly to the f_μ 's, one defines dual functions

$$g_\mu(x_1, \dots, x_n) = \begin{array}{cccccc} & 0 & 0 & 0 & \dots & \dots & \dots \\ & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \\ x_n \leftarrow n & \text{---} & \text{---} & \text{---} & \text{---} & \text{---} & \text{---} & 0 \\ \vdots & & & & & & & \vdots \\ \vdots & & & & & & & \vdots \\ x_2 \leftarrow 2 & \text{---} & \text{---} & \text{---} & \text{---} & \text{---} & \text{---} & 0 \\ x_1 \leftarrow 1 & \text{---} & \text{---} & \text{---} & \text{---} & \text{---} & \text{---} & 0 \\ & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \\ & A(0) & A(1) & A(2) & \dots & \dots & \dots & \end{array} \quad (2-24)$$

as matrix elements $\langle \mu | \mathcal{B}_1(x_1) \cdots \mathcal{B}_n(x_n) | \emptyset \rangle$ of $\mathcal{B}_1(x_1) \cdots \mathcal{B}_n(x_n)$. One proves (see [Borodin and Wheeler 2018, Proposition 5.6.1]) that these two families of functions are closely related:

$$g_{\tilde{\mu}}(x_n^{-1}, \dots, x_1^{-1}; q^{-1}, s^{-1}) = c_\mu(q, s) \prod_{i=1}^n x_i \cdot f_\mu(x_1, \dots, x_n; q, s), \quad \tilde{\mu} = (\mu_n, \dots, \mu_1), \quad (2-25)$$

where the multiplicative constant $c_\mu(q, s)$ is given by

$$c_\mu(q, s) = \frac{s^n (q-1)^n q^{\#\{i < j : \mu_i \leq \mu_j\}}}{\prod_{j \geq 0} (s^2; q)_{m_j(\mu)}}, \quad m_j(\mu) := \#\{1 \leq k \leq n : \mu_k = j\}, \quad j \geq 0. \quad (2-26)$$

It will be convenient for us to use a slightly different normalization of the dual functions:

$$g_\mu^*(x_1, \dots, x_n) := q^{n(n+1)/2} (q-1)^{-n} \cdot g_\mu(x_1, \dots, x_n).$$

Finally, let us introduce a third family of symmetric rational functions parametrized by a pair of compositions μ, ν , or rather by a skew composition μ/ν , by

$$G_{\mu/\nu}(x_1, \dots, x_p) = \begin{array}{ccccccc} & & \mathbf{B}(0) & \mathbf{B}(1) & \mathbf{B}(2) & \dots & \dots & \dots \\ & & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \\ x_p \leftarrow & 0 & \leftarrow & \leftarrow & \leftarrow & \leftarrow & \leftarrow & 0 \\ & \vdots & \leftarrow & \leftarrow & \leftarrow & \leftarrow & \leftarrow & \vdots \\ & \vdots & \leftarrow & \leftarrow & \leftarrow & \leftarrow & \leftarrow & \vdots \\ x_2 \leftarrow & 0 & \leftarrow & \leftarrow & \leftarrow & \leftarrow & \leftarrow & 0 \\ x_1 \leftarrow & 0 & \leftarrow & \leftarrow & \leftarrow & \leftarrow & \leftarrow & 0 \\ & & \mathbf{A}(0) & \mathbf{A}(1) & \mathbf{A}(2) & \dots & \dots & \dots \end{array} \quad (2-27)$$

(where $p = 1, 2, \dots$ is arbitrary) with $\mathbf{A}(k) = \sum_{j=1}^n \mathbf{1}_{\mu_j=k} \mathbf{e}_j$, $\mathbf{B}(k) = \sum_{j=1}^n \mathbf{1}_{\nu_j=k} \mathbf{e}_j$, for all $k \in \mathbb{Z}_{\geq 0}$. These are matrix elements $\langle \mu | \mathcal{B}_0(x_1) \cdots \mathcal{B}_0(x_p) | \nu \rangle$ of the operator $\mathcal{B}_0(x_1) \cdots \mathcal{B}_0(x_p)$, and their symmetry with respect to the x_i 's is a direct consequence of the commutativity of $\mathcal{B}_0(x_i)$'s noted after (2-20).

2D. Colour-blindness. For any integer $k \in \{0, 1, \dots, n\}$ define its colour-blind projection

$$\theta(k) := \mathbf{1}_{k \geq 1} = \begin{cases} 0, & k = 0, \\ 1, & k \geq 1. \end{cases}$$

Also, denote the set of compositions of a fixed length (number of entries) with a given weight (sum of entries) k as

$$\mathcal{W}(k) := \{\mathbf{K} \in \mathbb{Z}_{\geq 0}^n : |\mathbf{K}| = k\}.$$

Then one proves (see [Borodin and Wheeler 2018, Proposition 2.4.2]) that

$$\sum_{\mathbf{K} \in \mathcal{W}(k)} L_x(\mathbf{I}, j; \mathbf{K}, 0) = L_x^{(1)}(|\mathbf{I}|, \theta(j); k, 0), \quad \sum_{\mathbf{K} \in \mathcal{W}(k)} \sum_{1 \leq \ell \leq n} L_x(\mathbf{I}, j; \mathbf{K}, \ell) = L_x^{(1)}(|\mathbf{I}|, \theta(j); k, 1), \quad (2-28)$$

where $L^{(1)}$ refers to the weights (2-13) with $n = 1$. Similarly, one has

$$\sum_{\mathbf{K} \in \mathcal{W}(k)} M_x(\mathbf{I}, j; \mathbf{K}, 0) = M_x^{(1)}(|\mathbf{I}|, \theta(j); k, 0), \quad \sum_{\mathbf{K} \in \mathcal{W}(k)} \sum_{1 \leq \ell \leq n} M_x(\mathbf{I}, j; \mathbf{K}, \ell) = M_x^{(1)}(|\mathbf{I}|, \theta(j); k, 1).$$

This has been observed in a number of earlier publications [Foda and Wheeler 2013; Garbali et al. 2017; Kuan 2018], and used to different effects within those works. This property allows certain linear combinations of higher-rank partition functions to be computable as rank-1, or *colour-blind* partition functions.

As an example, one derives the symmetrization identities (see [Borodin and Wheeler 2018, Propositions 3.4.4 and 4.4.3])

$$\sum_{\mu: \mu^+ = \nu} f_\mu(x_1, \dots, x_n) = F_\nu^c(x_1, \dots, x_n), \quad \sum_{\mu: \mu^+ = \kappa} G_{\nu/\mu}(x_1, \dots, x_p) = q^{-np} G_{\nu^+/\kappa}(x_1, \dots, x_p), \quad (2-29)$$

with the sums taken over all compositions with a given dominant reordering denoted by the superscript “+”,

and where the symmetric rational functions F^c and G are colour-blind objects that were considered at length in [Borodin 2017; Borodin and Petrov 2017; 2018a].

2E. Recursive relations. While explicit formulas representing the functions f_μ as sums of monomials are rather involved (see Chapters 6 and 7 of [Borodin and Wheeler 2018] for two different versions) there exist concise recursive relations for them.

First, for *anti-dominant* compositions $\delta = (\delta_1 \leq \dots \leq \delta_n)$, the functions f_δ are completely factorized (see [Borodin and Wheeler 2018, Proposition 5.1.1]):

$$f_\delta(x_1, \dots, x_n) = \frac{\prod_{j \geq 0} (s^{2j}; q)_{m_j(\delta)}}{\prod_{i=1}^n (1 - sx_i)} \prod_{i=1}^n \left(\frac{x_i - s}{1 - sx_i} \right)^{\delta_i}, \quad m_j(\delta) = \#\{1 \leq k \leq n : \delta_k = j\}, \quad j \geq 0. \quad (2-30)$$

This happens because the partition function (2-23) has only one configuration of paths that contributes nontrivially.

Second, the following recursion allows one to move step by step from antidominant compositions to the dominant ones; see [Borodin and Wheeler 2018, Theorem 5.3.1]:

Let $\mu = (\mu_1, \dots, \mu_n)$ be a composition with $\mu_i < \mu_{i+1}$ for some $1 \leq i \leq n - 1$. Then

$$T_i \cdot f_\mu(x_1, \dots, x_n) = f_{(\mu_1, \dots, \mu_{i+1}, \mu_i, \dots, \mu_n)}(x_1, \dots, x_n), \quad (2-31)$$

where

$$T_i \equiv q - \frac{x_i - qx_{i+1}}{x_i - x_{i+1}}(1 - s_i), \quad 1 \leq i \leq n - 1, \quad (2-32)$$

with elementary transpositions $s_i \cdot h(x_1, \dots, x_n) := h(x_1, \dots, x_{i+1}, x_i, \dots, x_n)$, are the Demazure–Lusztig operators of the polynomial representation of the Hecke algebra of type A_{n-1} .

2F. Summation identities. The functions f_μ , g_μ , and $G_{\mu/\nu}$ satisfy several summation identities that can be found in [Borodin and Wheeler 2018, Section 4]; in what follows we will need a couple of them. The first one bears a certain similarity to a summation identity proved by Sahi [1996] and Mimachi and Noumi [1998] for nonsymmetric Macdonald polynomials [Mimachi and Noumi 1998]; it was proved as [Borodin and Wheeler 2018, Theorem 4.3.1].

Let (x_1, \dots, x_n) and (y_1, \dots, y_n) be two sets of complex parameters such that (see (2-21))

$$\left| \frac{x_i - s}{1 - sx_i} \cdot \frac{y_j - s}{1 - sy_j} \right| < 1 \quad \text{for all } 1 \leq i, j \leq n. \quad (2-33)$$

Then

$$\sum_{\mu} f_{\mu}(x_1, \dots, x_n) g_{\mu}^*(y_1, \dots, y_n) = \prod_{i=1}^n \frac{1}{1 - x_i y_i} \prod_{n \geq i > j \geq 1} \frac{1 - qx_i y_j}{1 - x_i y_j}, \quad (2-34)$$

where the summation is over all compositions μ (with nonnegative coordinates).

This identity is proved by evaluating the matrix element $\langle \emptyset | \mathcal{C}_1(x_1) \cdots \mathcal{C}_n(x_n) \mathcal{B}_1(y_1) \cdots \mathcal{B}_n(y_n) | \emptyset \rangle$ in two different ways, by inserting the partition of unity $\sum_{\mu} |\mu\rangle \langle \mu|$ between the groups of \mathcal{B} - and \mathcal{C} -operators, and by using the commutation relations (2-22).




Figure 2. Admissible contours $\{C_1, \dots, C_n\}$ with respect to (q, s) .

A similar argument applied to the matrix element $\langle \emptyset | C_1(x_1) \dots C_n(x_n) \mathcal{B}_0(y_1) \dots \mathcal{B}_0(y_p) | \nu \rangle$ leads to the following summation identity that is reminiscent of the skew-Cauchy identities known in the theory of symmetric functions (see [Borodin and Wheeler 2018, Proposition 4.5.1] for a detailed proof):

Let (x_1, \dots, x_n) and (y_1, \dots, y_p) be two sets of complex parameters satisfying the constraints (2-33), and fix a composition $\nu = (\nu_1, \dots, \nu_n)$. Then one has the identity

$$\sum_{\mu} f_{\mu}(x_1, \dots, x_n) G_{\mu/\nu}(y_1, \dots, y_p) = \prod_{i=1}^n \prod_{j=1}^p \frac{1 - qx_i y_j}{q(1 - x_i y_j)} \cdot f_{\nu}(x_1, \dots, x_n), \tag{2-35}$$

where the summation is taken over all length- n compositions $\mu = (\mu_1, \dots, \mu_n)$.

2G. Orthogonality and integral representations. Let $\{C_1, \dots, C_n\}$ be a collection of contours in the complex plane. We say that the set $\{C_1, \dots, C_n\}$ is admissible with respect to a pair of complex parameters (q, s) if the following conditions are met:

- The contours $\{C_1, \dots, C_n\}$ are closed, positively oriented and pairwise nonintersecting.
- The contours C_i and $q \cdot C_i$ are both contained within contour C_{i+1} for all $1 \leq i \leq n - 1$, where $q \cdot C_i$ denotes the image of C_i under multiplication by q .
- All contours surround the point s .

An illustration of such admissible contours is given in Figure 2.

Often when we integrate rational functions over $\{C_1, \dots, C_n\}$, the integrals can also be computed as sums of residues of the integrand inside the contours. Such sums also make sense for values of parameters that prevent admissible contours from existing, and thus the integrals could also be defined via the residue sums. Therefore, we will tacitly assume that we perform such a replacement should the admissible contours not exist, and we will also use a similar convention for other contour integrals below; see Remark 5.3.

Let $\mu = (\mu_1, \dots, \mu_n)$ and $\nu = (\nu_1, \dots, \nu_n)$ be two compositions. We have the following orthonormality of nonsymmetric spin Hall–Littlewood functions (see [Borodin and Wheeler 2018, Theorem 8.2.1]):

$$\frac{1}{(2\pi\sqrt{-1})^n} \oint_{C_1} \frac{dx_1}{x_1} \dots \oint_{C_n} \frac{dx_n}{x_n} \prod_{1 \leq i < j \leq n} \frac{x_j - x_i}{x_j - qx_i} f_{\mu}(x_1^{-1}, \dots, x_n^{-1}) g_{\nu}^*(x_1, \dots, x_n) = \mathbf{1}_{\mu=\nu}. \tag{2-36}$$

Coupled with (2-34) (which is easily shown to converge uniformly provided that the left-hand sides of (2-33) are bounded by a uniform constant < 1), this leads to the integral formula

$$f_\mu(x_1, \dots, x_n) = \frac{1}{(2\pi\sqrt{-1})^n} \oint_{C_1} \frac{dy_1}{y_1} \dots \oint_{C_n} \frac{dy_n}{y_n} \prod_{1 \leq i < j \leq n} \frac{y_j - y_i}{y_j - qy_i} f_\mu(y_1^{-1}, \dots, y_n^{-1}) \\ \times \prod_{i=1}^n \frac{1}{1 - x_i y_i} \prod_{n \geq i > j \geq 1} \frac{1 - qx_i y_j}{1 - x_i y_j}. \quad (2-37)$$

A similar integral representation for $G_{\mu/\nu}$ originating from (2-35) can be found in [Borodin and Wheeler 2018, Section 9.5].

2H. Coloured compositions. This section closely follows [Borodin and Wheeler 2018, Sections 3.3–3.4], where a more detailed exposition can be found.

Let $\lambda = (\lambda_1, \dots, \lambda_n)$ be a composition of length n and weight m : $|\lambda| = \sum_{i=1}^n \lambda_i = m$. Denote the partial sums of λ by $\sum_{i=1}^k \lambda_i = \ell_k$. We introduce the set \mathcal{S}_λ of λ -coloured compositions as follows:

$$\mathcal{S}_\lambda = \left\{ \mu = (\mu_1 \geq \dots \geq \mu_{\ell_1} | \mu_{\ell_1+1} \geq \dots \geq \mu_{\ell_2} | \dots | \mu_{\ell_{n-1}+1} \geq \dots \geq \mu_{\ell_n}) \right\}. \quad (2-38)$$

That is, the elements of \mathcal{S}_λ are length- m compositions μ , which have been subdivided into blocks of length λ_k , $1 \leq k \leq n$. These blocks demarcate the colouring of μ . Within any given block, the parts of μ have the same colouring and are weakly decreasing.

Two special cases of λ -coloured compositions play special roles. The first is when $\lambda = (n, 0, \dots, 0)$, when compositions $\mu \in \mathcal{S}_\lambda$ consist of a single block whose parts are weakly decreasing; i.e., one simply recovers partitions. As was noted in (2-29), reducing to this case recovers the symmetric rational functions F_* , G_* from [Borodin 2017; Borodin and Petrov 2017; Borodin and Petrov 2018a].

The second one is when $\lambda = (1, 1, \dots, 1) = (1^n)$. Then compositions $\mu \in \mathcal{S}_\lambda$ consist of n blocks, each of a different colour. Thus, the parts of μ are not bound by any inequalities; accordingly, one recovers the set of all length- n compositions. We will refer to these as *rainbow compositions*, or as composition in the *rainbow sector*. The functions f_μ , g_μ and $G_{\mu/\nu}$ introduced above were all defined under the assumption that the participating compositions were in the rainbow sector.

Let $\mu \in \mathcal{S}_\lambda$ be a λ -coloured composition, with ℓ_k denoting the partial sums of λ , as above. We associate to each such μ a vector $|\mu\rangle_\lambda \in \mathbb{V}$, defined as

$$|\mu\rangle_\lambda := \bigotimes_{k=0}^{\infty} |A(k)\rangle_k, \quad A(k) = \sum_{j=1}^n A_j(k) \mathbf{e}_j, \quad A_j(k) = \#\{i : \mu_i = k, \ell_{j-1} + 1 \leq i \leq \ell_j\}, \quad (2-39)$$

where by agreement $\ell_0 = 0$. In other words, the component $A_j(k)$ enumerates the number of parts in the j -th block of μ (these are the parts of colour j) which are equal to k . Further, we define vector subspaces $\mathbb{V}(\lambda)$ of \mathbb{V} which provide a natural grading of \mathbb{V} :

$$\mathbb{V} = \bigoplus_{m=0}^{\infty} \bigoplus_{|\lambda|=m} \mathbb{V}(\lambda), \quad \mathbb{V}(\lambda) := \text{Span}_{\mathbb{C}}\{|\mu\rangle_\lambda\}_{\mu \in \mathcal{S}_\lambda}. \quad (2-40)$$

The grading (2-40) splits \mathbb{V} into subspaces with fixed particle content: $\mathbb{V}(\lambda)$ is the linear span of all states

consisting of λ_i particles of colour i , for all $i \geq 1$. We refer to these subspaces as *sectors* of \mathbb{V} , thus generalizing the “rainbow sector” terminology.

The definitions of the rational functions f_μ , g_ν , and $G_{\mu/\nu}$ naturally lift to coloured composition labels. Concretely, let $\lambda = (\lambda_1, \dots, \lambda_n)$ be a composition of weight m with partial sums ℓ_k as above, and fix a λ -coloured composition $\mu = (\mu_1, \dots, \mu_m) \in \mathcal{S}_\lambda$. In analogy with (2-23), set

$$f_\mu(\lambda; x_1, \dots, x_m) := \langle \emptyset | \left(\prod_{i=1}^{\ell_1} \mathcal{C}_1(x_i) \right) \left(\prod_{i=\ell_1+1}^{\ell_2} \mathcal{C}_2(x_i) \right) \cdots \left(\prod_{i=\ell_{n-1}+1}^{\ell_n} \mathcal{C}_n(x_i) \right) | \mu \rangle_\lambda, \quad (2-41)$$

where $|\mu\rangle_\lambda \in \mathbb{V}(\lambda)$ is given by (2-39), and $\langle \emptyset | \in \mathbb{V}^*$ denotes the (dual) vacuum state $\langle \emptyset | = \bigotimes_{k=0}^{\infty} \langle \mathbf{0} |_k$, which is completely devoid of particles, and \mathcal{C}_i 's are the row-operators (2-19). Graphically, this is the partition function of the form (2-23), with the incoming paths in the bottom ℓ_1 rows having colour 1, having colour 2 in the ℓ_2 rows right above those, etc.

One similarly defines, in analogy with (2-24) and using the row operators (2-20),

$$g_\mu(\lambda; x_1, \dots, x_m) := \langle \mu |_\lambda \left(\prod_{i=1}^{\ell_1} \mathcal{B}_1(x_i) \right) \left(\prod_{i=\ell_1+1}^{\ell_2} \mathcal{B}_2(x_i) \right) \cdots \left(\prod_{i=\ell_{n-1}+1}^{\ell_n} \mathcal{B}_n(x_i) \right) | \emptyset \rangle, \quad (2-42)$$

where $\langle \mu |_\lambda \in \mathbb{V}^*(\lambda)$ is the dual of the vector (2-39), and $|\emptyset\rangle \in \mathbb{V}$ denotes the vacuum state $|\emptyset\rangle = \bigotimes_{k=0}^{\infty} |\mathbf{0}\rangle_k$. Again, here the exiting paths in the bottom ℓ_1 rows have colour 1, in the next ℓ_2 rows they have colour 2, etc.

Finally, in analogy with (2-27), for $\mu, \nu \in \mathcal{S}_\lambda$ we define

$$G_{\mu/\nu}(\lambda; x_1, \dots, x_p) = \langle \mu |_\lambda \mathcal{B}_0(x_1) \cdots \mathcal{B}_0(x_p) | \nu \rangle_\lambda, \quad (2-43)$$

with

$$\begin{aligned} \langle \mu |_\lambda &:= \bigotimes_{k=0}^{\infty} \langle \mathbf{A}(k) |_k, & \mathbf{A}(k) &= \sum_{j=1}^n A_j(k) \mathbf{e}_j, & | \nu \rangle_\lambda &:= \bigotimes_{k=0}^{\infty} | \mathbf{B}(k) \rangle_k, & \mathbf{B}(k) &= \sum_{j=1}^n B_j(k) \mathbf{e}_j, \\ A_j(k) &= \#\{i : \mu_i = k, \ell_{j-1} + 1 \leq i \leq \ell_j\}, & B_j(k) &= \#\{i : \nu_i = k, \ell_{j-1} + 1 \leq i \leq \ell_j\}, \end{aligned}$$

and the graphical depiction (2-27) does not require any modifications. Since the labelling composition λ only participates in this definition as a record of the colours in the boundary states μ and ν , we will often omit it from the notation of $G_{\mu/\nu}$.

3. Extensions

In this section we provide a few straightforward extensions of the results from Section 2, most of which have very similar proofs.

3A. Column inhomogeneities. In (2-23) we defined the functions f_μ by utilizing the weights L_x of (2-13) with $x = x_i$ and a fixed spin parameter s for all the vertices in the i -th row, $1 \leq i \leq n$. It is meaningful, however, to extend the definition when we take $x = x_i \xi_j$ and $s = s_j$ for the vertex in the i -th row and j -th column, where $1 \leq i \leq n$, $0 \leq j < +\infty$, and $\{\xi_j\}_{j \geq 0}$ and $\{s_j\}_{j \geq 0}$ are two infinite sequences of complex parameters. Correspondingly, in the definitions (2-24) and (2-27) of g_μ and $G_{\mu/\nu}$ one needs to use the

weights M_x of (2-14) with $x = x_i \xi_j^{-1}$ and $s = s_j$ for the vertex in the i -th row and j -th column, with the same sequences $\{\xi_j\}_{j \geq 0}$ and $\{s_j\}_{j \geq 0}$.

Many of the results cited in Section 2 and their proofs extend to such an inhomogeneous setup almost *verbatim*. In the colour-blind case of $n = 1$ this can be seen by comparing [Borodin and Petrov 2018a] and [Borodin and Petrov 2017].

The basic reason for such an easy extension lies in the fact that the Yang–Baxter equation (2-15) has a suitable extension. More exactly, the needed deformed equation has the form

$$\sum_{0 \leq k_1, k_3 \leq n} \sum_{\mathbf{K} \in \mathbb{N}^n} L_{x\xi}(\mathbf{I}, i_1; \mathbf{K}, k_1) R_{(qxz)^{-1}}(i_3, k_1; k_3, j_1) M_{z\xi^{-1}}(\mathbf{K}, k_3; \mathbf{J}, j_3) \\ = \sum_{0 \leq k_1, k_3 \leq n} \sum_{\mathbf{K} \in \mathbb{N}^n} M_{z\xi^{-1}}(\mathbf{I}, i_3; \mathbf{K}, k_3) R_{(qxz)^{-1}}(k_3, i_1; j_3, k_1) L_{x\xi}(\mathbf{K}, k_1; \mathbf{J}, j_1), \quad (3-1)$$

where the important thing to notice is that the parameters of the R -weights in the middle remain independent of ξ , and they are also independent of s by their definition (see Table 2).⁶

Let us quickly go through inhomogeneous analogues of the results from Section 2 that we will need.

The commutation relations (2-22) remain unchanged, as they are related to the R -matrix in (3-1), but the convergence condition (2-21) needs to be modified to

$$\lim_{L \rightarrow \infty} \prod_{j=0}^L \left| \frac{\xi_j x - s_j}{1 - s_j \xi_j x} \cdot \frac{y - s_j \xi_j}{\xi_j - s_j y} \right| = 0;$$

see [Borodin and Petrov 2018a, Proposition 4.8] for an explanation in the $n = 1$ case. In what follows, we will only need the situation of finitely many (in fact, exactly one) ξ_j 's different from 1 and s_j 's different from a certain fixed s , in which case, we can clearly continue using (2-21).

Correspondingly, the summation identities (2-34) and (2-35) also remain unchanged, modulo a similar comment about the convergence condition (2-33).

The relation (2-25) between f 's and g 's remains valid with the following modifications — the inversion of s for the g in the left-hand side needs to be applied to all the s_j 's, and the multiplicative constant c_μ needs to be read as

$$\frac{\prod_{j \geq 0} s_j \cdot (q-1)^n q^{\#\{i < j; \mu_i \leq \mu_j\}}}{\prod_{j \geq 0} (s_j^2; q)_{m_j(\mu)}}.$$

The proof remains the same.

The colour-blindness results (2-29) remain in place with identical proofs and inhomogeneous F and G understood as in [Borodin and Petrov 2018a].

The factorization (2-30) for the antidominant compositions looks very similar (and the same argument works):

$$f_\delta(x_1, \dots, x_n) = \prod_{j \geq 0} (s_j^2; q)_{m_j(\delta)} \cdot \prod_{i=1}^n \left(\frac{1}{1 - s_{\delta_i} \xi_{\delta_i} x_i} \prod_{j=0}^{\delta_i - 1} \frac{\xi_j x_i - s_j}{1 - s_j \xi_j x_i} \right). \quad (3-2)$$

⁶This version of the Yang–Baxter equation is also a consequence of the “master” Yang–Baxter equation (2-3).

The recurrence relation (2-31) remains unchanged, with the action of T_i 's given by the same Demazure–Lusztig operators (2-32). Its proof was based on the commutation relations between C_i 's given in [Borodin and Wheeler 2018, Theorem 3.2.1] and on [Borodin and Wheeler 2018, Proposition 5.3.3], both of which remain intact in the inhomogeneous setup.

Finally, the orthogonality relation (2-36) remains literally the same, with the contours surrounding all the points $\{\xi_j s_j\}_{j \geq 0}$ instead of just s in the homogeneous case. It is a bit more difficult to convince oneself that this is so because this proof relies on many ingredients. In addition to the facts already mentioned above, one needs monomial expansions of [Borodin and Wheeler 2018, Chapter 6] and a certain explicit contour integral computation. The latter in the inhomogeneous setup is exactly [Borodin and Petrov 2018a, Lemma 7.1], while the proof of the former carries over to the inhomogeneous case in the same spirit as all the other above-mentioned facts.

The principal reason for our carrying the column inhomogeneity of the model throughout Sections 3-6 is to be able to perform a limit transition in column 1 in Section 6, which will then give us access to averages of observables in the fused models and, as a consequence discussed in Section 7, in integrable models of directed random polymers.

3B. A simplifying specialization of $G_{\nu/\mu}$. Our next goal is to prove the following statement.

Proposition 3.1. *Fix generic complex parameters q and s , a composition λ , and two λ -coloured compositions μ, ν of length $n \geq 1$. Then for $L \in \mathbb{N}$ and generic $\epsilon \in \mathbb{C}$, $G_{\nu/\mu}(\lambda; \epsilon, q\epsilon, \dots, q^{L-1}\epsilon)$, with G defined as in (2-43), is a rational function in q^L , and there exists a limit*

$$\mathcal{G}_{\nu/\mu} := \lim_{\epsilon \rightarrow 0} (q^{nL} \cdot G_{\nu/\mu}(\lambda; \epsilon, q\epsilon, \dots, q^{L-1}\epsilon))|_{q^L=(s\epsilon)^{-1}}, \tag{3-3}$$

where in the right-hand side we substituted a particular value into a rational function. Explicitly, $\mathcal{G}_{\nu/\mu}$ has the form

$$\frac{(-s)^{|\mu|}}{(-s)^{|\nu|}} \mathcal{G}_{\nu/\mu} = \begin{cases} s^{-2n} \prod_{x=0}^{\infty} \left((s^2; q)_{|m^{(x)}|} q^{-\sum_{i>j} m_i^{(x)} m_j^{(x)}} \prod_{i=1}^n q^{m_i^{(x)} H_{>i}^{\nu/\mu}(x+1)} \binom{H_i^{\nu/\mu}(x+1)}{m_i^{(x)}}_q \right), & \text{if all } \nu_i \neq 0, \\ 0, & \text{otherwise,} \end{cases} \tag{3-4}$$

where we defined, for each $x \geq 0$, an n -component vector $\mathbf{m}^{(x)}$ whose i -th component $m_i^{(x)}$, $1 \leq i \leq n$, is equal to the number of parts of μ of colour i that are equal to x (symbolically $\mu = 0^{m^{(0)}} 1^{m^{(1)}} 2^{m^{(2)}} \dots$), and also **coloured height functions**

$$H_i^x(x) = \#\{j : \text{colour}(x_j) = i, x_j \geq x\}, \quad H_{>i}^x = \sum_{k>i} H_k^x(x), \quad H_*^{\nu/\mu} \equiv H_*^\nu - H_*^\mu. \tag{3-5}$$

The same conclusion holds in the inhomogeneous setting of Section 3A under the condition that for each column $x \geq 0$ such that there exists a part of μ equal to x (i.e., $|m^{(x)}| > 0$), the column rapidity and spin

parameter in that column are still equal to 1 and s , respectively, as in the homogeneous case, and the factor $(-s)^{|\mu|-|\nu|}$ in the left-hand side of (3-4) is replaced by $\prod_{i=1}^n \prod_{j=\mu_i}^{v_i-1} (-s_j)^{-1}$.

Remark 3.2. We tacitly follow the convention that the q -binomial coefficients vanish unless their arguments are nonnegative, and the top one is at least as large as the bottom one. Thus, the top line of (3-4) can also produce a zero outcome.

Proof. Let us first switch from using the weights M_x of (2-14) in the partition function of the form (2-27) to using the weights M_x^{stoch} of (2-12) instead. The product of the correcting factors $(-s)^{\mathbf{1}_{j \geq 1}}$ in (2-14) over all vertices gives $(-s)^{|\nu|-|\mu|}$ (the exponent counts the number of horizontal steps of all the paths, which is exactly $|\nu| - |\mu|$). Note that this matches the inverse of $(-s)^{|\mu|-|\nu|}$ in the left-hand side of (3-4), and in the inhomogeneous setup, the product of these correcting factors gives

$$\prod_{i=1}^n \prod_{j=\mu_i}^{v_i-1} (-s_j)$$

instead.

With the weights M_x^{stoch} , it is a bit more convenient to reflect the partition function with respect to a vertical axis so that paths move to the right horizontally. Then, noticing that the left and right boundary conditions of (2-27) consist of having no entering or exiting paths, we recognize that using the geometric progression $(\epsilon, q\epsilon, \dots, q^{L-1}\epsilon)$ for horizontal rapidities is equivalent to performing the stochastic fusion in the vertical direction (corresponding to the outer sum in (2-10)). Hence, we obtain that $(-s)^{|\mu|-|\nu|} G_{\nu/\mu}(\lambda; \epsilon, q\epsilon, \dots, q^{L-1}\epsilon)$ is a one-row partition function with the incoming paths from the bottom parametrizing ν , outgoing paths on the top parametrizing μ , no paths entering or exiting on either side, and the vertex weights given by

$$W_{L,M} \left(\begin{matrix} s & 1 \\ \epsilon & q \end{matrix}; \begin{matrix} C \\ B \uparrow \downarrow A \\ D \end{matrix} \right) \Big|_{q^M \rightarrow s^{-2}}. \tag{3-6}$$

Here s and ϵ need to be replaced by s_j and $\xi_j^{-1}\epsilon$, if the vertex is in a column j that carries inhomogeneities (s_j, ξ_j) . Pictorially,

$$(-s)^{|\mu|-|\nu|} G_{\nu/\mu}(\lambda; \epsilon, q\epsilon, \dots, q^{L-1}\epsilon) = \mathbf{0} \begin{matrix} \dots & \dots & \dots & I(2) & I(1) & I(0) \\ \uparrow & \uparrow & \uparrow & \uparrow & \uparrow & \uparrow \\ \dots & \dots & \dots & J(2) & J(1) & J(0) \end{matrix} \mathbf{0} \tag{3-7}$$

with $I(k) = \sum_{i=1}^n \mathbf{1}_{\mu_i=k} \mathbf{e}_i$, $J(k) = \sum_{j=1}^n \mathbf{1}_{\nu_j=k} \mathbf{e}_j$, for all $k \in \mathbb{Z}_{\geq 0}$. This partition function is a rational function in q^L because every vertex weight (3-6) is; see (2-6).

Assume we are in the homogeneous setting first. Observe that if in the right-hand side of (2-6) we have $q^{L-M}x = 1$, then the only nontrivially contributing term comes from $\mathbf{P} = \mathbf{C}$, for which the first Φ -factor turns into 1. In terms of our weights (3-6) this corresponds to $q^{M-L}s/\epsilon = 1$, which is realized by setting $q^L = (s\epsilon)^{-1}$ (recall that $q^M = s^{-2}$).

Writing out the term with $\mathbf{P} = \mathbf{C}$ we obtain

$$\mathbf{1}_{\mathbf{A}+\mathbf{B}=\mathbf{C}+\mathbf{D}} \cdot \left(\frac{s}{\epsilon}\right)^{|\mathbf{D}|-|\mathbf{B}|} (s\epsilon)^{|\mathbf{A}|} s^{-2|\mathbf{D}|} \times \frac{(s^{-2}; q^{-1})_{|\mathbf{C}|} (s\epsilon^{-1}; q^{-1})_{|\mathbf{B}|-|\mathbf{C}|}}{((s\epsilon)^{-1}; q^{-1})_{|\mathbf{B}|}} \left(\frac{s}{\epsilon}\right)^{|\mathbf{C}|} q^{-\sum_{i<j} (B_i - C_i) C_j} \prod_{i=1}^n \binom{B_i}{C_i}_{q^{-1}}. \quad (3-8)$$

There is an additional factor of $q^{nL} = (s\epsilon)^{-n}$ in (3-3). Since our compositions have n parts, there is a total of n paths exiting (3-7) from the top, and we can distribute $(s\epsilon)^{-n}$ as $(s\epsilon)^{-|\mathbf{C}|}$ over all vertices in (3-7) (with \mathbf{C} corresponding to the paths exiting the vertex at the top). Hence, we need to take the limit as $\epsilon \rightarrow 0$ of (3-8) multiplied by $(s\epsilon)^{-|\mathbf{C}|}$. This is a straightforward calculation which yields

$$\mathbf{1}_{\mathbf{A}+\mathbf{B}=\mathbf{C}+\mathbf{D}} \cdot s^{-2|\mathbf{C}|} (s^2; q)_{|\mathbf{C}|} \cdot q^{\sum_{i>j} (B_i - C_j) C_j} \prod_{i=1}^n \binom{B_i}{C_i}_q. \quad (3-9)$$

For the inhomogeneous setting, our hypothesis implies that for any vertex in a column with inhomogeneity parameters (\tilde{s}, ξ) we must have $\mathbf{C} = \mathbf{0}$. This turns the sum in (2-6) into a single term with $\mathbf{P} = \mathbf{0}$, which leads to the replacement of (3-8) by

$$\mathbf{1}_{\mathbf{A}+\mathbf{B}=\mathbf{C}+\mathbf{D}} \cdot \left(\xi^{-1} \frac{s}{\epsilon}\right)^{|\mathbf{D}|-|\mathbf{B}|} (s\epsilon)^{|\mathbf{A}|} \tilde{s}^{-2|\mathbf{D}|} \frac{((s\epsilon)^{-1}; q^{-1})_{|\mathbf{D}|}}{(\xi^{-1} \tilde{s}^{-2} s\epsilon^{-1}; q^{-1})_{|\mathbf{D}|}} \frac{(\xi^{-1} s\epsilon^{-1}; q^{-1})_{|\mathbf{B}|}}{((s\epsilon)^{-1}; q^{-1})_{|\mathbf{B}|}}. \quad (3-10)$$

This expression has the exact same $\epsilon \rightarrow 0$ asymptotics as (3-8) with $\mathbf{C} = \mathbf{0}$, thus confirming (3-9) for the inhomogeneous setup as well.

It remains to interpret \mathbf{B} and \mathbf{C} in (3-9) in terms of the height functions. Assume that our vertex is located in the column with coordinate $x \geq 0$ (which means, in particular, that $\mathbf{A} = \mathbf{J}(x)$, $\mathbf{C} = \mathbf{I}(x)$ in the notation of (3-7)). Then B_i is the number of paths of colour i in (3-7) that enter from the bottom at a location strictly to the left of x minus the number of paths of colour i that exit through the top at a location strictly to the left of x , and this equals $H_i^{v/\mu}(x+1)$ as defined in (3-5). Furthermore, C_i is exactly $m_i^{(x)}$. Hence, the part of (3-9) past the indicator function takes the form

$$s^{-2|\mathbf{m}^{(x)}|} (s^2; q)_{|\mathbf{m}^{(x)}|} \cdot q^{\sum_{j \geq 1} m_j^{(x)} H_{>j}^{v/\mu}(x+1) - \sum_{i > j} m_i^{(x)} m_j^{(x)}} \prod_{i=1}^n \binom{H_i^{v/\mu}(x+1)}{m_i^{(x)}}_q, \quad (3-11)$$

and the product of these expressions over $x \geq 0$ gives the first line of the right-hand side of (3-4).

On the other hand, the role of the indicator function $\mathbf{1}_{\mathbf{A}+\mathbf{B}=\mathbf{C}+\mathbf{D}}$ in (3-9) is in providing a recipe for uniquely assigning the paths along the horizontal edges of (3-7) inductively: \mathbf{D} is assigned the value of $\mathbf{A} + \mathbf{B} - \mathbf{C}$, starting from the far left where no paths are present. This works smoothly until column 0, in which \mathbf{D} must be equal to $\mathbf{0}$, enforcing $\mathbf{A} + \mathbf{B} = \mathbf{C}$. Since the q -binomial coefficients require $B_i \geq C_i$ for a nonzero outcome, we conclude that \mathbf{A} must be $\mathbf{0}$ in the 0-th column, giving us the second line of the right-hand side of (3-4). \square

An important special case of $\mathcal{G}_{v/\mu}$ is when μ has only zero parts. A direct inspection of (3-4) leads to the following:

Corollary 3.3. For $\mu = 0^n = (0, 0, \dots, 0)$, with notation $\mathcal{G}_\nu := \mathcal{G}_{\nu/0^n}$, the limit (3-3) takes the form

$$(-s)^{-|\nu|} \mathcal{G}_\nu = \begin{cases} s^{-2n}(s^2; q)_n, & \text{if all } \nu_i \neq 0, \\ 0, & \text{otherwise.} \end{cases} \quad (3-12)$$

In the inhomogeneous setting, assuming that $(s_0, \xi_0) = (s, 1)$, a similar formula holds, where one needs to replace $(-s)^{-|\nu|}$ in the left-hand side by $\prod_{i=1}^n \prod_{j=0}^{\nu_i-1} (-s_j)^{-1}$.

The second symmetrization relation of (2-29) implies that \mathcal{G}_ν has to equal

$$\lim_{\epsilon \rightarrow 0} \mathbf{G}_{\nu^+}(\epsilon, q\epsilon, \dots, q^{L-1}\epsilon) |_{q^L=(s\epsilon)^{-1}}$$

with ν^+ denoting the dominant reordering of ν , and \mathbf{G} as in [Borodin and Petrov 2018a]. This limit was evaluated in [Borodin and Petrov 2018a, Proposition 6.7], and was shown to be equivalent to (3-12).⁷

Let us also record for the future what happens to (3-4) when μ and ν are rainbow compositions (equivalently, $\lambda = (1, 1, \dots, 1)$).

Corollary 3.4. Under the assumption that μ and ν are rainbow compositions, (3-4) takes the form

$$\frac{(-s)^{|\mu|}}{(-s)^{|\nu|}} \mathcal{G}_{\nu/\mu} = \begin{cases} s^{-2n} \prod_{x=0}^{\infty} \left((s^2; q)_{|\mathbf{m}^{(x)}|} q^{-\binom{|\mathbf{m}^{(x)}|}{2}} \prod_{1 \leq i \leq n; m_i^{(x)}=1} \mathbf{1}_{H_i^{\nu/\mu}(x+1)=1} q^{H_{>i}^{\nu/\mu}(x+1)} \right), & \text{if all } \nu_i \neq 0, \\ 0, & \text{otherwise.} \end{cases} \quad (3-13)$$

The same formula holds in the inhomogeneous setting of Section 3A under the condition that for each column $x \geq 0$ such that there exists a part of μ equal to x (i.e., $|\mathbf{m}^{(x)}| > 0$), the column rapidity and spin parameter in that column are still equal to 1 and s , respectively, and the factor $(-s)^{|\mu|-|\nu|}$ in the left-hand side is replaced by $\prod_{i=1}^n \prod_{j=\mu_i}^{\nu_i-1} (-s_j)^{-1}$.

Proof. Direct inspection of (3-4). Note that in the rainbow sector, for any given i , $1 \leq i \leq n$, the i -th colour height function H_i^* defined as in (3-5), can only take values 0 or 1. \square

3C. Colour merging. Different versions of colour merging properties of vertex weights have been previously observed and studied in several works including [Foda and Wheeler 2013; Garbali et al. 2017; Kuan 2018; Borodin and Wheeler 2018; Borodin et al. 2019]. We use this section to formulate the statements we need in suitable notation.

Let n_1, n_2 be two positive integers, and let $\theta : \{1, \dots, n_1\} \rightarrow \{1, \dots, n_2\}$ be an arbitrary monotone map. It induces a map θ_* that turns a n_1 -dimensional vector into an n_2 -dimensional one as follows:

$$\theta_* : \mathbf{I} = (I_1, \dots, I_{n_1}) \mapsto \mathbf{J} = (J_1, \dots, J_{n_2}) \quad \text{if } J_j = \sum_{i \in \theta^{-1}(j)} I_i. \quad (3-14)$$

In other words, we sum the coordinates of \mathbf{I} that have the same θ -image and turn the result into a coordinate of \mathbf{J} whose index is that image. Empty sums are interpreted as having value 0.

⁷In fact, [Borodin and Petrov 2018a] provided two different evaluations for this limit, neither of which coincides with the color-blind version of the proof of Proposition 3.1.

Proposition 3.5. *Denote the weights $W_{L,M}$ given by (2-6) with n -dimensional vector arguments as $W_{L,M}^{(n)}$. Then for any $n_1, n_2 \geq 1$ and a map θ as above, we have the following colour merging relation: For any $\mathbf{A}, \mathbf{B} \in \mathbb{Z}_{\geq 0}^{n_1}$ and $\tilde{\mathbf{C}}, \tilde{\mathbf{D}} \in \mathbb{Z}_{\geq 0}^{n_2}$ with $|\mathbf{A}|, |\tilde{\mathbf{C}}| \leq M, |\mathbf{B}|, |\tilde{\mathbf{D}}| \leq L$,*

$$\sum_{\substack{C \in \mathbb{Z}_{\geq 0}^{n_1} : \theta_*(C) = \tilde{C} \\ D \in \mathbb{Z}_{\geq 0}^{n_1} : \theta_*(D) = \tilde{D}}} W_{L,M}^{(n_1)} \left(x; q; \begin{array}{c} C \\ \leftarrow \text{B} \text{---} \text{A} \rightarrow \\ D \end{array} \right) = W_{L,M}^{(n_2)} \left(x; q; \begin{array}{c} \tilde{C} \\ \leftarrow \theta_*(B) \text{---} \theta_*(A) \rightarrow \\ \tilde{D} \end{array} \right). \tag{3-15}$$

Proof. For $L = M = 1$, when the W -weights turn into matrix elements of the R -matrix (see (2-7)) the statement coincides with [Borodin et al. 2019, Proposition 4.3] (and it is also easy to check directly from the formulas given in Table 2 for the weights). For general $L, M \geq 1$, (3-15) readily follows from the $L = M = 1$ case and the stochastic fusion (2-10). \square

Corollary 3.6. *Colour merging statements similar to Proposition 3.5 hold for the vertex weights $L_x, M_x, L_x^{\text{stoch}}, M_x^{\text{stoch}}$ and q -Hahn weights (2-16), as well as for the vertex weights defined by (3-9).*

Proof. This follows from the fact that all these weights are obtained from $W_{L,M}$ by specializations, analytic continuation, multiplication by factors that give the same contribution to the two sides of the merging relation (3-15), and a limit transition $\epsilon \rightarrow 0$ in (3-6) in the case of (3-9). \square

The colour-blindness statements of Section 2D correspond, in the notation of Proposition 3.5, to $n_2 = 1$. In particular, applying the colour-blindness relation to either the q -Hahn weights (2-16) or to the weights (3-9) and removing common factors on the two sides, one obtains the q -identity (3-16).

Corollary 3.7. *For any $Q \in \mathbb{C}, (\alpha_1, \dots, \alpha_n) \in \mathbb{Z}_{\geq 0}^n$, and a fixed $|\beta| \in \mathbb{Z}_{\geq 0}, |\beta| \leq |\alpha|$, one has*

$$\sum_{\substack{0 \leq \beta_i \leq \alpha_i, 1 \leq i \leq n, \\ \beta_1 + \dots + \beta_n = |\beta|}} Q^{\sum_{1 \leq i < j \leq n} \beta_i (\alpha_j - \beta_j)} \prod_{i=1}^n \binom{\alpha_i}{\alpha_i - \beta_i}_Q = \binom{|\alpha|}{|\alpha| - |\beta|}_Q. \tag{3-16}$$

Since compositions are vectors, the map θ_* from (3-14) sends any composition of length n_1 to a composition of length n_2 . This can be naturally extended to coloured compositions as follows.

Let λ be a composition with $\ell(\lambda) = n_1$, weight $|\lambda| = m$, and partial sums $\ell_k = \sum_{i=1}^k \lambda_i$. Then $\rho := \theta_*(\lambda)$ is a composition with $\ell(\rho) = n_2$, same weight $|\rho| = m$, and partial sums that we denote as $r_k = \sum_{i=1}^k \rho_i$. Further, let μ be a λ -coloured composition of length m ; see Section 2H for a definition. As in (2-39), we can encode μ by a sequence of n_1 -dimensional vectors $\{\mathbf{A}(k)\}_{k \geq 0}$:

$$\mathbf{A}(k) = \sum_{j=1}^{n_1} A_j(k) \mathbf{e}_j, \quad A_j(k) = \#\{i : \mu_i = k, \ell_{j-1} + 1 \leq i \leq \ell_j\}.$$

It is not difficult to see that the sequence of n_2 -dimensional vectors $\mathbf{B}(k) := \theta_*(\mathbf{A}(k)), k \geq 0$, corresponds, via

$$\theta_*(\mathbf{A}(k)) = \mathbf{B}(k) = \sum_{j=1}^{n_2} B_j(k) \mathbf{e}_j, \quad B_j(k) = \#\{i : v_i = k, r_{j-1} + 1 \leq i \leq r_j\},$$

to a ρ -coloured composition ν of length m that we will define to be the image of μ under θ_* :

$$\theta^*(\mu) = \nu. \quad (3-17)$$

In less formal terms, if we view a coloured composition μ as positions of finitely many paths coloured by $\{1, \dots, n_1\}$, then $\nu = \theta_*(\mu)$ represents positions of the same paths that have been recoloured according to the map θ . Note that we required θ to be monotone, which means that the order of colours is being preserved.

Proposition 3.8. *Let λ be a composition with $\ell(\lambda) = n_1$ of weight $|\lambda| = m$, ρ be a composition with $\ell(\rho) = n_2$ and same weight $|\rho| = m$ such that $\theta_*(\lambda) = \rho$, ν' by a λ -coloured composition, and ν'' be a ρ -coloured composition (both of length m). Then for any $p \geq 1$ and complex parameters x_1, x_2, \dots , we have*

$$\sum_{\mu: \theta_*(\mu) = \nu''} f_\mu(\lambda; x_1, \dots, x_m) = f_{\nu''}(\rho; x_1, \dots, x_m), \quad (3-18)$$

$$\sum_{\mu: \theta_*(\mu) = \nu''} G_{\nu'/\mu}(\lambda; x_1, \dots, x_p) = G_{\theta_*(\nu')/\nu''}(\rho; x_1, \dots, x_p). \quad (3-19)$$

Remark 3.9. When $\lambda = (1, 1, \dots, 1)$ and $n_2 = 1$, one recovers the symmetrization formulas (2-29).

Proof. In complete analogy with proofs of (2-29) in [Borodin and Wheeler 2018, Propositions 3.4.4 and 4.4.3], the argument consists in multiple applications of Proposition 3.5, or rather its version for the weights L_x and M_x (Corollary 3.6) used in the definitions of f 's and G 's. In the case of (3-18), one starts with the top-right nontrivial vertex of the partition function (2-23) (with appropriate, not necessarily rainbow colours of entering paths on the left), while in the case of (3-19) one starts with the top-left nontrivial vertex of the partition function (2-27), and then moves step by step into the bulk of the partition function. Once the colour-merging summation has been performed for all nontrivial vertices, one recovers the right-hand sides of (3-18) and (3-19). \square

4. Cauchy identities

The goal of this section is to show how the skew-Cauchy identity (2-35) leads to formulas for averages of certain observables for stochastic vertex models.

For this section let us assume that we are either in the column-homogeneous situation, or, slightly more generally, the number of columns in which inhomogeneity parameters (s_j, ξ_j) of Section 3A are different from $(s, 1)$ is finite. We start by extending the skew-Cauchy identity (2-35) to limiting versions of the G -functions from Section 3B.

Proposition 4.1. *Let μ be a rainbow composition of length $n \geq 1$, and x_1, \dots, x_n be complex parameters satisfying*

$$\left| \frac{s(x_i - s)}{1 - sx_i} \right| < 1, \quad 1 \leq i \leq n. \quad (4-1)$$

Then we have

$$\sum_v f_v(x_1, \dots, x_n) \mathcal{G}_{v/\mu} = \prod_{i=1}^n \left(1 - \frac{x_i}{s}\right) \cdot f_\mu(x_1, \dots, x_n), \tag{4-2}$$

where $\mathcal{G}_{v/\mu}$ is as in [Corollary 3.4](#), and f_* 's are as in [\(2-23\)](#).

Proof. We start with the summation identity [\(2-35\)](#), multiply both sides by q^{np} , and substitute $p = L$, $(y_1, \dots, y_p) = (\epsilon, q\epsilon, \dots, q^{L-1}\epsilon)$ with $\epsilon = s^{-1}q^{-L}$ to eventually match with [\(3-3\)](#). The identity takes the form

$$\sum_v f_v(x_1, \dots, x_n) \cdot q^{nL} G_{v/\mu}(\epsilon, q\epsilon, \dots, q^{L-1}\epsilon) = \prod_{i=1}^n \frac{1 - x_i/s}{1 - x_i/(sq^L)} \cdot f_\mu(x_1, \dots, x_n), \tag{4-3}$$

where the convergence conditions [\(2-33\)](#) read

$$\left| \frac{x_i - s}{1 - sx_i} \cdot \frac{y_j - s}{1 - sy_j} \right| = \left| \frac{x_i - s}{1 - sx_i} \cdot \frac{s^{-1}q^{j-L-1} - s}{1 - q^{j-L-1}} \right| = \left| \frac{x_i - s}{1 - sx_i} \cdot \frac{1 - s^2q^{L-j+1}}{s(q^{L-j+1} - 1)} \right| < 1$$

for $1 \leq i \leq n$ and $1 \leq j \leq L$. They will hold for all $L = 1, 2, \dots$ as long as the x_i 's range over a sufficiently small neighbourhood of s (we assume $|q| < 1$).


As in the proof of [Proposition 3.1](#), $q^{nL}G_{v/\mu}(\epsilon, q\epsilon, \dots, q^{L-1}\epsilon)$ can be represented as a one-row partition function of the form [\(3-7\)](#) with fused weights [\(3-6\)](#) multiplied by $(s\epsilon)^{-|C|}$. Denoting $\mathfrak{l} := q^L$, we can rewrite those vertex weights as follows (see [\(3-8\)](#) and recalling that $\epsilon = (s\mathfrak{l})^{-1}$),

$$\mathbf{1}_{A+B=C+D} \cdot \frac{s^{-2|B|-|C|}(s^{-2}; q^{-1})_{|C|}(s^2\mathfrak{l}; q^{-1})_{|B|-|C|} \left(s^2\mathfrak{l}\right)^{|C|}}{(\mathfrak{l}; q^{-1})_{|B|}} q^{-\sum_{i < j} (B_i - C_i)C_j} \prod_{i=1}^n \binom{B_i}{C_i}_{q^{-1}}. \tag{4-4}$$

Using the fact that $|B|, |C| \leq n$, one readily sees that this expression remains uniformly bounded as \mathfrak{l} varies in the extended complex plane while staying uniformly bounded away from the potential poles at $\{1, q, \dots, q^{n-1}\}$. Hence, we obtain an estimate of the form $|q^{nL}G_{v/\mu}(\epsilon, q\epsilon, \dots, q^{L-1}\epsilon)| \leq \text{const}^{|\nu|}$ for such \mathfrak{l} . On the other hand, we also have

$$|f_v(x_1, \dots, x_n)| \leq \text{const} \prod_{i=1}^n \left| \frac{x_i - s}{1 - sx_i} \right|^{|\nu|}, \tag{4-5}$$

which follows from the fact that in the partition function [\(2-23\)](#) only vertices of the form



have the number of appearances that is not a priori bounded, and the weight of such a vertex in the j -th row, according to [\(2-13\)](#), is $(x_j - s)/(1 - sx_j)$ (at least sufficiently far to the right, even if there are finitely many column inhomogeneities). We conclude that for the x_i 's in a sufficiently small neighbourhood of s and \mathfrak{l} staying away from $\{1, q, \dots, q^{n-1}\}$, the series in the left-hand side of [\(4-3\)](#) converges uniformly, yielding an analytic function of \mathfrak{l} . Clearly, the right-hand side of [\(4-3\)](#) is also analytic in \mathfrak{l} , which implies that [\(4-3\)](#) holds under the same conditions on the x_i 's and \mathfrak{l} . Substituting $\mathfrak{l} = \infty$ (equivalently, $\epsilon = 0$) into [\(4-3\)](#) leads to [\(4-2\)](#).

Once (4-2) is proved for x_i 's sufficiently close to s and $|q| < 1$, we can relax the assumptions by further analytic continuation in these parameters (although we will not need $|q| \geq 1$ below). In particular, since the explicit formula (3-4) for $\mathcal{G}_{v/\mu}$ shows $\text{const} \cdot |s|^{|\nu|}$ behaviour of $\mathcal{G}_{v/\mu}$ for large ν , using (4-5) we can extend the equality to x_i 's satisfying (4-1). \square

Taking $\mu = (0, 0, \dots, 0)$ in Proposition 4.1, we obtain the following:

Corollary 4.2. For $x_1, \dots, x_n \in \mathbb{C}$ satisfying (4-1), we have

$$\prod_{i=1}^n \frac{1 - sx_i}{s(s - x_i)} \cdot \sum_{\nu: \text{all } \nu_i > 0} (-s)^{|\nu|} f_\nu(x_1, \dots, x_n) = 1, \quad (4-6)$$

where the summation is over all compositions ν of length n with no zero parts. In the inhomogeneous setting of Section 3A, assuming that $(s_0, \xi_0) = (s, 1)$, a similar formula holds, where one needs to replace $(-s)^{|\nu|}$ in the left-hand side by $\prod_{i=1}^n \prod_{j=0}^{\nu_i-1} (-s_j)$.

Proof. This is given by straightforward substitution of $\mu = (0, \dots, 0)$ and (3-12) into (4-2), with the evaluation

$$f_{(0, \dots, 0)}(x_1, \dots, x_n) = \prod_{i=1}^n \frac{1 - s^2 q^{i-1}}{1 - sx_i}, \quad (4-7)$$

which follows from the fact that the corresponding partition function (2-23) is the product of L -weights (2-13) for the unique path configuration that gives a nonzero contribution. \square

It is not difficult to extend Proposition 4.1 and Corollary 4.2 to partially merged colours.

Proposition 4.3. Let λ be a composition with $|\lambda| = n$, μ be a λ -coloured composition, and $x_1, \dots, x_n \in \mathbb{C}$ satisfy (4-1). Then

$$\sum_{\nu \text{ is } \lambda\text{-coloured}} f_\nu(\lambda; x_1, \dots, x_n) \mathcal{G}_{\nu/\mu} = \prod_{i=1}^n \left(1 - \frac{x_i}{s}\right) \cdot f_\mu(\lambda; x_1, \dots, x_n), \quad (4-8)$$

which in the case of μ having only zero parts, can be rewritten as

$$\prod_{i=1}^n \frac{1 - sx_i}{s(s - x_i)} \cdot \sum_{\substack{\nu \text{ is } \lambda\text{-coloured;} \\ \text{all } \nu_i > 0}} (-s)^{|\nu|} f_\nu(\lambda; x_1, \dots, x_n) = 1. \quad (4-9)$$

In the inhomogeneous setting of Section 3A, relation (4-9) also holds under assumption that $(s_0, \xi_0) = (s, 1)$ and with $(-s)^{|\nu|}$ in the left-hand side replaced by $\prod_{i=1}^n \prod_{j=0}^{\nu_i-1} (-s_j)$.

Proof. Let $\theta : \{1, \dots, n\} \rightarrow \{1, \dots, \ell(\lambda)\}$ be the unique colour merging monotone map such that $\theta_*((1, \dots, 1)) = \lambda$; see Section 3C. Then we can sum (4-2) over rainbow μ with a given image $\theta_*(\mu)$. The right-hand side is immediately computed via (3-18), and in the left-hand side we use (3-19) and subsequently perform, using (3-18), a partial summation over ν 's with the same image $\theta_*(\nu)$. The result is (4-8), with $\theta_*(\mu)$ and $\theta_*(\nu)$ replaced back by μ and ν . The second relation (4-9) follows from (4-8) in the same way as in the rainbow case of Corollary 4.2. \square

Definition 4.4. For any $n \geq 1$, composition λ with $|\lambda| = n$, and $q, s, x_1, \dots, x_n \in \mathbb{C}$ satisfying (4-1), we define a (generally speaking, complex valued) probability measure on the subset of \mathcal{S}_λ (defined in (2-38)) consisting of λ -coloured compositions ν with no zero parts by

$$P(\{\nu\}) = \prod_{i=1}^n \frac{1 - sx_i}{s(s - x_i)} \cdot (-s)^{|\nu|} f_\nu(\lambda; x_1, \dots, x_n). \tag{4-10}$$


Graphically, the weight of this measure could be seen as partition functions of the form (2-23) with incoming colours on the left partially identified according to λ , with vertex weights L^{stoch} given by (2-13), and conditioned to have no exiting paths in the 0-th column (i.e., $A(0) = \mathbf{0}$ in terms of (2-23)).

In what follows we will also use the notation

$$f_\nu^{\text{stoch}}(\lambda; x_1, \dots, x_n) = (-s)^{|\nu|} f_\nu(\lambda; x_1, \dots, x_n). \tag{4-11}$$

In the inhomogeneous setting of Section 3A, we will assume that $(s_0, \xi_0) = (s, 1)$, and in the right-hand sides of (4-10) and (4-11) replace $(-s)^{|\nu|}$ by $\prod_{i=1}^n \prod_{j=0}^{\nu_i-1} (-s_j)$.

The graphical interpretation is based on the observation that the prefactor of the sums in (4-6) and (4-9) is exactly the inverse of the product of L^{stoch} -weights of vertices



in the 0-th column of a partition function of the form (2-23) with no turns in the 0-th column, and (4-11) is the result of computing the partition function for f_ν with L -weights replaced by the L^{stoch} -weights. Together with the stochasticity of the L^{stoch} -weights, this also implies (4-6) and (4-9).

Definition 4.5. In the context of Definition 4.4, for any λ -coloured composition $\mu = 0^{m^{(0)}} 1^{m^{(1)}} 2^{m^{(2)}} \dots$, introduce an observable \mathcal{O}_μ , whose values on λ -coloured compositions ν with no zero parts are given by

$$\mathcal{O}_\mu(\nu) = \prod_{x \geq 1} \prod_{i \geq 1} q^{m_i^{(x)} H_{>i}^{\nu/\mu}(x+1)} \binom{H_i^{\nu/\mu}(x+1)}{m_i^{(x)}}_q, \tag{4-12}$$

where we use the coloured height functions (3-5). Note that only nonzero parts of μ play a role in this definition.

For rainbow compositions, i.e., when $\lambda = (1, \dots, 1)$, the observables take a simpler form; see Corollary 3.4:

$$\begin{aligned} \mathcal{O}_\mu^{\text{rainbow}}(\nu) &= \prod_{\substack{x \geq 1, i \geq 1 \\ m_i^{(x)} = 1}} \mathbf{1}_{H_i^{\nu/\mu}(x+1)=1} q^{H_{>i}^{\nu/\mu}(x+1)} \\ &= \prod_{\substack{x \geq 1, i \geq 1 \\ m_i^{(x)} = 1}} \frac{q^{H_{>i+1}^{\nu/\mu}(x+1)} - q^{H_{>i}^{\nu/\mu}(x+1)}}{q - 1}. \end{aligned} \tag{4-13}$$

Example 4.6. In the colour-blind case $\lambda = (n)$, thinking of the coordinates of μ as being ordered: $\mu_1 \geq \mu_2 \geq \dots \geq \mu_n$, we can rewrite (4-12) as a shifted q -moment of the colourless height function:

$$\mathcal{O}_\mu^{\text{colour-blind}}(v) = \frac{(1 - q^{H^v(\mu_1+1)})(1 - q^{H^v(\mu_2+1)-1}) \dots (1 - q^{H^v(\mu_k+1)-k+1})}{\prod_{j \geq 1} (q; q)_{\text{mult}_j(\mu)}}, \quad (4-14)$$

where $\text{mult}_j(\mu) = \#\{i : \mu_i = j\}$ and $k = \max\{j : \mu_j > 0\}$.

We are now in position to formulate the main result of this section.

Theorem 4.7. *With notation of Definitions 4.4 and 4.5 above, we have*

$$\mathbb{E} \mathcal{O}_\mu = \mathbb{E}_v [\mathcal{O}_\mu(v)] = \frac{q^{\sum_{x \geq 1} \sum_{i > j} m_i^{(x)} m_j^{(x)}}}{\prod_{x \geq 0} (s^2; q)_{|\mathbf{m}^{(x)}|}} \prod_{i=1}^n (1 - s x_i) \cdot f_\mu^{\text{stoch}}(\lambda; x_1, \dots, x_n), \quad (4-15)$$

where the expectation is taken with respect to the weights (4-10), and the observables are given by (4-12) in the general case, or by (4-13) in the rainbow case.

The formula (4-15) also holds in the inhomogeneous setting of Section 3A under the assumption that $(s_x, \xi_x) = (s, 1)$ for any x such that $|\mathbf{m}^{(x)}| > 0$, and for $x = 0$.

Proof. Let us take the ratio of two skew-Cauchy identities (4-8) with μ and with $\mu = (0, \dots, 0)$. This yields

$$\frac{\sum_v f_v(\lambda; x_1, \dots, x_n) \mathcal{G}_{v/\mu}}{\sum_v f_v(\lambda; x_1, \dots, x_n) \mathcal{G}_v} = \frac{f_\mu(\lambda; x_1, \dots, x_n)}{f_{(0, \dots, 0)}(\lambda; x_1, \dots, x_n)},$$

which can be rewritten, via (4-11) and (4-7) (stated in the rainbow case, but also holding in the non-rainbow one for the same reasons), as

$$\frac{\sum_v f_v(\lambda; x_1, \dots, x_n) \mathcal{G}_v \cdot (\mathcal{G}_{v/\mu} / \mathcal{G}_v)}{\sum_v f_v(\lambda; x_1, \dots, x_n) \mathcal{G}_v} = \frac{\prod_{i=1}^n (1 - s x_i)}{(-s)^{|\mu|} (s^2; q)_n} f_\mu^{\text{stoch}}(\lambda; x_1, \dots, x_n). \quad (4-16)$$

The summations are taken over λ -coloured compositions v with no zero parts.

Observe that the left-hand side of (4-16) is exactly $\mathbb{E}_v(\mathcal{G}_{v/\mu} / \mathcal{G}_v)$; see Definition 4.4. The expression (3-4) for $\mathcal{G}_{v/\mu}$ (which is also the source of our assumption in the inhomogeneous setting) implies

$$\begin{aligned} \frac{\mathcal{G}_{v/\mu}}{\mathcal{G}_v} &= \frac{(s^2; q)_{|\mathbf{m}^{(0)}|}}{(-s)^{|\mu|} (s^2; q)_n} q^{\sum_{i=1}^n (m_i^{(0)} \sum_{n \geq j > i} (H_j^{v/\mu}(1) - m_j^{(0)}))} \prod_{i \geq 1} \binom{H_i^{v/\mu}(1)}{m_i^{(0)}}_q \\ &\quad \times \prod_{x \geq 1} (s^2; q)_{|\mathbf{m}^{(x)}|} q^{-\sum_{i > j} m_i^{(x)} m_j^{(x)}} \cdot \mathcal{O}_\mu(v). \end{aligned}$$

The only v -dependent part of this expression is $\mathcal{O}_\mu(v)$, and it stays under the expectation; the other factors can be moved to the right-hand side of (4-16). It only remains to notice that for $\mathcal{G}_{v/\mu}$ not to vanish, we must have $H_i^{v/\mu}(1) = m_i^{(0)}$ for all colours $i \geq 1$. To see that, it might be easiest to return to the expression (3-9) for the vertex weights in the one-row representation (3-7) of $\mathcal{G}_{v/\mu}$, and note that in the 0-th column we have $\mathbf{D} = \mathbf{0}$, which together with $\mathbf{A} + \mathbf{B} = \mathbf{C} + \mathbf{D}$ and $\mathbf{B} \geq \mathbf{C}$ (enforced by the q -binomial coefficients) implies $\mathbf{B} = \mathbf{C}$, which is exactly the statement we are making, as $B_i = H_i^{v/\mu}(1)$ and $C_i = m_i^{(0)}$. Taking these equalities into account and cancelling out common factors gives (4-15). \square

5. Integral representations for f_μ

The result of averaging the observable \mathcal{O}_μ in [Theorem 4.7](#) is the function f_μ , up to simple prefactors. The coloured composition μ here has the same dimension (i.e., number of parts) as the coloured compositions ν over which we are averaging (see [Definition 4.4](#)), and thus does not offer much reduction in complexity. The goal of this section is to show that f_μ has an integral representation of the dimension equal to the number of *nonzero* parts of μ . Hence, by choosing μ consisting of mostly zeros, we will be able to obtain tangible formulas for averages over ν 's of a growing dimension.

Definition 5.1. For a coloured composition μ (defined as in [Section 2H](#)), we will denote by $\mu^{\geq 1}$ the coloured composition obtained from μ by removing all of its zero parts. The colouring of $\mu^{\geq 1}$ will be the one naturally inherited from μ , and we will denote it by $\lambda^{\geq 1}$ if μ is λ -coloured, or by $\text{colour}(\mu^{\geq 1})$ if λ is not explicit.

Our first goal is to prove the following integral representation of f_μ for rainbow compositions.

Proposition 5.2. *Let $\mu = 0^{m^{(0)}} 1^{m^{(1)}} 2^{m^{(2)}} \dots$ be a rainbow composition of length $n \geq 1$ not consisting entirely of zeros, let $\mu^{\geq 1}$ be as in [Definition 5.1](#) with $m := n - |m^{(0)}| \geq 1$ being the length of $\mu^{\geq 1}$, and let $c_1 < \dots < c_m$ be the colours of the (necessarily nonzero) parts of $\mu^{\geq 1}$. Then*

$$f_\mu(x_1, \dots, x_n) = \frac{(s^2; q)_{|m^{(0)}|}}{(1 - sx_1) \dots (1 - sx_n)} \frac{(-1)^m}{(2\pi\sqrt{-1})^m} \oint \dots \oint_{\text{around}\{x_j^{-1}\}} \prod_{1 \leq i < j \leq m} \frac{y_j - y_i}{y_j - qy_i} \times f_{\mu^{\geq 1}}(y_1^{-1}, \dots, y_m^{-1}) \prod_{j=1}^m \left(\frac{y_j - s}{y_j} \frac{x_{c_j}}{1 - x_{c_j} y_j} \prod_{i>j}^n \frac{1 - qx_i y_j}{1 - x_i y_j} dy_j \right), \tag{5-1}$$

where (positively oriented) integration contours are chosen to encircle all points $\{x_j^{-1}\}_{j=1}^n$ and no other singularities of the integrand, or as *q-nested* closed simple curves with y_i -contour containing $q^{-1} \cdot (y_j\text{-contour})$ for all $i < j$, and all of the contours encircling $\{x_j^{-1}\}_{j=1}^n$.

The formula also holds in the column inhomogeneous setting under the assumption that in the 0-th column $(s_0, \xi_0) = (s, 1)$.

Remark 5.3. Contour integration around a specific set of singularities can be viewed formally as the sum of residues at those singularities, and such a sum would make sense if the parameters are such that required contours are not possible to construct. This gives a slightly different way of interpreting [\(5-1\)](#), as well as all the other integral representations below.

Remark 5.4. According to [Definition 5.1](#), $\mu^{\geq 1}$ is a coloured composition, and thus $f_{\mu^{\geq 1}}(y_1^{-1}, \dots, y_m^{-1})$ should really be written as $f_{\mu^{\geq 1}}(\lambda; y_1^{-1}, \dots, y_m^{-1})$, where λ is the colouring composition for $\mu^{\geq 1}$. However, since all parts of $\mu^{\geq 1}$ have different colours, and our vertex weights always depend on the colours only through their ordering, we could replace the colours $c_1 < \dots < c_m$ represented in $\mu^{\geq 1}$ by $1, 2, \dots, m$, and denoting the resulting rainbow composition by $\tilde{\mu}^{\geq 1}$, we would have $f_{\mu^{\geq 1}}(\lambda, y_1^{-1}, \dots, y_m^{-1}) = f_{\tilde{\mu}^{\geq 1}}(y_1^{-1}, \dots, y_m^{-1})$. It is this function that we denoted as $f_{\mu^{\geq 1}}(y_1^{-1}, \dots, y_m^{-1})$, thus slightly abusing the notation.

Remark 5.5. In the case $s_0 = 0$, there exists a fairly straightforward argument leading to a formula similar to (5-1). Namely, in that case vertices of the form $j \nearrow i$ with $j > i$ have weight zero (in the 0-th column) due to vanishing of the bottom right entry in (2-13). This means that the paths of colours c_1, \dots, c_m are not allowed make any vertical steps in column 0, as otherwise they would have no way of exiting this column to the right (because of the ordering of entering colours along the left boundary). This completely determines the configuration of paths in column 0, and the contribution of the remaining columns can be encoded as the matrix element

$$\langle \emptyset | \mathcal{C}_0(x_1) \cdots \mathcal{C}_0(x_{c_1-1}) \mathcal{C}_{c_1}(x_{c_1}) \mathcal{C}_0(x_{c_1+1}) \cdots \mathcal{C}_0(x_{c_m-1}) \mathcal{C}_{c_m}(x_{c_m}) \mathcal{C}_0(x_{c_m+1}) \cdots \mathcal{C}_0(x_n) | \mathcal{X} \rangle,$$

with $\mathcal{X} = \mu^{\geq 1} - 1^m$ and \mathcal{C} -row operators from Section 2C. The summation over \mathcal{X} of such expressions multiplied by $g_{\mathcal{X}}^*(y_1, \dots, y_m)$ is evaluated as an explicit product similar to (2-34) via the commutation relations (2-22), and then the coefficients of $g_{\mathcal{X}}^*$ are extracted by the orthogonality (2-36).

It is not clear, however, how to extend this argument to $s_0 \neq 0$, and we need to employ a different idea in the proof below.

Proof of Proposition 5.2. Our argument consists of two steps: We will first prove the formula for $(c_1, \dots, c_m) = (n - m + 1, \dots, n)$, and then show that the formula continues to hold when we reduce one of the c_j 's by 1. Iterations of such reductions will cover all possible choices of $1 \leq c_1 < \dots < c_m \leq n$.

Both steps work identically in the column homogeneous and inhomogeneous settings, and we will give the argument in the homogeneous case. The inhomogeneous analogs of several statements from Section 2 used below can be found in Section 3A.

Assume that $(c_1, \dots, c_m) = (n - m + 1, \dots, n)$. This means that paths that enter the partition function of the form (2-23) from the left in rows $1, \dots, n - m$ must immediately turn up and exit on top, while paths that enter in rows $n - m + 1, \dots, n$ must move to the right across the 0-th column (recall that no horizontal edge can carry more than one path). Hence, the configuration of paths in the 0-th column is completely determined, and the product of the L -weights (2-13) in this column gives

$$\frac{(s^2; q)_{|m^{(0)}|}}{\prod_{i=1}^{n-m} (1 - sx_i)} \prod_{j=n-m+1}^n \frac{x_j - s}{1 - sx_j}.$$

On the other hand, the contribution of the remaining columns can be written as

$$\prod_{j=n-m+1}^n \frac{1 - sx_i}{x_i - s} \cdot f_{\mu^{\geq 1}}(x_{n-m+1}, \dots, x_n),$$

where the prefactor is responsible for the fact that the partition function for $f_{\mu^{\geq 1}}(x_{n-m+1}, \dots, x_n)$ starts with column 0, and we only had weights of vertices in columns ≥ 1 remaining. We now need to show that the product of the last two expressions agrees with the right-hand side of (5-1). With our choice of c_i 's, the integrand is independent of x_1, \dots, x_{n-m} , and its only poles are at $y_i = x_j^{-1}$ with $n - m + 1 \leq j \leq n$. The number of integration variables thus coincides with the number of potential pole locations, and no two variables can have nonvanishing residues at the same location because of $\prod_{i < j} (y_i - y_j)$ in the integrand. Hence, the residue locations cover all points $x_{n-m+1}^{-1}, \dots, x_n^{-1}$ exactly once, and $\prod_{j=1}^m (y_j - s) x_{c_j}$ in the

integrand necessarily evaluates to $\prod_{j=n-m+1}^n (1 - sx_j)$ in every set of nontrivial residues taken. Moving this factor out of the integral, we are led to the following desired equality:

$$f_{\mu \geq 1}(x_{n-m+1}, \dots, x_n) = \frac{(-1)^m}{(2\pi\sqrt{-1})^m} \oint \cdots \oint_{\text{around } \{x_j^{-1}\}} \prod_{1 \leq i < j \leq m} \frac{y_j - y_i}{y_j - qy_i} \\ \times f_{\mu \geq 1}(y_1^{-1}, \dots, y_m^{-1}) \prod_{j=n-m+1}^n \left(\frac{1}{1 - x_j y_j} \prod_{i>j}^n \frac{1 - qx_i y_j}{1 - x_i y_j} \frac{dy_j}{y_j} \right). \quad (5-2)$$

Comparing to (2-37), we see that the difference is in $(-1)^m$ and the choice of contours, which both come from the same origin. Namely, let us take the right-hand side of (2-37) (with n replaced by m and (x_1, \dots, x_n) replaced by (x_{n-m+1}, \dots, x_n)), and deform the outermost y_m -contour in the outside direction, moving it through ∞ and closing around $\{x_j^{-1}\}$. The recursive construction of (rainbow) f_μ 's from Section 2E readily shows that $(y_1 \cdots y_m)^{-1} f_{\mu \geq 1}(y_1^{-1}, \dots, y_m^{-1})$ is a ratio of two polynomials in y 's with the denominator consisting only of factors of the form $(y_i - s)$, and $f_{\mu \geq 1}(y_1^{-1}, \dots, y_m^{-1})$ viewed as such a ratio has the numerator degree 1 less than the denominator degree in each of the variables y_i , $1 \leq i \leq m$. Hence, our deformation of the y_m collects no residues along the way and yields the factor (-1) for changing the contour direction.

Next, we do the same deformation with the y_{m-1} -contour. Following the same reasoning, there is only one possible pole along the way: $y_{m-1} = q^{-1}y_m$. If we leave this potential singularity inside the y_{m-1} -contour, and proceed similarly for subsequent deformations, then we end up with two q -nested contours surrounding $\{x_j^{-1}\}$. It turns out that the q -nestedness is not necessary. Indeed, a direct inspection of the integrand shows that the only possible pole of y_m inside its current contour is x_m^{-1} , and then the factor $(1 - qx_m y_{m-1})$ makes the residue at $y_{m-1} = q^{-1}y_m$ vanish. Hence, we can close the y_{m-1} -contour around $\{x_j^{-1}\}$, orient it positively, and acquire another (-1) .

Continuing with this procedure for $y_{m-2}, y_{m-3}, \dots, y_1$ -contours (in this order), we turn (2-37) into (5-2), thus completing the first step of the proof.

Let us now see why lowering of the c_j 's in (5-1) keeps the formula intact. From the point of view of the (rainbow) composition μ , replacing $c_j \mapsto c_j - 1$ (assuming there is no $i \neq j$ such that $c_i = c_j - 1$) is equivalent to swapping $\mu_{c_j-1} = 0$ and $\mu_{c_j} > 0$. This can be done with the help of the exchange relations (2-31) by acting on the right-hand side of (5-1) by

$$T_{c_j-1} = q - \frac{x_{c_j-1} - qx_{c_j}}{x_{c_j-1} - x_{c_j}} (1 - s_{c_j-1});$$

see (2-32). The only part of the right-hand side of (5-1) that is not symmetric in (x_{c_j-1}, x_{c_j}) is the factor $x_{c_j}/(1 - x_{c_j}y_j)$ (note that the contours are symmetric too), and applying T_{c_j-1} to it we read

$$\frac{qx_{c_j}}{1 - x_{c_j}y_j} - \frac{x_{c_j-1} - qx_{c_j}}{x_{c_j-1} - x_{c_j}} \left(\frac{x_{c_j}}{1 - x_{c_j}y_j} - \frac{x_{c_j-1}}{1 - x_{c_j-1}y_j} \right) = \frac{x_{c_j-1}}{1 - x_{c_j-1}y_j} \frac{1 - qx_{c_j}y_j}{1 - x_{c_j}y_j}.$$

This recovers the integrand of (5-1) with $c_j \mapsto c_j - 1$ and completes the proof. □

Remark 5.6. While the integral representation (2-37) allowed for moving the contours through ∞ , this is no longer true for (5-1), as the added factors $(y_i - s)$ create poles at ∞ .

The main result of this section is a generalization of Proposition 5.2 to coloured nonrainbow compositions.

Theorem 5.7. *Let λ be a composition of weight $|\lambda| = n$ with partial sums $\sum_{i=1}^k \lambda_i = \ell_k$, $k \geq 1$; $\ell_0 := 0$. Further, let $\mu = 0^{m^{(0)}} 1^{m^{(1)}} 2^{m^{(2)}} \dots$ be a λ -coloured composition of length $\ell(\mu) = |\lambda| = n$ not consisting entirely of zeros, let $\mu^{\geq 1}$ be as in Definition 5.1 with inherited colouring $\lambda^{\geq 1}$ and $m := n - |\mathbf{m}^{(0)}|$ being the length of $\mu^{\geq 1}$, and let $c_1 < \dots < c_\alpha$ be the colours of the (necessarily nonzero) parts of $\mu^{\geq 1}$. Finally, let $m_1, \dots, m_\alpha \geq 1$ be the number of parts of $\mu^{\geq 1}$ of colours c_1, \dots, c_α , respectively, and denote $\mathbf{m}[a, b] = m_a + m_{a+1} + \dots + m_b$. Then*

$$\begin{aligned}
 f_\mu(\lambda; x_1, \dots, x_n) = & \frac{(s^2; q)_{|\mathbf{m}^{(0)}|}}{(1 - sx_1) \cdots (1 - sx_n)} \frac{1}{(2\pi\sqrt{-1})^m} \\
 & \times \oint \cdots \oint_{\text{around}\{x_j^{-1}\}} \prod_{1 \leq i < j \leq m} \frac{y_j - y_i}{y_j - qy_i} \\
 & \times \prod_{k=1}^{\alpha} \left(\sum_{j=0}^{m_k} \frac{(-1)^j q^{\binom{m_k-j}{2}}}{(q; q)_j (q; q)_{m_k-j}} \prod_{p > \mathbf{m}[1, k-1]}^{j + \mathbf{m}[1, k-1]} \prod_{a > \ell_{c_k-1}}^n \frac{1 - qx_a y_p}{1 - x_a y_p} \prod_{r > j + \mathbf{m}[1, k-1]}^{\mathbf{m}[1, k]} \prod_{b > \ell_{c_k}}^n \frac{1 - qx_b y_r}{1 - x_b y_r} \right) \\
 & \times f_{\mu^{\geq 1}}(\lambda^{\geq 1}; y_1^{-1}, \dots, y_m^{-1}) \prod_{i=1}^m \frac{(y_i - s) dy_i}{y_i^2}, \quad (5-3)
 \end{aligned}$$

where (positively oriented) integration contours are chosen to encircle all points $\{x_j^{-1}\}_{j=1}^n$ and no other singularities of the integrand, or as q -nested closed simple curves with y_i -contour containing $q^{-1} \cdot (y_j\text{-contour})$ for all $i < j$, and all of the contours encircling $\{x_j^{-1}\}_{j=1}^n$. The contours can also be chosen to either encircle or not encircle the point 0.

The formula also holds in the column inhomogeneous setting under the assumption that in the 0-th column $(s_0, \xi_0) = (s, 1)$.

Remark 5.8. For $\lambda = (1, \dots, 1)$ and μ a rainbow composition, we have $\ell_k \equiv k$, $\alpha = m$, $m_1 = \dots = m_\alpha = 1$, $\mathbf{m}[1, k] \equiv k$, and the middle line of (5-3) evaluates to

$$(-1) \left(\frac{1}{1-q} - \frac{1}{1-q} \frac{1 - qx_{c_k} y_k}{1 - x_{c_k} y_k} \right) \prod_{b > c_k} \frac{1 - qx_b y_k}{1 - x_b y_k} = \frac{x_{c_k} y_k}{1 - x_{c_k} y_k} \prod_{b > c_k} \frac{1 - qx_b y_k}{1 - x_b y_k},$$

thus reproducing (5-1).

Proof. The central role in the argument is played by the following:

Lemma 5.9. *For any positive integers ℓ, l and m which satisfy $\ell + m < l$, and any symmetric function $\Phi(z_1, \dots, z_m)$ that is holomorphic in a neighbourhood of the domain encircled by the integration contours,*

one has

$$\begin{aligned} & \sum_{\{-1 \geq i_1 > \dots > i_m \geq \ell\}} \oint \cdots \oint_{\text{around}\{\mathfrak{x}_j\}} \prod_{i < j} \frac{z_j - z_i}{z_j - qz_i} \Phi(z_1, \dots, z_m) \prod_{p=1}^m \left(\frac{1}{z_p - \mathfrak{x}_{i_p+1}} \prod_{j=1}^{i_p} \frac{qz_p - \mathfrak{x}_j}{z_p - \mathfrak{x}_j} dz_p \right) \\ &= \oint \cdots \oint_{\text{around}\{\mathfrak{x}_j\}} \prod_{i < j} \frac{z_j - z_i}{z_j - qz_i} \Phi(z_1, \dots, z_m) \\ & \quad \times \left(\sum_{j=0}^m \frac{(-1)^j q^{\binom{m-j}{2}}}{(q; q)_j (q; q)_{m-j}} \prod_{p>0}^j \prod_{a=1}^{\ell} \frac{qz_p - \mathfrak{x}_a}{z_p - \mathfrak{x}_a} \prod_{r>j}^m \prod_{b=1}^{\ell} \frac{qz_r - \mathfrak{x}_b}{z_r - \mathfrak{x}_b} \right) \prod_{p=1}^m \frac{dz_p}{z_p}, \end{aligned} \quad (5-4)$$

where (positively oriented) integration contours are chosen (independently in the two sides of (5-4)) to encircle all points $\{\mathfrak{x}_j\}_{j=1}^{\ell} \subset \mathbb{C}$ and no other singularities of the integrand, or as q -nested closed simple curves with z_i -contour containing $q^{-1} \cdot (z_j$ -contour) for all $i < j$, and all of the contours encircling $\{\mathfrak{x}_j\}_{j=1}^{\ell}$. The contours can also be chosen to either encircle or not encircle the point 0.

Let us postpone the proof of Lemma 5.9 and use it for the proof of Theorem 5.7 first.

Let $\theta : \{1, \dots, n\} \rightarrow \{1, \dots, \ell(\lambda)\}$ be the unique monotone map such that

$$\theta^{-1}(k) = \{\ell_{k-1} + 1, \dots, \ell_k\} \quad \text{for all } 1 \leq k \leq \ell(\lambda). \quad (5-5)$$

We can then use the colour merging relation (3-18) to obtain a formula for λ -coloured μ 's from the formula (5-1) of Proposition 5.2 for the rainbow ones. This means that we need to sum the right-hand sides of (5-1), written for a rainbow composition $\tilde{\mu}$, over all $\tilde{\mu}$ with $\theta_*(\tilde{\mu}) = \mu$.

Such a summation can be performed in two steps. At the first step we choose, for each colour c_k represented in $\mu^{\geq 1}$, $1 \leq k \leq \alpha$, the m_k colours in $\theta^{-1}(c_k)$ that are represented in $\tilde{\mu}^{\geq 1}$. At the second step we choose, for each k , $1 \leq k \leq \alpha$, different assignments of the chosen m_k colours in $\theta^{-1}(c_k)$ to the m_k parts of $\mu^{\geq 1}$ that have colour c_k . The summation of the second step is exactly the colour merging applied to $f_{\tilde{\mu}^{\geq 1}}$ (no other factors of the integrand of (5-1) depend on the choices in the second step), and (3-18) shows that the second step summation results in replacing $f_{\tilde{\mu}^{\geq 1}}$ in the integrand by $f_{\mu^{\geq 1}}$.

Returning to the first step summation, we see that $f_{\mu^{\geq 1}}$ in the integrand is now independent of the choices involved, and we can focus on the rest of the integrand. For each colour c_k , $1 \leq k \leq \alpha$, we are choosing $c_k^{(1)} < \dots < c_k^{(m_k)}$ in $\theta^{-1}(c_k)$, or, according to (5-5), $\ell_{c_k-1} < c_k^{(1)} < \dots < c_k^{(m_k)} \leq \ell_{c_k}$. Now the sum over such m_k -tuple of indices can be computed, for each $1 \leq k \leq \alpha$, using Lemma 5.9, where one needs to make substitutions

$$\begin{aligned} m &\mapsto m_k, & \mathfrak{k} &\mapsto (n - \ell_{c_k}), & \mathfrak{l} &\mapsto (n - \ell_{c_k-1}), & (z_1, \dots, z_m) &\mapsto (y_{m[1,k-1]+1}, \dots, y_{m[1,k]}), \\ (\mathfrak{x}_1, \mathfrak{x}_2, \dots, \mathfrak{x}_{\mathfrak{l}}) &\mapsto (x_n^{-1}, x_{n-1}^{-1}, \dots, x_{\ell_{c_k-1}+1}^{-1}), & (\mathfrak{x}_1, \mathfrak{x}_2, \dots, \mathfrak{x}_{\mathfrak{k}}) &\mapsto (x_n^{-1}, x_{n-1}^{-1}, \dots, x_{\ell_{c_k}+1}^{-1}), \\ (\mathfrak{l} > i_1 > \dots > i_m \geq \mathfrak{k}) &\mapsto (n - \ell_{c_k-1} > n - c_k^{(1)} > \dots > n - c_k^{(m_k)} \geq n - \ell_{c_k}), \end{aligned}$$

and collect the factors that do not correspond to the ones in the left-hand side of (5-4) into a $\Phi(z_1, \dots, z_m)$. The holomorphicity of $\Phi(z_1, \dots, z_m)$ in a neighbourhood of $\{\mathfrak{x}_j\}_{j=1}^{\ell}$ is readily visible; and the fact that it

is symmetric in z_1, \dots, z_m follows from the fact that $f_{\mu \geq 1}(y_1^{-1}, \dots, y_m^{-1})$, which enters $\Phi(z_1, \dots, z_m)$ as the only factor that is not manifestly symmetric, is symmetric in

$$(y_{m[1,k-1]+1}, \dots, y_{m[1,k]})$$

thanks to the commutativity of the $C_i(x)$ operators (for a fixed i and varying x) in the definition (2-41) of f -functions for coloured compositions.

Comparing the right-hand sides of (5-3) and (5-4), we see that this completes the proof of Theorem 5.7 modulo the proof of Lemma 5.9. □

Proof of Lemma 5.9. While a direct book-keeping of residues of the two sides of (5-4) might be possible, we will use the theory of Hall–Littlewood processes as a shortcut, with the work [Borodin et al. 2016a, Section 2] as our main reference; a more detailed description can be found in [Borodin and Corwin 2014, Section 2]. See also Remark 5.10 on the origin of the argument given below.

Consider an ascending Hall–Littlewood process with weights on sequences of partitions $\lambda^{(1)}, \dots, \lambda^{(n)}$ proportional to

$$P_{\lambda^{(1)}}(x_1) P_{\lambda^{(2)}/\lambda^{(1)}}(x_2) P_{\lambda^{(n)}/\lambda^{(n-1)}}(x_n) Q_{\lambda^{(n)}}(\rho), \quad \ell(\lambda^{(k)}) \leq k, \quad 1 \leq k \leq n, \quad (5-6)$$

with, generally speaking, complex parameters $\{x_i\}_{i=1}^n$, ρ being the specialization of the algebra of symmetric functions into a sequence of variables (b_1, b_2, \dots) , and P_* and Q_* being the Hall–Littlewood symmetric functions.

Our argument is based on [Borodin et al. 2016a, Proposition 2.2], see also [Borodin and Corwin 2014, Proposition 2.2.14], which says that for any $n \geq m_1 \geq \dots \geq m_n \geq 1$, one has

$$\begin{aligned} \mathbb{E}_{HL}(q^{m_1 - \ell(\lambda^{(m_1)})} \dots q^{m_n - \ell(\lambda^{(m_n)})}) \\ = \frac{q^{\binom{n}{2}}}{(2\pi\sqrt{-1})^n} \oint \dots \oint \prod_{i < j} \frac{z_j - z_i}{z_j - qz_i} \prod_{l=1}^n \left(\prod_{j \geq 1} \frac{1 - z_l b_j}{1 - qz_l b_j} \prod_{i=1}^{m_l} \frac{qz_l - x_i}{z_l - x_i} \frac{dz_l}{z_l} \right), \end{aligned} \quad (5-7)$$

where (positively oriented) z_j -contours are such that they surround $\{x_j\}_{j=1}^{m_1}$ and 0, and they are also q -nested in the sense that z_i -contour contains $q \cdot (z_j$ -contour) for all $i < j$; no other poles are taken into account.

Let us fix $1 \leq \ell < l \leq n$, and consider the sum

$$\begin{aligned} \sum_{\ell \leq i_m < i_{m-1} < \dots < i_1 < l} \prod_{p=1}^m \frac{q^{i_p - \ell(\lambda^{(i_p)})} - q^{i_p + 1 - \ell(\lambda^{(i_p+1)})}}{1 - q} \\ = q^{\binom{m}{2}} \cdot q^{m(\ell - \ell(\lambda^{(\ell)}))} \cdot \binom{l - \ell(\lambda^{(l)}) - \ell + \ell(\lambda^{(\ell)})}{m}_q. \end{aligned} \quad (5-8)$$

The equality between the two sides of (5-8) is a (nonobvious) special case of Corollary 3.7. More exactly, choosing $Q = q$, $\alpha_i \equiv 1$, $|\beta| = m$, and reversing the order of indices of α_i 's and β_i 's, turns (3-16) into

$$q^{-\binom{m}{2}} \cdot \sum_{\substack{\beta_1, \dots, \beta_n \in \{0,1\} \\ \beta_1 + \dots + \beta_n = m}} \prod_{i=1}^n q^{\beta_i \cdot (i-1)} = \binom{n}{m}_q. \quad (5-9)$$

Let $1 \leq k_m < k_{m-1} < \dots < k_1 \leq n$ be the set of index values k for which $\beta_k = 1$ in (5-9). Observe that

$$\frac{q^{j-\ell(\lambda(j))} - q^{j+1-\ell(\lambda(j+1))}}{1 - q} = \begin{cases} q^{j-\ell(\lambda(j))} = q^{\xi-\ell(\lambda(\xi))} \cdot q^{j-\ell(\lambda(j))-\xi+\ell(\lambda(\xi))}, & \ell(\lambda(j)) = \ell(\lambda(j+1)), \\ 0, & \text{otherwise,} \end{cases}$$

where ‘‘otherwise’’ refers to the only possible alternative $\ell(\lambda(j)) + 1 = \ell(\lambda(j+1))$.

Set $n = l - \ell(\lambda(l)) - \xi + \ell(\lambda(\xi))$, and let $\xi \leq j_1 < j_2 < \dots < j_m < l$ be all the values of $j \in [\xi, l)$ such that $\ell(\lambda(j)) = \ell(\lambda(j+1))$; equivalently $(j+1 - \ell(\lambda(j+1))) - (j - \ell(\lambda(j)))$ equals 1 rather than 0. These are all possible values that summation indices $i_1 > \dots > i_m$ in the left-hand side of (5-8) can take to produce a nonzero term; let $j_{l_1} > \dots > j_{l_m}$ be the corresponding choices. Then matching $(k_1, \dots, k_m) \equiv (l_1, \dots, l_m)$ establishes the equivalence of (5-8) and (5-9), thus proving (5-8).

Our next step is to compute the averages, with respect to the ascending Hall–Littlewood process, of both sides of (5-8) using (5-7).

The left-hand side of (5-8) is a simple linear combination of those from (5-7). Moving that linear combination inside the integrand and observing that

$$\frac{1}{1 - q} \left(\prod_{i=1}^{i_p} \frac{qz_p - \xi_i}{z_p - \xi_i} - \prod_{i=1}^{i_p+1} \frac{qz_p - \xi_i}{z_p - \xi_i} \right) = \frac{z_p}{z_p - \xi_{i_p+1}} \prod_{i=1}^{i_p} \frac{qz_p - \xi_i}{z_p - \xi_i},$$

we obtain

$$\begin{aligned} \mathbb{E}_{HL} \left(\sum_{\xi \leq i_m < i_{m-1} < \dots < i_1 < l} \prod_{p=1}^m \frac{q^{i_p - \ell(\lambda(i_p))} - q^{i_p + 1 - \ell(\lambda(i_p + 1))}}{1 - q} \right) \\ = \sum_{\xi \leq i_m < i_{m-1} < \dots < i_1 < l} \frac{q^{\binom{m}{2}}}{(2\pi\sqrt{-1})^m} \oint \dots \oint \prod_{i < j} \frac{z_j - z_i}{z_j - qz_i} \\ \times \prod_{p=1}^m \left(\prod_{j \geq 1} \frac{1 - z_p b_j}{1 - qz_p b_j} \frac{1}{z_p - \xi_{i_p+1}} \prod_{i=1}^{i_p} \frac{qz_p - \xi_i}{z_p - \xi_i} dz_p \right), \end{aligned} \tag{5-10}$$

with the same integration contours as in (5-7).

Note that 0 is no longer a potential singularity of the integrand; thus, the contour may or may not contain it. Let us also explain why the presence or absence of the potential poles at $z_i = q^{-1}z_j$ does not affect the value of the integral. Generally speaking, the integral with q -nested contours is equal to the sum of residues at $z_p = q^{-k_p}\xi_{l_p}$, $1 \leq p \leq m$, for certain values of $k_p \geq 0$ and $l_p \geq 1$, that arises by sequential evaluation of residues inside the z_m, z_{m-1}, \dots, z_1 -contours in that order. Let p^* be the maximal index such that $k_{p^*} > 0$. Since the z_m -contour encircles only the poles at ξ_j ’s, we must have $p^* \leq m - 1$. Also, since this pole must have come from a denominator factor $z_{j^*} - qz_{p^*}$ with $j^* > p^*$, due to the maximality of p^* , the pole must be at $z_{p^*} = q^{-1}\xi_{l^*}$ with $1 \leq l^* \leq i_{j^*}$. But the integrand contains the factor $(qz_{p^*} - \xi_{l^*}^*)$, which will turn the residue into 0 (note that we need the fact that $i_{p^*} \geq i_{j^*}$ to guarantee the presence of this factor). We conclude that the q -nestedness is irrelevant for the value of the integral.

Let us proceed to computing the Hall–Littlewood expectation of the right-hand side of (5-8).

Using the q -binomial theorem

$$\sum_{j \geq 0} \frac{(a; q)_j}{(q; q)_j} z^j = \frac{(az; q)_\infty}{(z; q)_\infty} = \left(1 - \frac{z}{q}\right) \cdots \left(1 - \frac{z}{q^m}\right) \quad \text{for } a = q^{-m},$$

we obtain

$$\binom{A}{m}_q = \frac{(1-q^A)(1-q^{A-1}) \cdots (1-q^{A-m+1})}{(q; q)_m} = \sum_{j=0}^m \frac{(q^{-m}; q)_j}{(q; q)_j (q; q)_m} q^{(A+1)j} = \sum_{j=0}^m \frac{(-1)^j q^{(A+1)j - mj + \binom{j}{2}}}{(q; q)_j (q; q)_{m-j}}.$$

Hence, the right-hand side of (5-8) can be written as

$$q^{\binom{m}{2}} \cdot q^{m(\ell - \ell(\lambda(\ell)))} \cdot \binom{\ell - \ell(\lambda(\ell)) - \ell + \ell(\lambda(\ell))}{m}_q = \sum_{j=0}^m \frac{q^{\binom{m-j}{2}}}{(q; q)_j (q; q)_{m-j}} q^{j(\ell - \ell(\lambda(\ell))) + (m-j)(\ell - \ell(\ell))},$$

where for powers of q we used $\binom{m}{2} - mj + j + \binom{j}{2} = \binom{m-j}{2}$. Employing (5-7), we now obtain

$$\begin{aligned} \mathbb{E}_{HL} \left(q^{\binom{m}{2}} \cdot q^{m(\ell - \ell(\lambda(\ell)))} \cdot \binom{\ell - \ell(\lambda(\ell)) - \ell + \ell(\lambda(\ell))}{m}_q \right) \\ = \sum_{j=0}^m \frac{q^{\binom{m-j}{2}}}{(q; q)_j (q; q)_{m-j}} \frac{q^{\binom{m}{2}}}{(2\pi\sqrt{-1})^m} \oint \cdots \oint \prod_{i < j} \frac{z_j - z_i}{z_j - qz_i} \\ \times \prod_{p>0} \prod_{a=1}^{\ell} \frac{qz_p - \varkappa_a}{z_p - \varkappa_a} \prod_{r>j} \prod_{b=1}^{\ell} \frac{qz_r - \varkappa_b}{z_r - \varkappa_b} \prod_{p=1}^m \prod_{j \geq 1} \frac{1 - z_p b_j}{1 - qz_p b_j} \frac{dz_p}{z_p}, \end{aligned} \quad (5-11)$$

where the integration contours are as for (5-7).

Since the two of sides of (5-8) are equal, the right-hand sides of (5-10) and (5-11) are also equal. This is literally the desired statement of Lemma 5.9, equation (5-4), with a specific choice of

$$\Phi(z_1, \dots, z_m) = \phi(z_1) \cdots \phi(z_m), \quad \phi(z) = \text{const} \prod_{j \geq 1} \frac{1 - zb_j}{1 - qzb_j}, \quad (5-12)$$

and a specific choice of contours for the right-hand side. The equality (of the right-hand sides of (5-10) and (5-11)) has been thus proven for generic $\{\varkappa_i\}$ and $\{b_j\}$, although certain inequalities on them are needed to make sure that the weights (5-6) are summable (see [Borodin et al. 2016a, Section 2] for details). However, the equality itself is an identity of finite sums (of residues), and the restrictions on $\{\varkappa_i\}$ and $\{b_j\}$ can thus be removed by analytic continuation.

The two sides of (5-4) computed as (finite) sums of residues are linear combinations of values of the function $\Phi(z_1, \dots, z_m)$ at m -tuples of distinct (because of $\prod_{i < j} (z_j - z_i)$ in the integrand) points from the list of possible singularities, consisting of $\{\varkappa_i\}$, their q -multiples, and 0. Values of $\Phi(z_1, \dots, z_m)$ with permuted z_j 's are equal (due to the symmetry of Φ) and can be grouped together; their coefficients are certain explicit rational functions of $\{\varkappa_i\}$ and q . Fix an m -tuple of distinct possible poles. Using the freedom in the choice of $\{b_j\}$ and the constant prefactor in (5-12), we can make $\phi(z)$ to be arbitrarily close to 1 at the points of the chosen m -tuple, and arbitrarily close to 0 at all the other potential singularities (we

are assuming that $\{x_i\}$ are generic, we only need to prove an identity between rational functions in them).⁸ This will lead to Φ being close to 1 at the chosen m -tuple and its permutations, and close to 0 at all other possible m -tuples. Since we already proved the equality of the two sides of (5-4) for all Φ 's of the form (5-12), this implies the equality of the coefficients of Φ evaluated at the chosen m -tuple (together its permutations) in residue expansions of the two sides of (5-4). This proves (5-4) for arbitrary Φ , but still with the particular choice of integration contours in the right-hand side, which are q -nested and include 0.

However, we already know that the declared freedom for the choice of contours is valid for the left-hand side; see the argument after (5-10). This implies that the values of Φ at m -tuples that include either 0 or a nontrivial q -multiple of a x_i do not contribute to the left-hand side. Since we already know that such coefficients are the same on both sides, these values must not contribute to the right-hand side either, which means that we can use the same freedom of contours in the right-hand side without changing the value of the integral. \square

Remark 5.10. Let us comment on the origin of our proof of Lemma 5.9 that might have looked somewhat cryptic. If, in the setting of Theorem 5.7, μ has no zero parts and only one colour, then f_μ is colour-blind, according to (2-29). Theorem 4.7 then implies that it is given by an average of a q -moment type observable (4-14) over a colour-blind stochastic vertex model. The height function of the colour-blind stochastic vertex models can be interpreted, along any down-right path in a quadrant (which includes horizontal lines), as lengths of partitions distributed according to Hall–Littlewood processes; this was the main result of [Borodin et al. 2016a]. Thus, we get an expression for f_μ in the form of an average over a Hall–Littlewood process.

On the other hand, the symmetrization of colours in (2-29) can also be taken after the computation of the expectation, with respect to a rainbow coloured model, in the left-hand side of (4-15). As was shown in [Borodin and Wheeler 2018, Chapter 10], the distribution of coloured height functions at a single observation point can also be described via lengths of partitions in a Hall–Littlewood process, and this is where the computation of that expectation can take place. Once the corresponding average over the Hall–Littlewood process is computed, one can perform its colour symmetrization.

The two resulting expressions must be the same - the operations of averaging over our measure and symmetrizing over colours commute. Understanding the reason for that in the language of the Hall–Littlewood processes is not too difficult, this is essentially (5-8). Rewriting what it means in terms of integral representations for averages of Hall–Littlewood observables results in a general identity for symmetric functions, which is exactly our Lemma 5.9.

6. Observables of stochastic lattice models

The purpose of this section is to combine the results of Sections 4-5 in order to obtain integral representations for averages of the observables introduced in Definition 4.5, as well as to explore corollaries thereof.

⁸Indeed, $\phi(z) = \text{const} \cdot \psi(x)/\psi(qz)$ with ψ being an arbitrary polynomial that can be chosen to approximate any values at any finite set of points; the constant in front is needed to control $\phi(0)$.

6A. The main result. Recalling graphical interpretation of [Definition 4.4](#), we consider a stochastic lattice model in the quadrant $\mathbb{Z}_{\geq 1} \times \mathbb{Z}_{\geq 1}$, with vertex weights given by $L_{x_i}^{\text{stoch}}$ in [\(2-11\)](#) (see also [\(2-13\)](#) and [Table 3](#)) for the vertices in row i from the bottom. The boundary conditions are as follows: No paths (equivalently, only paths of colour 0) enter the quadrant through its bottom boundary; and along the left boundary we have a single path entering at every row, with the bottom λ_1 paths having colour 1, next λ_2 paths having colour 2, and so on. We focus on the state of the model between row n and row $n+1$; that is, we record the locations where the paths of colour 1, 2, \dots exit the n -th row upwards as a λ -coloured composition ν , where λ is the composition with parts $\lambda_1, \lambda_2, \dots$, and we truncate this sequence so that $|\lambda| = n$.⁹ The partial sums of λ are denoted as $\sum_{i=1}^k \lambda_i = \ell_k$, $k \geq 1$; $\ell_0 := 0$.

The model is also allowed to have column inhomogeneities $\{s_j, \xi_j\}_{j \geq 1}$, which replace the (s, x_i) parameters in the L^{stoch} -weight of the vertex in row i and column j by $(s_j, \xi_j x_i)$; see [Section 3A](#). For convenience, we assume that the number of column inhomogeneities is finite. (For our results this assumption is not restrictive as the state of the model far enough to the right will not play any role, and thus, due to stochasticity, the column inhomogeneities there can be chosen freely.)

Also recall the observables \mathcal{O}_μ of [Definition 4.5](#), defined for any λ -coloured composition $\mu = 0^{m^{(0)}} 1^{m^{(1)}} 2^{m^{(2)}} \dots$, that take values

$$\begin{aligned} \mathcal{O}_\mu(\nu) &= \prod_{x \geq 1} \prod_{i \geq 1} q^{m_i^{(x)} H_{>i}^{\nu/\mu}(x+1)} \binom{H_i^{\nu/\mu}(x+1)}{m_i^{(x)}}_q \\ &= \prod_{x \geq 1} \prod_{i \geq 1} \frac{(q^{H_{>i}^{\nu/\mu}(x+1)} - q^{H_{\geq i}^{\nu/\mu}(x+1)}) (q^{H_{>i}^{\nu/\mu}(x+1)} - q^{H_{\geq i}^{\nu/\mu}(x+1)-1}) \dots (q^{H_{>i}^{\nu/\mu}(x+1)} - q^{H_{\geq i}^{\nu/\mu}(x+1)-m_i^{(x)}+1})}{(q; q)_{m_i^{(x)}}} \end{aligned} \quad (6-1)$$

given in terms of the coloured height functions [\(3-5\)](#).

For any λ -coloured μ , we make a new coloured composition $\mu^{\geq 1}$ that consists of its nonzero parts coloured in the same way with the labelling composition of $\mu^{\geq 1}$ denoted by $\lambda^{\geq 1}$; see [Definition 5.1](#). Clearly, given the colouring λ of μ , μ is uniquely reconstructed from $\mu^{\geq 1}$. We use the notation $m := n - |\mathbf{m}^{(0)}|$ for the length of $\mu^{\geq 1}$, denote by $c_1 < \dots < c_\alpha$ the colours of parts of $\mu^{\geq 1}$, and denote by $\mathbf{m}_1, \dots, \mathbf{m}_\alpha \geq 1$ the number of parts of $\mu^{\geq 1}$ of colours c_1, \dots, c_α , respectively. Set $\mathbf{m}[a, b] = \mathbf{m}_a + \mathbf{m}_{a+1} + \dots + \mathbf{m}_b$.

We can now state the main result of this paper.

Theorem 6.1. *With the above notation, we have*

$$\begin{aligned} \mathbb{E} \mathcal{O}_\mu &= \frac{q^{\sum_{u \geq 1} \sum_{i > j} m_i^{(u)} m_j^{(u)}}}{\prod_{j \geq 1} (s^2; q)_{|\mathbf{m}^{(j)}|}} \frac{1}{(2\pi\sqrt{-1})^m} \oint \dots \oint_{\text{around}\{x_j^{-1}\}} \prod_{1 \leq i < j \leq m} \frac{y_j - y_i}{y_j - qy_i} \\ &\times \prod_{k=1}^{\alpha} \left(\sum_{j=0}^{\mathbf{m}_k} \frac{(-1)^j q^{\binom{\mathbf{m}_k - j}{2}}}{(q; q)_j (q; q)_{\mathbf{m}_k - j}} \prod_{p > \mathbf{m}[1, k-1]}^{j + \mathbf{m}[1, k-1]} \prod_{a > \ell_{c_k-1}}^n \frac{1 - qx_a y_p}{1 - x_a y_p} \prod_{r > j + \mathbf{m}[1, k-1]}^{\mathbf{m}[1, k]} \prod_{b > \ell_{c_k}}^n \frac{1 - qx_b y_r}{1 - x_b y_r} \right) \\ &\times f_{\mu^{\geq 1}}^{\text{stoch}}(\lambda^{\geq 1}; y_1^{-1}, \dots, y_m^{-1}) \prod_{i=1}^m \frac{(y_i - s) dy_i}{y_i^2}, \end{aligned} \quad (6-2)$$

⁹Without loss of generality, for convenience of notation, we assume incoming paths in rows n and $n+1$ have different colours.

where (positively oriented) integration contours are chosen to encircle all points $\{x_j^{-1}\}_{j=1}^n$ and no other singularities of the integrand, or as q -nested closed simple curves with y_i -contour containing $q^{-1} \cdot (y_j\text{-contour})$ for all $i < j$, and all of the contours encircling $\{x_j^{-1}\}_{j=1}^n$. The contours can also be chosen to either encircle or not encircle the point 0.

The formula also holds in the column inhomogeneous setting under the assumption that $(s_x, \xi_x) = (s, 1)$ for any $x \geq 1$ such that $|\mathbf{m}^{(x)}| > 0$.

Proof. If the parameters satisfy the inequalities (4-1), then the statement is a substitution of Theorem 5.7 into Theorem 4.7. Observe that \mathcal{O}_μ is independent of the state of the model to the right of the maximal coordinate of μ . Thus, both sides of (6-2) are actually rational functions in x_i 's, and the extra assumption (4-1) can thus be removed by analytic continuation. \square

Remark 6.2. The factor $\prod_{j \geq 1} (s^2; q)_{|\mathbf{m}^{(j)}|}^{-1}$ in the right-hand side of (6-2) does not make the expression singular in the finite spin situation $s^2 = q^{-J}$, $J = 1, 2, \dots$, because the same factor appears in the numerator when one writes $f_{\mu \geq 1}^{\text{stoch}} = (-s)^{|\mu|} f_{\mu \geq 1}$ explicitly by taking the factorization (2-30), where this factor is manifest, and acting on it by difference operators (2-31)–(2-32) and colour merging (3-18) as needed.

6B. The colour-blind case. Let us see how Theorem 6.1 works in the colour-blind situation. The observables then simplify to (4-14), and we obtain:

Corollary 6.3. *In the colour-blind case $\lambda = (n)$, with ordered coordinates $\theta_1 \geq \dots \geq \theta_m \geq 1$ of $\mu^{\geq 1}$ and no column inhomogeneities, we have*

$$\begin{aligned} & \mathbb{E} \left[(1 - q^{H^v(\theta_1+1)})(1 - q^{H^v(\theta_2+1)-1}) \dots (1 - q^{H^v(\theta_m+1)-m+1}) \right] \\ &= \frac{(-1)^m (-s)^{|\theta|}}{(2\pi \sqrt{-1})^m} \oint \dots \oint_{\text{around}\{x_j^{-1}\}} \prod_{1 \leq i < j \leq m} \frac{y_i - y_j}{y_i - qy_j} \prod_{p=1}^m \left(\frac{1 - sy_p}{y_p - s} \right)^{\theta_i} \prod_{a=1}^n \frac{1 - qx_a y_p}{1 - x_a y_p} \frac{dy_p}{y_p}, \end{aligned} \quad (6-3)$$

where (positively oriented) integration contours are chosen to encircle all points $\{x_j^{-1}\}_{j=1}^n$ and no other singularities of the integrand, and $H^v(x) = \#\{i \in \{1, \dots, n\} : v_i \geq x\}$.

Remark 6.4. Equation (6-3) is readily seen to coincide with [Borodin and Petrov 2017, Lemma 9.10], which is essentially equivalent to the integral representation of the most general multipoint moments for the colourless stochastic vertex model along a single line.

It is not hard to extend this result, with a very similar proof to the one given below, to the column inhomogeneous case under the condition that $(s_x, \xi_x) = (s, 1)$ for any $x \geq 1$ such that no parts of μ are equal to x . However, [Borodin and Petrov 2018a, Lemma 9.11] gives such an extension to the fully column inhomogeneous situation. It remains unclear how to achieve this level of generality in the coloured case.

Remark 6.5. While the freedom in choosing the contours as q -nested is still there (it can be checked directly as in the proof of Lemma 5.9 or carried over from Theorem 6.1), the contours cannot include the point 0 anymore, and its inclusion (together with q -nestedness) would actually change the q -shifted moments in the left-hand side into unshifted moments of [Borodin and Petrov 2017, Theorem 9.8]; see [Borodin and Petrov 2017, Section 9] for a detailed explanation of that transition.

Proof of Corollary 6.3. In the colour-blind case $\lambda = (n)$, using (4-11), (2-29) and (4-14), we write (6-2) as

$$\begin{aligned} \mathbb{E} & \left[\frac{(1 - q^{H^v(\theta_1+1)})(1 - q^{H^v(\theta_2+1)-1}) \dots (1 - q^{H^v(\theta_m+1)-m+1})}{\prod_{j \geq 1} (q; q)_{\text{mult}_j(\theta)}} \right] \\ &= \frac{(-s)^{|\theta|}}{\prod_{j \geq 1} (s^2; q)_{\text{mult}_j(\theta)}} \frac{1}{(2\pi\sqrt{-1})^m} \oint \dots \oint_{\text{around}\{x_j^{-1}\}} \prod_{1 \leq i < j \leq m} \frac{y_j - y_i}{y_j - qy_i} \\ & \quad \times \sum_{j=0}^m \left(\frac{(-1)^j q^{\binom{m-j}{2}}}{(q; q)_j (q; q)_{m-j}} \prod_{p=1}^j \prod_{a=1}^n \frac{1 - qx_a y_p}{1 - x_a y_p} \right) F_{\theta}^c(y_1^{-1}, \dots, y_m^{-1}) \prod_{i=1}^m \frac{(y_i - s) dy_i}{y_i^2}, \quad (6-4) \end{aligned}$$

with contours around $\{x_j^{-1}\}$ and no other singularities. Observe that if the summation index j in the integrand above takes any value $j < m$, then the integrand, viewed as a function in y_m , has no singularities at $\{x_*^{-1}\}$, and the integral vanishes. Hence, we can set $j = m$.¹⁰

Further we write

$$\prod_{i < j} \frac{y_j - y_i}{y_j - qy_i} = \prod_{i \neq j} \frac{y_j - y_i}{y_j - qy_i} \cdot \prod_{i > j} \frac{y_j - qy_i}{y_j - y_i},$$

note that the first factor in the right-hand side is symmetric in y_i 's, and the same is true about all other parts of the integrand. Hence, we can sum over the second factor over all permutations of the y_i 's, which yields $(q; q)_m / (1 - q)^m$ in the integrand and multiplies the value of the integral by $m!$.

Finally, from [Borodin and Petrov 2017, Theorem 4.12] we read

$$F_{\theta}^c(y_1^{-1}, \dots, y_m^{-1}) = \frac{\prod_{j \geq 1} (s^2; q)_{\text{mult}_j(\theta)}}{\prod_{j \geq 1} (q; q)_{\text{mult}_j(\theta)}} \frac{(1 - q)^m}{\prod_{i=1}^m (1 - sy_i^{-1})} \sum_{\sigma \in \mathfrak{S}_m} \sigma \left(\prod_{i < j} \frac{y_i^{-1} - qy_j^{-1}}{y_i^{-1} - y_j^{-1}} \prod_{i=1}^m \left(\frac{y_i^{-1} - s}{1 - sy_i^{-1}} \right)^{\theta_i} \right),$$

where the sum is over all permutations σ in the symmetric group \mathfrak{S}_m on m symbols, and σ 's permute the variables y_1, \dots, y_m in the expression that they are applied to. Since the rest of the integrand is symmetric in the y_i 's, we can remove the sum over permutation leaving only the term with $\sigma = \text{id}$, and divide the integral by $m!$.

Implementing the above transformation and cancelling common factors yields (6-3). \square

6C. Duality. One convenient feature of the integral representation (6-2) for $\mathbb{E}\mathcal{O}_{\mu}$ is that the dependence on values of coordinates of μ is concentrated in the factor $f_{\mu \geq 1}(\lambda^{\geq 1}; y_1^{-1}, \dots, y_m^{-1})$ of the integrand. This allows us to easily derive certain difference equations that $\mathbb{E}\mathcal{O}_{\mu}$ must satisfy. Let us see how this works.

The skew-Cauchy identity (2-35) was stated for the rainbow case but is readily extended to the colour-merged case with the help of Proposition 3.8 (in fact, this colour merging argument was already used in the proof of Proposition 4.3 for G_* 's replaced by their limits \mathcal{G}_* 's). In particular, it implies

$$\sum_{\mu \geq 1 \text{ is } \lambda^{\geq 1} \text{ coloured}} f_{\mu \geq 1}(\lambda^{\geq 1}; y_1^{-1}, \dots, y_m^{-1}) G_{\mu \geq 1/x}((qX)^{-1}) = \prod_{i=1}^m \frac{1 - y_i X}{1 - qy_i X} \cdot f_x(\lambda^{\geq 1}; y_1^{-1}, \dots, y_m^{-1}),$$

¹⁰It is at this moment that we lose the freedom to have 0 inside the contours, as the terms that we remove may have nontrivial residues at 0; see Remark 6.5.

where \varkappa is an arbitrary $\lambda^{\geq 1}$ -coloured composition, X is a complex parameter. Multiplying both sides by $(-s)^{|\varkappa|} = (-s)^{|\varkappa|-|\mu|}(-s)^{|\mu|}$, we can also replace f by f^{stoch} and G by G^{stoch} in this identity, where $G_{\mu^{\geq 1}/\varkappa}^{\text{stoch}} = (-s)^{|\varkappa|-|\mu|}G_{\mu^{\geq 1}/\varkappa}$ is a stochastic kernel with matrix elements built from stochastic M^{stoch} -weights (2-12).

Observe that the prefactor is of the form of the exact inverse of the factors in the integrand of (6-2) that depend on x_i 's. Hence, if we compute the expectation $\mathbb{E}_{n+1}\mathcal{O}_\mu$ along the $(n+1)$ -st row of the stochastic vertex model, with left entering colour $(n+1)$ and rapidity X used in that row, and subsequently sum against the stochastic kernel $G_{\mu^{\geq 1}/\varkappa}^{\text{stoch}}((qX)^{-1})$, the result will coincide (subject to certain convergence conditions that we are ignoring here) with the expectation of $\mathbb{E}_n\mathcal{O}_\mu$ computed along the n -th row:

$$\sum_{\mu^{\geq 1}} \mathbb{E}_{n+1}\mathcal{O}_\mu \cdot G_{\mu^{\geq 1}/\varkappa}^{\text{stoch}}((qX)^{-1}) = \mathbb{E}_n\mathcal{O}_\mu. \tag{6-5}$$

As the transition from row n to row $n+1$ is also realized by a stochastic kernel, we observe a *duality* of the action of two stochastic operators on the *duality functional* \mathcal{O}_μ .

Of course, the above arguments only verify the duality relation (6-5) for a specific class of distributions on row n . However, this class is sufficiently general ($n \geq m$ and parameters x_1, \dots, x_n are arbitrary), and it is quite plausible that the duality relation will hold for generic distributions on coloured compositions $\mu^{\geq 1}$ on the n -th row, and also for the n -th row being the whole lattice \mathbb{Z} , rather than $\mathbb{Z}_{\geq 1}$ with a path entering from the left.

It would be very interesting to see an independent verification of (6-5), possibly by a reduction to duality functional constructed in [Kuan 2018]. Given that spectral decomposition of stochastic kernels G^{stoch} are known (see [Borodin and Wheeler 2018, Section 9.5]) this would likely lead to an alternative proof of Theorem 6.1, apart from other possible applications.

For a colour-blind version of the above discussion see [Borodin and Petrov 2017, Section 8.5] and [Borodin and Petrov 2018a, Section 8.5].

6D. The rainbow case. Let us now focus on the rainbow sector, with the labelling composition λ being $(1, 1, \dots, 1)$. The simplification of the observables \mathcal{O}_μ in this case was given in (4-13), and this leads us to:

Corollary 6.6. *In the notation of Section 6A, assume that $\lambda = (1, 1, \dots, 1)$. Then*

$$\begin{aligned} \mathbb{E} \prod_{\substack{i, j \geq 1 \\ m_i^{(j)} = 1}} \frac{q^{H_{\geq i}^{v/\mu}(j+1)} - q^{H_{> i}^{v/\mu}(j+1)}}{q-1} \\ = \frac{q^{\sum_{j \geq 1} \binom{m(j)}{2}}}{\prod_{j \geq 1} (s^2; q)_{|m(j)|}} \frac{(-1)^m}{(2\pi\sqrt{-1})^m} \oint \dots \oint_{\text{incl.}\{x_j^{-1}\}} \prod_{1 \leq i < j \leq m} \frac{y_j - y_i}{y_j - qy_i} \\ \times f_{\mu^{\geq 1}}^{\text{stoch}}(y_1^{-1}, \dots, y_m^{-1}) \prod_{j=1}^m \left(\frac{y_j - s}{y_j} \frac{x_{c_j}}{1 - x_{c_j}y_j} \prod_{i > c_j}^n \frac{1 - qx_iy_j}{1 - x_iy_j} dy_j \right), \tag{6-6} \end{aligned}$$

where (positively oriented) integration contours are chosen to encircle all points $\{x_j^{-1}\}_{j=1}^n$ and no

other singularities of the integrand, or as q -nested closed simple curves with y_i -contour containing $q^{-1} \cdot (y_j\text{-contour})$ for all $i < j$, and all of the contours encircling $\{x_j^{-1}\}_{j=1}^n$.

The formula also holds in the column inhomogeneous setting under the assumption that $(s_x, \xi_x) = (s, 1)$ for any $x \geq 1$ such that $|\mathbf{m}^{(x)}| > 0$.

Proof. This can be obtained by either a direct substitution $\lambda = (1, \dots, 1)$ into [Theorem 6.1](#), or a substitution of [Proposition 5.2](#) into the rainbow case of [Theorem 4.7](#). \square

Remark 6.7. Denote the parts of $\mu^{\geq 1}$ by $(\theta_1, \dots, \theta_m)$, where θ_1 carries the smallest colour c_1 of those represented in $\mu^{\geq 1}$, θ_2 carries the next smallest colour c_2 , etc. Then in the antidominant case, when $\theta_1 \leq \theta_2 \leq \dots \leq \theta_m$, $f_{\mu}^{\text{stoch}}(y_1^{-1}, \dots, y_m^{-1})$ in the integrand of (6-6) completely factorizes — and hence so does the whole integrand — as (in the column homogeneous case)

$$f_{\mu^{\geq 1}}^{\text{stoch}}(y_1^{-1}, \dots, y_m^{-1}) = \prod_{j \geq 1} (s^2; q)_{|\mathbf{m}^{(j)}|} \prod_{i=1}^n \frac{y_i}{y_i - s} \prod_{i=1}^n \left(\frac{1 - sy_i}{y_i - s} \right)^{\theta_i}; \quad (6-7)$$

see (2-30), (3-2), and [Remark 5.4](#). In this case there is also another path to (6-6). Namely, using the shift invariance property established in [[Borodin et al. 2019](#)], one can rewrite the left-hand side of (6-6) in terms of an average for a combination of q -moments of the height function for a colour-blind vertex on a quadrant taken at points along a down-right path in the quadrant. In their turn, such moments possess explicit integral representations, see [[Borodin et al. 2016a](#)] for details. See also [Remark 7.10](#) below for a related observation.

Remark 6.8. The identity (6-6) admits two colour-position symmetries, one for each side.

For the left-hand side, [[Borodin and Bufetov 2019](#), Theorem 7.3] shows that the joint distributions of the coloured height functions of the (rainbow) coloured stochastic vertex model along boundaries of a certain class of down-right domains are symmetric with respect to rotations of the domains by 180 degrees that also swap the roles of colour and position of the entering/exiting paths. In our case, the domain is the rectangle, and the application of this symmetry will swap colours and positions in the observables, as well as the roles of rows and columns. This will give integral formulas for averages of the new observables, as well as indicate that those observables are also likely to be duality functionals for the same reasons as those in [Section 6C](#).

For the right-hand side, if we represent f_{μ}^{stoch} as a result of a sequence of applications of the difference operators T_i given by (2-31)–(2-32) to a factorized expression of the type (6-7), then we can use self-adjointness of the T_i 's with respect to the integral scalar product with weight $\prod_{i < j} (y_j - y_i)/(y_j - qy_j)$ (see [[Borodin and Wheeler 2018](#), Proposition 8.1.3]) to move the application of the T_i 's to the fully factorized part of the integrand. Treating that part as an analogue of (2-30), or rather of (3-2), at the specific value of $s = q^{-1/2}$, we will see a new f -like function appearing in the integrand, that will utilize suitably permuted colours c_1, \dots, c_m for its coordinates, and horizontal rapidities x_1, \dots, x_n as its inhomogeneities. Thus, we will see a similar formula with positions and colours, as well as rows and columns, swapped.

It would be interesting to see if applying both symmetries, at least in the case $s = q^{-1/2}$, returns one to the original formula, but we will not pursue that here.

6E. Fusion. The goal of this section is to fuse rows of the stochastic vertex model from Section 6A and to see what Theorem 6.1 turns into in that situation.

First, one can fuse finitely many rows, which in terms of individual vertices corresponds to the outer sum in (2-10) (one can think of the inner sum in that relation as already performed, with our spin parameter s being $q^{-M/2}$). This starts by replacing a single row of colour c with rapidity x by $L \geq 1$ rows of the same colour c and rapidities forming a finite geometric progression $x, \dots, q^{L-1}x$. Since our left boundary condition in these L rows consists of all incoming edges occupied by paths of the same colour, no summation along that boundary is necessary. Hence, the L rows of vertices with L^{stoch} -weights can be collapsed into a single row with the weight of a vertex in that row and column j being

$$W_{L,M}(s_j^{-1}\xi_j x; q; *)|_{q^{-M}=s_j^2},$$

where (s_j, ξ_j) are the column inhomogeneities as in Section 3A, and the appearance of s_j^{-1} in front of $\xi_j x_j$ is due to the argument x/s in the expression (2-11) of L^{stoch} -weights in terms of W -weights.

The right-hand side of (6-2) also behaves well with respect to such fusion. More exactly, it leads to a simple replacement of all factors of the form $(1 - qxy_k)/(1 - xy_k)$, for various k and $x = x_*$ being the rapidity of the fused row, by

$$\frac{1 - qxy_k}{1 - xy_k} \cdot \frac{1 - q^2xy_k}{1 - qxy_k} \cdots \frac{1 - q^Lxy_k}{1 - q^{L-1}xy_k} = \frac{1 - q^Lxy_k}{1 - xy_k}. \tag{6-8}$$

Thus, the right-hand side can be immediately analytically continued in q^L . It is not clear, however, what that would mean on the side of the stochastic vertex model as the left boundary condition in the fused row consists of L paths.

To remedy this situation, we will perform a special limit transition with vertical inhomogeneity in column 1; this is parallel to what was done in [Borodin et al. 2019, Section 6]. More exactly, we will rely on the limiting relation (2-17). According to the left-hand side of that relation, we will take $\xi_1 = \zeta/s_1$ to turn the weight in the first column of the fused row into $W_{L,M}(s_1^{-2}\zeta x; q; *)|_{q^{-M}=s_1^2}$, and then take the limit $s_1 \rightarrow 0$. Here z is an additional parameter that remains finite in the limit $s_1 \rightarrow 0$; it regulates the strength of the left boundary. As we are about to make the rest of the model homogeneous, let us also set ζ to s , as this value will make the first column “blend in” with the other ones. Thus, our limit results, by virtue of (2-17), a random number of paths of colour c passing horizontally from column 1 to column 2 in the fused row, with the distribution of this random number given by

$$\text{Prob}\{k\} = \frac{(sq^Lx; q)_\infty}{(sx; q)_\infty} \frac{(q^{-L}; q)_k}{(q; q)_k} (sq^Lx)^k, \quad k \geq 0.$$

Note that for $L \in \mathbb{Z}_{\geq 1}$, this distribution is supported by $\{0, 1, \dots, L\}$, as it should be, as an L -fused row cannot carry more than L paths. But this distribution is also suitable for analytic continuation in q^L , which we will use momentarily.

Let us now discuss vertices in the fused row and other columns. Their weights $W_{L,M}(s_j^{-1}\xi_j x; q; *)$ are given by the right-hand side of (2-6), which is explicit but rather complicated. We will consider a simpler situation instead, governed by the q -Hahn specialization of Section 2B. Hence, we will specialize $(x, s_j, \xi_j) \mapsto (s, s, 1)$ to turn the weights into more tangible expressions as in the right-hand side of (2-16).

Finally, rather than performing fusion for a particular row of colour c as we have done above (replacing it by L columns first and then fusing them together), let us do it for every row of the stochastic vertex model of Section 6A.

Gathering all the pieces together, we obtain a stochastic vertex model in the quadrant $\mathbb{Z}_{\geq 2} \times \mathbb{Z}_{\geq 1}$, depending on three parameters q , $s(=q^{-M/2})$, and $z(=q^{-L/2})$ satisfying $|q|, |s|, |z|, |s/z| < 1$, defined as follows:

- Along the boundary of the quadrant, no paths enter the quadrant through its bottom boundary; and along the left boundary we have a random number of paths entering at every row, with the bottom λ_1 rows hosting entering paths of colour 1, next λ_2 rows hosting paths of colour 2, and so on; the sequence $\{\lambda_j\}_{j \geq 1}$ of nonnegative integers is given. As before, the partial sums of λ are denoted as $\sum_{i=1}^k \lambda_i = \ell_k$, $k \geq 1$; $\ell_0 := 0$. The distribution of the number of paths that enter in any row is given by

$$\text{Prob}\{k\} = \frac{(s^2/z^2; q)_{\infty}}{(s^2; q)_{\infty}} \frac{(z^2; q)_k}{(q; q)_k} \left(\frac{s^2}{z^2}\right)^k, \quad k \geq 0. \quad (6-9)$$

- The vertex weights for the model are given by

$$\text{weight}_{s,z} \left(\begin{array}{c} c \\ B \rightarrow A \rightarrow D \end{array} \right) = \left(\frac{s^2}{z^2}\right)^{|D|} \frac{(s^2/z^2; q)_{|A|-|D|} (z^2; q)_{|D|}}{(s^2; q)_{|A|}} q^{\sum_{i < j} D_i (A_j - D_j)} \prod_{i \geq 1} \binom{A_i}{A_i - D_i}_q. \quad (6-10)$$

This model is more general than the one in Section 6A in the sense that it can carry any number of paths along any edges, not just the vertical ones. However, it is less general in that there are no remaining row and column inhomogeneities.¹¹

As before, we focus on the paths that cross upwards from row n to row $n + 1$ for some $n \geq 1$.¹² Encoding colours and horizontal positions of these crossings by a coloured composition ν , we can define observables \mathcal{O}_{μ} on the set of possible ν 's by the same formula (6-1), where $\mu = 2^{m^{(2)}} 3^{m^{(3)}} \dots$ is an arbitrary coloured composition with no parts smaller than 2; this is what used to be $\mu^{\geq 1}$. Let us use the familiar notation $m = \ell(\mu)$ for the length of μ , $c_1 < \dots < c_{\alpha}$ for the colours represented in μ , and $m_1, \dots, m_{\alpha} \geq 1$ for the number of parts of μ of colours c_1, \dots, c_{α} , respectively. Finally, let $\text{colour}(\mu)$ be the composition that encodes the colouring of μ , and let $\mu - 1^m$ denote the composition obtained from μ by subtracting 1 from each part.

¹¹In fact, we could have left s and z parameters column and row dependent, respectively, but chose not to do so for the sake of simplicity. On the other hand, we could not have left row and column rapidities x_* and ξ_* generic and still had the same factorized form of the weights.

¹²Since rows above n do not matter to us, we assume, as in Section 6A, that colours of left-entering arrows in rows n and $n + 1$ are different.

Corollary 6.9. *With the above notation for the fused stochastic vertex model, we have*


$$\begin{aligned} \mathbb{E} \mathcal{O}_\mu &= \frac{q^{\sum_{u \geq 2} \sum_{i > j} m_i^{(u)} m_j^{(u)}} (-s)^m}{\prod_{j \geq 2} (s^2; q)_{|m^{(j)}|} (2\pi\sqrt{-1})^m} \\ &\quad \times \oint \cdots \oint \prod_{1 \leq i < j \leq m} \frac{y_j - y_i}{y_j - qy_i} \\ &\quad \times \prod_{k=1}^\alpha \left(\sum_{j=0}^{m_k} \frac{(-1)^j q^{\binom{m_k-j}{2}}}{(q; q)_j (q; q)_{m_k-j}} \prod_{p > m[1, k-1]}^{j+m[1, k-1]} \left(\frac{1-z^{-2}sy_p}{1-sy_p} \right)^{n-\ell_{c_k-1}} \prod_{r > j+m[1, k-1]}^{m[1, k]} \left(\frac{1-z^{-2}sy_r}{1-sy_r} \right)^{n-\ell_{c_k}} \right) \\ &\quad \times f_{\mu-1^m}^{\text{stoch}}(\text{colour}(\mu); y_1^{-1}, \dots, y_m^{-1}) \prod_{i=1}^m \frac{dy_i}{y_i^2}, \end{aligned} \tag{6-11}$$

where the integration contours are either positively oriented and q -nested around s^{-1} with y_i -contour containing $q^{-1} \cdot (y_j\text{-contour})$ for all $i < j$, or negatively oriented and q -nested around s , with y_j -contour containing $q \cdot (y_i\text{-contour})$ for all $i < j$. The point 0 can be either inside or outside the contours in either case.

Proof. The starting point is [Theorem 6.1](#) with μ from [\(6-11\)](#) being $\mu^{\geq 1}$ there.

The first step is to turn each row of the quadrant into L rows of the same colour, and fuse them as was described above. This does not affect the integral representation much, except for the change described around [\(6-8\)](#).

The next step is the limit transition in column 1 described above. To see how it affects the factor $f_{\mu^{\geq 1}}^{\text{stoch}}(y_1^{-1}, \dots, y_m^{-1})$, recall the partition function definition [\(2-23\)](#) (and its coloured modification [\(2-41\)](#)) of the f 's. Since all parts of $\mu^{\geq 1}$ are assumed to be at least 2, the first column contains only vertices of the form



that have L^{stoch} -weights

$$\frac{(-s_1)(\xi_1 y_j^{-1} - s_1)}{1 - s_1 \xi_1 y_j^{-1}} = \frac{(-s_1)(\xi_1 - s_1 y_j)}{y_j - s_1 \xi_1} = \frac{(-s_1)(s/s_1 - s_1 y_j)}{y_j - s} = \frac{s_1^2 y_j - s}{y_j - s}, \quad 1 \leq j \leq m,$$

where we used the value $\xi_1 = s/s_1$ as above. The limit $s_1 \rightarrow 0$ of this expression is $-s/(y_j - s)$, and the product over all $1 \leq j \leq m$ gives $(-s)^m \prod_{j=1}^m (y_j - s)^{-1}$. Adding that to the integrand of [\(6-2\)](#) yields the integrand of [\(6-11\)](#), together with the replacement $\mu^{\geq 1} \mapsto \mu - 1^m$ in the index of f^{stoch} , where the subtraction of 1^m is responsible for removing a column from the partition function representation of f^{stoch} that we just performed.

Let us now look at the contours. For the application of [Theorem 6.1](#) we could choose them to q -nest around $\{x_i, qx_i, \dots, q^{L-1}x_i\}_{i=1}^n = \{s, qs, \dots, q^{L-1}s\}$ and either contain 0 or not. (We could not choose them to encircle $\{x_i, qx_i, \dots, q^{L-1}x_i\}_{i=1}^n$ and no other singularities as this choice of horizontal rapidities

forces the inclusion of at least some singularities of $\prod_{i < j} (y_j - qy_i)^{-1}$ into the contours.) Since the integrand is manifestly nonsingular at $qs, \dots, q^{L-1}s$, we can remove the condition of encircling those points. As the integrand is readily seen to not have poles at $y_* = \infty$, we can also move the contours through ∞ and have them q -nest around the only other singularity, which is at s (again, 0 can be either inside or outside).

Thus, we have now proved (6-11) for $L = 1, 2, \dots$, and the final step consists in analytic continuation in $q^L = z^{-2}$ from the set of points $\{q, q^2, q^3, \dots\}$ accumulating at 0. This, however, is straightforward: The only dependence on z of the right-hand side is through factors $(1 - z^{-2}y_*)$, and the left-hand side is readily seen to be given by uniformly convergent series with rational terms at least as long as $|q|, |s|, |sz^{-1}| < \text{const} < 1$. \square

7. Limit to polymers

The goal of this section is to explore the consequences of Corollary 6.9 for a few models of directed polymers in (1+1) dimensions. The exposition of the limit transitions from fused coloured stochastic models to directed polymers follows [Borodin et al. 2019, Section 7].

7A. Continuum stochastic vertex model. Let us start by introducing a vertex model that will serve as a limiting object for the fused vertex model of Section 6E described by weights (6-9)–(6-10).

As in the fused case, the vertices of the continuum model will be parametrized by points of a quadrant, and to keep the notation parallel to that of Section 6E, we will use the quadrant $\mathbb{Z}_{\geq 2} \times \mathbb{Z}_{\geq 1}$.

Each vertex will have a certain *mass* of each colour ≥ 1 entering from the bottom and from the left, and exiting through the top and to the right. The mass is a real number in $[0, \infty)$, and for each vertex the total number of colours that have nonzero mass entering the vertex will always be finite. The mass of each colour passing through a vertex will always be preserved – the sum of incoming mass from the bottom and from the left must be equal to the sum of exiting mass to the right and through the top.

Let us denote the masses of colours $1, 2, \dots$ entering through the bottom of a vertex by $\boldsymbol{\alpha} = (\alpha_1, \alpha_2, \dots)$, entering from the left by $\boldsymbol{\beta} = (\beta_1, \beta_2, \dots)$, exiting through the top by $\boldsymbol{\gamma} = (\gamma_1, \gamma_2, \dots)$, and exiting to the right by $\boldsymbol{\delta} = (\delta_1, \delta_2, \dots)$, respectively. The mass preservation means

$$|\boldsymbol{\alpha}| + |\boldsymbol{\beta}| = |\boldsymbol{\gamma}| + |\boldsymbol{\delta}|.$$

The notation is chosen to be in parallel with $(\mathbf{A}, \mathbf{B}; \mathbf{C}, \mathbf{D})$ notation for the vertex models, as in (2-1).

Recall that a random variable with values in $(0, 1)$ is said to be *Beta-distributed* with parameters $a, b > 0$ if it has a density, with respect to the Lebesgue measure, given by

$$\frac{\Gamma(a+b)}{\Gamma(a)\Gamma(b)} x^{a-1}(1-x)^{b-1}, \quad 0 < x < 1. \quad (7-1)$$

Given the coloured masses $\boldsymbol{\alpha}, \boldsymbol{\beta}$ entering a vertex of our continuum vertex model, the coloured masses $\boldsymbol{\gamma}, \boldsymbol{\delta}$ exiting the vertex are random and determined as follows. The procedure has two parameters $a, b > 0$, as in the Beta-distribution (7-1). If all coordinates $\boldsymbol{\alpha}$ are zero, i.e., $\boldsymbol{\alpha} = \mathbf{0}$, then we set $\boldsymbol{\gamma} = \boldsymbol{\beta}$ and $\boldsymbol{\delta} = \mathbf{0}$. If $\boldsymbol{\alpha} \neq \mathbf{0}$, let $n \geq 1$ be the maximal natural number such that $\alpha_n \neq 0$, and let ζ be an (a, b) -Beta

distributed random variable. Then we set $\delta_{n+1}, \delta_{n+2}, \dots$ to 0 and define $\delta_n, \delta_{n-1}, \dots, \delta_1$ recursively via

$$\begin{aligned} \delta_n &= -\log(e^{-\alpha_n} + \zeta(1 - e^{-\alpha_n})), \\ \delta_{n-1} + \delta_n &= -\log(e^{-\alpha_{n-1}-\alpha_n} + \zeta(1 - e^{-\alpha_{n-1}-\alpha_n})), \\ &\vdots \\ \delta_1 + \dots + \delta_{n-1} + \delta_n &= -\log(e^{-\alpha_1-\dots-\alpha_{n-1}-\alpha_n} + \zeta(1 - e^{-\alpha_1-\dots-\alpha_{n-1}-\alpha_n})). \end{aligned} \tag{7-2}$$

One can show that this implies $0 < \delta_j < \alpha_j$ for $1 \leq j \leq n$. Finally, we set $\boldsymbol{\gamma} = \boldsymbol{\beta} + (\boldsymbol{\alpha} - \boldsymbol{\delta})$, thus enforcing mass conservation.

In addition to defining what happens at the vertices of the quadrant, we need to specify boundary conditions. As before, we will assume that no mass enters the quadrant from the bottom, i.e., $\boldsymbol{\alpha} = \mathbf{0}$ for all vertices in the bottom row. On the other hand, along the left boundary we will assume that for the left-most vertex in row i , the left-entering coloured mass $\boldsymbol{\beta}$ has all but one coordinates equal to 0, with the exception of the i -th one, which is (a, b) -Beta distributed.

As usual, we think of the randomness as having no space dependency, which means that the Beta-distributed random variables at different vertices, as well as those used to define the left boundary condition, are independent.

The following statement was proved in [Borodin et al. 2019, Corollary 6.22].

Proposition 7.1. *Consider the fused coloured vertex model defined around (6-9)-(6-10) and set*

$$q = \exp(-\epsilon), \quad s^2 = q^\sigma, \quad z^2 = q^\rho, \tag{7-3}$$

for some $\sigma > \rho > 0$ and $\epsilon > 0$. Then as $\epsilon \rightarrow 0$, the fused coloured vertex models scaled by ϵ converges to the continuum vertex model defined above with parameters $(a, b) = (\sigma - \rho, \rho)$, in the sense that any finite collection of numbers of paths of arbitrary fixed colours entering/exiting any fixed set of vertices in fixed directions, when multiplied by ϵ , weakly converges to the collection of corresponding colour masses entering/exiting the corresponding vertices of the continuum model.

This immediately implies the convergence of the averages of the observables from Corollary 6.9 as well, but we will postpone the limiting statement until we reformulate the continuum vertex model as a directed random polymer in the next section.

7B. Random Beta-polymer. The Beta-polymer was first introduced in [Barraquand and Corwin 2017]. In order to define it, let $\{\eta_{t,m}\}_{t,m \geq 1}$ be a family of independent identically Beta-distributed random variables with parameters $a, b > 0$; see (7-1). The partition function $\mathfrak{Z}_{t,m}$ of the Beta-polymer, with $(t, m) \in \mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 1}$ and $t \geq m - 1$, is determined by the recurrence relation

$$\mathfrak{Z}_{t,m} = \eta_{t,m} \mathfrak{Z}_{t-1,m} + (1 - \eta_{t,m}) \mathfrak{Z}_{t-1,m-1}$$

and boundary conditions

$$\mathfrak{Z}_{t,t+1} \equiv 1, \quad \mathfrak{Z}_{t,1} = \eta_{1,1} \eta_{2,1} \cdots \eta_{t,1}.$$

Pictorially, $\mathfrak{Z}_{t,m}$ is a sum over all directed lattice paths with $(0, 1)$ and $(1, 1)$ steps that join $(0, 1)$ and (t, m) ,

of products of edge weights that all have the form η_* or $1 - \eta_*$; see [Barraquand and Corwin 2017; Borodin et al. 2019].

We also define *delayed* partition functions $\mathfrak{Z}_{t,m}^{(k)}$, $k \geq 1$, $t \geq m + k - 1$, by the same recurrence relation

$$\mathfrak{Z}_{t,m}^{(k)} = \eta_{t,m} \mathfrak{Z}_{t-1,m}^{(k)} + (1 - \eta_{t,m}) \mathfrak{Z}_{t-1,m-1}^{(k)},$$

where we are using the same family of random variable $\{\eta_{t,m}\}_{t,m \geq 1}$ to evaluate the coefficients, and shifted boundary conditions

$$\mathfrak{Z}_{t,t-k+2}^{(k)} \equiv 1, \quad \mathfrak{Z}_{t,1}^{(k)} = \eta_{k,1} \eta_{k+1,1} \cdots \eta_{t,1}.$$

Clearly, $\mathfrak{Z}_{t,m}^{(1)} \equiv \mathfrak{Z}_{t,m}$, and for any $k \geq 1$, $\mathfrak{Z}_{t,m}^{(k)}$ can be interpreted graphically in a similar way to $\mathfrak{Z}_{t,m}$, but with paths joining $(k-1, 1)$ and (t, m) .

As was shown in [Borodin et al. 2019, Section 7.1], there is a way to identify the continuum vertex model of Section 7A and the family of Beta-polymer partition functions $\{\mathfrak{Z}_{t,m}^{(k)}\}$. In order to see the equivalence, let us introduce the coloured height functions $\{h^{(\geq k)}(x, y) \mid x \geq 2; y, k \geq 1\}$ that count the total mass of colours $\geq k$ that exit vertices $(x, 1), (x, 2), \dots, (x, y)$, either upward or rightward, in the continuum model. Then one has the identification

$$-\log \mathfrak{Z}_{t,m}^{(k)} = h^{(\geq k)}(m+1, t), \quad t, m, k \geq 1, \quad t \geq m + k - 1. \quad (7-4)$$

For $t, m, k \geq 1$ that do not satisfy the inequality $t \geq m + k - 1$, the right-hand side of (7-4) is readily seen to vanish, and we set $\mathfrak{Z}_{t,m}^{(k)}$ to 1 for these values as well; then (7-4) holds for any $t, m, k \geq 1$.

Together with Proposition 7.1, this allows us to obtain a limiting version of Corollary 6.9 for the Beta-polymer. But in order to take the corresponding limit of the integral representation, we need to introduce limiting versions of the rational functions f_μ .

Lemma 7.2. *Take $q = \exp(-\epsilon)$, and $s^2 = q^\sigma$ (as in (7-3)), and fix a λ -coloured composition $\mu = 0^{m^{(0)}} 1^{m^{(1)}} 2^{m^{(2)}} \cdots$ of length $m \geq 1$, where λ is a composition of weight $|\lambda| = m$. Then there exists a limit*

$$\mathfrak{f}_\mu(\lambda; u_1, \dots, u_m) = \lim_{\epsilon \rightarrow 0} \epsilon^m \cdot \frac{f_\mu(\lambda; 1 + \epsilon u_1, \dots, 1 + \epsilon u_m)}{\prod_{j \geq 0} (s^{2j}; q)_{|m^{(j)}|}}, \quad (7-5)$$

where the convergence is uniform for complex u_1, \dots, u_m varying in compact sets that do not include $\sigma/2$, and the rational function $\mathfrak{f}_\mu(\lambda; u_1, \dots, u_m)$ can be characterized as follows.

- (i) For a rainbow μ , i.e., for $\lambda = (1, \dots, 1)$, in the antidominant sector $\mu_1 \leq \mu_2 \leq \dots \leq \mu_m$ one has (omitting λ from the notation)

$$\mathfrak{f}_\mu(u_1, \dots, u_m) = \prod_{i=1}^m \frac{1}{\sigma/2 - u_i} \left(\frac{\sigma/2 + u_i}{\sigma/2 - u_i} \right)^{\mu_i}, \quad (7-6)$$

and in the case $\mu_i < \mu_{i+1}$ for some $1 \leq i \leq m-1$, one has

$$\begin{aligned} \mathfrak{T}_i \cdot \mathfrak{f}_\mu(u_1, \dots, u_m) &= \mathfrak{f}_{(\mu_1, \dots, \mu_{i+1}, \mu_i, \dots, \mu_m)}(u_1, \dots, u_m), \\ \mathfrak{T}_i &\equiv 1 - \frac{u_i - u_{i+1} + 1}{u_i - u_{i+1}} (1 - \mathfrak{s}_i), \quad 1 \leq i \leq m-1, \end{aligned} \quad (7-7)$$

with elementary transpositions $\mathfrak{s}_i \cdot h(u_1, \dots, u_m) := h(u_1, \dots, u_{i+1}, u_i, \dots, u_m)$; see (2-31)–(2-32).

(ii) For a general labelling composition λ , let $\theta : \{1, \dots, m\} \rightarrow \{1, \dots, n\}$ be a monotone map (as in Section 3C) with $\theta_*((1, \dots, 1)) = \lambda$. Then

$$\sum_{\text{rainbow } \nu : \theta_*(\nu) = \mu} \mathfrak{f}_\nu(u_1, \dots, u_m) = \mathfrak{f}_\mu(\lambda; u_1, \dots, u_m). \tag{7-8}$$

Proof. This limit is an immediate consequence of the recursive definition (2-30)–(2-32) of Section 2E and (3-18) of Proposition 3.8. □

We are now ready to take a $q \rightarrow 1$ limit in Corollary 6.9.

Proposition 7.3. Fix $\sigma > \rho > 0$, and consider the partition functions $\mathfrak{Z}_{t,m}^{(k)}$ of the Beta-polymer as defined above with the parameters of (7-1) given by $(a, b) = (\sigma - \rho, \rho)$. Let $\mu = 2^{m^{(2)}} 3^{m^{(3)}} \dots$ be a coloured composition with no parts smaller than 2, $m = \ell(\mu)$ be the length of μ , $c_1 < \dots < c_\alpha$ be the colours represented in μ , and $m_1, \dots, m_\alpha \geq 1$ be the number of parts of μ of colours c_1, \dots, c_α , respectively. Also, let $\text{colour}(\mu)$ be the composition that encodes the colouring of μ . Then for any $t \geq \max\{i : m_i^{(x)} > 0\}$ we have

$$\begin{aligned} \mathbb{E} \prod_{x \geq 1} \prod_{i \geq 1} \frac{(\mathfrak{Z}_{t,x}^{(i+1)} - \mathfrak{Z}_{t,x}^{(i)}) m_i^{(x)}}{m_i^{(x)}!} \\ = \frac{(-1)^{|\mu|}}{(2\pi\sqrt{-1})^m} \oint \dots \oint \prod_{1 \leq i < j \leq m} \frac{u_j - u_i}{u_j - u_i + 1} \\ \times \prod_{k=1}^\alpha \left(\sum_{j=0}^{m_k} \frac{(-1)^j}{j!(m_k - j)!} \prod_{p > m[1, k-1]}^{j+m[1, k-1]} \left(\frac{\sigma/2 - u_p - \rho}{\sigma/2 - u_p} \right)^{t-c_k+1} \prod_{r > j+m[1, k-1]}^{m[1, k]} \left(\frac{\sigma/2 - u_r - \rho}{\sigma/2 - u_r} \right)^{t-c_k} \right) \\ \times \mathfrak{f}_{\mu-1^m}(\text{colour}(\mu); -u_1, \dots, -u_m) \prod_{i=1}^m du_i, \end{aligned} \tag{7-9}$$

with the integration contours either positively oriented and nested around $\sigma/2$ with u_i -contour containing $(u_j\text{-contour}) + 1$ for all $i < j$, or negatively oriented and nested around $-\sigma/2$, with u_j -contour containing $(u_i\text{-contour}) - 1$ for all $i < j$.

Proof. We start with (6-11) and $(\lambda_1, \lambda_2, \dots) = (1, 1, \dots)$; equivalently, $\ell_j = j$. Let us make the substitution (7-3) and look at the asymptotics of both sides.

On the left-hand side we have averages of the observables \mathcal{O}_μ given by (6-1). The denominators are deterministic and asymptotically give $(q; q)_{m_i^{(x)}}^{-1} \sim \epsilon^{-m_i^{(x)}} m_i^{(x)}!$ as $\epsilon \rightarrow 0$. For the numerators, according to Proposition 7.1, we obtain, along row $t = n$, $\epsilon^{-1} H_{\geq i}^v(x+1) \rightarrow h^{(\geq i)}(x+1, t)$, and, changing ν to ν/μ with ϵ -independent μ , $\epsilon^{-1} H_{\geq k}^{\nu/\mu}(x+1) \rightarrow h^{(\geq k)}(x+1, t)$ weakly as $\epsilon \rightarrow 0$, where h denotes the coloured height functions of the Beta-polymer. Using (7-4) and the fact that all the observables are bounded, we see that $\mathbb{E}\mathcal{O}_\mu$ is asymptotically equivalent to ϵ^{-m} times the left-hand side of (7-9).

For the right-hand side of (6-11), we change the variables $y_i = 1 + \epsilon u_i$, use Lemma 7.2, and also $(q; q)_j^{-1} (q; q)_{m_k-j}^{-1} \sim \epsilon^{-m_k} j!^{-1} (m_k - j)!^{-1}$.

The powers of ϵ from the q -symbols on both sides cancel out, the powers of ϵ from the changes of variables and $f \rightarrow \mathfrak{f}$ limit also cancel out, and the prefactor $(-s)^m$ in the right-hand side of (6-11) together with $(-s)^{|\mu|-m}$ required to convert $f_{\mu-1^m}^{\text{stoch}}$ to $f_{\mu-1^m}$ give $(-1)^{|\mu|}$ in the limit. This concludes the proof. \square

7C. Strict-weak polymer. The strict-weak or gamma polymer was first introduced in [Corwin et al. 2015] and [O’Connell and Ortmann 2015]. Its partition functions are determined by a very similar recurrence as those for the Beta-polymer. Namely, let us define $Z_{t,m}^{(k)}$ for $(t, m, k) \in \mathbb{Z}_{\geq 0} \times \mathbb{Z}_{\geq 1} \times \mathbb{Z}_{\geq 1}$, $t \geq m + k - 1$, by

$$Z_{t,m}^{(k)} = \eta_{t,m} Z_{t-1,m}^{(k)} + Z_{t-1,m-1}^{(k)}$$

with boundary conditions

$$Z_{t,t-k+2}^{(k)} \equiv 1, \quad Z_{t,1}^{(k)} = \eta_{k,1} \eta_{k+1,1} \cdots \eta_{t,1},$$

where $\{\eta_{t,m}\}_{t,m \geq 1}$ is a family of independent identically distributed random variables with a Gamma distribution that has density

$$\frac{1}{\Gamma(\kappa)} x^{\kappa-1} \exp(-x), \quad x > 0,$$

with respect to the Lebesgue measure. Here $\kappa > 0$ is a parameter.

The strict-weak polymer is a limiting instance of the Beta-polymer of Section 7B, because a Beta-distributed random variable with density (7-1) and parameters $(a, b) = (\kappa, \epsilon^{-1})$, when multiplied by ϵ^{-1} , converges to a Gamma-distributed random variable with parameter κ as $\epsilon \rightarrow 0$, both in distribution and with all moments.

In order to argue the convergence of the partition functions and their moments, we will appeal to the following:

Lemma 7.4. *Let $\{X_n\}_{n \geq 1}$ and $\{Y_n\}_{n \geq 1}$ be two sequences of nonnegative random variables such that $\{(X_n, Y_n)\}_{n \geq 1}$ weakly converges to a two-dimensional random variable (X, Y) with finite moments and a continuous joint distribution function.¹³ Furthermore, assume that coordinate moments converge:*

$$\lim_{n \rightarrow \infty} \mathbb{E} X_n^k = \mathbb{E} X^k, \quad \lim_{n \rightarrow \infty} \mathbb{E} Y_n^k = \mathbb{E} Y^k, \quad k \geq 1.$$

Then the joint moments also converge:

$$\lim_{n \rightarrow \infty} \mathbb{E}(X_n^k Y_n^l) = \mathbb{E}(X^k Y^l), \quad k, l \geq 1.$$

Proof. Fix $C > 0$ and write

$$\mathbb{E}(X_n^k Y_n^l) = \mathbb{E}(X_n^k Y_n^l \cdot \mathbf{1}\{X_n < C, Y_n < C\}) + \mathbb{E}(X_n^k Y_n^l \cdot \mathbf{1}\{X_n \geq C \text{ or } Y_n \geq C\}),$$

where we use $\mathbf{1}\{A\}$ to denote the indicator function of an event A . In the first term, we have a bounded functional under the expectation, which converges to $\mathbb{E}(X^k Y^l \cdot \mathbf{1}\{X < C, Y < C\})$ by the distributional convergence of (X_n, Y_n) . (If the distribution of (X, Y) were not continuous, we would have needed to choose C as its continuity point.)

¹³This requirement of continuity can be easily removed by an extra step in the proof.

Further, let us show that the second term converges to 0 as $C \rightarrow \infty$ uniformly in n . We have

$$\begin{aligned} \mathbb{E}(X_n^k Y_n^l \cdot \mathbf{1}\{X_n \geq C \text{ or } Y_n \geq C\}) &\leq \mathbb{E}(X_n^k Y_n^l \cdot (X_n/C + Y_n/C)) = C^{-1} \mathbb{E}(X_n^k Y_n^l \cdot (X_n + Y_n)) \\ &\leq 2C^{-1} (\mathbb{E}X_n^{k+l+1} + \mathbb{E}Y_n^{k+l+1}), \end{aligned}$$

where we used the inequality $x^a y^b \leq x^{a+b} + y^{a+b}$ that holds for $x, y, a, b > 0$. Since the moments of X_n and Y_n are bounded (because they converge by the hypothesis), the final expression tends to 0 as $C \rightarrow \infty$ uniformly in n .

As $\lim_{C \rightarrow \infty} \mathbb{E}(X^k Y^l \cdot \mathbf{1}\{X < C, Y < C\}) = \mathbb{E}(X^k Y^l)$, the proof is complete.¹⁴ □

Lemma 7.4 implies, in particular, that by choosing $(\sigma - \rho, \rho) = (\kappa, \epsilon^{-1})$, we can ensure the convergence

$$\lim_{\epsilon \rightarrow 0} \epsilon^{m+k-t-2} \cdot \mathfrak{Z}_{t,m}^{(k)} = Z_{t,m}^{(k)}, \quad t \geq m+k-2,$$

together with all the joint moments. This will allow us to take such a limit in **Proposition 7.3** momentarily, after the following analogue of **Lemma 7.2**.

Lemma 7.5. *Fix a λ -coloured composition $\mu = 0^{m^{(0)}} 1^{m^{(1)}} 2^{m^{(2)}} \dots$ of length $m \geq 1$, where λ is a composition of weight $|\lambda| = m$. Then there exists a limit*

$$\mathfrak{p}_\mu(\lambda; v_1, \dots, v_m) = \lim_{\sigma \rightarrow \infty} (-1)^{|\mu|+m} \sigma^{-|\mu|} \cdot \mathfrak{f}_\mu(\lambda; \sigma/2 + v_1, \dots, \sigma/2 + v_m) \tag{7-10}$$

where the convergence is uniform for complex v_1, \dots, v_m varying in compact sets that do not include 0, and the function $\mathfrak{p}_\mu(\lambda; u_1, \dots, u_m)$ is a polynomial in $v_1^{-1}, \dots, v_m^{-1}$ that can be characterized as follows.

(i) *For a rainbow μ , in the antidominant sector $\mu_1 \leq \mu_2 \leq \dots \leq \mu_m$, one has (omitting λ from the notation)*

$$\mathfrak{p}_\mu(v_1, \dots, v_m) = \prod_{i=1}^m v_i^{-\mu_i-1}, \tag{7-11}$$

and in the case $\mu_i < \mu_{i+1}$ for some $1 \leq i \leq m-1$ one uses the exchange relations (7-7) with (\mathfrak{p}, v_*) instead of (\mathfrak{f}, u_*) .

(ii) *For a general labelling composition λ , let $\theta : \{1, \dots, m\} \rightarrow \{1, \dots, n\}$ be a monotone map with $\theta_*((1, \dots, 1)) = \lambda$. Then*

$$\sum_{\text{rainbow } \nu : \theta_*(\nu) = \mu} \mathfrak{p}_\nu(u_1, \dots, u_m) = \mathfrak{p}_\mu(\lambda; u_1, \dots, u_m). \tag{7-12}$$

The proof of this lemma is straightforward.

Proposition 7.6. *Fix $\kappa > 0$, and consider the partition functions $Z_{t,m}^{(k)}$ of the strict-weak polymer as defined above with κ being the parameter of the Gamma distribution. Let μ and the associated notation*

¹⁴We are very grateful to Vadim Gorin for providing this argument.

be as in [Proposition 7.3](#). Then for any $t \geq m + \max\{i : m_i^{(x)} > 0\} - 1$, we have

$$\begin{aligned} \mathbb{E} \prod_{x \geq 1} \prod_{i=1}^n \frac{(Z_{t,x}^{(i+1)})^{m_i^{(x)}}}{m_i^{(x)}!} &= \frac{1}{(2\pi\sqrt{-1})^m} \oint \cdots \oint \prod_{1 \leq i < j \leq m} \frac{v_j - v_i}{v_j - v_i - 1} \\ &\quad \times \prod_{k=1}^{\alpha} \prod_{r > m[1,k-1]}^{m[1,k]} \frac{(\kappa + v_r)^{t-c_k}}{m_k!} \cdot \mathbf{p}_{\mu-1^m}(\text{colour}(\mu); v_1, \dots, v_m) \prod_{i=1}^m dv_i, \end{aligned} \quad (7-13)$$

where the integration contours are positively oriented and nested around 0, with v_j -contour containing $(v_i\text{-contour}) + 1$ for all $i < j$.

Proof. We start with (7-9), set $\sigma - \rho$ to κ , change the integration variables via $u_i = -v_i - \sigma/2$, and take the limit. Since $\mathfrak{Z}_{t,x}^{(i)} \sim \sigma^{x+i-t-2} Z_{t,x}^{(i)}$ and $\mathfrak{Z}_{t,x}^{(i+1)} \sim \sigma^{x+i-t-1} Z_{t,x}^{(i)}$, the second term dominates, and employing [Lemma 7.4](#), we obtain the convergence of the left-hand side of (7-9) to that of (7-13), with an additional power of σ that has exponent

$$\sum_{i,x \geq 1} m_i^{(x)} (x + i - t - 1) = |\mu| + \sum_{k=1}^{\alpha} c_k m_k - (t + 1)m.$$

On the other hand, in the integrand we have factors of the form $\sigma/2 - u_* = \sigma + v_* \sim \sigma$ in the denominator, that make the terms with $j = 0$ dominate and produce the power of σ with the exponent

$$\sum_{k=1}^{\alpha} m_k (c_k - t) = \sum_{k=1}^{\alpha} c_k m_k - tm.$$

Finally, the limit relation (7-10) yields the power of σ with the exponent $|\mu| - m$, thus matching the powers of σ on both sides. All the signs cancel out, where we note that we used the second choice of the contours in [Proposition 7.3](#) and changed the negative orientation to the positive one for (7-13). \square

7D. O’Connell–Yor semidiscrete Brownian polymer. This model was first introduced in [[O’Connell and Yor 2001](#)]. It is defined using a family $\{B_n(t)\}_{n \geq 1, t \geq 0}$ of independent standard Brownian motions. For each $n \geq 1$ and $t \geq s \geq 0$ we define its point-to-point partition function (with one of the points situated on level 1) as

$$Z_{(1,s) \rightarrow (n,t)}^{OY} = \int_{s=\tau_0 < \tau_2 < \cdots < \tau_n=t} \exp\left(\sum_{i=1}^n (B_i(\tau_i) - B_i(\tau_{i-1}))\right) d\tau_1 \cdots d\tau_{n-1}. \quad (7-14)$$

The classical functional central limit theorem and the fact that a Gamma-distributed random variable with large parameter L divided by L is approximately equal to 1 plus a standard normal variable divided by \sqrt{L} , yield the convergence (see [[Borodin et al. 2019](#), Section 7.3])

$$\lim_{L \rightarrow \infty} \frac{Z_{L,t,n}^{(Ls+1)}(\kappa = L)}{L^{L(t-s)}} = \exp\left(\frac{s-t}{2}\right) \cdot Z_{(1,s) \rightarrow (n,t)}^{OY}, \quad t \geq s \geq 0, \quad n \geq 1, \quad (7-15)$$

where on the left we take the strict-weak polymer partition functions from the previous section with the parameter $\kappa = L$, and the convergence is in finite-dimensional distributions.

Proposition 7.7. *Let μ and the associate notation be as in Proposition 7.3, and fix $0 \leq s_1 < \dots < s_\alpha$. Then for any $t \geq s_\alpha$ we have*

$$\begin{aligned} \mathbb{E} \prod_{x \geq 1} \prod_{i \geq 1} \frac{(\exp(\frac{s_i - t}{2}) \cdot Z_{(1, s_i) \rightarrow (x, t)}^{OY})^{m_i^{(x)}}}{m_i^{(x)}!} \\ = \frac{1}{(2\pi\sqrt{-1})^m} \oint \cdots \oint \prod_{1 \leq i < j \leq m} \frac{v_j - v_i}{v_j - v_i - 1} \\ \times \prod_{k=1}^{\alpha} \frac{\exp((t - s_k) \cdot \sum_{r > m[1, k-1]}^{m[1, k]} v_r)}{m_k!} \cdot \mathfrak{p}_{\mu-1^m}(\text{colour}(\mu); v_1, \dots, v_m) \prod_{i=1}^m dv_i, \end{aligned} \quad (7-16)$$

where the integration contours are positively oriented and nested around 0, with v_j -contour containing $(v_i\text{-contour}) + 1$ for all $i < j$.

Proof. Let us take the limit $L \rightarrow \infty$ of (7-13) with $\kappa = L$ and the coloured composition μ replaced by $\mu^{(L)}$ that has exactly the same parts, but the colours of those parts are $[s_1 L], \dots, [s_\alpha L]$ instead of c_1, \dots, c_α , respectively.

Let us look at the left-hand side first. The weak joint convergence of the random variables

$$\lim_{L \rightarrow \infty} \frac{Z_{Lt, x}^{([s_i L] + 1)}(\kappa = L)}{L^{tL - [s_i L]}} = \exp(\frac{s_i - t}{2}) \cdot Z_{(1, s_i) \rightarrow (n, t)}^{OY}$$

for various i and x follows from the central limit theorem, as was mentioned above, and the convergence of moments of these random variables follows from the corresponding convergence of their integral representations; see [Corwin et al. 2015, Theorem 5.3] for moments of the left-hand side, and [Borodin and Corwin 2014, Proposition 5.2.8] for the moments of the right-hand side.¹⁵ Lemma 7.4 then shows that the left-hand side of (7-13) divided by the power of L with exponent

$$\sum_{i, x \geq 1} m_i^{(x)}(tL - [s_i L]) = \sum_{k=1}^{\alpha} m_k(tL - [s_k L])$$

converges to that of (7-16).

On the other hand, for the right-hand side the convergence of the L -dependent factors in the integrand is elementary:

$$\lim_{L \rightarrow \infty} \frac{(L + v_r)^{tL - [s_k L]}}{L^{tL - [s_k L]}} = \exp((t - s_k)v_r),$$

uniformly for bounded v_r 's, which leads to the right-hand side of (7-16). □

7E. Continuum Brownian polymer. One way to define partition functions of the continuum Brownian polymer in (1+1)-dimensions is through solving the stochastic heat equation with multiplicative

¹⁵That convergence is, in fact, a special case of the one we are about to observe for single-coloured μ .

two-dimensional white noise. More exactly, let $\tilde{\mathcal{Z}}^{(y)}(t, x)$ be the unique solution of the following stochastic partial differential equation with the initial condition

$$\mathcal{Z}_t^{(y)} = \frac{1}{2} \mathcal{Z}_{xx}^{(y)} + \eta(t, x) \mathcal{Z}^{(y)}, \quad t > 0, \quad x \in \mathbb{R}; \quad \mathcal{Z}^{(y)}(0, x) = \delta(x - y),$$

where $\eta = \eta(t, x)$ is the two-dimensional white noise. We refer to the survey [Quastel 2012] and references therein for an extensive literature on this equation and its close relation to continuum Brownian path integrals and the Kardar–Parisi–Zhang equation.

The solutions $\mathcal{Z}^{(y)}(t, x)$ arise naturally as limits of the partition functions of the semidiscrete Brownian polymer from the previous section:

$$\lim_{L \rightarrow \infty} \frac{\exp\left(\frac{y-t\sqrt{L}}{2}\right) \cdot Z_{(1,y) \rightarrow (tL-x\sqrt{L}, t\sqrt{L})}^{OY}}{\exp(tL) \cdot L^{(x\sqrt{L}-tL)/2}} = \mathcal{Z}^{(y)}(t, x). \quad (7-17)$$

This was essentially verified on the level of convergence of integral representations for moments in [Borodin and Corwin 2014], and a complete proof for convergence of finite-dimensional distributions and moments with varying x was given in [Nica 2016] (in different scalings). It is very likely that the methods of [Nica 2016] are sufficient to achieve the same result for varying y as well; we will not address that here but rather focus on convergence of integral representations for joint moments instead. We need to start with an appropriate analogue of Lemmas 7.2 and 7.5.

Lemma 7.8. *Take a λ -coloured composition $\mu = 0^{m^{(0)}} 1^{m^{(1)}} 2^{m^{(2)}} \dots$ of length $m \geq 1$, where λ is a composition of weight $|\lambda| = m$, and assume that*

$$\mu_i = tL - \kappa_i \sqrt{L} + o(\sqrt{L}) \quad \text{as } L \rightarrow \infty, \quad 1 \leq i \leq m,$$

for a fixed m -tuple of reals $\kappa = (\kappa_1, \dots, \kappa_m)$ and $t > 0$. Then there exists a limit

$$\lim_{L \rightarrow \infty} \frac{e^{t\sqrt{L}(w_1 + \dots + w_m)} \cdot \mathfrak{p}_{\mu_{-1^m}(\sqrt{L} + w_1, \dots, \sqrt{L} + w_m)}}{L^{-(\mu_1 + \dots + \mu_m)/2}} = e^{\frac{t}{2}(w_1^2 + \dots + w_m^2)} \cdot \mathfrak{e}_\kappa(\lambda; w_1, \dots, w_m), \quad (7-18)$$

uniformly for bounded w_i 's, where the function $\mathfrak{e}_\kappa(w_1, \dots, w_m)$ can be characterized as follows.

(i) For a rainbow μ , in the dominant sector $\kappa_1 \geq \kappa_2 \geq \dots \geq \kappa_m$ one has (omitting λ from the notation)

$$\mathfrak{e}_\mu(w_1, \dots, w_m) = \exp(\kappa_1 w_1 + \dots + \kappa_m w_m), \quad (7-19)$$

and in the case $\kappa_i > \kappa_{i+1}$ for some $1 \leq i \leq m-1$ one uses the exchange relations (7-7) with (\mathfrak{e}, w_*) instead of (\mathfrak{f}, u_*) .

(ii) For a general labelling composition λ , let $\theta : \{1, \dots, m\} \rightarrow \{1, \dots, n\}$ be a monotone map with $\theta_*((1, \dots, 1)) = \lambda$. Then

$$\sum_{\text{rainbow } \kappa' : \theta_*(\kappa') = \kappa} \mathfrak{e}_{\kappa'}(w_1, \dots, w_m) = \mathfrak{e}_\kappa(\lambda; w_1, \dots, w_m). \quad (7-20)$$

Proof. It suffices to check the convergence for rainbow antidominant μ 's, as the other cases follow from that one by finite linear combinations (note that w_i 's are shifted but not scaled in the left-hand side of (7-18)). Then we need to prove that

$$\lim_{L \rightarrow \infty} \frac{e^{t\sqrt{L}(w_1+\dots+w_m)}(\sqrt{L}+w_1)^{-\mu_1} \dots (\sqrt{L}+w_m)^{-\mu_m}}{L^{-(\mu_1+\dots+\mu_m)/2}} = e^{\frac{t}{2}(w_1^2+\dots+w_m^2)} \cdot e^{\kappa_1 w_1+\dots+\kappa_m w_m}.$$

As the relation splits into a product over w_i 's, it suffices to consider the case of a single variable. We have

$$\begin{aligned} -\mu_1 \log(\sqrt{L}+w_1) &= -\mu_1 \left(\log \sqrt{L} + \left(\frac{w_1}{\sqrt{L}} - \frac{w_1^2}{2L} + o(L^{-1}) \right) \right) \\ &= -\frac{\mu_1 \log(L)}{2} - (tL - \kappa_1 \sqrt{L} + o(\sqrt{L})) \left(\frac{w_1}{\sqrt{L}} - \frac{w_1^2}{2L} + o(L^{-1}) \right) \\ &= \log L^{-\mu_1/2} - t\sqrt{L}w_1 + \kappa_1 w_1 + \frac{tw_1^2}{2} + o(1), \end{aligned}$$

as required. □

We can now make a limiting statement for the moments of $\mathcal{Z}^{(y)}(t, x)$.

Proposition 7.9. *Let $\kappa = (\kappa_1, \dots, \kappa_m)$ be a coloured composition of length m with real coordinates, and let the colours $s_1 < \dots < s_\alpha$ of the parts of κ also take real values; denote*

$$m_i^{(x)} = \#\{j : \kappa_j = x \text{ and has colour } s_i\}, \quad m_i = \sum_x m_i^{(x)}, \quad 1 \leq i \leq \alpha, \quad x \in \mathbb{R}.$$

Then

$$\begin{aligned} \mathbb{E} \prod_{(i,x) : m_i^{(x)} > 0} \frac{(\mathcal{Z}^{(s_i)}(t, x))^{m_i^{(x)}}}{m_i^{(x)}!} &= \frac{1}{(2\pi\sqrt{-1})^m} \int \dots \int \prod_{1 \leq i < j \leq m} \frac{w_j - w_i}{w_j - w_i - 1} \\ &\quad \times \prod_{k=1}^{\alpha} \frac{\exp(-s_k \cdot \sum_{r>m[1,k-1]}^{m[1,k]} w_r)}{m_k!} \cdot \mathfrak{e}_\kappa(\text{colour}(\kappa); w_1, \dots, w_m) \prod_{i=1}^m e^{tw_i^2/2} dw_i, \quad (7-21) \end{aligned}$$

where the integration is over upwardly oriented lines $w_i = a_i + \sqrt{-1} \cdot \mathbb{R}$ with $\Re a_j > \Re a_i + 1$ for $j > i$.

Sketch of the proof. We obtain (7-21) as a limit of (7-16). The convergence of the left-hand sides was discussed below (7-17). The convergence of the right-hand sides is a standard steepest descent argument with the main contribution coming from a finite neighbourhood of the critical point $v = \sqrt{L}$; see the proof of [Borodin and Corwin 2014, Proposition 5.4.2] for a similar situation. The change of variables $v_i = \sqrt{L} + w_i$, $1 \leq i \leq m$, together with Lemma 7.8, leads to the convergence of the integrands. □

Remark 7.10. Assume that the string $\kappa = (\kappa_1, \dots, \kappa_m)$ can be split into three sequential (possibly empty) substrings $\kappa = (\kappa', \kappa'', \kappa''')$, with all the coordinates of κ'' being (weakly) smaller than those of κ' and (weakly) larger than those of κ''' , and with all the colours s_i of the coordinates of κ'' being (strictly) larger

than those of κ' and (strictly) smaller than those of κ''' . A special case of this situation is the dominant sector $\kappa_1 \geq \dots \geq \kappa_m$ with rainbow compositions served by (7-19).

It is not hard to show from the definition of the ϵ_κ -functions in Lemma 7.8, that under a simultaneous shift of all the coordinates of κ'' by Δ that does not change the ordering conditions above, $\epsilon_\kappa(w_1, \dots, w_m)$ is multiplied by $\exp(\Delta(w_a + \dots, w_b))$, where $\kappa'' = (\kappa_a, \dots, \kappa_b)$. Hence, if one simultaneously performs the shift of all the colours of the parts of κ'' by the same amount Δ , then the right-hand side of (7-21) is not going to change. Since this means that the moments in the left-hand side do not change either, it is natural to conjecture that the joint distribution of the participating \mathcal{Z} 's also does not change (the moments do not determine this distribution uniquely, though). When κ'' consists of one part, this conjecture was verified in [Borodin et al. 2019], along with its versions for higher models, up to coloured stochastic vertex models in general “down-right” domains. For general κ'' this conjecture was very recently proved in [Dauvergne 2020] and [Galashin 2020] by two different methods.

Acknowledgments

Borodin was partially supported by NSF grants DMS-1664619 and DMS-1853981. Wheeler was partially supported by ARC grant DP19010289.

References

- [Aggarwal et al. 2019] A. Aggarwal, A. Borodin, and A. Bufetov, “Stochasticization of solutions to the Yang–Baxter equation”, *Ann. Henri Poincaré* **20**:8 (2019), 2495–2554. [MR](#)
- [Amir et al. 2011] G. Amir, I. Corwin, and J. Quastel, “Probability distribution of the free energy of the continuum directed random polymer in $1 + 1$ dimensions”, *Comm. Pure Appl. Math.* **64**:4 (2011), 466–537. [MR](#)
- [Barraquand and Corwin 2017] G. Barraquand and I. Corwin, “Random-walk in beta-distributed random environment”, *Probab. Theory Related Fields* **167**:3-4 (2017), 1057–1116. [MR](#)
- [Borodin 2017] A. Borodin, “On a family of symmetric rational functions”, *Adv. Math.* **306** (2017), 973–1018. [MR](#)
- [Borodin 2018] A. Borodin, “Stochastic higher spin six vertex model and Macdonald measures”, *J. Math. Phys.* **59**:2 (2018), [art. id. 023301]. [MR](#)
- [Borodin and Bufetov 2019] A. Borodin and A. Bufetov, “Color-position symmetry in interacting particle systems”, 2019. [arXiv](#)
- [Borodin and Corwin 2014] A. Borodin and I. Corwin, “Macdonald processes”, *Probab. Theory Related Fields* **158**:1-2 (2014), 225–400. [MR](#)
- [Borodin and Olshanski 2017] A. Borodin and G. Olshanski, “The ASEP and determinantal point processes”, *Comm. Math. Phys.* **353**:2 (2017), 853–903. [MR](#)
- [Borodin and Petrov 2017] A. Borodin and L. Petrov, “Integrable probability: stochastic vertex models and symmetric functions”, pp. 26–131 in *Stochastic processes and random matrices*, edited by G. Schehr et al., Oxford Univ. Press, 2017. [MR](#)
- [Borodin and Petrov 2018a] A. Borodin and L. Petrov, “Higher spin six vertex model and symmetric rational functions”, *Selecta Math. (N.S.)* **24**:2 (2018), 751–874. [MR](#)
- [Borodin and Petrov 2018b] A. Borodin and L. Petrov, “Inhomogeneous exponential jump model”, *Probab. Theory Related Fields* **172**:1-2 (2018), 323–385. [MR](#)
- [Borodin and Wheeler 2018] A. Borodin and M. Wheeler, “Coloured stochastic vertex models and their spectral theory”, preprint, 2018. [arXiv](#)
- [Borodin et al. 2014] A. Borodin, I. Corwin, and T. Sasamoto, “From duality to determinants for q -TASEP and ASEP”, *Ann. Probab.* **42**:6 (2014), 2314–2382. [MR](#)

- [Borodin et al. 2015a] A. Borodin, I. Corwin, L. Petrov, and T. Sasamoto, “Spectral theory for interacting particle systems solvable by coordinate Bethe ansatz”, *Comm. Math. Phys.* **339**:3 (2015), 1167–1245. [MR](#)
- [Borodin et al. 2015b] A. Borodin, I. Corwin, L. Petrov, and T. Sasamoto, “Spectral theory for the q -Boson particle system”, *Compos. Math.* **151**:1 (2015), 1–67. [MR](#)
- [Borodin et al. 2016a] A. Borodin, A. Bufetov, and M. Wheeler, “Between the stochastic six vertex model and Hall–Littlewood processes”, preprint, 2016. To appear in *J. Comb. Theory, Series A*. [arXiv](#)
- [Borodin et al. 2016b] A. Borodin, I. Corwin, and V. Gorin, “Stochastic six-vertex model”, *Duke Math. J.* **165**:3 (2016), 563–624. [MR](#)
- [Borodin et al. 2019] A. Borodin, V. Gorin, and M. Wheeler, “Shift-invariance for vertex models and polymers”, preprint, 2019. [arXiv](#)
- [Bosnjak and Mangazeev 2016] G. Bosnjak and V. V. Mangazeev, “Construction of R -matrices for symmetric tensor representations related to $U_q(\widehat{\mathfrak{sl}}_n)$ ”, *J. Phys. A* **49**:49 (2016), 495204, 19. [MR](#)
- [Calabrese et al. 2010] P. Calabrese, P. L. Doussal, and A. Rosso, “Free-energy distribution of the directed polymer at high temperature”, *EPL (Europhysics Letters)* **90**:2 (2010), 20002.
- [Corwin and Petrov 2016] I. Corwin and L. Petrov, “Stochastic higher spin vertex models on the line”, *Comm. Math. Phys.* **343**:2 (2016), 651–700. [MR](#)
- [Corwin et al. 2015] I. Corwin, T. Seppäläinen, and H. Shen, “The strict-weak lattice polymer”, *J. Stat. Phys.* **160**:4 (2015), 1027–1053. [MR](#)
- [Dauvergne 2020] D. Dauvergne, “Hidden invariance of last passage percolation and directed polymers”, preprint, 2020. [arXiv](#)
- [Dotsenko 2010] V. Dotsenko, “Bethe ansatz derivation of the Tracy–Widom distribution for one-dimensional directed polymers”, *EPL (Europhysics Letters)* **90**:2 (2010), [20003].
- [Foda and Wheeler 2013] O. Foda and M. Wheeler, “Colour-independent partition functions in coloured vertex models”, *Nuclear Phys. B* **871**:2 (2013), 330–361. [MR](#)
- [Galashin 2020] P. Galashin, “Symmetries of stochastic colored vertex models”, 2020. [arXiv](#)
- [Garbali et al. 2017] A. Garbali, J. de Gier, and M. Wheeler, “A new generalisation of Macdonald polynomials”, *Comm. Math. Phys.* **352**:2 (2017), 773–804. [MR](#)
- [Gwa and Spohn 1992] L.-H. Gwa and H. Spohn, “Six-vertex model, roughened surfaces, and an asymmetric spin Hamiltonian”, *Phys. Rev. Lett.* **68**:6 (1992), 725–728. [MR](#)
- [Kardar 1987] M. Kardar, “Replica Bethe ansatz studies of two-dimensional interfaces with quenched random impurities”, *Nuclear Phys. B* **290**:4 (1987), 582–602. [MR](#)
- [Kardar et al. 1986] M. Kardar, G. Parisi, and Y.-C. Zhang, “Dynamic Scaling of Growing Interfaces”, *Phys. Rev. Lett.* **56** (1986), 889–892.
- [Kuan 2018] J. Kuan, “An algebraic construction of duality functions for the stochastic $U_q(A_n^{(1)})$ vertex model and its degenerations”, *Comm. Math. Phys.* **359**:1 (2018), 121–187. [MR](#)
- [Kuniba et al. 2016] A. Kuniba, V. V. Mangazeev, S. Maruyama, and M. Okado, “Stochastic R matrix for $U_q(A_n^{(1)})$ ”, *Nuclear Phys. B* **913** (2016), 248–277. [MR](#)
- [Mimachi and Noumi 1998] K. Mimachi and M. Noumi, “A reproducing kernel for nonsymmetric Macdonald polynomials”, *Duke Math. J.* **91**:3 (1998), 621–634. [MR](#)
- [Nica 2016] M. Nica, “Intermediate disorder limits for multi-layer semi-discrete directed polymers”, preprint, 2016. [arXiv](#)
- [O’Connell and Ortmann 2015] N. O’Connell and J. Ortmann, “Tracy–Widom asymptotics for a random polymer model with gamma-distributed weights”, *Electron. J. Probab.* **20** (2015), [art. id.] 25. [MR](#)
- [O’Connell and Yor 2001] N. O’Connell and M. Yor, “Brownian analogues of Burke’s theorem”, *Stochastic Process. Appl.* **96**:2 (2001), 285–304. [MR](#)
- [Quastel 2012] J. Quastel, “Introduction to KPZ”, pp. 125–194 in *Current developments in mathematics, 2011*, edited by D. Jerison et al., International Press, Somerville, MA, 2012. [MR](#)

- [Sahi 1996] S. Sahi, “A new scalar product for nonsymmetric Jack polynomials”, *Internat. Math. Res. Notices* 20 (1996), 997–1004. [MR](#)
- [Sasamoto and Spohn 2010] T. Sasamoto and H. Spohn, “One-Dimensional Kardar–Parisi–Zhang Equation: An Exact Solution and its Universality”, *Phys. Rev. Lett.* **104** (2010), 230602.
- [Tracy and Widom 2008a] C. A. Tracy and H. Widom, “A Fredholm determinant representation in ASEP”, *J. Stat. Phys.* **132**:2 (2008), 291–300. [MR](#)
- [Tracy and Widom 2008b] C. A. Tracy and H. Widom, “Integral formulas for the asymmetric simple exclusion process”, *Comm. Math. Phys.* **279**:3 (2008), 815–844. [MR](#)
- [Tracy and Widom 2009] C. A. Tracy and H. Widom, “Asymptotics in ASEP with step initial condition”, *Comm. Math. Phys.* **290**:1 (2009), 129–154. [MR](#)

Received 22 Mar 2020. Revised 6 Sep 2020. Accepted 21 Sep 2020.

ALEXEI BORODIN: borodin@math.mit.edu

Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA, United States

and

Institute for Information Transmission Problems, Moscow, Russia

MICHAEL WHEELER: wheelerm@unimelb.edu.au

School of Mathematics and Statistics, University of Melbourne, Parkville VIC, Australia

EDITORS-IN-CHIEF

Alexei Borodin	Massachusetts Institute of Technology, United States
Hugo Duminil-Copin	IHÉS, France & Université de Genève, Switzerland
Robert Seiringer	Institute of Science and Technology, Austria
Sylvia Serfaty	Courant Institute, New York University, United States

EDITORIAL BOARD

Nalini Anantharaman	Université de Strasbourg, France
Scott Armstrong	Courant Institute, New York University, United States
Roland Bauerschmidt	University of Cambridge, UK
Ivan Corwin	Columbia University, United States
Mihalis Dafermos	Princeton University, United States
Semyon Dyatlov	University of California Berkeley, United States
Yan Fyodorov	King's College London, UK
Christophe Garban	Université Claude Bernard Lyon 1, France
Alessandro Giuliani	Università degli studi Roma Tre, Italy
Alice Guionnet	École normale supérieure de Lyon & CNRS, France
Pierre-Emmanuel Jabin	Pennsylvania State University, United States
Mathieu Lewin	Université Paris Dauphine & CNRS, France
Eyal Lubetzky	Courant Institute, New York University, United States
Jean-Christophe Mourrat	Courant Institute, New York University, United States
Laure Saint Raymond	École normale supérieure de Lyon & CNRS, France
Benjamin Schlein	Universität Zürich, Switzerland
Vlad Vicol	Courant Institute, New York University, United States
Simone Warzel	Technische Universität München, Germany

PRODUCTION

Silvio Levy	(Scientific Editor) production@msp.org
-------------	---

See inside back cover or msp.org/pmp for submission instructions.

Probability and Mathematical Physics (ISSN 2690-1005 electronic, 2690-0998 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

PMP peer review and production are managed by EditFlow[®] from MSP.

PUBLISHED BY
 **mathematical sciences publishers**
nonprofit scientific publishing
<https://msp.org/>

© 2020 Mathematical Sciences Publishers

PROBABILITY and MATHEMATICAL PHYSICS

1:1

2020

Introducing PMP	1
Alexei Borodin, Hugo Duminil-Copin, Robert Seiringer and Sylvia Serfaty	
Sharp spectral asymptotics for nonreversible metastable diffusion processes	3
Dorian Le Peutrec and Laurent Michel	
Joint distribution of Busemann functions in the exactly solvable corner growth model	55
Wai-Tong Louis Fan and Timo Seppäläinen	
Optimal lower bound on the least singular value of the shifted Ginibre ensemble	101
Giorgio Cipolloni, László Erdős and Dominik Schröder	
New critical exponent inequalities for percolation and the random cluster model	147
Tom Hutchcroft	
Mean-field tricritical polymers	167
Roland Bauerschmidt and Gordon Slade	
Observables of coloured stochastic vertex models and their polymer limits	205
Alexei Borodin and Michael Wheeler	