

# Tunisian Journal of Mathematics

an international publication organized by the Tunisian Mathematical Society

**Local estimates for Hörmander's operators  
with Gevrey coefficients and application  
to the regularity of their Gevrey vectors**

Makhlouf Derridj

2019 vol. 1 no. 3



# Local estimates for Hörmander's operators with Gevrey coefficients and application to the regularity of their Gevrey vectors

Makhlouf Derridj

Given a general Hörmander's operator  $P = \sum_{j=1}^m X_j^2 + Y + b$  in an open set  $\Omega \subset \mathbb{R}^n$ , where  $Y, X_1, \dots, X_m$  are smooth real vector fields in  $\Omega$ ,  $b \in C^\infty(\Omega)$ , and given also an open, relatively compact set  $\Omega_0$  with  $\bar{\Omega}_0 \subset \Omega$ , and  $s \in \mathbb{R}$ ,  $s \geq 1$ , such that the coefficients of  $P$  are in  $G^s(\Omega_0)$  and  $P$  satisfies a  $\frac{1}{p}$ -Sobolev estimate in  $\Omega_0$ , our aim is to establish local estimates reflecting local domination of ordinary derivatives by powers of  $P$ , in  $\Omega_0$ . As an application, we give a direct proof of the  $G^{2ps}(\Omega_0)$ -regularity of any  $G^s(\Omega_0)$ -vector of  $P$ .

1. Introduction	321
2. Some notation and definitions	323
3. Preliminary lemmas and propositions	325
4. Local relations of domination by powers of $P$	333
5. Gevrey regularity for Gevrey vectors	340
Acknowledgements	344
References	344

## 1. Introduction

The study of the regularity of analytic vectors of partial differential operators goes back to the work of T. Kotake and N. S. Narasimhan [1962] who proved the (local) analyticity of (local) analytic vectors of elliptic operators with analytic coefficients (see also [Nelson 1959] for another related context). This property, called the “iteration property” or even the “Kotake–Narasimhan property” was further studied in the following decades in more general situations (such as systems, or nonelliptic operators) and also in the Gevrey categories  $G^s$ ,  $s \geq 1$  ( $s = 1$  corresponds to the analytic case). This was in particular the case for the class of differential operators of principal type with analytic coefficients and also, after the famous article by L. Hörmander on hypoelliptic operators of second order, the systems of real-analytic real vector fields satisfying the so-called Hörmander condition, and

*MSC2010:* 35B65, 35G99, 35J70, 35K65.

*Keywords:* Gevrey vectors, Degenerate elliptic-parabolic differential operators.

also for the Hörmander's operators themselves. A result of G. Métivier [1978] shows that the "iteration property" is not true for nonelliptic operators in the Gevrey category  $G^s$ ,  $s > 1$ , and another one by M.S. Baouendi and Métivier [1982] gave the "iteration property" for hypoelliptic operators of principal type in the analytic setting. Concerning analytic real vector fields, satisfying Hörmander's condition, that property was shown by M. Damak and B. Helffer [1980], followed by a more precise version by Helffer and C. Mattera [1980].

More recently, a series of papers studying the case of involutive systems of analytic complex vector fields, concerning analytic or more generally Gevrey vectors, have been published [Barostichi et al. 2011; Castellanos et al. 2013]. Even knowing that the "iteration property" is not true for nonelliptic operators, one can ask about the Gevrey regularity  $G^{s'}$  of an  $s$ -Gevrey vector ( $s \geq 1$ ) and even give the best  $s'$  one may obtain. Such a study is contained in the above mentioned papers. A more recent paper by N. Braun Rodrigues, G. Chinni, P. Cordaro and M. Jahnke [Braun Rodrigues et al. 2016] was partly devoted to the study of global analytic vectors for some sums of squares on a product of two tori. A little later, we studied in [Derridj  $\geq$  2019] the case of  $G^k(\Omega)$ -vectors of Hörmander's operators of the first kind (or degenerate elliptic) with  $G^k(\Omega)$  coefficients,  $\Omega$  an open set in  $\mathbb{R}^n$  (see the definitions in Section 2), and  $k \in \mathbb{N}^*$  (in particular analytic vectors for  $k = 1$ ), in which we proved an optimal result (the optimality following from the work in the global case by the authors of [Braun Rodrigues et al. 2016]).

In this paper, we consider general Hörmander's operators  $P$  (of the second kind or degenerate elliptic parabolic) for which we study the existence of local estimates giving local domination of the ordinary derivatives by powers of  $P$ , when the coefficients of  $P$  are in  $G^s(\Omega_0)$ ,  $\bar{\Omega}_0 \subset \Omega$ , and  $P$  satisfies a " $\frac{1}{p}$ -Sobolev estimate" on  $\Omega_0$  (see Section 2). This, with our main result Theorem 4.2, is used to obtain  $G^{2ps}(\Omega_0)$ -regularity for  $G^s(\Omega_0)$ -vectors of  $P$  ( $s \geq 1$ ), providing therefore, first a direct proof, without using the method of addition of an extra variable, and second the result, with any  $s \in \mathbb{R}$ ,  $s \geq 1$ . Let us remark that, in our preceding result, as we used the above method, our result was obtained there for  $s = k \in \mathbb{N}^*$ .

In a forthcoming paper, we will consider operators of the first kind, for which the integer  $p$  (giving the  $\frac{1}{p}$ -Sobolev estimate in  $\Omega_0$ ) is intimately related to the vector fields  $(X_1, \dots, X_m)$  and prove finer local estimates of domination by powers of  $P$ , giving, as an application, the optimal  $G^{ps}(\Omega_0)$  regularity for any  $G^s(\Omega_0)$ -vector of  $P$ . A complete survey on results in this field until 1987 may be found in [Bolley et al. 1987] and a more very recent short one may be found in [Derridj 2017].

We recall in Section 2 some definitions and elementary facts about the operators  $P$ , Gevrey functions and Gevrey vectors. Section 3 will be devoted to preliminary lemmas and propositions as a preparation for the proof of our main theorem.

Our main theorem will be proved in [Section 4](#) and the last section is devoted to the application of the main result to the regularity of Gevrey vectors of  $P$ .

### 2. Some notation and definitions

We consider a system  $(Y, X_1, \dots, X_m)$  of real vector fields with smooth coefficients on an open set  $\Omega \subset \mathbb{R}^n$ , and

$$P = \sum_{j=1}^m X_j^2 + Y + b, \quad b \in C^\infty(\Omega), \tag{2-1}$$

see [\[Hörmander 1967; Kohn 1978; Rothschild and Stein 1976\]](#). Let us just recall Hörmander's condition for hypoellipticity in  $\Omega$  of the operator (2-1):

$$\text{The Lie algebra, } \text{Lie}(Y, X_1, \dots, X_m), \text{ generated by the} \tag{2-2}$$

$$\text{vector fields } Y, X_1, \dots, X_m, \text{ is of maximal rank in } \Omega.$$

Concretely, the family of brackets of all lengths of the vector fields  $Y, X_1, \dots, X_m$  span at any point  $x \in \Omega$  the tangent space at  $x$ .

Under the condition (2-2), Hörmander proved the following a priori subelliptic estimate which we briefly describe. The  $L^2$ -norm and Sobolev  $H^\sigma$ -norm are denoted by  $\|\cdot\|$  and  $\|\cdot\|_\sigma$ .

Let us recall, below, some norms introduced by Hörmander [\[1967\]](#):

$$\begin{aligned} \|v\| &= \|v\| + \sum_{j=1}^m \|X_j v\|, \quad v \in \mathcal{D}(\Omega), \\ \|v\|' &= \sup\{\|(v, w)\| : \|w\| < 1, w \in \mathcal{D}(\Omega)\} \leq \|v\|. \end{aligned} \tag{2-3}$$

**Theorem 2.1** [\[Hörmander 1967\]](#). *Let  $\Omega_0 \Subset \Omega$  and assume (2-2). Then there exist  $\sigma > 0$  and  $C > 0$  such that*

$$\|v\|_\sigma \leq C(\|v\| + \|Yv\|'), \quad \text{for all } v \in \mathcal{D}(\Omega_0). \tag{2-4}$$

We call (2-4) a subelliptic estimate for  $P$ .

The constants  $\sigma$  and  $C$  depend on  $\Omega_0$  and  $Y, X_1, \dots, X_m$ . More specifically,  $\sigma$  depends on the length of the brackets needed in order to span the tangent space at every point of  $\bar{\Omega}_0$ .

Now, elementary technical manipulations give, with  $C_0 > 0$ ,

$$\|v\|_\sigma \leq C_0(\|Pv\|' + \|v\|), \quad \text{for all } v \in \mathcal{D}(\Omega_0). \tag{2-5}$$

A particular ingredient in order to get (2-5) and which we need in the sequel is the set of obvious inequalities:

$$\|X_j v\|' \leq C\|v\|, \quad \forall v \in \mathcal{D}(\Omega), \text{ for some } C > 0, j = 1, \dots, m. \tag{2-6}$$

We want to say a word on the case  $Y = 0$ , or more generally the case where in (2-2) one considers the Lie algebra  $\text{Lie}(X_1, \dots, X_m)$  generated by  $X_1, \dots, X_m$ ; in that case, one has a more precise estimate. We considered that case in our preceding work, obtaining an optimal result for the Gevrey regularity of  $k$ -Gevrey vectors of  $P$  (named in that case Hörmander's operator of the first kind),  $k \in \mathbb{N}^*$ .

Coming back to our general case (named operator of the second kind), we need to write a more precise a priori estimate than (2-5) which we need to consider in the sequel:

$$\|v\|_\sigma + \sum_{j=1}^m \|X_j v\| \leq C_0(\|Pv\|' + \|v\|), \quad \text{for all } v \in \mathcal{D}(\Omega_0). \quad (2-7)$$

Again this is easily obtained, using (2-4).

Let us recall, in order to be complete, definitions of Gevrey functions and Gevrey vectors of a differential operator of order  $m$  (here it will be  $m = 2$ ).

**Definition 2.2.** Let  $s \geq 1$ . The space of Gevrey functions of order  $s$ ,  $G^s(\Omega)$ , is defined as

$$G^s(\Omega) := \left\{ u \in C^\infty(\Omega) : \forall K \Subset \Omega, \exists C_K > 0 \text{ s.t. } |\partial^\alpha u|_K \leq C_K^{|\alpha|+1} |\alpha|!^s \right. \\ \left. \forall \alpha \in \mathbb{N}^n, |\alpha| = \sum_{j=1}^n \alpha_j, \partial^\alpha = \partial_{x_1}^{\alpha_1} \dots \partial_{x_n}^{\alpha_n} \right\}. \quad (2-8)$$

**Remark 2.3.** It is known that for  $s > 1$ , one has the property of partition of unity in  $G^s(\Omega)$ : in particular, given two open sets  $\Omega_1, \Omega_2$ , with  $\bar{\Omega}_1$  compact and  $\bar{\Omega}_1 \subset \Omega_2$ , there exists a function  $\varphi \in \mathcal{D}(\Omega_2)$ ,  $\varphi \equiv 1$  on  $\bar{\Omega}_1$ ,  $\varphi \in G^s(\Omega_2)$ .

**Definition 2.4.** Let  $P$  be a differential operator of order  $m$  in  $\Omega$ . The space  $G^s(\Omega, P)$  of  $s$ -Gevrey vectors of  $P$  in  $\Omega$ ,  $s \geq 1$ , is defined as

$$G^s(\Omega, P) := \left\{ u \in L^2_{\text{loc}}(\Omega) : \forall K \Subset \Omega, \exists C_K > 0 \text{ s.t. } \forall k \in \mathbb{N}, \right. \\ \left. P^k u \in L^2(K) \text{ and } \|P^k u\|_{L^2(K)} \leq C_K^{k+1} (mk)!^s \right\}. \quad (2-9)$$

As in our case,  $P$  is of order 2 and hypoelliptic, with a subelliptic estimate (2-5) or (2-7), (2-9) reduces to

$$G^s(\Omega, P) := \left\{ u \in C^\infty(\Omega) : \forall K \Subset \Omega, \exists C_K > 0 \right. \\ \left. \text{s.t. } \forall k \in \mathbb{N}, \|P^k u\|_{L^2(K)} \leq C_K^{k+1} (2k)!^s \right\}. \quad (2-10)$$

**Remark 2.5.** We used in our definitions (2-9), (2-10), the commonly used  $L^2$ -norm, but in some specific situations, such as for systems of complex vectors, other norms are used [Barostichi et al. 2011; Castellanos et al. 2013].

### 3. Preliminary lemmas and propositions

When trying to get estimates for derivatives  $\partial^\alpha u$  of a function  $u$ , knowing  $Pu$ , we are faced in particular with the study of commutators  $[P, \partial^\alpha]$ ,  $\alpha \in \mathbb{N}^n$ , so, to the study of commutators  $[X_j^2, \partial^\alpha]$ ,  $[Y + b, \partial^\alpha]$ . Now, one has the following equality:

$$[X_j^2, \partial^\alpha] = 2X_j[X_j, \partial^\alpha] - [X_j, [X_j, \partial^\alpha]]. \quad (3-1)$$

So we have to look closely at the commutators  $[X_j, \partial^\alpha]$ ,  $[Y + b, \partial^\alpha]$  and the double commutators  $[X_j, [X_j, \partial^\alpha]]$ . In order to get an estimate for the coefficients of the differential operators above, it is sufficient to consider the following basic commutators

$$\begin{aligned} [a_\ell \partial_\ell, \partial^\alpha], \quad \ell = 1, \dots, n, \quad a_\ell \in G^s(\Omega), \\ [a_\ell \partial_\ell, [a_k \partial_k, \partial^\alpha]], \quad \ell, k = 1, \dots, n, \quad a_\ell, a_k \in G^s(\Omega), \end{aligned} \quad (3-2)$$

as  $Y$  and  $X_j$  are linear combinations of the basic vector fields  $a_\ell \partial_\ell$ ,  $a_\ell \in G^s(\Omega)$ .

Of course, the commutator  $[b, \partial^\alpha]$  is elementary. Then

$$\begin{aligned} [b, \partial^\alpha] &= - \sum_{\beta < \alpha} \binom{\alpha}{\beta} b^{(\alpha-\beta)} \partial^\beta = \sum_{\beta < \alpha} b_{\alpha\beta} \partial^\beta, \\ &\quad \binom{\alpha}{\beta} = \prod_j \binom{\alpha_j}{\beta_j}, \quad \beta < \alpha \Leftrightarrow \beta_j \leq \alpha_j, \quad \beta \neq \alpha, \\ [a_\ell \partial_\ell, \partial^\alpha] &= - \sum_{\beta < \alpha} \binom{\alpha}{\beta} a_\ell^{(\alpha-\beta)} \partial^{\beta+\ell}, \\ &\quad \beta + \ell = \{(\beta + \ell)_i = \beta_i, \quad i \neq \ell, \quad (\beta + \ell)_\ell = \beta_\ell + 1\}. \end{aligned} \quad (3-3)$$

Let us, now, give the expression of  $[a_\ell \partial_\ell, [a_k \partial_k, \partial^\alpha]]$  or, in view of (3-3),

$$\sum_{\beta < \alpha} \binom{\alpha}{\beta} [a_k^{(\alpha-\beta)}, \partial^{\beta+k}, a_\ell \partial_\ell].$$

But for  $\beta < \alpha$ ,

$$[a_k^{(\alpha-\beta)} \partial^{\beta+k}, a_\ell \partial_\ell] = \sum_{\gamma < \beta+k} \binom{\beta+k}{\gamma} a_k^{(\alpha-\beta)} a_\ell^{(\beta+k-\gamma)} \partial^{\gamma+\ell} - a_\ell a_k^{(\alpha-\beta+\ell)} \partial^{\beta+k}.$$

Hence we get

$$\begin{aligned} [a_\ell \partial_\ell, [a_k \partial_k, \partial^\alpha]] \\ = \sum_{\substack{\gamma < \beta+k, \\ \beta < \alpha}} \binom{\alpha}{\beta} \binom{\beta+k}{\gamma} a_k^{(\alpha-\beta)} a_\ell^{(\beta+k-\gamma)} \partial^{\gamma+\ell} - \sum_{\beta < \alpha} \binom{\alpha}{\beta} a_\ell a_k^{(\alpha-\beta+\ell)} \partial^{\beta+k}. \end{aligned} \quad (3-4)$$

In the first sum in the second member of (3-4), we distinguish two families of  $\gamma$ 's such that  $\gamma < \beta + k$ :

(i) If  $\gamma_k = 0$ , then  $\gamma \leq \beta$ , so  $\gamma < \alpha$  (as  $\beta < \alpha$ ). Hence in that case

$$\begin{aligned} \sum_{\beta < \alpha, \gamma < \beta + k, \gamma_k = 0} \binom{\alpha}{\beta} \binom{\beta + k}{\gamma} a_k^{(\alpha - \beta)} a_\ell^{(\beta + k - \gamma)} \partial^{\gamma + \ell} \quad \text{is a part of} \\ \sum_{\beta < \alpha, \gamma \leq \beta} \binom{\alpha}{\beta} \binom{\beta + k}{\gamma} a_k^{(\alpha - \beta)} a_\ell^{(\beta + k - \gamma)} \partial^{\gamma + \ell}. \end{aligned} \quad (3-5)$$

(ii) If  $\gamma_k \neq 0$ , we set  $\delta = (\delta_1, \dots, \delta_n)$  with  $\delta_\rho = \gamma_\rho$  if  $\rho \neq k$  and  $\delta_k = \gamma_k - 1$ . So  $\delta < \beta < \alpha$ , in particular  $|\delta| \leq |\alpha| - 2$  (easy to see) and  $\gamma = \delta + k$ . Hence

$$\begin{aligned} \sum_{\beta < \alpha, \gamma < \beta + k, \gamma_k \neq 0} \binom{\alpha}{\beta} \binom{\beta + k}{\gamma} a_k^{(\alpha - \beta)} a_\ell^{(\beta + k - \gamma)} \partial^{\gamma + \ell} \quad \text{is a part of} \\ \sum_{\beta < \alpha, \delta \leq \beta} \binom{\alpha}{\beta} \binom{\beta + k}{\delta + k} a_k^{(\alpha - \beta)} a_\ell^{(\beta - \delta)} \partial^{\delta + \ell + k}. \end{aligned} \quad (3-6)$$

Setting  $I_{(\ell, k)} = (I_1, \dots, I_n)$  with  $I_\rho = 0$  if  $\rho \notin \{\ell, k\}$  and  $I_\rho = 1$  if  $\rho \in \{\ell, k\}$ , the sum in the second line of (3-6) can be written as

$$\sum_{\substack{\beta < \alpha \\ \delta < \beta}} \binom{\alpha}{\beta} \binom{\beta + k}{\delta + k} a_k^{(\alpha - \beta)} \partial^{\delta + I_{(\ell, k)}} \quad (3-7)$$

So, looking at (3-4) and in view of (3-5), (3-6) and taking as 0 the other coefficients in  $\sum_{\beta < \alpha, \gamma \leq \beta}$  (in (3-5)) and  $\sum_{\beta < \alpha, \delta < \beta}$  (in (3-6)) which are not in (3-4), we may write

$$\begin{aligned} [a_\ell \partial_\ell, [a_k \partial_k, \partial^\alpha]] \\ = - \sum_{\beta < \alpha} a_{\ell k \alpha \beta} \partial^{\beta + k} + \sum_{\gamma < \alpha} b_{\ell k \alpha \gamma} \partial^{\gamma + \ell} + \sum_{\substack{\delta < \alpha \\ |\delta| \leq |\alpha| - 2}} c_{\ell k \alpha \delta} \partial^{\delta + I_{(\ell, k)}}, \end{aligned} \quad (3-8)$$

$$\text{where } a_{\ell k \alpha \beta} = \binom{\alpha}{\beta} a_\ell a_k^{(\alpha - \beta + \ell)},$$

$$b_{\ell k \alpha \gamma} = \sum_{\gamma \leq \beta < \alpha} \binom{\alpha}{\beta} \binom{\beta + k}{\gamma} a_k^{(\alpha - \beta)} a_\ell^{(\beta + k - \gamma)},$$

$$c_{\ell k \alpha \delta} = \sum_{\delta < \beta < \delta} \binom{\alpha}{\beta} \binom{\beta + k}{\delta + k} a_k^{(\alpha - \beta)} a_\ell^{(\beta - \delta)}.$$

Now considering a vector field  $X_j$  with smooth coefficients  $a_{j\ell}$ , i.e.,

$$X_j = \sum_{\ell=1}^n a_{j\ell} \partial_\ell = \sum_{\ell=1}^n Z_{j\ell},$$

we obtain from (3-3)

$$[X_j, \partial^\alpha] = \sum_{\ell=1}^n [Z_{j\ell}, \partial^\alpha] = - \sum_{\substack{\beta < \alpha \\ \ell=1, \dots, n}} \binom{\alpha}{\beta} a_{j\ell}^{(\alpha - \beta)} \partial^{\beta + \ell} = \sum_{\substack{\beta < \alpha \\ \ell=1, \dots, n}} a_{j\alpha\beta\ell} \partial^{\beta + \ell}, \quad (3-9)$$

$$[Y, \partial^\alpha] = \sum_{\ell=1}^n [b_\ell \partial_\ell, \partial^\alpha] = - \sum_{\substack{\beta < \alpha \\ \ell=1, \dots, n}} \binom{\alpha}{\beta} b_\ell^{(\alpha - \beta)} \partial^{\beta + \ell} = \sum_{\substack{\beta < \alpha \\ \ell=1, \dots, n}} b_{\alpha\beta\ell} \partial^{\beta\ell}.$$

Concerning the double brackets, we obtain:

$$[X_j, [X_j, \partial^\alpha]] = \sum_{\ell, k} [Z_{j\ell}, [Z_{jk}, \partial^\alpha]], \quad \text{with } [Z_{j\ell}, [Z_{k\ell}, \partial^\alpha]] \text{ given in (3-8),}$$

where  $a_{\ell k\alpha\beta}$  is replaced by  $a_{j\ell k\alpha\beta}$ , and so on. (3-10)

Now we assume that the coefficients of  $P$  are in  $G^s(\Omega)$ . So  $a_{j\ell}$ ,  $b_\ell$  and  $b$  are in  $G^s(\Omega)$ . So we have that for any compact  $K$  in  $\Omega$ , there exists  $C_K > 0$  such that

$$\text{for all } \nu \in \mathbb{N}^n,$$

$$|a_{j\ell}^{(\nu)}|_K + |b_\ell^{(\nu)}|_K + |b^{(\nu)}|_K \leq C_K^{|\nu|+1} |\nu|!^s, \quad \ell \in \{1, \dots, n\}, j \in \{1, \dots, m\}. \quad (3-11)$$

Writing (3-10) more concretely, we get using (3-8)

$$[X_j, [X_j, \partial^\alpha]] = \sum_{\substack{\beta < \alpha \\ k=1, \dots, n}} d_{jk\alpha\beta} \partial^{\beta+k} + \sum_{\substack{\beta < \alpha \\ k, \ell=1, \dots, n}} c_{j\ell k\alpha\beta} \partial^{\beta+\ell+k},$$

where  $d_{jk\alpha\beta} = -\sum_{\ell=1}^n a_{j\ell k\alpha\beta} + \sum_{\ell=1}^n b_{jk\ell\alpha\beta}$ . (3-12)

We want now to get estimates for the coefficients of the brackets and double brackets in (3-9) and (3-12) and the operators associated to these brackets, when the coefficients are in  $G^s(\Omega)$ .

**Proposition 3.1.** *Assume the coefficients of  $P$  are in  $G^s(\Omega)$ . Given any compact set  $K$  in  $\Omega$ , there exists  $B = B_K > 0$  such that the coefficients of the operators  $[b, \partial^\alpha]$  (given in (3-3)),  $[X_j, \partial^\alpha]$ ,  $[Y, \partial^\alpha]$  (given in (3-9)),  $[X_j, [X_j, \partial^\alpha]]$  (given in (3-12)) satisfy the following estimates:*

$$|b_{\alpha\beta}|_K + |b_{\alpha\beta\ell}|_K + |a_{j\alpha\beta\ell}|_K + |\nabla b_{\alpha\beta}|_K$$

$$+ |\nabla b_{\alpha\beta\ell}|_K + |\nabla a_{j\alpha\beta\ell}|_K \leq B^{|\alpha-\beta|} \left(\frac{\alpha!}{\beta!}\right)^s,$$

$$|c_{j\ell\alpha\beta}|_K + |\nabla c_{j\ell k\alpha\beta}|_K \leq B^{|\alpha-\beta|} \left(\frac{(\alpha+k)!}{(\beta+k)!}\right)^s, \quad (3-13)$$

$$|d_{jk\alpha\beta}|_K + |\nabla d_{jk\alpha\beta}|_K \leq B^{|\alpha-\beta|} (|\alpha| + 1) \left(\frac{\alpha!}{\beta!}\right)^s.$$

*Proof.* We recall first that  $\beta < \alpha$  and  $(\alpha+k)_\ell = \alpha_\ell$  for  $\ell \neq k$  and  $(\alpha+k)_k = \alpha_k + 1$ . The first line comes easily from the expression of the functions  $b_{\alpha\beta}$ ,  $b_{\alpha\beta\ell}$ ,  $a_{j\alpha\beta\ell}$ , and their derivatives. Note that we took  $B^{|\alpha-\beta|}$  in place of  $B^{|\alpha-\beta|+1}$  as  $|\alpha-\beta| \geq 1$  and used (3-11). The proof for the other functions needs more work. We first use the following estimate for  $a_{jk}^{(\alpha-\beta)} a_{j\ell}^{(\beta-\delta)}$ , which is in the expression of  $c_{j\ell k\alpha\delta}$  (see last line in (3-8)):

$$\left| a_{jk}^{(\alpha-\beta)} a_{j\ell}^{(\beta-\delta)} \binom{\alpha}{\beta} \binom{\beta+k}{\delta+k} \right| \leq B^{|\alpha-\beta|} (\lambda B)^{|\beta-\delta|} \left(\frac{(\alpha+k)!}{(\delta+k)!}\right)^s, \quad \lambda \geq 1. \quad (3-14)$$

For that we used

$$\frac{\alpha! (\beta+k)!}{\beta! (\delta+k)!} = \frac{\alpha! (\beta_k+1)}{(\delta+k)!} \leq \frac{(\alpha+k)!}{(\delta+k)!}.$$



Hence

$$|c_{j\ell k\alpha\delta}|_K \leq \left( \sum_{\delta < \beta < \alpha} B^{|\alpha-\beta|} (\lambda B)^{|\beta-\delta|} \right) \left( \frac{(\alpha+k)!}{(\delta+k)!} \right)^s, \quad \lambda \geq 1.$$

As  $(\lambda B)^{|\beta-\delta|} = (\lambda B)^{|\alpha-\delta|} (\lambda B)^{-|\alpha-\beta|}$ , ( $\delta < \beta < \alpha$ ), we get:

$$|c_{j\ell k\alpha\delta}|_K \leq \left( \frac{(\alpha+k)!}{(\delta+k)!} \right)^s (\lambda B)^{|\alpha-\delta|} \sum_{\beta < \alpha} \lambda^{-|\alpha-\beta|}.$$

Now we use the following easy lemma:

**Lemma 3.2.** *There exists  $\epsilon_0 > 0$  (independent from  $\alpha$ ) such that if  $0 < \epsilon \leq \epsilon_0$  then  $\sum_{\beta < \alpha} \epsilon^{|\alpha-\beta|} \leq 1$ .*

This comes from the fact that if  $\beta_j \leq \alpha_j$ ,  $j = 1, \dots, n$ , and  $\alpha \neq \beta$  then  $\sum_{j=1}^n \beta_j < \sum_{j=1}^n \alpha_j$ . In fact, setting  $\lambda_j = \alpha_j - \beta_j \in \mathbb{N}$ ,  $\sum_{j=1}^n \lambda_j \geq 1$ ,

$$\begin{aligned} \sum \epsilon^{\sum \lambda_j} &\leq \sum_{\lambda_1 \geq 1} \epsilon^{\sum \lambda_j} + \dots + \sum_{\lambda_n \geq 1} \epsilon^{\sum \lambda_j} \\ &\leq \epsilon \left( \sum_{(\lambda_2, \dots, \lambda_n) \in \mathbb{N}^{n-1}} \epsilon^{\lambda_2 + \dots + \lambda_n} + \dots + \sum_{(\lambda_1, \dots, \lambda_{n-1}) \in \mathbb{N}^{n-1}} \epsilon^{\lambda_1 + \dots + \lambda_{n-1}} \right) \\ &\leq \epsilon n 2^{n-1} \quad \text{if } \epsilon \leq \frac{1}{2}. \end{aligned}$$

The lemma follows by taking  $\epsilon_0 = (n 2^{n-1})^{-1}$ .

Coming back to our proof, we consider  $\lambda_0 = n 2^{n-1} = \epsilon_0^{-1}$  and replace  $B$  by  $\lambda_0 B$ . We get the estimate for  $c_{jk\alpha\beta}$ . As  $d_{jk\alpha\beta} = -\sum_{\ell=1}^n (a_{j\ell k\alpha\beta} - b_{j\ell k\alpha\beta})$ , we just need to bound on  $K$  the functions  $a_{j\ell k\alpha\beta}$  and  $b_{j\ell k\alpha\beta}$  for  $\ell = 1, \dots, n$ . The worst term is  $b_{j\ell k\alpha\gamma}$ . As we did above, we use, with  $\lambda \geq 1$ ,

$$\left| \binom{\alpha}{\beta} \binom{\beta+\ell}{\gamma} a_\ell^{(\alpha-\beta)} a_k^{(\beta+\ell-\gamma)} \right| \leq B^{|\alpha-\beta|+1} (\lambda B)^{|\beta-\gamma|} \left( \frac{\alpha!}{\gamma!} (\beta_\ell + 1) \right)^s.$$

So,

$$\begin{aligned} |b_{j\ell k\alpha\gamma}| &\leq \left( \sum_{\gamma \leq \beta < \alpha} B^{|\alpha-\beta|+1} (\lambda B)^{|\beta-\gamma|} \right) \left( \frac{(\alpha+\ell)!}{\gamma!} \right)^s \\ &\leq \left( \frac{(\alpha+\ell)!}{\gamma!} \right)^s B (\lambda B)^{|\alpha-\gamma|} \sum_{\beta < \alpha} \lambda^{-|\alpha-\beta|} \\ &\leq B \left( \frac{\alpha!}{\gamma!} \right)^s (|\alpha|+1)^s (\lambda B)^{|\alpha-\gamma|} \sum_{\beta < \alpha} \lambda^{-|\alpha-\beta|}. \end{aligned}$$

Now taking  $\lambda \geq \lambda_0$ ,  $B$  large enough, we obtain what we want (more precisely we take  $\tilde{B}$  such that (as  $|\alpha-\gamma| \geq 1$ ),  $B(\lambda B)^{|\alpha-\gamma|} \leq (\lambda \tilde{B})^{|\alpha-\gamma|}$ , and then choose the final  $B$  as  $\lambda \tilde{B}$ ). Concerning the derivatives of first order, we just have to apply what we did, using bounds on  $K$ , not only for the coefficients of  $P$ , but also bounds of

their derivatives of first order. In order to be rigorous and complete, let us bound a derivative of  $b_{j\ell k\alpha\gamma}$ , which we denote by  $b'_{j\ell k\alpha\gamma}$ . Then we have (see (3-8))

$$b'_{j\ell k\alpha\beta} = \sum_{\gamma \leq \beta < \alpha} \binom{\alpha}{\beta} \binom{\beta+k}{\gamma} ((a_{jk}^{(\alpha-\beta)})' a_{j\ell}^{(\beta+k-\gamma)} + a_{jk}^{(\alpha-\beta)} (a_{j\ell}^{(\beta+k-\gamma)})').$$

So we just have to do the same as before to get the bounds on  $K$  for the functions  $a'_{jk}$ ,  $a_{j\ell}$ ,  $a_{jk}$  and  $a'_{j\ell}$ . Hence taking  $B$ , greater if necessary, we obtain (3-13).  $\square$

As a consequence, we obtain the following:

**Proposition 3.3.** *Assume that the coefficients of  $P$  are in  $G^s(\Omega)$ . Then for every relatively compact open set  $\Omega_0$  in  $\Omega$  ( $\Omega_0 \Subset \Omega$ ), there exists  $B = B(\Omega_0, P) > 0$  such that, for every  $\tau$ ,  $0 \leq \tau \leq 1$ , one has for  $j = 1, \dots, m$ ,  $\beta < \alpha$ ,  $\ell, k \in \{1, \dots, n\}$ ,*

$$\begin{aligned} \|b_{\alpha\beta} v\|_{\tau} + \|b_{\alpha\beta\ell} v\|_{\tau} + \|a_{j\alpha\beta\ell} v\|_{\tau} &\leq B^{|\alpha-\beta|} \left(\frac{\alpha!}{\beta!}\right)^s \|v\|_{\tau}, \quad \beta < \alpha, \ell = 1, \dots, n, \\ \|c_{j\ell k\alpha\beta} v\|_{\tau} &\leq B^{|\alpha-\beta|} \left(\frac{(\alpha+k)!}{(\beta+k)!}\right)^s \|v\|_{\tau}, \\ \|d_{jk\alpha\beta} v\|_{\tau} &\leq B^{|\alpha-\beta|} \left(\frac{\alpha!(|\alpha|+1)}{\beta!}\right)^s \|v\|_{\tau}, \quad \forall v \in \mathcal{D}(\Omega_0). \end{aligned} \quad (3-15)$$

*Proof.* From inequalities (3-13), one obtains estimates (3-15) for  $\tau = 0$  and  $\tau = 1$ . Then (3-15) follows from the cases  $\tau = 0$  and  $\tau = 1$  by interpolation between Sobolev spaces  $L^2$  and  $H^1$ .  $\square$

**Remark 3.4.** Our Proposition 3.1 is a refinement of our Lemma 5.3 in [Derridj  $\geq$  2019]. We need this refinement in order to prove our local estimates of ordinary derivatives in terms of powers of  $P$ .

In order to begin to state what we need for our results, we make the following assumption:

$$\text{Estimate (2-4) is true with } \sigma = \frac{1}{p}, \quad p \in \mathbb{N}^*. \quad (3-16)$$

Then our local estimates will use the equality  $\sigma = \frac{1}{p}$ . In this section, we want to give and prove the basic ones which we will use in another section in order to give a sequence of local estimates for  $P$ , assuming (2-7), (3-16) and  $P$  with coefficients in  $G^s(\Omega)$ , for some  $s \geq 1$ .

**Proposition 3.5.** *Assume (2-4), (3-16) and that  $P$  has smooth coefficients. Let  $\Omega_1$  be a relatively compact open subset of  $\Omega_0$ . Then there exists a constant  $C = C(\Omega_0, P) > 0$  such that for all  $(u, \varphi) \in C^\infty(\overline{\Omega}_1) \cap \mathcal{D}(\Omega_1)$ ,*

$$\|\varphi u\|_{\sigma} \leq C \left( \|\varphi P u\|' + \sum_{|\beta| \leq 2} \|\varphi^{(\beta)} u\| \right). \quad (3-17)$$

*Proof.* This proposition has a simple proof. Taking  $v = \varphi u$  in (2-7), we get

$$\|\varphi u\|_\sigma + \sum_{j=1}^m \|X_j \varphi u\| \leq C_0(\|P\varphi u\|' + \|\varphi u\|),$$

taking  $\Omega_0 = \Omega_1$  and  $C_0$  as in (2-7) related to  $\bar{\Omega}_0 \Subset \Omega$ , we have

$$\|P\varphi u\|' \leq \|\varphi P u\|' + \|[P, \varphi]u\|', \quad (3-18)$$

with

$$\begin{aligned} \|[P, \varphi]u\|' &\leq 2 \sum_{j=1}^m (\|X_j [X_j, \varphi]u\|' + \|X_j^2(\varphi)u\|') + \|Y(\varphi)u\|' \\ &\leq 2 \sum_{j=1}^m (\|X_j(\varphi)u\| + \|X_j^2(\varphi)u\|) + \|Y(\varphi)u\|, \quad \text{from (2-3), (2-6)}. \end{aligned}$$

Now as the  $X_j, Y$  are smooth in  $\Omega$  and  $\bar{\Omega}_0 \subset \Omega$ ,

$$\|X_j(\varphi)u\| \leq C_1 \sum_{|\beta| \leq 1} \|\varphi^{(\beta)}u\|, \quad \|X_j^2(\varphi)u\| \leq C_1 \sum_{|\beta| \leq 2} \|\varphi^{(\beta)}u\|. \quad (3-19)$$

So, with a suitable  $C$ , (3-17) is obtained from (3-18), (3-19).  $\square$

In order to reach estimates for ordinary derivatives, one way is to try to obtain estimates for  $\varphi u$  in the Sobolev spaces  $H^{\ell\sigma}$ ,  $\ell = 1, \dots, p$ , (3-17) corresponding to  $\ell = 1$ . As the basic estimate (2-6) is with  $H^\sigma$  (i.e.,  $\ell = 1$ ), it is natural to use the following, recalling some notation and definitions in [Derridj  $\geq$  2019, Section 2]: let  $\Omega_0$ , be such that  $\bar{\Omega}_1 \subset \Omega_0 \subset \bar{\Omega}_0 \Subset \Omega$ . So we consider  $\psi \in \mathcal{D}(\Omega_0)$ ,  $\psi = 1$  on  $\bar{\Omega}_1$ ; one way to estimate  $\|v\|_{\ell\sigma}$ , knowing (2-7) is to consider  $\psi T_{\ell\sigma} v$ ,  $v \in \mathcal{D}(\Omega_1)$ ; in fact, for all  $v \in \mathcal{D}(\Omega_1)$ ,

$$\|v\|_{(\ell+1)\sigma} = \|\psi v\|_{(\ell+1)\sigma} = \|T_{(\ell+1)\sigma} \psi v\| = \|T_{\ell\sigma} \psi v\|_\sigma,$$

(see [Derridj  $\geq$  2019, 2.11-2.13]),

where  $\widehat{T_\rho w}(\xi) = (1 + \|\xi\|^2)^{\rho/2} \widehat{w}(\xi)$ , for all  $w \in \mathcal{D}(\mathbb{R}^n)$ . Hence

$$\|v\|_{(\ell+1)\sigma} \leq \|[T_{\ell\sigma}, \psi]v\|_\sigma + \|\psi T_{\ell\sigma} v\|_\sigma \leq C_2 \|v\|_{(\ell+1)\sigma-1} + \|\psi T_{\ell\sigma} v\|_\sigma.$$

Now for  $\ell = 1, \dots, p-1$ ,  $(\ell+1)\sigma - 1 \leq 0$ . So

$$\|v\|_{(\ell+1)\sigma} \leq \tilde{C} \|v\| + \|\psi T_{\ell\sigma} v\|_\sigma, \quad \text{for all } v \in \mathcal{D}(\Omega_1). \quad (3-20)$$

So this boils down to applying (2-7) to  $\psi T_{\ell\sigma} v \in \mathcal{D}(\Omega_0)$  using the constant  $C_0$ . Then we have

$$\|v\|_{(\ell+1)\sigma} \leq \tilde{C} (\|v\| + \|\psi T_{\ell\sigma} v\|_\sigma) \leq \tilde{C} \left( \|\psi T_{\ell\sigma} v\| + \sum_j \|X_j \psi T_{\ell\sigma} v\| + \|v\| \right).$$

So, we get:

$$\|\varphi u\|_{(\ell+1)\sigma} \leq C_0 \tilde{C} (\|P\psi T_{\ell\sigma}\varphi u\|' + \|\psi T_{\ell\sigma}\varphi u\| + \|\varphi u\|),$$

yielding

$$\|\varphi u\|_{(\ell+1)\sigma} \leq C_3 (\|P\psi T_{\ell\sigma}\varphi u\|' + \|\varphi u\|_{\ell\sigma}),$$

for all  $(u, \varphi) \in C^\infty(\bar{\Omega}_1) \times \mathcal{D}(\Omega_1)$ . (3-21)

Now we provide a suitable bound for  $\|P\psi T_{\ell\sigma}\varphi u\|'$  as follows:

$$\begin{aligned} & \|P\psi T_{\ell\sigma}\varphi u\|' \\ & \leq \| [P, \psi T_{\ell\sigma}] \varphi u \|' + \| \psi T_{\ell\sigma} [P, \varphi] u \|' + \| \psi T_{\ell\sigma} \varphi P u \|' \\ & \leq \sum_{j=1}^m (2 \| X_j [X_j, \psi T_{\ell\sigma}] \varphi u \|' + \| [X_j, [X_j, \psi T_{\ell\sigma}]] \varphi u \|) + \| [Y + b, \psi T_{\ell\sigma}] \varphi u \| \\ & \quad + \sum_{j=1}^m (2 \| \psi T_{\ell\sigma} X_j [X_j, \varphi] u \|' + \| \psi T_{\ell\sigma} [X_j [X_j, \varphi]] u \|) \\ & \quad \quad \quad + \| \psi T_{\ell\sigma} [Y, \varphi] u \|' + \| \psi T_{\ell\sigma} \varphi P u \|'. \end{aligned}$$

Using again inequalities in (2-3), (2-6), we get, with some constant  $C$  which may vary from line to another,

$$\begin{aligned} & \|P\psi T_{\ell\sigma}\varphi u\|' \\ & \leq C \left( \|\varphi u\|_{\ell\sigma} + \sum_{j=1}^m \left( 2 \| [\psi T_{\ell\sigma}, X_j] [X_j, \varphi] u \|' + \| X_j \psi T_{\ell\sigma} [X_j, \varphi] u \|' \right. \right. \\ & \quad \quad \quad \left. \left. + \| X_j^2(\varphi) u \|_{\ell\sigma} \right) + \| Y(\varphi) u \|_{\ell\sigma} \right) + \| \psi T_{\ell\sigma} \varphi P u \|' \\ & \leq C \left( \| \psi T_{\ell\sigma} \varphi P u \|' + \sum_{j=1}^m \left( \| X_j(\varphi) u \|_{\ell\sigma} + \| X_j^2(\varphi) u \|_{\ell\sigma} + \| \varphi u \|_{\ell\sigma} \right. \right. \\ & \quad \quad \quad \left. \left. + \| Y(\varphi) u \|_{\ell\sigma} \right) \right). \end{aligned} \quad (3-22)$$

Hence, in fact, we proved the following with  $\Omega_0$  as above:

**Proposition 3.6.** *Under the hypotheses in Proposition 3.5, there exists a constant  $C = C(\Omega_0, P) > 0$  such that*

$$\|\varphi u\|_{(\ell+1)\sigma} \leq C \left( \|\varphi P u\|_{\ell\sigma} + \sum_{|\beta| \leq 2} \|\varphi^{(\beta)} u\|_{\ell\sigma} \right), \quad \ell = 0, \dots, p-1. \quad (3-23)$$

The next step is to obtain such estimates for couples  $(\partial^\alpha u, \varphi)$  for  $(u, \varphi) \in C^\infty(\bar{\Omega}_1) \times \mathcal{D}(\Omega_1)$ ,  $\alpha \in \mathbb{N}^n$ . In order to get such an estimate, let us first consider the case  $\ell = 0$ . We use (3-17). So

$$\|\varphi \partial^\alpha u\|_\sigma \leq C \|\varphi P \partial^\alpha u\|' + \sum_{|\beta| \leq 2} \|\varphi^{(\beta)} \partial^\alpha u\|.$$

Here, what is new is to use  $\varphi P \partial^\alpha u = \varphi [P, \partial^\alpha] u + \varphi \partial^\alpha P u$ :

$$\begin{aligned} \|\varphi P \partial^\alpha u\|' &\leq \|\varphi [P, \partial^\alpha] u\|' + \|\varphi \partial^\alpha P u\|' \\ &\leq \sum_{j=1}^m (2\|\varphi X_j [X_j, \partial^\alpha] u\|' + \|\varphi [X_j, [X_j, \partial^\alpha]] u\|') \\ &\quad + \|\varphi [Y + b, \partial^\alpha] u\|' + \|\varphi \partial^\alpha P u\|'. \end{aligned} \tag{3-24}$$

Writing  $\varphi X_j [X_j, \partial^\alpha] u = [\varphi, X_j] [X_j, \partial^\alpha] u + X_j \varphi [X_j, \partial^\alpha] u$  and using again (2-3) and (2-6), we get

$$\begin{aligned} \|\varphi P \partial^\alpha u\| &\leq \sum_{j=1}^m (2\|X_j(\varphi) [X_j, \partial^\alpha] u\| + \|\varphi [X_j, [X_j, \partial^\alpha]] u\|) \\ &\quad + \|\varphi [Y + b, \partial^\alpha] u\| + \|\varphi \partial^\alpha P u\|, \end{aligned}$$

and then we have to use expressions in (3-9) and (3-12). As we have to do that in order to bound  $\|\varphi \partial^\alpha u\|_{(\ell+1)\sigma}$ , we will write the step after the use of (3-9), (3-12) just in this general case; replacing  $u$  by  $\partial^\alpha u$  in (3-22), we get, with some constant  $C > 0$ , which may vary from line to line,

$$\|\varphi \partial^\alpha u\|_{(\ell+1)\sigma} \leq C \left( \|\psi T_{\ell\sigma} \varphi P \partial^\alpha u\|' + \sum_{|\beta| \leq 2} \|\varphi^{(\beta)} \partial^\alpha u\|_{\ell\sigma} \right). \tag{3-25}$$

Now, as we did above with (3-24), we get

$$\begin{aligned} \|\varphi \partial^\alpha u\|_{(\ell+1)\sigma} &\leq C \left( \|\psi T_{\ell\sigma} \varphi \partial^\alpha P u\|' + \sum_{j=1}^m (\|X_j(\varphi) [X_j, \partial^\alpha] u\|_{\ell\sigma} + \|\varphi [X_j, [X_j, \partial^\alpha]] u\|_{\ell\sigma}) \right. \\ &\quad \left. + \|\varphi [Y + b, \partial^\alpha] u\|_{\ell\sigma} + \sum_{|\beta| \leq 2} \|\varphi^{(\beta)} \partial^\alpha u\|_{\ell\sigma} \right). \end{aligned}$$

As  $\ell\sigma \leq 1$ , ( $\ell = 0, \dots, p - 1$ ), we finally obtain:

**Proposition 3.7.** *There exists a constant  $C > 0$  such that*

$$\begin{aligned} &\text{for all } (u, \varphi) \in C^\infty(\bar{\Omega}_1) \times \mathcal{D}(\Omega_1), \quad \text{and all } \alpha \in \mathbb{N}^n, \\ \|\varphi \partial^\alpha u\|_{(\ell+1)\sigma} &\leq C \left( \|\varphi \partial^\alpha P u\|_{\ell\sigma} + \sum_{|\beta| \leq 2} \|\varphi^{(\beta)} \partial^\alpha u\|_{\ell\sigma} \right. \\ &\quad \left. + \sum_{j=1}^m \left( \sum_{|\beta|=1} \|\varphi^{(\beta)} [X_j, \partial^\alpha] u\|_{\ell\sigma} + \|\varphi [X_j, [X_j, \partial^\alpha]] u\|_{\ell\sigma} \right) \right. \\ &\quad \left. + \|\varphi [Y + b, \partial^\alpha] u\|_{\ell\sigma} \right). \end{aligned} \tag{3-26}$$

In what follows,  $s$  is given and  $s \geq 1$ .

#### 4. Local relations of domination by powers of $P$

We need to introduce, in this section, further notation:

For  $\epsilon > 0$ ,  $j \in \mathbb{N}$ ,  $\gamma \in \mathbb{N}^n$ , and  $(u, \varphi) \in C^\infty(\bar{\Omega}_1) \times \mathcal{D}(\Omega_1)$ , we set

$$N_{j,\gamma}^\epsilon(u, \varphi) = \epsilon^{|\gamma|+2j} |\gamma|!^{-s} (2j)!^{-s} \|\varphi^{(\gamma)} P^j u\|. \quad (4-1)$$

Once  $\epsilon$  is fixed, we often delete  $\epsilon$  and write  $N_{j,\gamma}^\epsilon = N_{j,\gamma}$ .

Before stating our main theorem, we give a simple useful lemma, which we will apply many times.

**Lemma 4.1.** *Let  $(k, \beta) \in \mathbb{N} \times \mathbb{N}^n$ , and  $\rho \in \mathbb{N}$ . Then*

$$\rho!^s N_{j,\gamma}^\epsilon(P^k u, \varphi^{(\beta)}) \leq \epsilon^{-(|\beta|+2k)} (\rho + |\beta| + 2k)!^s N_{j+k,\gamma+\beta}^\epsilon(u, \varphi),$$

if  $|\gamma| + 2j \leq \rho$ . (4-2)

*Proof.* From the definition in (4-1), we see that

$$\begin{aligned} N_{j,\gamma}^\epsilon(P^k u, \varphi^{(\beta)}) \\ = \epsilon^{-(|\beta|+2k)} (|\gamma| + 1)^s \cdots (|\gamma| + |\beta|)^s ((2j + 1) \cdots (2j + 2k))^s N_{j+k,\gamma+\beta}(u, \varphi). \end{aligned}$$

Then, observe that, for  $|\gamma| + 2j \leq \rho$ ,

$$\rho! (|\gamma| + 1) \cdots (|\gamma| + |\beta|) (2j + 1) \cdots (2j + k) \leq (\rho + |\beta| + 2k)!. \quad (4-3)$$

The proof is then finished by taking (4-3) to the power  $s \geq 1$ . □

**Theorem 4.2.** *Let  $s \geq 1$  be given. Assume that the coefficients of  $P$  are in  $G^s(\Omega)$  and properties (2-4) and (3-16) hold on  $\Omega_0$ ,  $\bar{\Omega}_0 \subset \Omega$ . Let  $\Omega_1$  be an open set with  $\bar{\Omega}_1 \subset \Omega_0$ . For every  $0 < \epsilon \leq 1$ , there exists  $M = M(\epsilon, \Omega_0, P) \geq 1$  such that for all  $\alpha \in \mathbb{N}^n$ ,  $\ell = 0, \dots, p$ ,  $(u, \varphi) \in C^\infty(\bar{\Omega}_1) \times \mathcal{D}(\Omega_1)$ ,*

$$\|\varphi \partial^\alpha u\|_{\ell\sigma} \leq M^{2p|\alpha|+\ell+1} (2(p|\alpha| + \ell))!^s \sum_{|\beta|+2j \leq 2(p|\alpha|+\ell)} N_{j,\beta}(u, \varphi). \quad (4-4)$$

*Proof.* It consists of a double induction on  $|\alpha| = r$  and on  $\ell$ . More precisely, in a first step, we prove the estimates (4-2) for  $\alpha = 0$ . In all the proof, we will specify (4-4) $_{\alpha,\ell}$  for (4-4) when we consider the couple  $(\alpha, \ell) \in \mathbb{N} \times \{0, \dots, p\}$ . So we want, in this first step, to prove (4-4) $_{0,\ell}$ ,  $\ell \in \{0, \dots, p\}$ .

(A) Proof of (4-4) $_{0,\ell}$ ,  $\ell \in \{0, \dots, p\}$ : As (4-4) $_{0,0}$  is trivial, all we have to do is to make an induction on  $\ell$ . So assume that (4-4) $_{0,i}$  is true for  $i \leq \ell$ ,  $\ell \in \{0, \dots, p-1\}$ . Then we want to prove (4-4) $_{0,\ell+1}$ . For that we use (3-23) in Proposition 3.6 for  $(u, \varphi) \in C^\infty(\bar{\Omega}_1) \times \mathcal{D}(\Omega_1)$ . So in order to bound  $\|\varphi u\|_{(\ell+1)\sigma}$ , we just have to suitably bound  $\|\varphi P u\|_{\ell\sigma}$  and  $\sum_{|\beta| \leq 2} \|\varphi^{(\beta)} u\|$ . Hence this reduces to applying (4-4) $_{0,\ell}$

respectively to the couples  $(Pu, \varphi)$  and  $(u, \varphi^{(\beta)})$  in  $C^\infty(\bar{\Omega}_1) \times \mathcal{D}(\Omega_1)$ . So

$$\|\varphi Pu\|_{\ell\sigma} \leq M^{\ell+1}(2\ell)!^s \sum_{|\beta|+2j \leq 2\ell} N_{j,\beta}(Pu, \varphi), \quad (u, \varphi) \in C^\infty(\bar{\Omega}_1) \times \mathcal{D}(\Omega_1). \tag{4-5}$$

Now we apply [Lemma 4.1](#) with  $\rho = 2\ell$  in (4-2). Then

$$\begin{aligned} \|\varphi Pu\|_{\ell\sigma} &\leq \epsilon^{-2}(2(\ell+1))!^s M^{\ell+1} \sum_{|\beta|+2j \leq 2(\ell+1)} N_{j,\beta}(u, \varphi) \\ &\leq \epsilon^{-2} M^{-1} M^{(\ell+1)+1} (2(\ell+1))!^s \sum_{|\beta|+2j \leq 2(\ell+1)} N_{j,\beta}(u, \varphi). \end{aligned}$$

We do exactly the same for  $\|\varphi^{(\beta)}u\|_{\ell\sigma}$ , as  $|\beta| \leq 2$ :

$$\|\varphi^{(\beta)}u\|_{\ell\sigma} \leq \epsilon^{-2} M^{-1} M^{(\ell+1)+1} (2(\ell+1))!^s \sum_{|\gamma|+2j \leq 2(\ell+1)} N_{j,\gamma}(u, \varphi). \tag{4-6}$$

So from (3-23), (4-5) and (4-6), we get

$$\begin{aligned} \|\varphi u\|_{(\ell+1)\sigma} \\ \leq C(2+n+n^2)\epsilon^{-2} M^{-1} M^{(\ell+1)+1} (2(\ell+1))!^s \sum_{|\beta|+2j \leq 2(\ell+1)} N_{j,\beta}(u, \varphi). \end{aligned} \tag{4-7}$$

So we see that under the condition

$$C(2+n+n^2)\epsilon^{-2} M^{-1} \leq 1 \quad (\text{equivalently, } M \geq C(2+n+n^2)\epsilon^{-2}), \tag{4-8}$$

[Equation \(4-4\)<sub>0,\ell+1</sub>](#) is satisfied for all  $(u, \varphi) \in C^\infty(\bar{\Omega}_1) \times \mathcal{D}(\Omega_1)$ .

(B) Proof of [\(4-4\)<sub>\alpha,\ell</sub>](#) for all  $(\alpha, \ell) \in \mathbb{N}^n \times \{0, \dots, p\}$ : All we have to do, as [\(4-4\)<sub>0,\ell</sub>](#) is true for  $\ell = 0, \dots, p$ , is to make an induction on  $|\alpha| = r$ . More precisely, if [\(4-4\)<sub>\alpha,\ell</sub>](#) is true for  $|\alpha| \leq r, \ell \in \{0, \dots, p\}$ , then it is true for  $|\alpha| = r+1, \ell = 0, \dots, p$ . We use the same kind of proof as before: given  $\alpha + i$ , with  $|\alpha| = r, i \in \{1, \dots, n\}$ , we want to suitably bound  $\|\varphi \partial_i \partial^\alpha u\|_{\ell\sigma}$  for  $\ell \in \{0, \dots, p\}$ .

(1)  $\ell = 0$ : We use  $\|\varphi \partial_i \partial^\alpha u\| \leq \|\partial_i(\varphi) \partial^\alpha u\| + \|\varphi \partial^\alpha u\|_1$ . Hence we just have to apply [\(4-4\)<sub>\alpha,0</sub>](#) with  $(u, \varphi^{(i)})$  and [\(4-4\)<sub>\alpha,p</sub>](#) with  $(u, \varphi)$ . We will obtain, directly, or applying also [Lemma 4.1](#),

$$\begin{aligned} \|\varphi^{(i)} \partial^\alpha u\| \\ \leq \epsilon^{-1} M^{-2p} M^{2p(|\alpha|+1)+1} (2p(|\alpha|+1))!^s \sum_{|\beta|+2j \leq 2p(|\alpha|+1)} N_{j,\beta}(u, \varphi), \end{aligned} \tag{4-9}$$

$$\|\varphi \partial^\alpha u\|_{p\sigma} \leq M^{-p} M^{2p(|\alpha|+1)+1} (2p(|\alpha|+1))!^s \sum_{|\beta|+2j \leq 2p(|\alpha|+1)} N_{j,\beta}(u, \varphi). \tag{4-10}$$

So, summing (4-9) and (4-10), we get that if

$$M^{-p}(1 + \epsilon^{-1}M^{-p}) \leq 1, \quad (4-11)$$

then (4-4) $_{\alpha+i,0}$  is satisfied, for  $i = 1, \dots, n$ .

(2) Proof of (4-4) $_{\alpha,\ell}$ , for  $|\alpha| = r + 1$ ,  $\ell > 0$ , i.e.,  $\ell \in \{1, \dots, p\}$ : So assuming that (4-4) $_{\alpha,\ell}$  is true for  $|\alpha| = r$ ,  $\ell = 0, \dots, p$ , and (4-4) $_{\alpha,\rho}$  is true for  $|\alpha| = r + 1$ ,  $\rho \in \{1, \dots, \ell\}$ , then, if  $\ell < p$ , we want to prove that (4-4) $_{\alpha,\ell+1}$  is true,  $|\alpha| = r + 1$ . Now we have to use (3-26) in Proposition 3.7. So in order to suitably bound  $\|\varphi \partial^\alpha u\|_{(\ell+1)\sigma}$ , we are led to bound  $\|\varphi \partial^\alpha Pu\|_{\ell\sigma}$ ,  $\|\varphi^{(\beta)} \partial^\alpha u\|_{\ell\sigma}$ ,  $|\beta| \leq 2$ , but also much more terms like simple brackets of  $X_j$ 's and  $Y$  with  $\partial^\alpha$  and double brackets of  $X_j$ 's with  $\partial^\alpha$ .

The proof will follow the lines of our proof for  $\alpha = 0$ , but here with more terms, and some are more difficult to handle than others, namely those coming from the brackets of the  $X_j$ 's with  $\partial^\alpha$  (simple and double brackets). The term  $\|\varphi[Y + b, \partial^\alpha]u\|_{\ell\sigma}$  is similar to the terms  $\|\varphi^{(\beta)}[X_j, \partial^\alpha]u\|$ ,  $\beta = 0$ .

(a) A bound on  $\|\varphi \partial^\alpha Pu\|_{\ell\sigma}$ : We apply (4-4) $_{\alpha,\ell}$  for the couple  $(Pu, \varphi)$ :

$$\|\varphi \partial^\alpha Pu\|_{\ell\sigma} \leq M^{2p|\alpha|+\ell+1}(2(p|\alpha| + \ell))!^s \sum_{|\beta|+2j \leq 2(p|\alpha|+\ell)} N_{j,\beta}(Pu, \varphi).$$

Applying Lemma 4.1, we get (here  $\rho = 2(p|\alpha| + \ell)$ )

$$\begin{aligned} & \|\varphi \partial^\alpha Pu\|_{\ell\sigma} \\ & \leq \epsilon^{-2} M^{2p|\alpha|+\ell+1}(2(p|\alpha| + \ell + 1))!^s \sum_{|\beta|+2j \leq 2(p|\alpha|+\ell+1)} N_{j,\beta}(u, \varphi) \\ & \leq (\epsilon^{-2} M^{-1}) M^{2p|\alpha|+(\ell+1)+1}(2(p|\alpha| + \ell + 1))!^s \\ & \quad \cdot \sum_{|\beta|+2j \leq 2(p|\alpha|+\ell+1)} N_{j,\beta}(u, \varphi). \end{aligned} \quad (4-12)$$

(b) A bound on  $\|\varphi^{(\beta)} \partial^\alpha u\|_{\ell\sigma}$ ,  $|\beta| \leq 2$ : We apply (4-4) $_{\alpha,\ell}$  to the couple  $(u, \varphi^{(\beta)})$ :

$$\|\varphi^{(\beta)} \partial^\alpha u\|_{\ell\sigma} M^{2p|\alpha|+\ell+1}(2(p|\alpha| + \ell))!^s \sum_{|\beta|+2j \leq 2(p|\alpha|+\ell)} N_{j,\beta}(u, \varphi^{(\beta)}).$$

Applying again Lemma 4.1 (as  $|\beta| \leq 2$ ), we get

$$\begin{aligned} & \|\varphi^{(\beta)} \partial^\alpha u\|_{\ell\sigma} \\ & \leq \epsilon^{-2} M^{2p|\alpha|+\ell+1}(2(p|\alpha| + \ell + 1))!^s \sum_{|\beta|+2j \leq 2(p|\alpha|+\ell+1)} N_{j,\beta}(u, \varphi) \\ & \leq \epsilon^{-2} M^{-1} M^{2p|\alpha|+(\ell+1)+1}(2(p|\alpha| + \ell + 1))!^s \\ & \quad \cdot \sum_{|\beta|+2j \leq 2(p|\alpha|+\ell+1)} N_{j,\beta}(u, \varphi). \end{aligned} \quad (4-13)$$



(c) A bound on  $\|\varphi^{(\beta)}[X_j, \partial^\alpha]u\|_{\ell\sigma}$ ,  $|\beta| \leq 1$ ,  $j = 1, \dots, m$ ,  $\|\varphi[Y + b, \partial^\alpha]u\|_{\ell\sigma}$ : (It is easy to see that the term  $\|\varphi[Y + b, \partial^\alpha]u\|_{\ell\sigma}$  is much simpler than the others, since it corresponds to  $\beta = 0$ .)

From expressions in (3-9) where we delete  $j \in \{1, \dots, m\}$ :

$$\|\varphi^{(\beta)}[X, \partial^\alpha]u\|_{\ell\sigma} \leq \sum_{\substack{\gamma < \alpha \\ i=1, \dots, n}} \|a_{\alpha\gamma i} \varphi^{(\beta)} \partial^{\gamma+i} u\|_{\ell\sigma}. \tag{4-14}$$

Then applying estimates in (3-15) (recall that  $j$  is deleted):

$$\|\varphi^{(\beta)}[X, \partial^\alpha]u\|_{\ell\sigma} \leq \sum_{\substack{\gamma < \alpha \\ i=1, \dots, n}} B^{|\alpha-\gamma|} \left(\frac{\alpha!}{(\gamma+i)!}\right)^s \|\varphi^{(\beta)} \partial^{\gamma+i} u\|_{\ell\sigma}. \tag{4-15}$$

As  $|\gamma + i| \leq |\alpha| = r + 1$ , we can apply (4-4) $_{\gamma+i, \ell}$  to  $(u, \varphi^{(\beta)})$ . So we get

$$\begin{aligned} &\|\varphi^{(\beta)}[X, \partial^\alpha]u\|_{\ell\sigma} \\ &\leq n \sum_{\substack{\gamma < \alpha \\ i=1, \dots, n}} \left( B^{|\alpha-\gamma|} M^{2p(|\gamma|+1)+\ell+1} (2(p(|\gamma|+1)+\ell))!^s \left(\frac{\alpha!}{\gamma!}\right)^s \right. \\ &\qquad \left. \cdot \sum_{|\delta|+2j \leq 2(p(|\gamma|+1)+\ell)} N_{j,\delta}(u, \varphi^{(\beta)}) \right). \end{aligned} \tag{4-16}$$

Using (4-2) in Lemma 4.1, with  $\rho = 2(p(|\gamma|+1)+\ell)$ , we find

$$(2(p(|\gamma|+1)+\ell))!^s N_{j,\delta}(u, \varphi^{(\beta)}) \leq (2(p(|\gamma|+1)+\ell)+1)!^s N_{j,\delta+\beta}(u, \varphi), \tag{4-17}$$

and hence

$$\begin{aligned} &(2(p(|\gamma|+1)+\ell))!^s \left(\frac{\alpha!}{\gamma!}\right)^s N_{j,\delta}(u, \varphi^{(\beta)}) \\ &\leq (2(p(|\alpha|+\ell+1))!^s N_{j,\delta+\beta}(u, \varphi), \end{aligned} \tag{4-18}$$

for all  $(j, \delta)$  such that  $|\delta| + 2j \leq 2(p(|\gamma|+1)+\ell)+1$ . So, coming back to the second member in (4-16),

$$\begin{aligned} &\|\varphi^{(\beta)}[X, \partial^\alpha]u\|_{\ell\sigma} \\ &\leq n \sum_{\gamma < \alpha} \left( B^{|\alpha-\gamma|} M^{2p(|\gamma|+1)+\ell+1} (2(p|\alpha|+\ell+1))!^s \right. \\ &\qquad \left. \cdot \epsilon^{-1} \sum_{|\delta|+2j \leq 2(p|\alpha|+\ell+1)} N_{j,\delta}(u, \varphi) \right), \end{aligned} \tag{4-19}$$

or

$$\begin{aligned} \|\varphi^{(\beta)}[X, \partial^\alpha]u\|_{\ell\sigma} &\leq n(2(p|\alpha| + \ell + 1))!^s \sum_{|\delta|+2j \leq 2(p|\alpha|+\ell+1)} N_{j,\delta}(u, \varphi) \\ &\quad \cdot \epsilon^{-1} \sum_{\gamma < \alpha} B^{|\alpha-\gamma|} M^{2p(|\gamma|+1)+\ell+1}. \end{aligned} \quad (4-20)$$

The last sum is bounded as follows:

$$\sum_{\gamma < \alpha} B^{|\alpha-\beta|} M^{2p(|\gamma|+1)+\ell+1} = BM^{2p|\alpha|+\ell+1} \sum_{\gamma < \alpha} \left(\frac{B}{M^{2p}}\right)^{|\alpha-\gamma|-1}. \quad (4-21)$$

Now we have the following lemma:

**Lemma 4.3.** *There exists  $\theta_0 > 0$ , independent of  $\alpha$ , such that*

$$\sum_{\gamma < \alpha} \lambda^{|\alpha-\gamma|-1} \leq n + 1, \quad \text{if } 0 \leq \lambda < \theta_0. \quad (4-22)$$

The proof of [Lemma 4.3](#) is similar to that of [Lemma 3.2](#) after noticing that  $\sum_{\gamma < \alpha, |\alpha-\gamma|=1} \lambda^{|\alpha-\gamma|-1} = n$ .

**Remark 4.4.** [Lemma 4.3](#) is not true when one works with the sum  $\sum_{|\gamma| < |\alpha|} \lambda^{|\alpha-\gamma|-1}$  as the sum  $\sum_{|\gamma| < |\alpha|, |\alpha-\gamma|=1} 1$  is not bounded by a constant independent of  $\alpha$ .

Applying [\(4-22\)](#) we see that under the condition

$$M^{2p} \geq \theta_0^{-1} B, \quad (4-23)$$

we obtain from [\(4-20\)](#), [\(4-21\)](#) and [\(4-22\)](#)

$$\begin{aligned} \|\varphi^{(\beta)}[X, \partial^\alpha]u\|_{\ell\sigma} &\leq n(n+1)\epsilon^{-1} BM^{-1} M^{2p|\alpha|+(\ell+1)+1} (2p|\alpha| + \ell + 1)!^s \\ &\quad \cdot \sum_{|\delta|+2j \leq 2(p|\alpha|+\ell+1)} N_{j,\delta}(u, \varphi). \end{aligned} \quad (4-24)$$

(d) A bound on  $\|\varphi[X_j, [X_j, \partial^\alpha]]u\|_{\ell\sigma}$ ,  $j = 1, \dots, m$ : As we did above we delete the index  $j$  and write  $\|\varphi[X, [X, \partial^\alpha]]u\|$ . Of course, we also delete  $j$  in the coefficients of the bracket. Looking at [\(3-12\)](#), we have two kinds of terms to study:

(i) A bound on  $\sum_{\beta < \alpha, k=1, \dots, n} \|\varphi d_{k\alpha\beta} \partial^{\beta+k} u\|_{\ell\sigma} = E_1$ : Using [Equation \(3-15\)](#) in [Proposition 3.3](#), as  $\ell\sigma \leq 1$ , we have

$$\begin{aligned} E_1 &\leq \sum_{\substack{\beta < \gamma \\ k=1, \dots, n}} B^{|\alpha-\beta|} \left(\frac{\alpha!}{\beta!} (|\alpha| + 1)\right)^s M^{2\beta(|\beta|+1)+\ell+1} (2(p(|\beta| + 1) + \ell))!^s \\ &\quad \cdot \sum_{|\gamma|+2j \leq 2(p(|\beta|+1)+\ell)} N_{j,\gamma}(u, \varphi). \end{aligned} \quad (4-25)$$

We want to remark here that there is a factor  $(|\alpha| + 1)^s$  in (4-25), but it is compensated by the fact that one has  $\varphi$ , not  $\varphi^{(\beta)}$ ,  $|\beta| = 1$ . Precisely, we have

$$E_1 \leq n \sum_{\beta < \alpha} B^{|\alpha - \beta|} \left( \frac{(|\alpha| + 1)!}{|\beta|!} \right)^s (2p(|\beta| + 1) + \ell)!^s M^{2p(|\beta| + 1) + \ell + 1} \cdot \sum_{|\gamma| + 2j \leq 2(p|\alpha| + \ell)} N_{j, \gamma}(u, \varphi), \quad \text{as } |\beta| + 1 \leq |\alpha|. \quad (4-26)$$

Now we have the following inequality:

$$\left( \frac{(|\alpha| + 1)!}{|\beta|!} \right)^s (2(p(|\beta| + 1) + \ell))!^s \leq (2(p|\alpha| + \ell + 1))!^s, \quad |\beta| + 1 \leq |\alpha|. \quad (4-27)$$

So, from (4-26) and (4-27), we get

$$E_1 \leq n(2(p|\alpha| + \ell + 1))!^s \sum_{\beta < \alpha} B^{|\alpha - \beta|} M^{2p(|\beta| + 1) + \ell + 1} \cdot \sum_{|\gamma| + 2j \leq 2(p|\alpha| + \ell)} N_{j, \gamma}(u, \varphi). \quad (4-28)$$

The sum in the second member in (4-28) is the same than the sum in (4-21), replacing  $\gamma$  by  $\beta$ . So from (4-22) in Lemma 4.3, we get, under condition (4-23),

$$E_1 \leq n(n + 1)BM^{-1} M^{2p|\alpha| + (\ell + 1) + 1} (2p|\alpha| + \ell + 1)!^s \cdot \sum_{|\gamma| + 2j \leq 2(p|\alpha| + \ell)} N_{j, \gamma}(u, \varphi). \quad (4-29)$$

(ii) A bound on  $\sum_{\beta < \alpha, i, k=1, \dots, n} \|\varphi c_{ik\alpha\beta} \partial^{\beta+i+k} u\|_{\ell\sigma} = E_2$ : We write the proof, for completeness, noticing however that  $\partial^{\beta+i+k}$  is of the form  $\partial^{\beta+I}$  with  $|I| = 2$ . Using estimates in (3-15), we get

$$E_2 \leq \sum_{\substack{\beta < \alpha \\ i, k=1, \dots, n}} B^{|\alpha - \beta|} \left( \frac{(\alpha + k)!}{(\beta + k)!} \right)^s \|\varphi \partial^{\beta+i+k} u\|_{\ell\sigma}. \quad (4-30)$$

Hence,

$$E_2 \leq \sum_{\substack{\beta < \alpha, |\beta| \leq |\alpha| - 2 \\ i, k=1, \dots, n}} B^{|\alpha - \beta|} M^{2p(|\beta| + 2) + \ell + 1} \left( \frac{(\alpha + k)!}{(\beta + k)!} \right)^s (2(p(|\beta| + 2) + \ell))!^s \cdot \sum_{|\gamma| + 2j \leq 2(p(|\beta| + 2) + \ell)} N_{j, \gamma}(u, \varphi). \quad (4-31)$$

Now we have the following:

$$\left( \frac{(\alpha + k)!}{(\beta + k)!} \right)^s (2(p(|\beta| + 2) + \ell))!^s \leq (2(p|\alpha| + \ell + 1))!^s. \quad (4-32)$$

(4-32) is a consequence of  $(\alpha + k)!/(\beta + k)! \leq (|\alpha| + 1)!/(|\beta| + 1)!$ . Hence

$$E_2 \leq n^2(2(p|\alpha| + \ell + 1))!^s \left( \sum_{\substack{\beta < \alpha \\ |\beta| \leq |\alpha| - 2}} B^{|\alpha - \beta|} M^{2p(|\beta| + 2) + \ell + 1} \right) \cdot \sum_{|\gamma| + 2j \leq 2(p|\alpha| + \ell + 1)} N_{j,\gamma}(u, \varphi), \quad (4-33)$$

as  $\text{Card}\{(i, k) \in \{1, \dots, n\}\} = n^2$ . Now, we bound

$$F = \sum_{\beta < \alpha, |\beta| \leq |\alpha| - 2} B^{|\alpha - \beta|} M^{2p(|\beta| + 2) + \ell + 1}$$

with

$$F \leq \sum_{\substack{\beta < \alpha \\ |\beta| \leq |\alpha| - 2}} \left( \frac{B}{M^{2p}} \right)^{|\alpha - \beta| - 2} M^{2p|\alpha| + (\ell + 1) + 1} B^2 M^{-1}. \quad (4-34)$$

Now using Lemma 4.3 and the fact that  $\sum_{\beta < \alpha, |\beta| \leq |\alpha| - 2} 1 \leq n^2$ , we obtain

$$E_2 \leq (n^2 + 1)^2 B^2 M^{-1} M^{2p|\alpha| + (\ell + 1) + 1} (2(p|\alpha| + \ell + 1))!^s \cdot \sum_{|\delta| + 2j \leq 2(p|\alpha| + \ell + 1)} N_{j,\delta}(u, \varphi). \quad (4-35)$$

We recall that we said in (c) that we do not consider the term

$$\|\varphi[Y + b, \partial^\alpha]\|_{\ell\sigma} \leq \|\varphi[Y, \partial^\alpha]u\|_{\ell\sigma} + \|\varphi[b, \partial^\alpha]u\|_{\ell\sigma}$$

as  $\|\varphi[Y, \partial^\alpha]u\|_{\ell\sigma}$  is like  $\|\varphi[X_j, \partial^\alpha]u\|_{\ell\sigma}$  and  $\|\varphi[b, \partial^\alpha]u\|_{\ell\sigma}$  much smaller. So in order to bound the term  $\|\varphi[X_j, [X_j, \partial^\alpha]]u\|_{\ell\sigma}$ , we have to collect (4-29) and (4-35), which are true under condition (4-23). We find

$$\|\varphi[X_j, [X_j, \partial^\alpha]]u\|_{\ell\sigma} \leq 2(n^2 + 1)^2 B^2 M^{-1} M^{2p|\alpha| + (\ell + 1) + 1} (2(p|\alpha| + \ell + 1))!^s \cdot \sum_{|\gamma| + 2j \leq 2(p|\alpha| + \ell)} N_{j,\gamma}(u, \varphi) \quad (4-36)$$

In order to give a bound for  $\|\varphi\partial^\alpha u\|_{(\ell+1)\sigma}$ , we have from (3-26) in Proposition 3.7 to take  $C$  times the bound in (a) plus  $C(n^2 + n + 1)$  times the bound in (b) plus  $Cm(n + 2)$  times the bound in (c) plus  $Cm$  times the bound in (d), under, of course, the conditions on  $M$  indicated in the proofs of (a), (b), (c) and (d).

Of course, we also have to take care of the conditions needed on  $M$  for the validity of these bounds. These may be summarized as follows:

- (a) (4-12) is satisfied, just under the induction hypothesis,
- (b) (4-13) is satisfied, under the induction hypothesis,
- (c) (4-24) is satisfied, under the induction hypothesis and (4-23),
- (d) (4-36) is satisfied, under the induction hypothesis and (4-23).

So adding all these estimates with the suitable factors yields

$$\begin{aligned} & \|\varphi \partial^\alpha u\|_{(\ell+1)\sigma} \\ & \leq CM^{-1}(2\epsilon^{-2} + n(n+1)\epsilon^{-1}B + 2(n^2 + 1)^2B^2) M^{2p|\alpha| + (\ell+1)+1} (2(p|\alpha| + \ell + 1))!^s \\ & \qquad \cdot \sum_{|\gamma|+2j \leq 2(p|\alpha| + \ell)} N_{j,\gamma}(u, \varphi) \end{aligned} \quad (4-37)$$

under the condition (4-23). From (4-37), we deduce that if  $M$  satisfies

$$M \geq \sup\{(\theta_0^{-1}B)^{1/2p}, 1, C(2\epsilon^{-2} + n(n+1)\epsilon^{-1}B + 2(n^2 + 1)^2B^2)\}, \quad (4-38)$$

then (4-4) $_{\alpha, \ell+1}$  is true. So (4-4) $_{\alpha, \ell}$  is true,  $|\alpha| = r + 1$ , for all  $\ell$ .

Now, let us finish the proof of the theorem. Since we proved (4-4) $_{0, \ell}$ ,  $\ell = 1, \dots, p$ , and the induction

$$\{(4-4)_{\alpha, \ell}, |\alpha| \leq r, \ell \in \{1, \dots, p\}\} \Rightarrow \{(4-4)_{\alpha, \ell}, |\alpha| = r + 1, \ell \in \{1, \dots, p\}\},$$

under respectively condition ((4-8) and (4-11)) and condition (4-38), we have  $M = M(\epsilon, \Omega_0, P) > 0$  so that the theorem is completely proved, when the conditions (4-8), (4-11) and (4-38) are satisfied. As  $\epsilon \leq 1$ , we see that  $M^p \geq 2\epsilon^{-1}$  implies that (4-11) holds. So, everything boils down to only the following condition:

$$\begin{aligned} M & \geq \sup\left\{(\theta_0^{-1}B)^{1/2p}, C(2 + n + n^2)\epsilon^{-2}, \right. \\ & \qquad \left. C(2\epsilon^{-2} + n(n+1)\epsilon^{-1}B + 2(n^2 + 1)^2B^2)\right\} \\ & = M(\epsilon, \theta_0, B). \end{aligned} \quad (4-39)$$

$B$  depends on  $P$  and  $\Omega_0$  and  $\theta_0$  depends on  $n$ . Hence  $M(\epsilon, \theta_0, B)$  can be written, as  $n$  is fixed,  $M(\epsilon, \Omega_0, P)$ . The proof of Theorem 4.2 is now complete.  $\square$

### 5. Gevrey regularity for Gevrey vectors

We want to give in this section an application of Theorem 4.2. In fact, we shall just use the estimates (4-4) for  $\ell = 0$ , which we rewrite here, for  $(u, \varphi) \in C^\infty(\bar{\Omega}_1) \times \mathcal{D}(\Omega_1)$ :

$$\|\varphi \partial^\alpha u\| \leq M_\epsilon^{2p|\alpha|+1} (2p|\alpha|)!^s \sum_{|\beta|+2j \leq 2p|\alpha|} N_{j,\beta}^\epsilon(u, \varphi). \quad (5-1)$$

Moreover, we want to state a theorem for operators of order  $m$ , satisfying estimates similar to (5-1), but with 2 replaced by  $m$  and with coefficients in  $G^s(\Omega)$ . For that purpose, we clarify the notation as  $m$  replaces 2.

Firstly,  $\Omega$  and  $\Omega_1$  are as in Section 4,  $\bar{\Omega}_1 \Subset \Omega$  and  $s \geq 1$ . Then define

$$N_{j,\beta}^\epsilon = \epsilon^{|\beta|+mj} |\beta|!^{-s} (m_j)!^{-s} \|\varphi^{(\beta)} P^j u\|, \quad (u, \varphi) \in C^\infty(\bar{\Omega}_1) \times \mathcal{D}(\Omega_1). \quad (5-2)$$

Now assume that  $P$  satisfies the following estimates:

$$\forall \epsilon, 0 < \epsilon \leq 1, \exists M_\epsilon \geq 1 \quad \text{such that} \quad \forall \alpha \in \mathbb{N}^n, \forall (u, \varphi) \in C^\infty(\bar{\Omega}_1) \times \mathcal{D}(\Omega_1)$$

$$\|\varphi \partial^\alpha u\| \leq M_\epsilon^{r|\alpha|+1} (r|\alpha|)!^s \sum_{|\beta|+mj \leq r|\alpha|} N_{j,\beta}(u, \varphi) \quad \text{for some } r \in \mathbb{N}^*. \quad (5-3)$$

Now, we provide a proposition on Gevrey regularity of Gevrey vectors of  $P$  satisfying (5-3).

**Proposition 5.1.** *Let  $P$  be a linear partial differential operator with  $G^s(\Omega)$  coefficients, of order  $m$ , satisfying (5-3) in  $\Omega_1$  with  $\bar{\Omega}_1 \subset \Omega$ . Then any  $G^s(\Omega_1)$ -vector of  $P$  which is  $C^\infty(\Omega)$  is in  $G^{rs}(\Omega_1)$ ,  $s \geq 1$ .*

*Proof.* We have to distinguish between the cases  $s > 1$  and  $s = 1$ .

(1) Case  $s > 1$ : Let  $u \in C^\infty(\Omega) \cap G^s(\Omega_1, P)$ . In order to prove that  $u \in G^{rs}(\Omega_1)$ , we have to show that given any open set  $\Omega_2$  with  $\bar{\Omega}_2 \subset \Omega_1$ , we have:

$$\exists C_{\Omega_2} > 0 \text{ s. t. } \forall \alpha \in \mathbb{N}^n, \quad \|\partial^\alpha u\|_{L^2(\Omega_2)} \leq C_{\Omega_2}^{|\alpha|+1} (r|\alpha|)!^s = C^{|\alpha|+1} (r|\alpha|)!^s. \quad (5-4)$$

First, we consider  $\varphi \in G^s(\Omega_1) \cap \mathcal{D}(\Omega_1)$ ,  $\varphi = 1$  on  $\Omega_2$ . Then

$$\|\partial^\alpha u\|_{L^2(\Omega_2)} \leq \|\varphi \partial^\alpha u\|. \quad (5-5)$$

As  $u \in C^\infty(\bar{\Omega}_1)$  and  $\varphi \in \mathcal{D}(\Omega_1)$ , we apply (5-3). So we get

$$\|\partial^\alpha u\|_{L^2(\Omega_2)} \leq (CM_\epsilon)^{r|\alpha|+1} (r|\alpha|)!^s \sum_{|\beta|+mj \leq r|\alpha|} N_{j,\beta}^\epsilon(u, \varphi). \quad (5-6)$$

Now as  $\varphi \in G^s(\Omega_1) \cap \mathcal{D}(\Omega_1)$ , there exists  $A = A_\varphi$  such that

$$\sup |\varphi^{(\beta)}| \leq A^{|\beta|+1} |\beta|!^s. \quad (5-7)$$

Also, from (5-2), we have

$$N_{j,\beta}^\epsilon = \epsilon^{|\beta|+mj} |\beta|!^{-s} (mj)!^{-s} \|\varphi^{(\beta)} P^j u\|. \quad (5-8)$$

Now as  $u \in G^s(\Omega_1, P)$ , taking  $K = \text{Supp}(\varphi) \subset \Omega_1$

$$\exists B = B_K > 0, \quad \text{such that} \quad \|P^j u\|_{L^2(K)} \leq B^{mj+1} (mj)!^s, \quad \forall j \in \mathbb{N}. \quad (5-9)$$

From (5-7), (5-8) and (5-9), we obtain

$$N_{j,\beta}^\epsilon(u, \varphi) \leq \epsilon^{|\beta|+mj} A^{|\beta|+1} B^{mj+1} \leq \epsilon^{|\beta|+mj} D^{|\beta|+mj+1} \quad (5-10)$$

for some constant  $D$  (for example  $A + B$ ). So, with (5-6),

$$\|\partial^\alpha u\|_{L^2(\Omega_2)} \leq D(CM_\epsilon)^{r|\alpha|+1} (r|\alpha|)!^s \sum_{|\beta|+mj \leq r|\alpha|} (\epsilon D)^{|\beta|+mj}. \quad (5-11)$$

Now let us choose  $\epsilon$  such that

$$\epsilon D = \frac{1}{2}. \tag{5-12}$$

Then with  $M = M_{1/(2D)}$ , we have

$$\|\partial^\alpha u\|_{L^2(\Omega_2)} \leq D(CM)^{r|\alpha|+1} (r|\alpha|)!^s \sum_{|\beta|+mj \leq r|\alpha|} \left(\frac{1}{2}\right)^{|\beta|+mj}. \tag{5-13}$$

We have to estimate the sum in the (5-13).

**Lemma 5.2.** *There exists a constant  $C_0 > 0$  such that*

$$\sum_{|\beta|+mj \leq r|\alpha|} \left(\frac{1}{2}\right)^{|\beta|+mj} \leq C_0, \quad \forall \alpha \in \mathbb{N}^n. \tag{5-14}$$

*Proof of Lemma 5.2.* This simple lemma has an elementary proof. For completeness, let us give it. For  $\alpha = 0$ , it is trivial. So assume  $|\alpha| \geq 1$ . It suffices to see that

$$\begin{aligned} \sum_{|\beta|+mj \leq r|\alpha|} \left(\frac{1}{2}\right)^{|\beta|+mj} &= \sum_{k=0}^{r|\alpha|} \left( \sum_{|\beta|+mj=k} 1 \right) \left(\frac{1}{2}\right)^k \leq \sum_{k=0}^{r|\alpha|} (k+1)^n \left(\frac{1}{2}\right)^k \\ &\leq \sum_{k=0}^{\infty} (k+1)^n \left(\frac{1}{2}\right)^k = C_0 < +\infty. \quad \square \end{aligned}$$

Coming back to the proof of Proposition 5.1, we have, with some constant  $C_1 > 0$ , using (5-13) and (5-14), the following estimate:

$$\|\partial^\alpha u\|_{L^2(\Omega_2)} \leq C_1(CM)^{r|\alpha|} (r|\alpha|)!^s \tag{5-15}$$

which shows that  $u$  is in  $G^{rs}(\Omega_2)$ , hence in  $G^{rs}(\Omega_1)$  as  $\Omega_2$  is any relatively compact set in  $\Omega_1$ .

(2) Case  $s = 1$ : In this case, as we have no  $\varphi \in \mathcal{D}(\Omega_1)$  which is in  $G^1(\Omega_1)$ , we proceed by using a sequence of functions of L. Ehrenpreis associated to the couple  $(\Omega_0, \Omega_1)$  with  $\bar{\Omega}_1 \subset \Omega_0$  and  $\bar{\Omega}_0 \subset \Omega$ . We state below a proposition due to Ehrenpreis, providing the precise details regarding the sequence.

**Proposition 5.3 [Ehrenpreis 1960].** *Let  $\Omega_0, \Omega_1$  be as above. Then there exists a constant  $\tilde{C} > 0$  such that:*

$$\forall N \in \mathbb{N}, \exists \varphi_N \in \mathcal{D}(\Omega_0), \varphi_N|_{\Omega_1} = 1,$$

$$\text{such that } |\varphi_N^{(\beta)}| \leq \tilde{C}^{|\beta|+1} N^\beta, \quad \text{for } |\beta| \leq N. \tag{5-16}$$

In our proof below, in order to bound  $\|\partial^\alpha u\|_{L^2(\Omega_2)}$ , we use, in place of  $\varphi$  used in case (1), the function  $\varphi_{r|\alpha|}$  given by taking  $N = r|\alpha|$  in (5-16). So, as  $\bar{\Omega}_2 \subset \Omega_1$ , we have

$$\|\partial^\alpha u\|_{L^2(\Omega_2)} \leq \|\varphi_{r|\alpha|} \partial^\alpha u\| \leq M_\epsilon^{r|\alpha|+1} (r|\alpha|)! \sum_{|\beta|+mj \leq r|\alpha|} N_{j,\beta}^\epsilon(u, \varphi_{r|\alpha|}). \tag{5-17}$$

Now, taking  $A$  given by:

$$A = \sup(B, \tilde{C}), \tag{5-18}$$

where  $B$  is given by (5-9) with  $K = \bar{\Omega}_0$ , we get

$$E_\alpha = \sum_{|\beta|+mj \leq r|\alpha|} N_{j,\beta}^\epsilon(u, \varphi_{r|\alpha|}) \leq A \sum_{|\beta|+mj \leq r|\alpha|} (\epsilon A)^{|\beta|+mj} |\beta|!^{-1} (r|\alpha|)^{|\beta|}, \tag{5-19}$$

since  $|\beta| \leq r|\alpha|$  in the sum. Looking at the second member in (5-19), we may bound by

$$E_\alpha \leq \sum_{mj \leq r|\alpha|} (\epsilon A)^{mj} \sum_{|\beta| \leq r|\alpha|} (\epsilon A)^{|\beta|} |\beta|!^{-1} (r|\alpha|)^{|\beta|}. \tag{5-20}$$

Hence

$$E_\alpha \leq \sum_{mj \leq r|\alpha|} (\epsilon A)^{mj} \sum_{k=0}^{r|\alpha|} \left( \sum_{|\beta|=k} 1 \right) (\epsilon A r|\alpha|)^k k!^{-1}. \tag{5-21}$$

If we choose  $\epsilon_0$  so that  $\epsilon_0 A = \frac{1}{2}$ , we get

$$E_\alpha \leq 2 \sum_{k=0}^{r|\alpha|} (k+1)^n \left( \frac{r|\alpha|}{2} \right)^k k!^{-1} \leq 2(r|\alpha|+1)^n \sum_k \left( \frac{r|\alpha|}{2} \right)^k k!^{-1} \tag{5-22}$$

So finally, with some constant  $C_1 > 0$ ,  $C_1 = C_1(n)$ ,

$$E_\alpha \leq 2(r|\alpha|+1)^n \exp\left(\frac{r|\alpha|}{2}\right) \leq C_1 \exp(r|\alpha|). \tag{5-23}$$

Then, coming back to (5-17), we obtain

$$\|\partial^\alpha u\|_{L^2(\Omega_2)} \leq M_{\epsilon_0}^{r|\alpha|+1} (r|\alpha|)! C_1 \exp(r|\alpha|) = C_1 M_{\epsilon_0} (e M_{\epsilon_0})^{r|\alpha|} (r|\alpha|)!. \tag{5-24}$$

As we took any  $\Omega_2$  with  $\bar{\Omega}_2 \subset \Omega_1$ ,  $\bar{\Omega}_1 \subset \Omega_0$ , we obtain that  $u \in G^r(\Omega_1)$ . The proof of Proposition 5.1 is complete.  $\square$

As a corollary of Theorem 4.2 and Proposition 5.1, we get:

**Theorem 5.4.** *Let  $P$  be a Hörmander's operator on an open set  $\Omega$  in  $\mathbb{R}^n$  and  $s \in \mathbb{R}$ ,  $s \geq 1$ . Assume that  $P$  satisfies the estimate (2-4) in some open subset  $\Omega_0$  with  $\bar{\Omega}_0 \subset \Omega$  with  $\sigma = 1/p$ ,  $p \in \mathbb{N}^*$  and that its coefficients are in  $G^s(\Omega_0)$ . Then  $G^s(\Omega_0, P) \subset G^{2ps}(\Omega_0)$ .*

We conclude this article with some final remarks.

(1) In the case  $s = 1$ , there is another proof, using the method of addition of an extra variable (see, for example, [Bolley et al. 1987] or [Lions and Magenes 1970]), by considering the operator  $\partial_t^2 + P$  in  $\mathbb{R} \times \Omega \subset \mathbb{R}^{n+1}$ , which is also a Hörmander's operator in  $\mathbb{R} \times \Omega$ , with analytic coefficients (case  $s = 1$ ), to which one can use the theorem of Gevrey hypoellipticity  $G^s$  for  $s \geq 2p$ , [Derridj and Zuily 1973].



(2) We know nothing on optimality of our result. In our preceding paper [Derridj  $\geq$  2019], in the case of operators of the first kind, the result was optimal:  $G^k(\Omega_0, P) \subset G^{pk}(\Omega_0)$ ,  $k \in \mathbb{N}$ .

(3) In a forthcoming paper, we will study the question of local relations of domination by powers of  $P$ , in the case where  $P$  is of the first kind, which will be finer, giving therefore the optimal result  $G^s(\Omega_0, P) \subset G^{ps}(\Omega_0)$ ,  $p$  being the type of  $\bar{\Omega}_0$ .

### Acknowledgements

I am greatly indebted to Nordine Mir for encouraging me to write the present paper when he knew about the result and for his help.

### References

- [Baouendi and Métivier 1982] M. S. Baouendi and G. Métivier, “Analytic vectors of hypoelliptic operators of principal type”, *Amer. J. Math.* **104**:2 (1982), 287–319. [MR](#) [Zbl](#)
- [Barostichi et al. 2011] R. F. Barostichi, P. D. Cordaro, and G. Petronilho, “Analytic vectors in locally integrable structures”, pp. 1–14 in *Geometric analysis of several complex variables and related topics* (Marrakesh, 2010), edited by Y. Barkatou et al., *Contemp. Math.* **550**, Amer. Math. Soc., Providence, RI, 2011. [MR](#) [Zbl](#)
- [Bolley et al. 1987] P. Bolley, J. Camus, and L. Rodino, “Hypoellipticité analytique-Gevrey et itérés d’opérateurs”, *Rend. Sem. Mat. Univ. Politec. Torino* **45**:3 (1987), 1–61. [MR](#) [Zbl](#)
- [Braun Rodrigues et al. 2016] N. Braun Rodrigues, G. Chinni, P. D. Cordaro, and M. R. Jahnke, “Lower order perturbation and global analytic vectors for a class of globally analytic hypoelliptic operators”, *Proc. Amer. Math. Soc.* **144**:12 (2016), 5159–5170. [MR](#) [Zbl](#)
- [Castellanos et al. 2013] J. E. Castellanos, P. D. Cordaro, and G. Petronilho, “Gevrey vectors in involutive tube structures and Gevrey regularity for the solutions of certain classes of semilinear systems”, *J. Anal. Math.* **119** (2013), 333–364. [MR](#) [Zbl](#)
- [Damlakhi and Helffer 1980] M. Damlakhi and B. Helffer, “Analyticité et itères d’un système de champs non elliptique”, *Ann. Sci. École Norm. Sup. (4)* **13**:4 (1980), 397–403. [MR](#) [Zbl](#)
- [Derridj 2017] M. Derridj, “On Gevrey vectors of some partial differential operators”, *Complex Var. Elliptic Equ.* **62**:10 (2017), 1474–1491. [MR](#) [Zbl](#)
- [Derridj  $\geq$  2019] M. Derridj, “On Gevrey vectors of L. Hörmander’s operators”, preprint. To appear in *Trans. Amer. Math. Soc.*
- [Derridj and Zuily 1973] M. Derridj and C. Zuily, “Sur la régularité Gevrey des opérateurs de Hörmander”, *J. Math. Pures Appl. (9)* **52** (1973), 309–336. [MR](#) [Zbl](#)
- [Ehrenpreis 1960] L. Ehrenpreis, “Solution of some problems of division, IV: Invertible and elliptic operators”, *Amer. J. Math.* **82** (1960), 522–588. [MR](#) [Zbl](#)
- [Helffer and Mattera 1980] B. Helffer and C. Mattera, “Analyticité et itérés réduits d’un système de champs de vecteurs”, *Comm. Partial Differential Equations* **5**:10 (1980), 1065–1072. [MR](#) [Zbl](#)
- [Hörmander 1967] L. Hörmander, “Hypoelliptic second order differential equations”, *Acta Math.* **119** (1967), 147–171. [MR](#) [Zbl](#)
- [Kohn 1978] J. J. Kohn, “Lectures on degenerate elliptic problems”, pp. 89–151 in *Pseudodifferential operator with applications* (Bressanone, Italy, 1977), Liguori, Naples, 1978. [MR](#) [Zbl](#)

- [Kotake and Narasimhan 1962] T. Kotake and M. S. Narasimhan, “Regularity theorems for fractional powers of a linear elliptic operator”, *Bull. Soc. Math. France* **90** (1962), 449–471. [MR](#) [Zbl](#)
- [Lions and Magenes 1970] J.-L. Lions and E. Magenes, *Problèmes aux limites non homogènes et applications*, vol. 3, Travaux et Recherches Mathématiques **20**, Dunod, Paris, 1970. [MR](#) [Zbl](#)
- [Métivier 1978] G. Métivier, “Propriété des itérés et ellipticité”, *Comm. Partial Differential Equations* **3:9** (1978), 827–876. [MR](#) [Zbl](#)
- [Nelson 1959] E. Nelson, “Analytic vectors”, *Ann. of Math. (2)* **70** (1959), 572–615. [MR](#) [Zbl](#)
- [Rothschild and Stein 1976] L. P. Rothschild and E. M. Stein, “Hypoelliptic differential operators and nilpotent groups”, *Acta Math.* **137:3-4** (1976), 247–320. [MR](#) [Zbl](#)

Received 24 Nov 2017. Revised 11 Dec 2017.

MAKHLOUF DERRIDJ:

[makloul.derridj@numericable.fr](mailto:makloul.derridj@numericable.fr)

78350 Les Loges en Josas, France

# Tunisian Journal of Mathematics

[msp.org/tunis](http://msp.org/tunis)

## EDITORS-IN-CHIEF

Ahmed Abbes CNRS & IHES, France  
[abbes@ihes.fr](mailto:abbes@ihes.fr)  
Ali Baklouti Faculté des Sciences de Sfax, Tunisia  
[ali.baklouti@fss.usf.tn](mailto:ali.baklouti@fss.usf.tn)

## EDITORIAL BOARD

Hajer Bahouri CNRS & LAMA, Université Paris-Est Créteil, France  
[hajer.bahouri@u-pec.fr](mailto:hajer.bahouri@u-pec.fr)  
Arnaud Beauville Laboratoire J. A. Dieudonné, Université Côte d'Azur, France  
[beauville@unice.fr](mailto:beauville@unice.fr)  
Bassam Fayad CNRS & Institut de Mathématiques de Jussieu-Paris Rive Gauche, Paris, France  
[bassam.fayad@imj-prg.fr](mailto:bassam.fayad@imj-prg.fr)  
Benoit Fresse Université Lille 1, France  
[benoit.fresse@math.univ-lille1.fr](mailto:benoit.fresse@math.univ-lille1.fr)  
Dennis Gaitsgory Harvard University, United States  
[gaitsgde@gmail.com](mailto:gaitsgde@gmail.com)  
Emmanuel Hebey Université de Cergy-Pontoise, France  
[emmanuel.hebey@math.u-cergy.fr](mailto:emmanuel.hebey@math.u-cergy.fr)  
Mohamed Ali Jendoubi Université de Carthage, Tunisia  
[ma.jendoubi@gmail.com](mailto:ma.jendoubi@gmail.com)  
Sadok Kallel Université de Lille 1, France & American University of Sharjah, UAE  
[sadok.kallel@math.univ-lille1.fr](mailto:sadok.kallel@math.univ-lille1.fr)  
Minhyong Kim Oxford University, UK & Korea Institute for Advanced Study, Seoul, Korea  
[minhyong.kim@maths.ox.ac.uk](mailto:minhyong.kim@maths.ox.ac.uk)  
Toshiyuki Kobayashi The University of Tokyo & Kavli IPMU, Japan  
[toshi@kurims.kyoto-u.ac.jp](mailto:toshi@kurims.kyoto-u.ac.jp)  
Yanyan Li Rutgers University, United States  
[yyli@math.rutgers.edu](mailto:yyli@math.rutgers.edu)  
Nader Masmoudi Courant Institute, New York University, United States  
[masmoudi@cims.nyu.edu](mailto:masmoudi@cims.nyu.edu)  
Haynes R. Miller Massachusetts Institute of Technology, United States  
[hrm@math.mit.edu](mailto:hrm@math.mit.edu)  
Nordine Mir Texas A&M University at Qatar & Université de Rouen Normandie, France  
[nordine.mir@qatar.tamu.edu](mailto:nordine.mir@qatar.tamu.edu)  
Detlef Müller Christian-Albrechts-Universität zu Kiel, Germany  
[mueller@math.uni-kiel.de](mailto:mueller@math.uni-kiel.de)  
Mohamed Sifi Université Tunis El Manar, Tunisia  
[mohamed.sifi@fst.utm.tn](mailto:mohamed.sifi@fst.utm.tn)  
Daniel Tataru University of California, Berkeley, United States  
[tataru@math.berkeley.edu](mailto:tataru@math.berkeley.edu)  
Sundaram Thangavelu Indian Institute of Science, Bangalore, India  
[veluma@math.iisc.ernet.in](mailto:veluma@math.iisc.ernet.in)  
Saïd Zarati Université Tunis El Manar, Tunisia  
[said.zarati@fst.utm.tn](mailto:said.zarati@fst.utm.tn)

## PRODUCTION

Silvio Levy (Scientific Editor)  
[production@msp.org](mailto:production@msp.org)

The Tunisian Journal of Mathematics is an international publication organized by the Tunisian Mathematical Society (<http://www.tms.rnu.tn>) and published in electronic and print formats by MSP in Berkeley.


See inside back cover or [msp.org/tunis](http://msp.org/tunis) for submission instructions.

The subscription price for 2019 is US \$315/year for the electronic version, and \$370/year (+\$20, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Tunisian Journal of Mathematics (ISSN 2576-7666 electronic, 2576-7658 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

TJM peer review and production are managed by EditFlow<sup>®</sup> from MSP.

PUBLISHED BY

 **mathematical sciences publishers**

**nonprofit scientific publishing**

<http://msp.org/>

© 2019 Mathematical Sciences Publishers

# Tunisian Journal of Mathematics

2019

vol. 1

no. 3

Rigid local systems and alternating groups	295
ROBERT M. GURALNICK, NICHOLAS M. KATZ and PHAM HUU TIEP	
Local estimates for Hörmander's operators with Gevrey coefficients and application to the regularity of their Gevrey vectors	321
MAKHLOUF DERRIDJ	
Generic colourful tori and inverse spectral transform for Hankel operators	347
PATRICK GÉRARD and SANDRINE GRELLIER	
Ramification groups of coverings and valuations	373
TAKESHI SAITO	
Almost sure local well-posedness for the supercritical quintic NLS	427
JUSTIN T. BRERETON	