

Tunisian Journal of Mathematics

an international publication organized by the Tunisian Mathematical Society

Duality relations among multiple series with three parameters

Masahiro Igarashi

2020 vol. 2 no. 1



Duality relations among multiple series with three parameters

Masahiro Igarashi

We prove a duality relation among multiple series with three parameters. As special case of it, we obtain some new identities for multiple Hurwitz zeta values, which are relations among extensions of the multiple Hurwitz zeta value.

1. Introduction

The extended multiple zeta value (EMZV for short) is defined by the multiple series

$$\sum_{\substack{0 < m_1 < c_1 \cdots < c_{p-1} m_p < \infty \\ m_i \in \mathbb{Z}}} \frac{1}{m_1^{k_1} \cdots m_p^{k_p}}, \quad (1)$$

where $1 \leq p \in \mathbb{Z}$, $c_i \in \{\frac{1}{2}, 1\}$, $1 \leq k_i \in \mathbb{Z}$ ($i = 1, \dots, p-1$), $2 \leq k_p \in \mathbb{Z}$, and the symbols $<_{c_i}$ ($i = 1, \dots, p-1$) denote $<$ if $c_i = 1$ and \leq if $c_i = \frac{1}{2}$ (see [Ulanskiĭ 2011]). The case $c_i = 1$ ($i = 1, \dots, p-1$) and the case $c_i = \frac{1}{2}$ ($i = 1, \dots, p-1$) of (1) are the multiple zeta value (MZV for short) and the multiple zeta-star value (MZSV for short), respectively, which were studied by Euler [1776], Hoffman [1992], and Zagier [1994]. EMZVs were studied by Fischler and Rivoal [2016], Kawashima [2009] and Ulanskiĭ [2011]. Kawashima and Ulanskiĭ proved relations among them, and Fischler and Rivoal showed a connection between EMZVs and a solution to a Padé approximation problem involving multiple polylogarithms. Their works show that EMZVs are as useful and fruitful as MZ(S)V's.

In [Igarashi 2015], we also studied multiple series of the extended form (1), which generalize EMZVs. In fact, we proved duality relations among them which yield numerous relations. The duality relations were proved by using a combination of the method used in [Igarashi 2012] and the calculational techniques for the Pochhammer symbol $(a)_m$ used in [Igarashi 2018]. In the present paper, we shall show that the combinative method used in [Igarashi 2015] can be applied to derive

MSC2010: 11M32.

Keywords: duality, multiple series, multiple Hurwitz zeta value, extended multiple zeta value.

numerous relations among multiple series of the following type:

$$\sum_{\substack{0 \leq m_1 < c_1 \cdots < c_{p-1} m_p \\ m_p < c_p \cdots < c_{p+q-1} m_{p+q} < \infty \\ m_i \in \mathbb{Z}}} \frac{(y)_{m_p} (w)_{m_{p+q}}}{(x)_{m_p} (z)_{m_{p+q}}} \times \left\{ \prod_{i=1}^{p+q} \frac{1}{(m_i + x)^{a_i} (m_i + y)^{b_i} (m_i + z)^{c_i} (m_i + w)^{d_i}} \right\}, \quad (2)$$

where $1 \leq p \in \mathbb{Z}$, $0 \leq q \in \mathbb{Z}$, $x, y, z, w \in \mathbb{C} \setminus \mathbb{Z}_{\leq 0}$, $a_i, b_i, c_i, d_i \in \mathbb{Z}$ such that

$$\sum_{i=r}^{p+q} (a_i + b_i + c_i + d_i) + \delta_r \operatorname{Re}(x - y) + \operatorname{Re}(z - w) > p + q - r + 1$$

($r = 1, \dots, p+q$), where $\mathbb{Z}_{\leq 0} := \{0, -1, -2, \dots\}$; $\delta_r = 1$ if $r \leq p$ and $\delta_r = 0$ if $r > p$. (These conditions guarantee the absolute convergence of (2): see [Krattenthaler and Rivoal 2007, Lemmas 1 and 3, (3.12)].) The symbols $<_{c_i}$ ($i = 1, \dots, p+q-1$) are the same as in (1), and the symbol $(a)_m$ denotes the Pochhammer symbol defined by $(a)_m = a(a+1) \cdots (a+m-1)$ ($1 \leq m \in \mathbb{Z}$) and $(a)_0 = 1$. The Pochhammer symbol $(a)_m$ can be expressed as $(a)_m = \Gamma(a+m)/\Gamma(a)$ by using the gamma function $\Gamma(s)$. The case $x = y = z = w = 1$ and the case $x = y, z = w$ of (2) are EMZV and the following multiple Hurwitz zeta value (MHZV for short), respectively:

$$\sum_{\substack{0 \leq m_1 < c_1 \cdots < c_{p-1} m_p < \infty \\ m_i \in \mathbb{Z}}} \prod_{i=1}^p \frac{1}{(m_i + x)^{s_i} (m_i + z)^{t_i}}$$

($s_i, t_i \in \mathbb{Z}; i = 1, \dots, p$). Since I proved the results in [Igarashi 2007], one of my research subjects has been to find the extensions of (multiple) Hurwitz zeta values which satisfy various relations as MZ(S)V_s. As I showed in [Igarashi 2018], the multiple series (2) and (3) below give such extensions. The results in [Igarashi 2018] have given me a motivation to the further research on (2) and (3). In the present paper, I prove a class of relations among (2) which gives numerous relations among extensions of MHZV: see Theorem 1 and Corollary 8 below. I remark that the partial derivatives of (2) with respect to x, y, z, w can be expressed by \mathbb{Z} -linear combinations of (2). This suggests that, by partial differentiation, one relation among (2) yields further relations. The proof of Theorem 1 in the present paper gives an embodiment of this suggestion. In [Igarashi 2018] and its manuscripts (submitted in March and May 2015), I studied relations among multiple series of the types (2) and (3) by using the hypergeometric identities of [Andrews 1975, Theorem 4; Krattenthaler and Rivoal 2007, Proposition 1]. This work of mine is one of the bases of the present research.

1.1. Algebraic formulation. In a revised version of [Igarashi 2015], in order to describe the results, we followed the algebraic formulation for EMZVs given by Ulanskiĭ [2011] (see also [Hoffman 1997]). That formulation gave us concise descriptions of the results. For this reason, in the present paper, we also follow the formulation of Ulanskiĭ. The following formulation is parallel to that in the revised version of [Igarashi 2015].

Hereafter we assume that $m, n, p, q, r, k_i, k'_i, m_i, r_i, s_i, M_{ij} \in \mathbb{Z}$. We consider the three noncommutative variables $x_0, x_{\frac{1}{2}}$ and x_1 . For these variables, we use the expressions

$$x_1 x_0^{k_1-1} x_{c_1} x_0^{k_2-1} \cdots x_{c_{p-1}} x_0^{k_p-1} = z_1(k_1) z_{c_1}(k_2) \cdots z_{c_{p-1}}(k_p) = \prod_{i=1}^p z_{c_{i-1}}(k_i),$$

where $p \geq 1$ and $z_{c_{i-1}}(k_i) := x_{c_{i-1}} x_0^{k_i-1}$ ($c_0 = 1, k_p \geq 1, c_i \in \{\frac{1}{2}, 1\}, k_i \geq 1; i = 1, \dots, p-1$). In case $p = 0$, we regard all these expressions as $1 \in \mathbb{Q}$. Here we consider the following set of monomials:

$$B^0 := \left\{ \prod_{i=1}^p z_{c_{i-1}}(k_i) \mid p \geq 0, c_0 = 1, c_i \in \left\{ \frac{1}{2}, 1 \right\}, k_i \geq 1 (i = 1, \dots, p-1), k_p \geq 2 \right\},$$

and thus $1 \in B^0$ as the case $p = 0$. We denote by V^0 the \mathbb{Q} -vector space with the basis B^0 . For B^0 , we define the evaluation map $H = H_{(x,y,z)} : B^0 \rightarrow \mathbb{C}$ by $H(1; (x, y, z)) = 1$ and

$$\begin{aligned} & H(z_1(k_1) z_{c_1}(k_2) \cdots z_{c_{p-1}}(k_p); (x, y, z)) \\ &= \sum_{0 \leq m_1 < c_1 \cdots < c_{p-1} m_p < \infty} \frac{(y)_{m_p}}{(x)_{m_p+1}} \left\{ \prod_{i=1}^{p-1} \frac{1}{(m_i + z)^{k_i}} \right\} \frac{1}{(m_p + z)^{k_p-1}}, \quad (3) \end{aligned}$$

where $x, z \in \mathbb{C} \setminus \mathbb{Z}_{\leq 0}, y \in \mathbb{C}, k_p \in \mathbb{Z}$ such that $k_p + \text{Re}(x - y) > 1$ and $k_i \geq 1$ ($i = 1, \dots, p-1$). This map can be extended as a \mathbb{Q} -linear map onto the whole space V^0 . Multiple series of the type (3) are studied in [Coppo 2009; Coppo and Candelpergher 2010; Emery 2004; Hasse 1930; Igarashi 2018; 2015]. See also [Igarashi 2007, Examples]. For brevity, we put $\langle_{c_i}^* := \langle_{(2c_i)-1}$ ($c_i \in \{\frac{1}{2}, 1\}$). This is the inversion of \langle_{c_i} , namely

$$\langle_{c_i}^* = \begin{cases} \leq & \text{if } c_i = 1, \\ < & \text{if } c_i = \frac{1}{2}. \end{cases}$$

We also define the evaluation map

$$H_{((r_i)_{i=1}^n; \{s_i\}_{i=1}^q)}^* = H_{((r_i)_{i=1}^n; \{s_i\}_{i=1}^q), (x, y, z)}^* : B^0 \rightarrow \mathbb{C}$$

by $H^*_{((r_i)_{i=1}^n; \{s_i\}_{i=1}^q)}(1; (x, y, z)) = 1$ and

$$\begin{aligned}
& H^*_{((r_i)_{i=1}^n; \{s_i\}_{i=1}^q)}(z_1(k_1)z_{c_1}(k_2) \cdots z_{c_{q-1}}(k_q); (x, y, z)) \\
&= \sum_{\substack{0 \leq M_{11} < \cdots < M_{1n} < m_1 \\ m_1 < c_1 M_{21} \leq \cdots \leq M_{2s_2} < c_2^* m_2 \\ \vdots \\ m_{i-1} < c_{i-1} M_{i1} \leq \cdots \leq M_{is_i} < c_i^* m_i \\ \vdots \\ m_{q-1} < c_{q-1} M_{q1} \leq \cdots \leq M_{qs_q} < c_q^* m_q < \infty}} \frac{(x)_{m_1}}{m_1!} \frac{(y)_{m_q+1}}{(z)_{m_q+1}} \\
&\quad \times \left(\prod_{i=1}^n \frac{1}{(M_{1i} + x)(M_{1i} + z)^{r_i}} \right) \frac{1}{(m_1 + y)^{k_1} (m_1 + z)^{s_1}} \\
&\quad \times \left\{ \prod_{i=2}^q \left(\prod_{j=1}^{s_i} \frac{1}{M_{ij} + z} \right) \frac{1}{(m_i + y)^{k_i}} \right\}, \quad (4)
\end{aligned}$$

where $n, r_i \geq 0$ ($i = 1, \dots, n$); $q \geq 1, s_i \geq 0$ ($i = 1, \dots, q$), $c_q = 1$; $x, y, z \in \mathbb{C} \setminus \mathbb{Z}_{\leq 0}$ such that $\operatorname{Re}(2 - x - y + z), \operatorname{Re}(1 - y + z) > 0$. We regard $\{a_i\}_{i=m}^{m-1}$ as the empty set \emptyset . In case $s_i = 0$, we regard the inequalities $m_{i-1} < c_{i-1} M_{i1} \leq \cdots \leq M_{is_i} < c_i^* m_i$ under the summation sign in (4) as $m_{i-1} < c_{i-1} m_i$. For example, the case $n = 0$ and $s_i = 0$ ($i = 1, \dots, q$) of (4) becomes

$$\begin{aligned}
& H^*_{(\emptyset; \{0\}_{i=1}^q)}(z_1(k_1)z_{c_1}(k_2) \cdots z_{c_{q-1}}(k_q); (x, y, z)) \\
&= \sum_{0 \leq m_1 < c_1 \cdots < c_{q-1} m_q < \infty} \frac{(x)_{m_1}}{m_1!} \frac{(y)_{m_q+1}}{(z)_{m_q+1}} \left\{ \prod_{i=1}^q \frac{1}{(m_i + y)^{k_i}} \right\} \\
&=: H^*(z_1(k_1)z_{c_1}(k_2) \cdots z_{c_{q-1}}(k_q); (x, y, z)).
\end{aligned}$$

This map can also be extended as a \mathbb{Q} -linear map onto the whole space V^0 .

We define the map $\sigma_r^b : B^0 \rightarrow V^0$ by $\sigma_r^b(1) = 1$ and

$$\begin{aligned}
& \sigma_r^b(z_1(k_1)z_{c_1}(k_2) \cdots z_{c_{p-1}}(k_p)) \\
&= \sum_{\substack{r_1 + \cdots + r_p = r \\ r_i \geq 0}} \left\{ \prod_{i=1}^{p-1} \binom{k_i + r_i - 1}{r_i} \right\} \binom{k_p + r_p - 2}{r_p} \prod_{i=1}^p z_{c_{i-1}}(k_i + r_i),
\end{aligned}$$

where $r \geq 0$. This map corresponds to the partial derivative of (3) with respect to z . Following [Ulanskiĭ 2011, p. 106], we also define the dual map $\tau : B^0 \rightarrow B^0$ by $\tau(1) = 1$ and $\tau(x_1 x_{e_1} \cdots x_{e_{n-1}} x_0) = x_1 x_{1-e_{n-1}} \cdots x_{1-e_1} x_0$, where $n \geq 1$ and $e_i \in \{0, \frac{1}{2}, 1\}$ ($i = 1, \dots, n-1$). We call $\tau(v)$ the dual of v . By the definition, it is easy to see that $\tau^2(v) = v$. The dual $\tau(v)$ can also be written as $\tau(v) = \prod_{i=1}^q z'_{c_{i-1}}(k'_i)$,

where $q \geq 0$, $c'_0 = 1$, $c'_i \in \{\frac{1}{2}, 1\}$, $k'_i \geq 1$ ($i = 1, \dots, q - 1$), $k'_q \geq 2$. Hereafter we assume this expression for $\tau(v)$. The maps σ_r^b and τ can be extended as \mathbb{Q} -linear maps from the whole space V^0 to V^0 .

1.2. Main theorem and its examples. We define the symbol $\varepsilon(c_i)$ by

$$\varepsilon(c_i) = \begin{cases} 1 & \text{if } c_i = 1, \\ 0 & \text{if } c_i \neq 1. \end{cases}$$

For brevity, we put

$$\begin{aligned} (\mathbf{r}^n, \mathbf{s}^q, \varepsilon(c'_1)) &:= (\{r_i\}_{i=1}^n; \varepsilon(c'_1)s_1, \{s_i\}_{i=2}^q), \\ (\alpha, \beta, \gamma)^\tau &:= (1 - \beta + \gamma, \alpha - \beta + 1, \alpha - \beta + \gamma). \end{aligned}$$

The main theorem is as follows:

Theorem 1. *Let $v \in B^0$, and let $\tau(v)$ be its dual. Then the identity*

$$\begin{aligned} &H(\sigma_r^b(v); (\alpha, \beta, \gamma)) \\ &= \sum_{n=0}^r (1-\alpha)^n \sum_{\substack{\sum_{i=0}^n r_i + \varepsilon(c'_1)s_1 + \sum_{i=2}^q s_i = r \\ r_i \geq 1 \ (i=1, \dots, n) \\ r_0, s_i \geq 0 \ (i=1, \dots, q)}} G^{(r_0)}(\alpha, \beta, \gamma) H_{(\mathbf{r}^n, \mathbf{s}^q, \varepsilon(c'_1))}^*(\tau(v); (\alpha, \beta, \gamma)^\tau) \quad (5) \end{aligned}$$

holds for all $r \geq 0$, $\alpha, \beta, \gamma \in \mathbb{C}$ such that $\text{Re}(\alpha), \text{Re}(\beta), \text{Re}(\gamma), \text{Re}(1 - \beta + \gamma), \text{Re}(\alpha - \beta + 1), \text{Re}(\alpha - \beta + \gamma) > 0$, where c'_1 and q are those of the dual $\tau(v) = \prod_{i=1}^q z_{c'_{i-1}}(k'_i)$. (For the definition of $G^{(r_0)}(\alpha, \beta, \gamma)$, see Section 2.)

The identity (5) gives numerous relations among (2) which contain identities for MHZVs and EMZVs as special cases. For example, we consider the monomial $v_0 := \{\prod_{i=1}^{p-1} z_{c_{i-1}}(1)\} z_{c_{p-1}}(2)$ ($c_0 = 1, c_i \in \{\frac{1}{2}, 1\}; i = 1, \dots, p - 1$). This has the dual

$$\tau(v_0) = x_1 x_{1-c_{p-1}} \cdots x_{1-c_1} x_0 = z_1(k'_1) \left\{ \prod_{i=2}^q z_{\frac{1}{2}}(k'_i) \right\},$$

where $q, k'_i \geq 1$ ($i = 1, \dots, q - 1$), $k'_q \geq 2$. Taking $v = v_0$ in (5), we get the identity

$$\begin{aligned} &\sum_{\substack{r_1 + \dots + r_p = r \\ r_i \geq 0}} H\left(\left\{ \prod_{i=1}^{p-1} z_{c_{i-1}}(1 + r_i) \right\} z_{c_{p-1}}(2 + r_p); (\alpha, \beta, \gamma)\right) \\ &= \sum_{n=0}^r (1-\alpha)^n \sum_{\substack{\sum_{i=0}^n r_i + \varepsilon(q)s_1 + \sum_{i=2}^q s_i = r \\ r_i \geq 1 \ (i=1, \dots, n) \\ r_0, s_i \geq 0 \ (i=1, \dots, q)}} G^{(r_0)}(\alpha, \beta, \gamma) \\ &\quad \times H_{(\mathbf{r}^n, \mathbf{s}^q, \varepsilon(q))}^*\left(z_1(k'_1) \left\{ \prod_{i=2}^q z_{\frac{1}{2}}(k'_i) \right\}; (\alpha, \beta, \gamma)^\tau\right). \quad (6) \end{aligned}$$

The sum on the left-hand side of (6) has the same form as that for the sum formula for MZVs: see (8) below. By the definition of $G^{(r_0)}(\alpha, \beta, \gamma)$ in Section 2, we see that $G^{(r_0)}(\alpha, \alpha, \alpha) = 0$ ($r_0 \geq 1$) and $G^{(0)}(\alpha, \alpha, \alpha) = 1$. Therefore, taking $\alpha = \beta = \gamma$ and $c_i = 1$ ($i = 1, \dots, p - 1$) in (6), we get the following sum formula for MHZVs:

$$\sum_{\substack{r_1 + \dots + r_p = r \\ r_i \geq 0}} \zeta \left(\left\{ \prod_{i=1}^{p-1} z_1(1+r_i) \right\} z_1(2+r_p); \alpha \right) = \sum_{n=0}^r (1-\alpha)^n \sum_{\substack{\sum_{i=1}^n r_i + s_1 = r \\ r_i \geq 1 (i=1, \dots, n) \\ s_1 \geq 0}} H_{((r_i)_{i=1}^n; s_1)}^* (z_1(p+1); (1, 1, \alpha)) \quad (7)$$

for $p \geq 1, r \geq 0, \alpha \in \mathbb{C}$ with $\text{Re}(\alpha) > 0$. Here

$$\zeta(v; \alpha) := H(v; (\alpha, \alpha, \alpha)) = \sum_{0 \leq m_1 < c_1 \dots < c_{p-1} m_p < \infty} \prod_{i=1}^p \frac{1}{(m_i + \alpha)^{k_i}}$$

($v = \prod_{i=1}^p z_{c_{i-1}}(k_i) \in B^0$). If $v = \prod_{i=1}^p z_1(k_i)$, this becomes the usual MHZV, i.e., the case $c_i = 1$ ($i = 1, \dots, p - 1$) of $\zeta(v; \alpha)$. The identity (7) was shown in [Igarashi 2018, (R3)] as an explicit expression for the identity (2) in [Igarashi 2007]. The identity (2) in [Igarashi 2007] is one of the bases of my research on MHZVs. For other identities for MHZVs which can be derived from Theorem 1, see Corollary 8 below. Finally, taking $\alpha = 1$ in (7), we get the sum formula for MZVs, which may be one of the basic relations among MZVs:

$$\sum_{\substack{r_1 + \dots + r_p = r \\ r_i \geq 0}} \zeta \left(\left\{ \prod_{i=1}^{p-1} z_1(1+r_i) \right\} z_1(2+r_p) \right) = \zeta(z_1(p+1+r)) \quad (8)$$

for $p \geq 1$ and $r \geq 0$ ([Granville 1997], Zagier (unpublished)). Here $\zeta(v) := \zeta(v; 1)$ ($v \in B^0$), which is EMZV. If $v = \prod_{i=1}^p z_1(k_i)$, this becomes the usual MZV. The case $\alpha = \beta = \gamma$ and the case $\alpha = \beta = \gamma = 1$ of (6) are extensions of (7) and (8), respectively. For another example of Theorem 1, taking $\alpha = \beta = \gamma = 1$ in (5), we can get the following relation among EMZVs:

$$\sum_{\substack{r_1 + \dots + r_p = r \\ r_i \geq 0}} \left\{ \prod_{i=1}^{p-1} \binom{k_i + r_i - 1}{r_i} \right\} \binom{k_p + r_p - 2}{r_p} \zeta \left(\prod_{i=1}^p z_{c_{i-1}}(k_i + r_i) \right) = \sum_{\substack{s_1 + \dots + s_q = r \\ s_i \geq 0}} \sum_{\substack{0 < m_1 < c_1 M_{21} \leq \dots \leq M_{2s_2} < c_2^* m_2 \\ \vdots \\ m_{i-1} < c_{i-1} M_{i1} \leq \dots \leq M_{is_i} < c_i^* m_i \\ \vdots \\ m_{q-1} < c_{q-1} M_{q1} \leq \dots \leq M_{qs_q} < c_q^* m_q < \infty}} \frac{1}{m_1^{k'_1 + \varepsilon(c'_1) s_1}} \left\{ \prod_{i=2}^q \left(\prod_{j=1}^{s_i} \frac{1}{M_{ij}} \right) \frac{1}{m_i^{k'_i}} \right\} \quad (9)$$

for $r \geq 0$. The identity (9) gives numerous relations among EMZVs. In fact, in [Igarashi 2015], I showed that the identity (9) is equivalent to a certain new extension of Ohno’s relation for MZVs [Ohno 1999, Theorem 1]: Ohno’s relation is a large class of relations among MZVs. By the above examples, we see that Theorem 1 contains various interesting relations for multiple series. See also Remark 11 below.

We explain the idea of the proof of Theorem 1. It is proved by using a symmetry of (3) with respect to the parameters x, y and z . Indeed we derive Theorem 1 from the duality formula (12) below, which has a symmetry with respect to the parameters α, β and X , by partial differentiation. The symmetry of (12) can be found by applying a change of variables to an iterated integral representation of (3) (see (10) below and the proof of (12)). It is important that the change of variables also brings about a change of the positions of the parameters of (3) (compare, e.g., the positions of X on both sides of (12)). Consequently partial differential operators act on each side of (12) in different ways, and this gives the identity in Theorem 1. (See also [Igarashi 2012].) For the factor $G^{(r_0)}(\alpha, \beta, \gamma)$ in Theorem 1, which is a partial differential coefficient of the gamma factor in (12), we prove its two explicit expressions (see Lemma 6 below). Those expressions can be used for deriving various relations among (2) from Theorem 1 (see Remarks 7 and 11 below).

I remark that Theorem 1 is a variation of my former results written in [Igarashi 2015], but it yields some new identities for MHZVs, which are different from those in [Igarashi 2015]: see Corollary 8 and Remark 10 below. For example, taking $\alpha = \beta = \gamma$ in Theorem 1, we get the following new identity for MHZVs:

$$\zeta(\sigma_r^b(v); \alpha) = \sum_{n=0}^r (1 - \alpha)^n \sum_{\substack{\sum_{i=1}^n r_i + \varepsilon(c'_1) s_1 + \sum_{i=2}^q s_i = r \\ r_i \geq 1 \ (i=1, \dots, n) \\ s_i \geq 0 \ (i=1, \dots, q)}} H_{(r^n, s^q, \varepsilon(c'_1))}^*(\tau(v); (1, 1, \alpha))$$

for all $r \geq 0, \alpha \in \mathbb{C}$ such that $\text{Re}(\alpha) > 0$, where $\zeta(\sigma_r^b(v); \alpha) := H(\sigma_r^b(v); (\alpha, \alpha, \alpha))$. This identity gives an extension of (7). See also Corollary 8(i) and Remark 10 below.

2. Proof of Theorem 1

In this section, we prove Theorem 1. The proof is a variation of the proofs of the results in [Igarashi 2015], therefore several proofs in this section overlap those in [Igarashi 2015] (see also [Igarashi 2012, Section 2]). However we shall not omit the details for the sake of keeping the present paper self-contained. The calculational techniques for the Pochhammer symbol $(a)_m$ used in the present paper are based on those used in [Igarashi 2018].

We first prove some lemmas. We define the symbol $\omega_{e_i}(t)$ ($e_i \in \{0, \frac{1}{2}, 1\}$) by

$$\omega_0(t) = \frac{1}{t}, \quad \omega_{\frac{1}{2}}(t) = \frac{1}{t(1-t)}, \quad \omega_1(t) = \frac{1}{1-t}.$$

The multiple series (3) has the following iterated integral representation, which is one of the most important ingredients of the proof of Theorem 1:

Lemma 2. *Let $n \geq 1$ and $e_i \in \{0, \frac{1}{2}, 1\}$ ($i = 1, \dots, n - 1$). Then the identity*

$$\begin{aligned} & H(x_1 x_{e_1} \cdots x_{e_{n-1}} x_0; (\alpha, \beta, X)) \\ &= \frac{\Gamma(\alpha)}{\Gamma(\beta)\Gamma(\alpha - \beta + 1)} \\ & \times \int_{0 < t_0 < \cdots < t_n < 1} t_0^{X-1} \omega_1(t_0) \left\{ \prod_{i=1}^{n-1} \omega_{e_i}(t_i) \right\} \omega_0(t_n) t_n^{\beta-X} (1 - t_n)^{\alpha-\beta} dt_0 \cdots dt_n \quad (10) \end{aligned}$$

holds for all $\alpha, \beta, X \in \mathbb{C}$ such that $\text{Re}(\alpha), \text{Re}(\beta), \text{Re}(X), \text{Re}(\alpha - \beta + 1) > 0$, where $\Gamma(s)$ is the gamma function.

Proof. Using $\omega_0(t)$ and $\omega_{c_i}(t)$ ($c_i \in \{\frac{1}{2}, 1\}$), we can rewrite the integrand as

$$\omega_1 \left\{ \prod_{i=1}^{n-1} \omega_{e_i} \right\} \omega_0 = \prod_{i=1}^p \omega_{c_{i-1}} \omega_0^{k_i-1},$$

where $p \geq 1$, $c_0 = 1$, $c_i \in \{\frac{1}{2}, 1\}$, $k_i \geq 1$ ($i = 1, \dots, p - 1$), $k_p \geq 2$. This gives the following expression for the iterated integral in (10), which we denote by I :

$$\begin{aligned} I = & \int_{\substack{0 < t_{11} < \cdots < t_{1k_1} \\ \vdots \\ < t_{i1} < \cdots < t_{ik_i} \\ \vdots \\ < t_{p1} < \cdots < t_{pk_p} < 1}} t_{11}^{X-1} \left\{ \prod_{i=1}^p \omega_{c_{i-1}}(t_{i1}) \left(\prod_{j=2}^{k_i} \omega_0(t_{ij}) \right) \right\} t_{pk_p}^{\beta-X} (1 - t_{pk_p})^{\alpha-\beta} \\ & \times \left(\prod_{i=1}^p \prod_{j=1}^{k_i} dt_{ij} \right). \quad (11) \end{aligned}$$

Applying the expansions $(1 - t_{i1})^{-1} = \sum_{m=0}^{\infty} t_{i1}^m$ ($i = 1, \dots, p$) to the integrand in (11) and integrating term by term, we get the identities

$$\begin{aligned} I &= \sum_{0 \leq m_1 < c_1 \cdots < c_{p-1} m_p < \infty} \left\{ \prod_{i=1}^{p-1} \frac{1}{(m_i + X)^{k_i}} \right\} \frac{1}{(m_p + X)^{k_p-1}} \int_0^1 (1 - t_{pk_p})^{\alpha-\beta} t_{pk_p}^{\beta+m_p-1} dt_{pk_p} \\ &= \sum_{0 \leq m_1 < c_1 \cdots < c_{p-1} m_p < \infty} \left\{ \prod_{i=1}^{p-1} \frac{1}{(m_i + X)^{k_i}} \right\} \frac{1}{(m_p + X)^{k_p-1}} \frac{\Gamma(\alpha - \beta + 1)\Gamma(\beta + m_p)}{\Gamma(\alpha + m_p + 1)} \\ &= \sum_{0 \leq m_1 < c_1 \cdots < c_{p-1} m_p < \infty} \left\{ \prod_{i=1}^{p-1} \frac{1}{(m_i + X)^{k_i}} \right\} \frac{1}{(m_p + X)^{k_p-1}} \frac{(\beta)_{m_p}}{(\alpha)_{m_p+1}} \frac{\Gamma(\alpha - \beta + 1)\Gamma(\beta)}{\Gamma(\alpha)}, \end{aligned}$$

so

$$I = \frac{\Gamma(\alpha - \beta + 1)\Gamma(\beta)}{\Gamma(\alpha)} H(x_1 x_0^{k_1-1} x_{c_1} x_0^{k_2-1} \cdots x_{c_{p-1}} x_0^{k_p-1}; (\alpha, \beta, X))$$

for $\text{Re}(\alpha), \text{Re}(\beta), \text{Re}(X), \text{Re}(\alpha - \beta + 1) > 0$. We can also get the following expression for the above monomial:

$$x_1 x_0^{k_1-1} x_{c_1} x_0^{k_2-1} \cdots x_{c_{p-1}} x_0^{k_p-1} = x_1 x_{e_1} \cdots x_{e_{n-1}} x_0,$$

where $e_i \in \{0, \frac{1}{2}, 1\}$ ($i = 1, \dots, n - 1$), therefore we get (10). □

Lemma 2 gives the following duality formula for (3):

Lemma 3 (Duality formula). *Let $v \in B^0$, and let $\tau(v)$ be its dual. Then the identity*

$$H(v; (\alpha, \beta, X)) = \frac{\Gamma(\alpha)}{\Gamma(\beta)} \frac{\Gamma(X)}{\Gamma(\alpha - \beta + X)} H^*(\tau(v); (\alpha, \beta, X)^\tau) \tag{12}$$

holds for all $\alpha, \beta, X \in \mathbb{C}$ such that

$$\text{Re}(\alpha), \text{Re}(\beta), \text{Re}(X), \text{Re}(\alpha - \beta + 1), \text{Re}(\alpha - \beta + X) > 0.$$

Proof. The assertion follows by applying the change of variables $t_i = 1 - u_{n-i}$ ($i = 0, 1, \dots, n$) to the iterated integral on the right-hand side of (10). (This change of variables was used in [Zagier 1994, p. 510] for MZVs.) □

Remark 4. The cases $\alpha = \beta = X = 1$ of (10) and (12) are Ulanskiĭ’s results on EMZVs [Ulanskiĭ 2011, Corollary 2 and Theorem 1]. The case $\alpha = \beta$ and $v = z_1(1)^{p-1} z_1(2)$ ($p \geq 1$) of (12) was used in [Igarashi 2007] to prove a sum formula for MHZVs:

$$\sum_{0 \leq m_1 < \dots < m_p < \infty} \left(\prod_{i=1}^p \frac{1}{m_i + X} \right) \frac{1}{m_p + \alpha} = \sum_{m=0}^{\infty} \frac{(1 - \alpha + X)_m}{(X)_{m+1}} \frac{1}{(m + 1)^p}.$$

To use the partial differentiation, we need the following lemma:

Lemma 5. *Let $v = \prod_{i=1}^q z_{c_{i-1}}(k_i) \in B^0$, and let $x, y, z \in \mathbb{C}$ with $\text{Re}(x), \text{Re}(y), \text{Re}(z) > 0$. Then the multiple series*

$$H^*(v; (x + Y, y, z + Y)) = \sum_{0 \leq m_1 < c_1 < \dots < c_{q-1} < m_q < \infty} \frac{(x + Y)_{m_1}}{m_1!} \frac{(y)_{m_q+1}}{(z + Y)_{m_q+1}} \left\{ \prod_{i=1}^q \frac{1}{(m_i + y)^{k_i}} \right\} \tag{13}$$

converges uniformly in $\{Y \in \mathbb{C} \mid |Y| \leq \varepsilon\}$, where $\varepsilon \in \mathbb{R}$ such that

$$0 < \varepsilon < \min\{\text{Re}(x), \text{Re}(z), 1 - \text{Re}(x + y - z)/2, 1 - \text{Re}(y - z)\}.$$

Proof. By using the expression $(a)_m = \Gamma(a + m)/\Gamma(a)$, the Pochhammer symbol $(a)_m$ can be estimated as follows:

$$\begin{aligned} |(a)_m| &= \left| \frac{\Gamma(a + m)}{\Gamma(a)} \right| = \left| \frac{1}{\Gamma(a)} \int_0^\infty e^{-t} t^{a+m-1} dt \right| \\ &\leq \frac{1}{|\Gamma(a)|} \int_0^\infty e^{-t} t^{\operatorname{Re}(a)+m-1} dt = \frac{\Gamma(\operatorname{Re}(a) + m)}{|\Gamma(a)|} = \frac{\Gamma(\operatorname{Re}(a))}{|\Gamma(a)|} (\operatorname{Re}(a))_m \\ &\leq C_1 \frac{\Gamma(\operatorname{Re}(a))}{|\Gamma(a)|} (m + 1)^{\operatorname{Re}(a)} \end{aligned}$$

for $m \geq 0$, $a \in \mathbb{C}$ with $\operatorname{Re}(a) > 0$, where C_1 is a positive constant which does not depend on m . (The last inequality above follows by applying Stirling’s formula for $\Gamma(s)$ to $(\operatorname{Re}(a))_m$.) Using this estimate and $|(a)_m| \geq (\operatorname{Re}(a))_m > 0$ for $a \in \mathbb{C}$ with $\operatorname{Re}(a) > 0$, we get the estimate

$$\begin{aligned} \left| \frac{(x + Y)_{m_1}}{m_1!} \frac{(y)_{m_q+1}}{(z + Y)_{m_q+1}} \right| &\leq \frac{\Gamma(\operatorname{Re}(x + Y))\Gamma(\operatorname{Re}(y))}{|\Gamma(x + Y)\Gamma(y)|} \frac{(\operatorname{Re}(x + Y))_{m_1}}{m_1!} \frac{(\operatorname{Re}(y))_{m_q+1}}{(\operatorname{Re}(z + Y))_{m_q+1}} \\ &\leq C_2 \frac{(\operatorname{Re}(x) + \varepsilon)_{m_1}}{m_1!} \frac{(\operatorname{Re}(y))_{m_q+1}}{(\operatorname{Re}(z) - \varepsilon)_{m_q+1}} \\ &\leq C_3 \frac{1}{(m_1 + 1)^{1-\operatorname{Re}(x)-\varepsilon}} \frac{1}{(m_q + 1)^{-\operatorname{Re}(y-z)-\varepsilon}} \end{aligned}$$

for all $Y \in \{Y \in \mathbb{C} \mid |Y| \leq \varepsilon\}$, where C_2 and C_3 are positive constants which do not depend on Y . By this estimate, we can take the absolutely convergent multiple series

$$\sum_{0 \leq m_1 < c_1, \dots, c_{q-1} m_q < \infty} \frac{1}{(m_1 + 1)^{1-\operatorname{Re}(x)-\varepsilon}} \frac{1}{(m_q + 1)^{-\operatorname{Re}(y-z)-\varepsilon}} \left\{ \prod_{i=1}^q \frac{1}{(m_i + \operatorname{Re}(y))^{k_i}} \right\}$$

as a majorant of (13), and this implies the uniform convergence of (13). For conditions on the absolute convergence of the above majorant, see those for (2). \square

For the gamma factor in (12), we put

$$G^{(n)}(\alpha, \beta, \gamma) := \frac{\Gamma(\alpha)}{\Gamma(\beta)} \frac{(-1)^n}{n!} \frac{\partial^n}{\partial X^n} \left(\frac{\Gamma(X)}{\Gamma(\alpha - \beta + X)} \right) \Bigg|_{X=\gamma},$$

where $n \geq 0$, $\alpha, \beta, \gamma \in \mathbb{C}$ such that $\alpha, \gamma, \alpha - \beta + \gamma \notin \mathbb{Z}_{\leq 0}$, which appears in Theorem 1. We prove two explicit expressions for $G^{(n)}(\alpha, \beta, \gamma)$. For brevity, we

also put

$$\begin{aligned}
 P^{(n)}(z, w) &:= \frac{1}{n!} \frac{\partial^n}{\partial Y^n} (\psi(z - Y) - \psi(w - Y)) \Big|_{Y=0} \\
 &= \begin{cases} (z - w) \sum_{m=0}^{\infty} (m + z)^{-1} (m + w)^{-1} & \text{if } n = 0, \\ -\sum_{m=0}^{\infty} (m + z)^{-n-1} + \sum_{m=0}^{\infty} (m + w)^{-n-1} & \text{if } n \geq 1, \end{cases}
 \end{aligned}$$

where $z, w \in \mathbb{C} \setminus \mathbb{Z}_{\leq 0}$ and $\psi(z) := \Gamma'(z) / \Gamma(z)$ is the digamma function. This identity is an immediate consequence of the following two properties of the digamma function:

$$\psi(z) - \psi(w) = (z - w) \sum_{m=0}^{\infty} (m + z)^{-1} (m + w)^{-1}$$

($z, w \in \mathbb{C} \setminus \mathbb{Z}_{\leq 0}$) and $\psi^{(n)}(z) = (-1)^{n+1} n! \sum_{m=0}^{\infty} (m + z)^{-n-1}$ ($n \geq 1$), where $\psi^{(n)}(z)$ is the n -th derivative of $\psi(z)$. (For the property of the digamma function, see, e.g., [Srivastava and Choi 2001, Section 1.2].) Then we can prove the following:

Lemma 6. *Let $n \geq 1$. Then $G^{(n)}(\alpha, \beta, \gamma)$ has the following two expressions:*

(i)

$$\begin{aligned}
 &G^{(n)}(\alpha, \beta, \gamma) \\
 &= \frac{\Gamma(\alpha)}{\Gamma(\beta)} \frac{\Gamma(\gamma)}{\Gamma(\alpha - \beta + \gamma)} \sum_{i=1}^n \sum_{0=l_0 < l_1 < \dots < l_{i-1} < l_i = n} \prod_{j=1}^i l_j^{-1} P^{(l_j - l_{j-1} - 1)}(\alpha - \beta + \gamma, \gamma) \quad (14)
 \end{aligned}$$

for all $\alpha, \beta, \gamma \in \mathbb{C}$ such that $\alpha, \gamma, \alpha - \beta + \gamma \notin \mathbb{Z}_{\leq 0}$.

(ii)

$$G^{(n)}(\alpha, \beta, \gamma) = \frac{\Gamma(\alpha)}{\Gamma(\beta)\Gamma(\alpha - \beta)} \sum_{m=0}^{\infty} \frac{(\beta - \alpha + 1)_m}{m!} \frac{1}{(m + \gamma)^{n+1}} \quad (15)$$

for all $\alpha, \beta, \gamma \in \mathbb{C}$ such that $\alpha \notin \mathbb{Z}_{\leq 0}$, $\text{Re}(\gamma), \text{Re}(\alpha - \beta + 1) > 0$: this single series is $H^*(z_1(n + 1); (\beta - \alpha + 1, \gamma))$.

Proof. The expression (14) can be proved in the same way as in [Igarashi 2018, Proof of Lemma 2.18]. The expression (15) can be proved as follows: The quotient $\Gamma(X) / \Gamma(\alpha - \beta + X)$ can be expressed as

$$\begin{aligned}
 \frac{\Gamma(X)}{\Gamma(\alpha - \beta + X)} &= \frac{\alpha - \beta + X}{\Gamma(\alpha - \beta + 1)} \frac{\Gamma(X)\Gamma(\alpha - \beta + 1)}{\Gamma(\alpha - \beta + X + 1)} = \frac{\alpha - \beta + X}{\Gamma(\alpha - \beta + 1)} \int_0^1 t^{X-1} (1-t)^{\alpha-\beta} dt \\
 &= \frac{\alpha - \beta + X}{\Gamma(\alpha - \beta + 1)} \int_0^1 \sum_{m=0}^{\infty} \frac{(\beta - \alpha)_m}{m!} t^{m+X-1} dt \\
 &= \frac{\alpha - \beta + X}{\Gamma(\alpha - \beta + 1)} \sum_{m=0}^{\infty} \frac{(\beta - \alpha)_m}{m!} \frac{1}{m + X}
 \end{aligned}$$

for $\operatorname{Re}(X), \operatorname{Re}(\alpha - \beta + 1) > 0$. Using the last expression above, which is a special case of Gauss' formula for the hypergeometric series ${}_2F_1(a, b; c; 1)$, we get the identities

$$\begin{aligned} & \frac{(-1)^n}{n!} \frac{\partial^n}{\partial X^n} \left(\frac{\Gamma(X)}{\Gamma(\alpha - \beta + X)} \right) \Big|_{X=\gamma} \\ &= \frac{1}{\Gamma(\alpha - \beta + 1)} \sum_{m=0}^{\infty} \frac{(\beta - \alpha)_m}{m!} \frac{(-1)^n}{n!} \frac{\partial^n}{\partial X^n} \left(\frac{\alpha - \beta + X}{m + X} \right) \Big|_{X=\gamma} \\ &= \frac{1}{\Gamma(\alpha - \beta + 1)} \sum_{m=0}^{\infty} \frac{(\beta - \alpha)_m}{m!} \left(\frac{\alpha - \beta + \gamma}{(m + \gamma)^{n+1}} - \frac{1}{(m + \gamma)^n} \right) \\ &= \frac{1}{\Gamma(\alpha - \beta)} \sum_{m=0}^{\infty} \frac{(\beta - \alpha + 1)_m}{m!} \frac{1}{(m + \gamma)^{n+1}} \end{aligned}$$

for $n \geq 1, \operatorname{Re}(\gamma), \operatorname{Re}(\alpha - \beta + 1) > 0$. Thus we get (15). □

Remark 7. Both (14) and (15) can be applied to (5), in particular, to evaluating (5) in terms of MHZVs: see Corollary 8(ii) below. The expression (15) is simpler than (14).

Proof of Theorem 1. Let

$$v = \prod_{i=1}^p z_{c_{i-1}}(k_i) \in B^0 \quad \text{with the dual} \quad \tau(v) = \prod_{i=1}^q z_{c'_{i-1}}(k'_i).$$

For brevity, we put $a_0 := 1 - \beta$ and $b_0 := \alpha - \beta$. Differentiating the left-hand side of (12) r times with respect to X at $X = \gamma$ ($\operatorname{Re}(\gamma) > 0$) and by the Leibniz rule, we get the left-hand side of (5).

The right-hand side of (5) can be proved as follows: We remark that the identity

$$(b_0 + X)_{m_1 + \varepsilon(c'_1)} = (b_0 + X)_{m_1} (m_1 + b_0 + X)^{\varepsilon(c'_1)}$$

holds, because $\varepsilon(c_i) \in \{0, 1\}$. Using this identity, we get the identities

$$\begin{aligned} \frac{(a_0 + X)_{m_1}}{(b_0 + X)_{m_q+1}} &= \frac{(a_0 + X)_{m_1}}{(b_0 + X)_{m_1}} \frac{(b_0 + X)_{m_1}}{(b_0 + X)_{m_q+1}} \\ &= \frac{(a_0 + X)_{m_1}}{(b_0 + X)_{m_1}} \frac{1}{(m_1 + b_0 + X)^{\varepsilon(c'_1)}} \left(\prod_{i=2}^q \frac{(b_0 + X)_{m_{i-1} + \varepsilon(c'_{i-1})}}{(b_0 + X)_{m_i + \varepsilon(c'_i)}} \right) \quad (16) \end{aligned}$$

for $m_1, \dots, m_q \in \mathbb{Z}$ such that $0 \leq m_1 < c'_1 \cdots < c'_{q-1} m_q$, where $c'_q = 1$. We calculate the partial differential coefficients of each factor on the right-hand side of (16). By

direct calculation, we get the identities

$$\begin{aligned}
 & \frac{(b_0 + \gamma)_{m_1} (a_0 + X)_{m_1}}{(a_0 + \gamma)_{m_1} (b_0 + X)_{m_1}} \\
 &= \prod_{n=0}^{m_1-1} \frac{(b_0 + \gamma + n)(a_0 + X + n)}{(a_0 + \gamma + n)(b_0 + X + n)} \\
 &= \prod_{n=0}^{m_1-1} \left(1 + \frac{(1 - \alpha)(\gamma - X)}{(a_0 + \gamma + n)(b_0 + X + n)} \right) \\
 &= \sum_{n=0}^{m_1} \sum_{0 \leq M_{11} < \dots < M_{1n} < m_1} \prod_{i=1}^n \frac{(1 - \alpha)(\gamma - X)}{(a_0 + \gamma + M_{1i})(b_0 + X + M_{1i})} \tag{17}
 \end{aligned}$$

for $m_1 \geq 0$. Using (17), we can get the identity

$$\begin{aligned}
 & \frac{(-1)^{s_0}}{s_0!} \frac{\partial^{s_0}}{\partial X^{s_0}} \left(\frac{(a_0 + X)_{m_1}}{(b_0 + X)_{m_1}} \right) \Big|_{X=\gamma} \\
 &= \frac{(a_0 + \gamma)_{m_1}}{(b_0 + \gamma)_{m_1}} \sum_{n=0}^{m_1} \sum_{\substack{\sum_{i=1}^n r_i = s_0 \\ r_i \geq 1}} (1 - \alpha)^n \\
 &\quad \times \sum_{0 \leq M_{11} < \dots < M_{1n} < m_1} \prod_{i=1}^n \frac{1}{(a_0 + \gamma + M_{1i})(b_0 + \gamma + M_{1i})^{r_i}} \tag{18}
 \end{aligned}$$

for $m_1, s_0 \geq 0$. The partial differential coefficients of other factors can be calculated as follows:

$$\begin{aligned}
 & \frac{(-1)^{s_i}}{s_i!} \frac{\partial^{s_i}}{\partial X^{s_i}} \left(\frac{(b_0 + X)_{m_{i-1} + \varepsilon(c'_{i-1})}}{(b_0 + X)_{m_i + \varepsilon(c'_i)}} \right) \Big|_{X=\gamma} \\
 &= \frac{(b_0 + \gamma)_{m_{i-1} + \varepsilon(c'_{i-1})}}{(b_0 + \gamma)_{m_i + \varepsilon(c'_i)}} \sum_{m_{i-1} + \varepsilon(c'_{i-1}) \leq M_{i1} \leq \dots \leq M_{is_i} < m_i + \varepsilon(c'_i)} \prod_{j=1}^{s_i} \frac{1}{M_{ij} + b_0 + \gamma} \\
 &= \frac{(b_0 + \gamma)_{m_{i-1} + \varepsilon(c'_{i-1})}}{(b_0 + \gamma)_{m_i + \varepsilon(c'_i)}} \sum_{m_{i-1} < c'_{i-1} \leq M_{i1} \leq \dots \leq M_{is_i} < c'_i m_i} \prod_{j=1}^{s_i} \frac{1}{M_{ij} + b_0 + \gamma} \tag{19}
 \end{aligned}$$

for $s_i \geq 0$ ($i = 2, \dots, q$): By the definitions of the symbols $<_{c_i}$, $<_{c_i}^*$ and $\varepsilon(c_i)$, the inequalities $m_{i-1} + \varepsilon(c'_{i-1}) \leq M_{i1}$ and $M_{is_i} < m_i + \varepsilon(c'_i)$ under the summation sign in (19) can be rewritten as $m_{i-1} < c'_{i-1} M_{i1}$ and $M_{is_i} < c'_i m_i$, respectively. Indeed these can be verified directly. For example, if $c'_i = 1$, then we get $\varepsilon(c'_i) = 1$ and $<_{c'_i}^* = \leq$. Therefore the inequality $M_{is_i} < m_i + \varepsilon(c'_i) = m_i + 1$ can be rewritten as

$M_{i s_i} <_{c'_i}^* m_i$. Using (19) and the Leibniz rule, we get the identities

$$\begin{aligned}
 & \frac{(-1)^r}{r!} \frac{\partial^r}{\partial X^r} \left(\frac{1}{(m_1 + b_0 + X)^{\varepsilon(c'_1)}} \left(\prod_{i=2}^q \frac{(b_0 + X)_{m_{i-1} + \varepsilon(c'_{i-1})}}{(b_0 + X)_{m_i + \varepsilon(c'_i)}} \right) \right) \Big|_{X=\gamma} \\
 &= \sum_{\substack{s_1 + \dots + s_q = r \\ s_i \geq 0}} \frac{\binom{s_1 + \varepsilon(c'_1) - 1}{s_1}}{(m_1 + b_0 + \gamma)^{s_1 + \varepsilon(c'_1)}} \\
 & \quad \times \left(\prod_{i=2}^q \frac{(b_0 + \gamma)_{m_{i-1} + \varepsilon(c'_{i-1})}}{(b_0 + \gamma)_{m_i + \varepsilon(c'_i)}} \sum_{m_{i-1} <_{c'_{i-1}} M_{i1} \leq \dots \leq M_{i s_i} <_{c'_i}^* m_i} \prod_{j=1}^{s_i} \frac{1}{M_{ij} + b_0 + \gamma} \right) \\
 &= \sum_{\substack{s_1 + \dots + s_q = r \\ s_i \geq 0}} \frac{\binom{s_1 + \varepsilon(c'_1) - 1}{s_1}}{(m_1 + b_0 + \gamma)^{s_1}} \\
 & \quad \times \frac{(b_0 + \gamma)_{m_1}}{(b_0 + \gamma)_{m_q + 1}} \prod_{i=2}^q \left(\sum_{m_{i-1} <_{c'_{i-1}} M_{i1} \leq \dots \leq M_{i s_i} <_{c'_i}^* m_i} \prod_{j=1}^{s_i} \frac{1}{M_{ij} + b_0 + \gamma} \right) \\
 &= \sum_{\substack{\varepsilon(c'_1) s_1 + \sum_{i=2}^q s_i = r \\ s_j \geq 0}} \frac{1}{(m_1 + b_0 + \gamma)^{\varepsilon(c'_1) s_1}} \\
 & \quad \times \frac{(b_0 + \gamma)_{m_1}}{(b_0 + \gamma)_{m_q + 1}} \prod_{i=2}^q \left(\sum_{m_{i-1} <_{c'_{i-1}} M_{i1} \leq \dots \leq M_{i s_i} <_{c'_i}^* m_i} \prod_{j=1}^{s_i} \frac{1}{M_{ij} + b_0 + \gamma} \right) \quad (20)
 \end{aligned}$$

for $r \geq 0$. The last equality sign of (20) comes from the identity

$$\binom{s_1 + \varepsilon(c'_1) - 1}{s_1} = \begin{cases} 1 & \text{if } \varepsilon(c'_1) = s_1 = 0 \text{ or } \varepsilon(c'_1) = 1, s_1 \geq 0, \\ 0 & \text{if } \varepsilon(c'_1) = 0, s_1 \geq 1. \end{cases}$$

Letting

$$\begin{aligned}
 & g(\{m_i\}_{i=1}^q; \{r_i\}_{i=1}^n; \{s_i\}_{i=1}^q) \\
 &= \sum_{\substack{0 \leq M_{11} < \dots < M_{1n} < m_1 \\ m_1 <_{c'_1} M_{21} \leq \dots \leq M_{2 s_2} <_{c'_2}^* m_2 \\ \vdots \\ m_{i-1} <_{c'_{i-1}} M_{i1} \leq \dots \leq M_{i s_i} <_{c'_i}^* m_i \\ \vdots \\ m_{q-1} <_{c'_{q-1}} M_{q1} \leq \dots \leq M_{q s_q} <_{c'_q}^* m_q}} \frac{(a_0 + \gamma)_{m_1} (b_0 + 1)_{m_q + 1}}{m_1! (b_0 + \gamma)_{m_q + 1}} \\
 & \quad \times \left(\prod_{i=1}^n \frac{1}{(M_{i1} + a_0 + \gamma)(M_{i1} + b_0 + \gamma)^{r_i}} \right) \frac{1}{(m_1 + b_0 + 1)^{k'_1} (m_1 + b_0 + \gamma)^{\varepsilon(c'_1) s_1}} \\
 & \quad \times \left\{ \prod_{i=2}^q \left(\prod_{j=1}^{s_i} \frac{1}{M_{ij} + b_0 + \gamma} \right) \frac{1}{(m_i + b_0 + 1)^{k'_i}} \right\},
 \end{aligned}$$

and using (16), (18) and (20), we get the identity

$$\begin{aligned} & \frac{(-1)^r}{r!} \frac{\partial^r}{\partial X^r} \left(\frac{\Gamma(\alpha)}{\Gamma(\beta)} \frac{\Gamma(X)}{\Gamma(b_0+X)} \frac{(a_0+X)_{m_1}}{m_1!} \frac{(b_0+1)_{m_q+1}}{(b_0+X)_{m_q+1}} \left\{ \prod_{i=1}^q \frac{1}{(m_i+b_0+1)^{k'_i}} \right\} \right) \Big|_{X=\gamma} \\ &= \sum_{\substack{r_0+s_0+\varepsilon(c'_1)s_1+\sum_{i=2}^q s_i=r \\ r_0, s_j \geq 0}} \sum_{n=0}^{m_1} \sum_{\substack{\sum_{i=1}^n r_i=s_0 \\ r_i \geq 1}} (1-\alpha)^n G^{(r_0)}(\alpha, \beta, \gamma) \\ & \qquad \qquad \qquad \times g(\{m_i\}_{i=1}^q; \{r_i\}_{i=1}^n; \{s_i\}_{i=1}^q) \quad (21) \end{aligned}$$

for $r \geq 0$ and $m_1, \dots, m_q \in \mathbb{Z}$ such that $0 \leq m_1 < c'_1 \dots < c'_{q-1} m_q$, where $c'_q = 1$. Here we impose the conditions

$$\operatorname{Re}(\beta), \operatorname{Re}(\gamma), \operatorname{Re}(1 - \beta + \gamma), \operatorname{Re}(\alpha - \beta + 1), \operatorname{Re}(\alpha - \beta + \gamma) > 0.$$

Then, taking $x = 1 - \beta + \gamma$, $y = \alpha - \beta + 1$ and $z = \alpha - \beta + \gamma$ in Lemma 5, we see that the multiple series on the right-hand side of (12) converges uniformly in $\{X \in \mathbb{C} \mid |X - \gamma| \leq \varepsilon\}$, where $\varepsilon \in \mathbb{R}$ such that

$$0 < \varepsilon < \min\{\operatorname{Re}(1 - \beta + \gamma), \operatorname{Re}(\alpha - \beta + \gamma), \operatorname{Re}(\beta/2), \operatorname{Re}(\gamma)\}.$$

Thus, differentiating term by term and using (21), we can get the following identity for the right-hand side of (12):

$$\begin{aligned} & \frac{(-1)^r}{r!} \frac{\partial^r}{\partial X^r} \left(\frac{\Gamma(\alpha)}{\Gamma(\beta)} \frac{\Gamma(X)}{\Gamma(\alpha - \beta + X)} H^*(\tau(v); (\alpha, \beta, X)^\tau) \right) \Big|_{X=\gamma} \\ &= \sum_{\substack{r_0+s_0+\varepsilon(c'_1)s_1+\sum_{i=2}^q s_i=r \\ r_0, s_j \geq 0}} \sum_{n=0}^{s_0} \sum_{\substack{\sum_{i=1}^n r_i=s_0 \\ r_i \geq 1}} (1-\alpha)^n G^{(r_0)}(\alpha, \beta, \gamma) \\ & \qquad \qquad \qquad \times H_{(r^n, s^q, \varepsilon(c'_1))}^*(\tau(v); (\alpha, \beta, \gamma)^\tau) \quad (22) \end{aligned}$$

for $r \geq 0$, $\operatorname{Re}(\alpha)$, and

$$\operatorname{Re}(\beta), \operatorname{Re}(\gamma), \operatorname{Re}(1 - \beta + \gamma), \operatorname{Re}(\alpha - \beta + 1), \operatorname{Re}(\alpha - \beta + \gamma) > 0.$$

Finally we verify that the right-hand side of (22) is the same as that of (5). For brevity, we put the summand on the right-hand side of (22) in $h(n; \{r_i\}_{i=0}^n; \{s_i\}_{i=1}^q)$. Then the right-hand side of (22) can be rewritten as

$$\begin{aligned} & \sum_{\substack{r_0+s_0+\varepsilon(c'_1)s_1+\sum_{i=2}^q s_i=r \\ r_0, s_j \geq 0}} \sum_{n=0}^{s_0} \sum_{\substack{\sum_{i=1}^n r_i=s_0 \\ r_i \geq 1}} h(n; \{r_i\}_{i=0}^n; \{s_i\}_{i=1}^q) \\ &= \sum_{s_0=0}^r \sum_{n=0}^{s_0} \sum_{\substack{r_0+\varepsilon(c'_1)s_1+\sum_{i=2}^q s_i=r-s_0 \\ r_0, s_j \geq 0}} \sum_{\substack{\sum_{i=1}^n r_i=s_0 \\ r_i \geq 1}} h(n; \{r_i\}_{i=0}^n; \{s_i\}_{i=1}^q) \end{aligned}$$

$$\begin{aligned}
 &= \sum_{n=0}^r \sum_{s_0=n}^r \sum_{\substack{r_0+\varepsilon(c'_1)s_1+\sum_{i=2}^q s_i=r-s_0 \\ r_0, s_j \geq 0}} \sum_{\substack{\sum_{i=1}^n r_i=s_0 \\ r_i \geq 1}} h(n; \{r_i\}_{i=0}^n; \{s_i\}_{i=1}^q) \\
 &= \sum_{n=0}^r \sum_{\substack{r_0+s_0+\varepsilon(c'_1)s_1+\sum_{i=2}^q s_i=r-n \\ r_0, s_j \geq 0}} \sum_{\substack{\sum_{i=1}^n r_i=s_0+n \\ r_i \geq 1}} h(n; \{r_i\}_{i=0}^n; \{s_i\}_{i=1}^q) \\
 &= \sum_{n=0}^r \sum_{\substack{\sum_{i=0}^n r_i+\varepsilon(c'_1)s_1+\sum_{i=2}^q s_i=r \\ r_i \geq 1 (i=1, \dots, n) \\ r_0, s_i \geq 0 (i=1, \dots, q)}} h(n; \{r_i\}_{i=0}^n; \{s_i\}_{i=1}^q).
 \end{aligned}$$

The last rewrite is exactly the same as the right-hand side of (5). This completes the proof of Theorem 1. □

For $v = \prod_{i=1}^p z_{c_{i-1}}(k_i) \in B^0$, we put

$$\begin{aligned}
 \zeta(v; (x, z)) &:= H(v; (x, x, z)) \\
 &= \sum_{0 \leq m_1 < c_1 \dots < c_{p-1} m_p < \infty} \left(\prod_{i=1}^{p-1} \frac{1}{(m_i + z)^{k_i}} \right) \frac{1}{(m_p + x)(m_p + z)^{k_p-1}} \tag{23}
 \end{aligned}$$

and

$$\begin{aligned}
 &\zeta^*_{(\{r_i\}_{i=1}^n; \{s_i\}_{i=1}^p)}(v; y) \\
 &:= H^*_{(\{r_i\}_{i=1}^n; \{s_i\}_{i=1}^p)}(v; (1, y, y)) \\
 &= \sum_{\substack{0 \leq M_{11} < \dots < M_{1n} < m_1 \\ m_1 < c_1 M_{21} \leq \dots \leq M_{2s_2} < c_2^* m_2 \\ \vdots \\ m_{i-1} < c_{i-1} M_{i1} \leq \dots \leq M_{i s_i} < c_i^* m_i \\ \vdots \\ m_{p-1} < c_{p-1} M_{p1} \leq \dots \leq M_{p s_p} < c_p^* m_p < \infty}} \left(\prod_{i=1}^n \frac{1}{(M_{1i} + 1)(M_{1i} + y)^{r_i}} \right) \frac{1}{(m_1 + y)^{k_1+s_1}} \\
 &\qquad \times \left\{ \prod_{i=2}^p \left(\prod_{j=1}^{s_i} \frac{1}{M_{ij} + y} \right) \frac{1}{(m_i + y)^{k_i}} \right\}.
 \end{aligned}$$

These multiple series are MHZVs. We can derive the following two identities for these MHZVs from Theorem 1:

Corollary 8. *Let $v \in B^0$, and let $\tau(v)$ be its dual. Then the following two identities hold:*

(i)

$$\zeta(\sigma_r^b(v); (\alpha, \gamma)) = \sum_{n=0}^r (1-\alpha)^n \sum_{\substack{\sum_{i=1}^n r_i + \varepsilon(c'_1)s_1 + \sum_{i=2}^q s_i = r \\ r_i \geq 1 (i=1, \dots, n) \\ s_i \geq 0 (i=1, \dots, q)}} H_{(r^n, s^q, \varepsilon(c'_1))}^*(\tau(v); (\alpha, \alpha, \gamma)^\tau) \quad (24)$$

for all $r \geq 0, \alpha, \gamma \in \mathbb{C}$ such that

$$\operatorname{Re}(\alpha), \operatorname{Re}(\gamma), \operatorname{Re}(1 - \alpha + \gamma) > 0,$$

where $\zeta(\sigma_r^b(v); (x, z)) := H(\sigma_r^b(v); (x, x, z))$.

(ii)

$$H(\sigma_r^b(v); (\alpha, 1, 1)) = \sum_{n=0}^r (1-\alpha)^n \sum_{\substack{\sum_{i=0}^n r_i + \varepsilon(c'_1)s_1 + \sum_{i=2}^q s_i = r \\ r_i \geq 1 (i=1, \dots, n) \\ r_0, s_i \geq 0 (i=1, \dots, q)}} G^{(r_0)}(\alpha, 1, 1) \zeta_{(r^n, s^q, \varepsilon(c'_1))}^*(\tau(v); \alpha) \quad (25)$$

for all $r \geq 0, \alpha \in \mathbb{C}$ with $\operatorname{Re}(\alpha) > 0$. The factor $G^{(r_0)}(\alpha, 1, 1)$ ($r_0 \geq 1$) becomes $(\alpha - 1)\zeta(z_1(1)^{r_0-1}z_1(2); (\alpha, 1))$ or a \mathbb{Q} -polynomial of

$$(1 - \alpha)^c \sum_{m=0}^{\infty} (m + 1)^{-a} (m + \alpha)^{-b}$$

($a, b, c \in \mathbb{Z}$ such that $a, b, c \geq 0, a + b > 1$), therefore the right-hand side of (25) can be expressed by MHZVs and $(1 - \alpha)^n$ ($n \geq 0$).

Proof. By the definition of $G^{(n)}(\alpha, \beta, \gamma)$, we see that $G^{(n)}(\alpha, \alpha, \gamma) = 0$ ($n \geq 1$) and $G^{(0)}(\alpha, \alpha, \gamma) = 1$. By this fact and taking $\alpha = \beta$ in Theorem 1, we get (24). The identity (25) can be derived from Theorem 1 by taking $\beta = \gamma = 1$. In this case, Lemma 6(i) shows that the factor $G^{(r_0)}(\alpha, 1, 1)$ ($r_0 \geq 1$) becomes a \mathbb{Q} -polynomial of $(1 - \alpha)^c \sum_{m=0}^{\infty} (m + 1)^{-a} (m + \alpha)^{-b}$ ($a, b, c \in \mathbb{Z}, a, b, c \geq 0, a + b > 1$). On the other hand, Lemma 6(ii) gives the expressions

$$\begin{aligned} G^{(r_0)}(\alpha, 1, 1) &= (\alpha - 1) \sum_{m=0}^{\infty} \frac{(2 - \alpha)_m}{m!} \frac{1}{(m + 1)^{r_0+1}} \\ &= (\alpha - 1) H^*(z_1(r_0 + 1); (2 - \alpha, 1, 1)) \end{aligned}$$

for $r_0 \geq 1, \operatorname{Re}(\alpha) > 0$. Further, this single series can be rewritten as

$$\begin{aligned} H^*(z_1(r_0 + 1); (2 - \alpha, 1, 1)) &= H^*(z_1(r_0 + 1); (\alpha, \alpha, 1)^\tau) \\ &= H(z_1(1)^{r_0-1} z_1(2); (\alpha, \alpha, 1)) \quad (\text{by Lemma 3}) \\ &= \zeta(z_1(1)^{r_0-1} z_1(2); (\alpha, 1)) \quad (\text{by (23)}). \end{aligned}$$

Thus we get $G^{(r_0)}(\alpha, 1, 1) = (\alpha - 1)\zeta(z_1(1)^{r_0-1} z_1(2); (\alpha, 1))$ ($r_0 \geq 1, \operatorname{Re}(\alpha) > 0$), and this completes the proof of the assertion for $G^{(r_0)}(\alpha, 1, 1)$ stated in (ii). \square

Remark 9. The identity

$$\sum_{m=0}^{\infty} \frac{(2 - \alpha)_m}{m!} \frac{1}{(m + 1)^{p+1}} = \sum_{0 \leq m_1 < \dots < m_p < \infty} \left(\prod_{i=1}^p \frac{1}{m_i + 1} \right) \frac{1}{m_p + \alpha}$$

($p \geq 1, \operatorname{Re}(\alpha) > 0$), which we used in the proof of Corollary 8(ii), is proved by Hoffman [1992, Section 4] by using a theorem of Mordell. He used this identity to prove a duality formula for MZVs [Hoffman 1992, Theorem 4.4].

Remark 10. I give a remark on a connection between (24) and a former result of mine written in [Igarashi 2015]. Let $v = \prod_{i=1}^p z_{c_{i-1}}(k_i) \in B^0$, and let $\tau(v) = \prod_{i=1}^q z_{c'_{i-1}}(k'_i)$ be its dual. In [Igarashi 2015], I proved the following new identity for MHZVs:

$$\begin{aligned} \sum_{\substack{r_1 + \dots + r_p = r \\ r_i \geq 0}} \left\{ \prod_{i=1}^p \binom{k_i + r_i - 1}{r_i} \right\} \zeta \left(\prod_{i=1}^p z_{c_{i-1}}(k_i + r_i); \alpha \right) \\ = \sum_{\substack{s_1 + \dots + s_q = r \\ s_i \geq 0}} H^*_{(\{s_i\}_{i=1}^q)}(\tau(v); \alpha) \quad (26) \end{aligned}$$

for all $r \geq 0, \alpha \in \mathbb{C}$ with $\operatorname{Re}(\alpha) > 0$, where $\zeta(v; \alpha)$ is the extension of MHZV defined under (7) and

$$\begin{aligned} H^*_{(\{s_i\}_{i=1}^p)}(v; \alpha) \\ := \sum_{\substack{0 = m_0 \leq M_{11} \leq \dots \leq M_{1s_1} <^*_{c_1} m_1 \\ \vdots \\ m_{i-1} <_{c_{i-1}} M_{i1} \leq \dots \leq M_{is_i} <^*_{c_i} m_i \\ \vdots \\ m_{p-1} <_{c_{p-1}} M_{p1} \leq \dots \leq M_{ps_p} <^*_{c_p} m_p < \infty}} \frac{(m_p + 1)!}{(\alpha)_{m_p + 1}} \left\{ \prod_{i=1}^p \left(\prod_{j=1}^{s_i} \frac{1}{M_{ij} + \alpha} \right) \frac{1}{(m_i + 1)^{k_i}} \right\} \end{aligned}$$

($p \geq 1, s_i \geq 0$ ($i = 1, \dots, p$), $c_p = 1, \alpha \in \mathbb{C}$ with $\operatorname{Re}(\alpha) > 0$). By the identity $\binom{k_p + r_p - 1}{r_p} = \binom{k_p + r_p - 2}{r_p} + \frac{r_p}{k_p - 1} \binom{k_p + r_p - 2}{r_p}$, we see that the sum on the left-hand side of (24) with $\alpha = \gamma$ is a partial sum of that of (26), and therefore the identity (24) with $\alpha = \gamma$ is a decomposition of (26). The factors $(1 - \alpha)^n$ ($n = 0, 1, \dots, r$) in

(24) explicitly indicate vanishing terms at $\alpha = 1$ of the decomposition: compare the case $\alpha = 1$ of the right-hand side of (24) with that of (26). The identity (24) gives a two-parameter extension of (7) also. The identities for MHZVs (24), (25) and (26) are relations among extensions of MHZV, namely the multiple series (2).

Remark 11. As is shown in the present paper, the gamma factor in (12) contributes to deriving various relations among (2) from (12). Here we give another example of this sort: By dividing both sides of (12) by the gamma factor, the identity (12) can be modified as

$$\frac{\Gamma(\beta)}{\Gamma(\alpha)} \frac{\Gamma(\alpha - \beta + X)}{\Gamma(X)} H(v; (\alpha, \beta, X)) = H^*(\tau(v); (\alpha, \beta, X)^\tau). \tag{27}$$

In the same way as in the proof of Theorem 1, we can derive the following inversion formula for (5) from (27):

$$\begin{aligned} \sum_{r_0=0}^r G^{(r_0)}(\beta, \alpha, \alpha - \beta + \gamma) H(\sigma_{r-r_0}^b(v); (\alpha, \beta, \gamma)) \\ = \sum_{n=0}^r (1 - \alpha)^n \sum_{\substack{\sum_{i=1}^n r_i + \varepsilon(c'_1)s_1 + \sum_{i=2}^q s_i = r \\ r_i \geq 1 \ (i=1, \dots, n) \\ s_i \geq 0 \ (i=1, \dots, q)}} H_{(r^n, s^q, \varepsilon(c'_1))}^*(\tau(v); (\alpha, \beta, \gamma)^\tau) \end{aligned}$$

for all $v \in B^0$, $r \geq 0$, $\alpha, \beta, \gamma \in \mathbb{C}$ such that

$$\operatorname{Re}(\alpha), \operatorname{Re}(\beta), \operatorname{Re}(\gamma), \operatorname{Re}(1 - \beta + \gamma), \operatorname{Re}(\alpha - \beta + 1), \operatorname{Re}(\alpha - \beta + \gamma) > 0,$$

where $\tau(v) = \prod_{i=1}^q z_{c'_{i-1}}(k'_i)$ is the dual of v . This identity also yields numerous relations among (2). For this kind of application of gamma factors, see also [Igarashi 2018, Theorem 2.17] and its proof.

Acknowledgment

I am grateful to the referee for careful reading and his or her comments on this manuscript, which were useful for me to improve the presentation.

References

[Andrews 1975] G. E. Andrews, “Problems and prospects for basic hypergeometric functions”, pp. 191–224 in *Theory and application of special functions* (Madison, WI, 1975), edited by R. A. Askey, Academic, New York, 1975. MR Zbl

[Coppo 2009] M.-A. Coppo, “Nouvelles expressions des formules de Hasse et de Hermite pour la fonction zêta d’Hurwitz”, *Expo. Math.* **27**:1 (2009), 79–86. MR Zbl

[Coppo and Candelpergher 2010] M.-A. Coppo and B. Candelpergher, “The Arakawa–Kaneko zeta function”, *Ramanujan J.* **22**:2 (2010), 153–162. MR Zbl

- [Emery 2004] M. Emery, “On a multiple harmonic power series”, preprint, 2004. arXiv:math.NT/0411267
- [Euler 1776] L. Euler, “Meditationes circa singulare serierum genus”, *Novi Comm. Acad. Sci. Petropol.* **20** (1776), 140–186. Reprinted in *Opera Omnia*, Ser. I, Vol. 15, Berlin, 1927, pp. 217–267.
- [Fischler and Rivoal 2016] S. Fischler and T. Rivoal, “Multiple zeta values, Padé approximation and Vasilyev’s conjecture”, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **15** (2016), 1–24. MR Zbl
- [Granville 1997] A. Granville, “A decomposition of Riemann’s zeta-function”, pp. 95–101 in *Analytic number theory* (Kyoto, 1996), edited by Y. Motohashi, London Math. Soc. Lecture Note Ser. **247**, Cambridge Univ. Press, 1997. MR Zbl
- [Hasse 1930] H. Hasse, “Ein Summierungsverfahren für die Riemannsche ξ -Reihe”, *Math. Z.* **32**:1 (1930), 458–464. MR Zbl
- [Hoffman 1992] M. E. Hoffman, “Multiple harmonic series”, *Pacific J. Math.* **152**:2 (1992), 275–290. MR Zbl
- [Hoffman 1997] M. E. Hoffman, “The algebra of multiple harmonic series”, *J. Algebra* **194**:2 (1997), 477–495. MR Zbl
- [Igarashi 2007] M. Igarashi, *On generalizations of the sum formula for multiple zeta values*, master’s thesis, Nagoya University, 2007. In Japanese; see also arXiv:1110.4875 and arXiv:0908.2536v5.
- [Igarashi 2012] M. Igarashi, “A generalization of Ohno’s relation for multiple zeta values”, *J. Number Theory* **132**:4 (2012), 565–578. MR Zbl
- [Igarashi 2015] M. Igarashi, “On the duality formula for parametrized multiple series”, preprint, 2015. Revised in 2018.
- [Igarashi 2018] M. Igarashi, “Note on relations among multiple zeta(-star) values”, *Italian J. Pure Appl. Math.* **39** (2018), 710–756. Zbl
- [Kawashima 2009] G. Kawashima, “Multiple series expressions for the Newton series which interpolate finite multiple harmonic sums”, preprint, 2009. arXiv:0905.0243v1
- [Krattenthaler and Rivoal 2007] C. Krattenthaler and T. Rivoal, “An identity of Andrews, multiple integrals, and very-well-poised hypergeometric series”, *Ramanujan J.* **13**:1-3 (2007), 203–219. MR Zbl
- [Ohno 1999] Y. Ohno, “A generalization of the duality and sum formulas on the multiple zeta values”, *J. Number Theory* **74**:1 (1999), 39–43. MR Zbl
- [Srivastava and Choi 2001] H. M. Srivastava and J. Choi, *Series associated with the zeta and related functions*, Kluwer, Dordrecht, Netherlands, 2001. MR Zbl
- [Ulanskiĭ 2011] E. A. Ulanskiĭ, “Multiple zeta values”, *Vestnik Moskov. Univ. Ser. I Mat. Mekh.* **66**:3 (2011), 14–19. In Russian; translated in *Moscow Univ. Math. Bull.* **66**:03 (2011), 105–109. MR Zbl
- [Zagier 1994] D. Zagier, “Values of zeta functions and their applications”, pp. 497–512 in *First European Congress of Mathematics, II* (Paris, 1992), Progr. Math. **120**, Birkhäuser, Basel, 1994. MR Zbl

Received 25 Oct 2018. Revised 31 Jan 2019.

MASAHIRO IGARASHI:

masahiro.igarashi2018@gmail.com

12-6, Hayato-cho, Showa-ku, Nagoya 466-0833, Japan

Tunisian Journal of Mathematics

msp.org/tunis

EDITORS-IN-CHIEF

- Ahmed Abbes CNRS & IHES, France
abbes@ihes.fr
- Ali Baklouti Faculté des Sciences de Sfax, Tunisia
ali.baklouti@fss.usf.tn

EDITORIAL BOARD

- Hajer Bahouri CNRS & LAMA, Université Paris-Est Créteil, France
hajer.bahouri@u-pec.fr
- Arnaud Beauville Laboratoire J. A. Dieudonné, Université Côte d'Azur, France
beauville@unice.fr
- Bassam Fayad CNRS & Institut de Mathématiques de Jussieu-Paris Rive Gauche, Paris, France
bassam.fayad@imj-prg.fr
- Benoit Fresse Université Lille 1, France
benoit.fresse@math.univ-lille1.fr
- Dennis Gaitsgory Harvard University, United States
gaitsgde@gmail.com
- Emmanuel Hebey Université de Cergy-Pontoise, France
emmanuel.hebey@math.u-cergy.fr
- Mohamed Ali Jendoubi Université de Carthage, Tunisia
ma.jendoubi@gmail.com
- Sadok Kallel Université de Lille 1, France & American University of Sharjah, UAE
sadok.kallel@math.univ-lille1.fr
- Minhyong Kim Oxford University, UK & Korea Institute for Advanced Study, Seoul, Korea
minhyong.kim@maths.ox.ac.uk
- Toshiyuki Kobayashi The University of Tokyo & Kavli IPMU, Japan
toshi@kurims.kyoto-u.ac.jp
- Yanyan Li Rutgers University, United States
yyli@math.rutgers.edu
- Nader Masmoudi Courant Institute, New York University, United States
masmoudi@cims.nyu.edu
- Haynes R. Miller Massachusetts Institute of Technology, United States
hrm@math.mit.edu
- Nordine Mir Texas A&M University at Qatar & Université de Rouen Normandie, France
nordine.mir@qatar.tamu.edu
- Detlef Müller Christian-Albrechts-Universität zu Kiel, Germany
mueller@math.uni-kiel.de
- Mohamed Sifi Université Tunis El Manar, Tunisia
mohamed.sifi@fst.utm.tn
- Daniel Tataru University of California, Berkeley, United States
tataru@math.berkeley.edu
- Sundaram Thangavelu Indian Institute of Science, Bangalore, India
veluma@math.iisc.ernet.in
- Saïd Zarati Université Tunis El Manar, Tunisia
said.zarati@fst.utm.tn

PRODUCTION

- Silvio Levy (Scientific Editor)
production@msp.org

The Tunisian Journal of Mathematics is an international publication organized by the Tunisian Mathematical Society (<http://www.tms.rnu.tn>) and published in electronic and print formats by MSP in Berkeley.

See inside back cover or msp.org/tunis for submission instructions.

The subscription price for 2020 is US \$320/year for the electronic version, and \$380/year (+\$20, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Tunisian Journal of Mathematics (ISSN 2576-7666 electronic, 2576-7658 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840 is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

TJM peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY
 **mathematical sciences publishers**
nonprofit scientific publishing
<http://msp.org/>

© 2020 Mathematical Sciences Publishers

Tunisian Journal of Mathematics

2020 vol. 2 no. 1

Looijenga line bundles in complex analytic elliptic cohomology CHARLES REZK	1
Fronts d'onde des représentations tempérées et de réduction unipotente pour $SO(2n + 1)$ JEAN-LOUP WALDSPURGER	43
Spectral Mackey functors and equivariant algebraic K -theory, II CLARK BARWICK, SAUL GLASMAN and JAY SHAH	97
Twisted Calabi–Yau ring spectra, string topology, and gauge symmetry RALPH L. COHEN and INBAR KLANG	147
Semiclassical approximation of the magnetic Schrödinger operator on a strip: dynamics and spectrum MOUEZ DIMASSI	197
Duality relations among multiple series with three parameters MASAHIRO IGARASHI	217